

Supporting information

Phytotoxic dioxolanones are potential virulence factors in the infection process of *Guignardia bidwellii*

Buckel, Iris; Andernach, Lars; Schüffler, Anja; Piepenbring, Meike; Opatz, Till; Thines,
Eckhard

Toxin detection *in planta*




Fig. 1: HPLC-ESIMS run data of the HPLC peak (retention time 8.7 min). (gradient: 70:30 \rightarrow 65:35 H₂O/MeCN in 4 minutes, then in 2 minutes 65:35 \rightarrow 10:90 H₂O/MeCN)




Fig 2: HPLC-ESIMS data of isolated partially ¹³C-labeled guignardic acid. (gradient: 70:30 \rightarrow 65:35 H₂O/MeCN in 4 minutes, then in 2 minutes 65:35 \rightarrow 10:90 H₂O/MeCN)




Fig. 3: UV spectrum at 1.3 minutes of the collected HPLC peak (retention time 8.7 minutes).




Fig. 4: UV spectrum at 1.3 minutes of isolated partially ^{13}C -labeled guignardic acid.

Elucidation of the precursor for the biosynthesis of phytotoxic dioxolanones




Fig. 5: HPLC-ESIMS data of isolated partially ¹³C-labeled alaguignardic acid. (gradient: 90:10 → 10:90 H₂O/MeCN in 4 minutes, then 2 minutes 10:90 H₂O/MeCN)




Fig. 6: HPLC-ESIMS data of isolated partially ^{13}C -labeled guignardic acid. (gradient: 90:10 \rightarrow 10:90 $\text{H}_2\text{O}/\text{MeCN}$ in 4 minutes, then 2 minutes 10:90 $\text{H}_2\text{O}/\text{MeCN}$)




Fig. 7: HPLC-ESIMS data of isolated partially ^{13}C -labeled phenguignardic acid. (gradient: 90:10 \rightarrow 10:90 $\text{H}_2\text{O}/\text{MeCN}$ in 4 minutes, then 2 minutes 10:90 $\text{H}_2\text{O}/\text{MeCN}$)




Fig. 8: ^1H -NMR of partially ^{13}C labeled guignardic acid (600 MHz, CDCl_3).




Fig. 9: Igated ^{13}C -NMR of partially ^{13}C labeled guignardic acid (151 MHz, CDCl_3).




Fig. 10: ^1H -NMR of partially ^{13}C labeled alaguignardic acid (600 MHz, CDCl_3).




Fig. 11: gated ^{13}C -NMR of partially ^{13}C labeled alaguignardic acid (151 MHz, CDCl_3).




Fig. 12: ^1H -NMR of partially ^{13}C labeled phenguignardic acid (600 MHz, CDCl_3).




Fig. 13: I gated ^{13}C -NMR of partially ^{13}C labeled phenguignardic acid (151 MHz, CDCl_3).