

A REVIEW OF THE INDIAN SPECIES OF THE GENUS
OLIGODON SUPPRESSING THE GENUS *SIMOTES*
(*OPHIDIA.*)

By F. WALL, Colonel, C.M.G., I.M.S.

Last year at my request all the available specimens of *Simotes* and *Oligodon* in the Indian Museum were submitted to me for study, also all those available in the Bombay Natural History Society's collection. I also had an opportunity while in England of examining some of the specimens in the British Museum. To notes from this material I have added notes from the many scores of specimens examined in smaller museums and of my own collecting. I propose in the following paper to review this genus and put on record all that is known of the various species up to date.

In a previous paper (*Journ. Bombay Nat. Hist. Soc.*, XIX, p. 556, foot-note) I showed from a study of skulls in my collection that Duméril and Bibron had established a genus *Oligodon* distinct from *Simotes* on the erroneous observation that species to which they applied the generic title *Oligodon* had no teeth in the palate. Later Boulenger discovered that some of these had palatine teeth, but he retained the genus as distinct from *Simotes* on the erroneous assumption that these same species had no pterygoid teeth. All those whose skulls I have prepared, except doubtfully *templetoni*, have pterygoid teeth so that the grounds for separating *Oligodon* from *Simotes* on dentition do not exist.

As *Oligodon* has priority (1827) over *Simotes* (1853) the former title must be retained to designate the whole.

A study of the genitalia show that there are two very different varieties to be met with among the species, and it is possible the genus, as now represented, may be divided hereafter on this basis.

However, as I have genitalia of only four species, it is wisest provisionally to let them all remain under the one generic title *Oligodon*.


IDENTIFICATION.—Distinctive as are all the species of this genus, there is no easily defined guide to their recognition. The following associated characters concerning lepidosis appear to me to afford the most important basis for identification. *Rostral.* In the median line the portion visible from above is twice or more than twice the length of the suture between the internasal fellows (except in *cruentatus* and *theobaldi*). *Internasals.* In species where two are present, as is usually the case, the suture between them is half or less than half the internaso-præ-frontals. *Frontal.* As long as the parietals (except in *cruentatus* and *theobaldi*). *Temporals.* Two shields border the parietals, the posterior subequal to, or longer than, the anterior (except in *splendidus* where three border the parietals). *Infralabials.* The last of the series (the 4th or 5th) is as long as, and rather broader than the posterior sub-linguals, and touches only two scales behind.

HABITS.—Terrestrial. Diurnal. Active. Courageous.

FOOD.—Chiefly lizard's (or snake's) eggs and frog's spawn. Sometimes small mammals, lizards, and other reptiles.

BREEDING.—Oviparous in all the species whose habits I have investigated.

The male genitalia vary in the species here included in one genus, and it is possible that this variation may be used as a basis for a subdivision of the genus into two. I have dried genitalia of four species in my collection. In *taeniatus* and *sublineatus* the organ is relatively small, cylindrical and beset with recurved cartilaginous processes from base to extremity, just as one sees in so many other species of the Colubridæ. In *albocinctus* and *juglandifer*, however, the organ is relatively very large, cylindrical, and increases in girth from base to extremity, being cleft at the extreme tip. There are no recurved processes. The proximal third has feeble longitudinal rugæ, the middle third a honeycomb network of rugæ, and the distal third concentric and somewhat imbricate rugæ (see figures).



A

A Genitalia of *Oligodon albocinctus* ($\times 2$).
B " " *sublineatus* ($\times 5$).!

B

LEPIDOSIS.—For Indian species. *Rostral*. Touching six shields (eight in *splendidus*). Portion visible above as long as or nearly as long as its distance to the frontal. *Internasals*. A pair. (Absent in *brevicauda* and *herberti*, four in *splendidus*. Separated by the rostral in *planiceps*.) *Praefrontals*. A pair. *Frontal*. Touching six shields. Length greater than its distance to the end of the snout (except in *cruentatus*, where it is subequal or less); equal to or almost equal to the parietals (two-thirds to three-fourths in *theobaldi* and *cruentatus*). *Nasals*. Entire, semidivided, or divided. *Loreal*. Present or absent. *Pracocular*. One. *Postoculars*. Two. (One in *herberti*, *mcdougalli* and *dorsalis*.) *Temporals*. $1+2$, or $2+2$, well developed (small and hardly differentiated in *splendidus*). Two border the parietals, the posterior being as long as or longer than the anterior (except in *splendidus*, where three border the parietals). Two shields between the parietals and the last supralabial (except in *splendidus*, where there are three). *Supralabials*. Usually 7 or 8. 4 in *planiceps*, 6 in *herberti*, *melanozonatus*, and *woodmasoni* (where the 3rd is divided into an upper and lower part) and in *violaceus*, *purpurascens*, and *splendidus*, where the 4th is similarly

divided. Two touching the eye (except in *planiceps*, where only the 3rd touches). *Infralabials*. 4 or 5. The last as long as, or longer than the posterior sublinguals, as broad or broader than those shields, touching two scales behind. *Sublinguals*. Two pairs, the posterior shorter than the anterior. Two or three pairs of scales between the posterior and the 1st ventral.

OSTEOLOGICAL CHARACTERS.—*Skull*. *Praemaxilla*.—Broader than high. *Nasals*. Whole breadth of posterior border forming a ligamentous suture with the frontals. *Praefrontals*. The suture with the frontal equal to the orbital rim of the frontal. *Frontals*. Longer than broad, slightly constricted opposite the middle of the orbit. Contributing about one-third to two-fifths to the orbital rim. *Postfrontal*. Touching the frontal in some species, not in others. *Parietal*. Contributing a little to the orbital rim in some species, nothing in others. *Supratemporal*. Projecting beyond the quadrate. *Maxilla*. Projecting beyond the palatine anteriorly, and posteriorly. As long as the dentary. Teeth 6 to 16; much compressed; anododont, syncranterian, strongly coryphodont. With or without an edentulous space anteriorly. *Ectopterygoid*. Rather shorter than the maxilla. *Palatine*. Slender; teeth 0 to 10; anododont, isodont or feebly kumatodont. With or without an edentulous space anteriorly. *Pterygoid*. Teeth 1 to 23 (? none in *sublineatus*). Anododont, kumatodont. With or without an edentulous space anteriorly. *Mandible*. Angular present. Sphenoid present. Coronoid absent. Dentary about as long as its distance to the quadrate. Teeth 6 to 20; anododont, feebly kumatodont. With or without an edentulous space anteriorly. *Quadrate*. About as long as its distance to the orbit. Oblique from above backwards. *Columella auris*. Short, ending in cartilage which is attached to about the middle of the quadrate.

Vertebræ. *Neural spines*.—Wanting on the atlas. On the axis as long as the body. On the 3rd somewhat oblique backwards. On the succeeding vertebrae the depth is about three-fourths the body, length rather shorter than the body. *Hypapophyses*. On the atlas a small tubercle. On the axis bifid, the posterior part oblique backwards. Well developed in succeeding vertebrae, disappearing in the anterior part of the second eighth of the body, where they are replaced by a shallow, even ridge. Bifid to form two laterally disposed laminæ on the 2nd and succeeding caudal vertebrae.

Costae.—1st rib articulated to the 3rd vertebra, rather shorter than the 2nd. Penultimate bifid, the outer ramus about one-fifth the inner. Ultimate bifid, the outer ramus about two-thirds the inner.

Pseudo-costal processes.—Bifid to form outer and inner rami on the first three caudal vertebrae, single in succeeding vertebrae.

***Oligodon planiceps* (Boulenger).**

Simotes planiceps, Boulenger, Cat. II, 1894, p. 232.

COLOUR.—Pale brown dorsally, some of the scales with blackish lower borders. A blackish bar behind the parietal shields. A black oblique subocular streak. Ventrally white (pinkish, Boulenger), with lateral, squarish, black spots, usually in pairs.

FOOD AND BREEDING.—Nothing known.

GROWTH.—Maximum Length. 130 mm. ($5\frac{1}{4}$ inches.) Tail 15 mm. ($\frac{3}{5}$ of an inch).

LEPIDOSIS.—As detailed in the synopsis. *Rostral*. Entirely separating the internasals, and partially separating the praefrontals. In the specimen in the Indian Museum the supralabials are 4. The 4th is a very long shield and possibly a confluence of two.

DISTRIBUTION.—*Burma*. The type is from Minhlā, Burma (Lat. 20° . Long 95°). Another specimen in the Indian Museum killed on the Yunnan Expedition is dubiously from Upper Burma or Yunnan.

Oligodon herberti Boulenger.

O. herberti, Boulenger, *Bomb. N. H. J.*, XVI, p. 235.

TYPE.—In the British Museum, from Mogok, N. Shan States, Burma.

COLOUR.—Dark grey dorsally. A buff, black-edged vertebral stripe involving the vertebral and upper half of the next row from the nape to the tip of the tail, and constricted at intervals to form elongate spindle-shaped segments. An ill-defined, blackish line on the confines of the second and third rows above the ventrals, and ending at the vent. Ventrally whitish (orange-red, Boulenger) with squarish, black, lateral spots, usually on alternate shields. Head with an indistinct praefrontal blackish bar, re-appearing below the eye, and an ill-defined blackish bar on the parietals, sending forward an angular process to the frontal, behind which is a buff oblique bar on the neck, separated from its fellow vertebrally.

FOOD AND BREEDING.—Nothing known.

GROWTH.—Maximum Length. 560 mm. (1 foot, 10 inches).

LEPIDOSIS.—As detailed in the synopsis.

DENTITION.—Maxillary. 7? with an edentulous space anteriorly.

DISTRIBUTION.—*Burma*. Mogok, N. Shan States. Sinlum Kaba (Lat. 24° . Long. $97^{\circ}5$). 6,000 ft. (Bombay colln.)

NOTE.—I have seen four specimens.

Oligodon mcdougalli Wall.

O. mcdougalli, Wall, *Bomb. N. H. J.*, XVI, p. 251.

TYPE.—From Sandoway, in the Bombay Natural History collection.

COLOUR.—Dusky black dorsally with a rufous-brown, vertebral stripe from the nape to tail tip, bordered interruptedly with black and involving the vertebral and half the next row. A linear black line on the confines of the 2nd and 3rd rows above the ventrals, interrupted anteriorly and ending at the vent. A supra-anal and a subterminal caudal black bar. Ventrally black mottled with fawn. Tail beneath black laterally, and crimson centrally. Head blackish. Rostral rufous yellow blotched with black below. Labials mottled black and rufous-yellow. A rufous collar, incomplete on the nape.

FOOD AND BREEDING.—Nothing known.

GROWTH.—Maximum Length. 350 mm. (1 foot, $1\frac{3}{4}$ inches).

LEPIDOSIS.—As detailed in the synopsis.

DISTRIBUTION.—*Burma.* Sandoway.

NOTE.—Known from a single specimen.

Oligodon torquatus (Boulenger).

Simotes torquatus, Boulenger, Cat. II, 1894, p. 232.

COLOUR.—Dorsally brown with or without four indistinct longitudinal stripes. The anterior superior edge of the 3rd row above the ventrals, and both anterior edges of the superjacent rows with a whitish speck. Ventrally white with a few irregularly disposed black spots on a few shields before the vent. Tail immaculate beneath. Head with obscure blackish marks. A more or less distinct blackish bar behind the parietals. A subocular blackish streak.

FOOD.—Nothing known.

BREEDING.—A gravid ♀ killed at Myitkyina by Captain Venning on the 1st of May 1911 was sent to me which was found to contain two (dubiously three) eggs. One of the eggs was protruding from the cloaca proving that the species is oviparous. No mention was made of any eggs already discharged.

GROWTH.—Maximum Length. 292 mm. (11½ inches). Tail 37 mm. (1½ inches).

LEPIDOSIS.—As detailed in the synopsis.

DENTITION.—As detailed in the synopsis. I have two skulls in my collection.

DISTRIBUTION.—*Burma.* Hills between Lat. 24° and 26°, and East of Long. 96°. Bhamo. Myitkyina.

NOTE.—I have seen four specimens including the types.

Oligodon erythrorhachis Wall.

O. erythrorhachis, Wall, Bomb. N. H. J., XIX, p. 923.

TYPE.—From Jaipur, Assam, in the British Museum.

COLOUR.—Dorsally pale grey with a powdering of black, giving a dark grey effect extending to the edges of the ventrals. A conspicuous red vertebral stripe from the nape to tail tip, involving the vertebral and next one and a half rows, 29 narrow, black, light-edged cross bars on the body and 7 on the tail. Ventrally white with square black lateral spots in the whole length. A pinkish suffusion in the middle from tail tip, disappearing in the fore-body. Head with a blackish transverse bar across the praefrontals, reappearing below the eye. A broad oblique black temporal streak. An elongate, black, median arrow head on the nape with the point confluent with the praefrontal bar, but not united with the temporal streaks.

FOOD AND BREEDING.—Nothing known.

GROWTH.—Maximum Length. 375 mm. (1 foot, 2¾ inches). Tail 62 mm. (2¾ inches).

LEPIDOSIS.—As detailed in the synopsis.

DISTRIBUTION.—*Assam.* Namsang near Jaipur, at the foot of the Naga Hills.

NOTE.—Only one specimen is known.

Oligodon dorsalis (Gray).

O. dorsalis, Boulenger, Cat. II, 1894, p. 241. Venning, *Bomb. N. H. J.*, XX, pp. 338 and 772. Wall, *Bomb. N. H. J.*, XVIII, p. 327; *l. c.* XIX, p. 831.

COLOUR.—Dorsally brown the colour deepening in the flanks. A light vertebral stripe from the nape to the tip of the tail, outlined with black, involves the vertebral and half the next row. A black linear stripe on the confines of the second and third rows above the ventrals. A basal and a sub-terminal caudal blackish bar. Ventrally parti-coloured black and white, some shields being entirely black or white, others half black and half white. Beneath the tail is crimson, unspotted and unbanded. Head powdered with black. A more or less distinct transverse bar on the praefrontals, and a chevron on the nape sending forward a median streak to meet the praefrontal bar. These may be very obscure, losing definition as age advances.

FOOD.—Nothing known.

BREEDING. (A) The Sexes.—The female appears to grow longer than the male. In females the body is rather longer, and the tail rather shorter than in the males as seen by the ventral and subcaudal shields. (Ventrals ♂ 162 to 182, ♀ 170 to 189. Subcaudals ♂ 35 to 51, ♀ 27 to 40.)

(B) Method of Reproduction.—From the size and character of eggs I have seen "in abdomina" there is no doubt that the species is oviparous.

(C) Season.—From available data eggs appear to be hatching from June to about the end of September. Venning's six inch specimen was killed on the 17th of June. A gravid female with eggs in an advanced stage of development was acquired by me on the cartroad below Shillong on the 20th of August 1907. Supposing these eggs were fit for discharge they would not probably have hatched for six or eight weeks.

(D) The eggs.—The specimen just referred to contained two eggs, but as it had been mutilated by cartwheels it is possible there were more.

GROWTH.—(a) The Hatchling.—The smallest specimen known to me was 152 mm. (6 inches) in length, but it is uncertain whether this was a hatchling.

(b) Maximum Length. 560 mm. (1 foot, 8 inches). Tail 68 mm. ($2\frac{3}{4}$ inches).

LEPIDOSIS.—(a) Typical.—As detailed in the synopsis.

(b) Anomalies.—In a specimen from Haka the 3rd and 4th rows above the ventrals blend, redivide, and reunite repeatedly on the left side in the anterior part of the body. I have seen two postoculars on one side in one specimen.

DENTITION.—As indicated in the synopsis. I have two skulls.

DISTRIBUTION.—Assam. Garo Hills. (Tura, Ind. Mus.). Naga Hills. (Samaguting. Ind. Mus. Namsang, near Jaipur, F. W.). Khasi Hills. (Ind. Mus., Shillong, F.W.). *Bengal.* Chittagong Hills. (Ind. Mus.). *Burma.* Chin Hills. (Haka 6,500 ft. Bombay Colln.). Mansi, Katha Dist. (Lat. 24° 1'. Long. 97° 4'. Bombay Colln.).

NOTE.—I have now examined eighteen examples.

***Oligodon brevicauda* Günther.**

O. brevicauda, Boulenger, Cat. II, 1894, p. 240.

COLOUR.—Light brown dorsally with two lateral darker stripes. The upper involves the lower half of the seventh, and the sixth rows above the ventrals, and extends from the nape to the tip of tail. It is bordered with a black line above which is interrupted anteriorly and renders the light vertebral stripe increasingly conspicuous from the fore body posteriorly. The lower stripe involves the lower half of the third, and the second rows, and extends from the neck to the vent. Ventrally pink, half or the whole of several ventrals blackish. Tail pink beneath, immaculate or with a few spots. Head with a transverse dark bar across the praefrontals reappearing below the eye. An oblique, dark, temporal stripe from the gape to the parietals. A dark frontal spot which may be confluent with the praefrontal bar. A broad dark chevron on the nape.

FOOD AND BREEDING.—Nothing known.

GROWTH.—Maximum Length. 480 mm. (1 foot, 7 inches). Tail 50 mm. (2 inches).

LEPIDOSIS.—(a) *Typical*.—As detailed in the synopsis. Peculiar in that the internasals are wanting as in *herberti*.

(b) *Anomalies*.—There is sometimes a segment detached from the parietals which forms a spurious upper anterior temporal.

DISTRIBUTION.—*Western Ghats*.—South of the Goa Gap. (Nilgiris. Anamalais. Travancore Hills.)

***Oligodon taeniatus* (Jerdon).**

Coronella taeniata, Jerdon, J. A. S., Bengal, XXII, 1853, p. 258.

Oligodon subgriseus, Boulenger, Cat. II, 1894, p. 243. Ferguson, Bomb. N. H. J., X, p. 71. Millard, Bomb. N. H. J., XV, p. 348. Pearless, Spol. Zeylan, 1909, p. 54. Sarasin, Zool. Jahr. Jena, 1910, p. 130. Slater, List. Sn. Ind. Mus., 1891, p. 25. Wall, Bomb. N. H. J., XVI, p. 298; l.c., XIX, p. 556; l.c., XXVI, p. 568. Spol. Zeylan, 1921, p. 406. Willey, Spol. Zeylan, 1906, p. 233.

Oligodon taeniatus, Wall, Oph. Tap., 1921, p. 239.

COLOUR.—Subject to much variation. Dorsally the ground colour is tawny or light brown. The anterior inferior edges of some of the scales are edged with a lighter and some a darker shade so as to form a fine variegation. The darker streaks show a more or less decided tendency to form crossbars of which there may be from 28 to 42 on the body and 7 to 11 on the tail. In some specimens there is an intermediate series of less obvious bars. In most specimens there are four more or less distinct darker longitudinal stripes. The upper and broader stripes pass from the nape where they are confluent, to the tip of the tail, and involve the edge of the vertebral, and two and half adjoining rows. The lower stripes pass from the neck to the vent, and involve the contiguous halves of the 2nd and 3rd rows above the ventrals. A pale line occupies the middle of the vertebral row, and expands anteriorly to the edge of the uppermost row. The belly is pearly white and usually quite immaculate but some specimens exhibit a few scattered brownish spots in the posterior part of the body. The head is marked with three dark bars.

The anterior crosses the praefrontals and reappears below the eye ; the median is chevron-shaped with its apex on the frontal and its limbs pass obliquely to the gape ; the third and broadest is chevron-shaped and situated on the nape, its apex extending forwards to the parietals. These marks which are so characteristic of the species of this genus are usually complete and discrete, but may be more or less confluent in the median line, or disintegrate. In some old specimens they become effaced, and replaced by a fine mottling.

Four varieties may be recognised of which the first three are completely connected.

Variety (A) (a). In this the variegations are light and sparse, the bars are not or hardly suggested, and the stripes are wanting or very faint.

Variety (A) (b) *taeniolatus* (Jerdon). The variegations are more pronounced than in (A) (a) and both bars and stripes are fairly conspicuous.

Variety (C) (c) *fasciatus* (Günther). Like the last but with the bars replaced by from 23 to 37 round, indented, or completely bisected spots on the body, and from 5 to 11 on the tail. I have examined specimens intermediate between (A) (b) and (A) (c), i.e., with spots anteriorly and crossbars posteriorly.

Variety (C) *spilonotus* (Günther). Differs from the previous forms in having large, light-edged marks, shaped like a walnut kernel, and reminding one of the ornamentation seen in *splendidus*, *venustus* and *juglansifer*. There are from 13 to 23 of these on the body, and 4 to 6 on the tail.

FOOD.—I have found lizard's (or snake's ?) eggs in the stomach many times and also frog's spawn.

BREEDING.—(a) *Method of reproduction*.—I have examined a gravid female with eggs of such a size and character as to make it fairly certain that the species is oviparous.

(b) *Season*.—Two specimens found in company at Bangalore, on the 23rd of April 1920, proved to be ♂ and ♀. The latter showed no enlargement of the ovarian follicles. A young specimen, apparently a hatchling which measured 122 mm. (4 $\frac{7}{8}$ inches), was taken by me at Cannanore in March. None of the 18 females collected by me in the Nilgiris and Wynad in the months of June to September 1917 were gravid, but 14 young of the year varying from 150 to 232 mm. (6 to 9 $\frac{1}{2}$ inches) were brought in during that period.

(c) *The Brood*.—The only gravid female that has come into my hands contained four large eggs. The date of capture is unfortunately not on record.

GROWTH.—(a) *The Hatchling*.—The smallest specimens I have seen were 118 mm. (4 $\frac{3}{4}$ inches), date unknown, and 121 mm. (4 $\frac{7}{8}$ inches) in March. I have seen another 4 $\frac{7}{8}$ inches long, date and locality not on record.

(b) *Early Life*.—From rather meagre figures at my disposal it would appear to grow about 5 inches a year in the first three years of life, when it would be about 20 inches long.

(c) *Maximum Length*.—The longest I know taped 585 mm. (1 foot 11 inches).

LEPIDOSIS.—(a) *Typical*.—As detailed in the synopsis.

(b) *Anomalies*.—*Praefrontals*.—Rarely quite separated by the frontal. *Parietals*. Sometimes a small detached portion forms a spurious upper anterior temporal. *Loreal*. Rarely absent, being confluent with the praefrontal. *Praeocular*. I have seen two in one specimen. *Postoculars*. Rarely three. *Supralabials*. The 3rd and 4th are rarely confluent. I have seen the 3rd divided in one specimen, as one sees the 4th divided in *purpurascens* and *violaceus*. In one specimen the 6th was cuneate and failed to reach the edge of the lip. Rarely there is a supernumerary making the total 8. *Ventrals*. I have seen the last divided like the anal in one example. *Anal*. Rarely entire. *Subcaudals*. Rarely some at the base of the tail are entire.

DENTITION.—From five skulls in my collection.

Maxillary.—An edentulous space anteriorly followed by 6 to 7 teeth. *Palatine*. Edentulous anteriorly and posteriorly; a single tooth (or none?) in the middle. *Pterygoid*. Edentulous anteriorly and posteriorly; 6 to 10 teeth in the middle. *Mandibular*. A short edentulous space anteriorly followed by 12 teeth.

DISTRIBUTION.—*Ceylon*. *Peninsular India* to Sind and Baluchistan in the North-West, the Western Himalayas, and to Bengal (Purnea) in the North-East.

Variety (A) (a).—Baluchistan (Khila Abdulla). Sind. Punjab. Western Himalayas (Dhikala, Garhwal District). Bengal (Purnea District). Throughout the Peninsula. Ceylon.

Variety (A) (b). Sind. N.-W. Frontier. Punjab. Western Himalayas (Dhikala, Garhwal District). Bengal (Barakur) to South India and Ceylon.

Variety (A) (c).—Western Ghats (Matheran. Mysore Hills. Wynad. Nilgiris. Anamalais. High Range, Travancore. Tinnevelly Hills). Ceylon.

I have examined specimens intermediate between (A) (b) and (A) (c) from Mysore Hills (Koppa), the Anamalais, Tinnevelly Hills and Bellary.

Variety (B).—Western Ghats (Castle Rock, Canara. Mysore Hills. Anamalais. Nelliampathi Hills. Travancore.)

Oligodon ellioti Günther.

O. ellioti, Sarasin, *Zool. Jahr. Jena*, 1910, p. 138. Wall, *Bomb. N. H. J.*, XIX p. 533.

COLOUR.—Very like *taeniolatus* (Jerdon). Dorsally brown, with the anterior inferior borders of some scales edged lighter and others darker, so forming a variegation. There are from 36 to 40 darker, light-edged crossbars on the body (or spots with a median indentation anteriorly and posteriorly), and 10 to 11 on the tail. The spots are completely bisected in the forebody and on the tail in one specimen, which thus resembles a specimen of variety *fasciatus* of *taeniolatus*. Three darker ill-defined stripes are more or less in evidence. The median stripes begin on the nape, and end at the tail tip, and involve the vertebral, and the next one and a half rows. A narrower lateral stripe begins in the neck and passes to the vent, and involves the lower half of the fourth, the third and the upper half of the second rows above the ventrals. Ventrally

yellowish and unspotted, or with the basal parts of the ventrals dusky. A dark transverse bar crosses the praefrontals and edge of the frontal to reappear below the eye. A dark chevron with its apex on the frontal and its arms passing to the gape curves forward to the throat, and nearly meets its fellow behind the posterior sublinguals. A dark broad mark on the nape is projected forwards in the middle line to the posterior angle of the frontal.

FOOD AND BREEDING.—Nothing known.

GROWTH.—*Maximum Length.*—343 mm. (1 foot, 1½ inches). Tail 47 mm. (1¾ inches).

LEPIDOSIS.—As detailed in the synopsis.

DISTRIBUTION.—South India. Ceylon ?

NOTE.—Only three specimens are known, the type in the British Museum, labelled "Madras Presidency," and two discovered by me in the Bombay collection labelled *O. subgriseus*.

One of these is from Nasik, and the other said to be from Ceylon, but before finally accepting the last locality it would be advisable to await confirmation of this.

***Oligodon sublineatus* Dumeril and Bibron.**

O. sublineatus, Abercromby, *Sn. of Ceylon*, 1910, p. 72. *Spol. Zeylan.* 1911, p. 206. Annandale, *J. A. S. Beng.*, 1905, p. 175. Boulenger, *Cat. II.* 1894, p. 242; *I. c. III.* 1896, p. 640. Pearloss, *Spol. Zeylan.* 1909, p. 54. Sarasin, *Zool. Jahr. Jena*, 1910, p. 134. Selator, *List Sn. Ind. Mus.* 1891, p. 25. Wall, *Spol. Zeylan.* 1910, p. 37; *I. c. 1921*, p. 400. *Oph. Tap.* 1921, p. 248. Willey, *Spol. Zeylan.* 1906, p. 233.

COLOUR.—Dorsally brown, the anterior inferior borders of many scales edged darker and lighter to form a variegation. A series of from 12 to 15 paired dark brown, light-edged spots on the body, 3 to 4 on the tail. These sometimes alternate with those of the opposite side. Ventrally whitish with three series of brown spots, sometimes confluent into lines. The lateral pass from the neck to the tip of the tail. The median ceases at the vent. Head with a dark praefrontal bar re-appearing below the eye. A similar transverse bar on the posterior part of the frontal, and anterior edge of the parietals. A dark bar on the nape, sometimes interrupted in the median line, sometimes confluent in the median line with the frontal bar. These marks tend to become obscure with age. They are sometimes very distinct and discrete, sometimes more or less confluent, or disintegrated.

FOOD.—On two occasions I have found the eggs of a lizard (or snake ?) in the stomach. In one there were two soft shelled eggs 17 mm. (5/8 of an inch) long, and in the other a single egg about 13 mm. (1/2 an inch) long.

BREEDING.—(a) *Method of Reproduction.*—The single gravid female that has come into my hands contained an egg of such dimensions as to leave no doubt that the species is oviparous.

(b) *Season.*—The gravid specimen was killed on the 4th of October 1920, and must have been very near the climax of her cyoësis.

(c) *The Brood.*—My one gravid female contained but one egg which measured 26 × 5 mm. (1 inch × ⅛ of an inch).

GROWTH.—(a) *The Hatchling*.—The length at exovation is not known, but one is justified in assuming that it is appreciably smaller than in *taeniolatus*, probably less than 100 mm. (4 inches).

(b) *Early Life*.—Two specimens obtained in 1920, one in January 200 mm. (8 inches) and the other in February 178 mm. ($7\frac{1}{2}$ inches) were probably hatched in the preceding year, and these seem to indicate that the young as in so many other snakes double their length in the first year of life. Eight specimens obtained between August and October and varying from 225 to 263 mm. (9 to $10\frac{1}{2}$ inches) would make it appear that these were about eighteen months old.

(c) *Maturity*.—My egg-bound female taped 412 mm. ($9\frac{1}{4}$ inches), a length suggesting that she was about eighteen months old.

(d) *Maximum Length*.—The largest known to me was 324 mm. ($12\frac{3}{4}$ inches) long, the tail 38 mm. ($1\frac{1}{2}$ inches).

LEPIDOSIS.—(a) *Typical*.—As detailed in the synopsis.

(b) *Anomalies. Loreal*.—I have seen this absent on one side in three specimens. *Supralabials*. The 2nd and 3rd were confluent on one side in two specimens, and on both sides in one example. *Costals*. In one specimen the 6th and 7th rows above the ventrals blended and redivided several times anteriorly so that the costals were alternately 15 and 13. Rarely the 2nd and 3rd rows above the ventrals blend posteriorly, so that the rows count 13 at a point two heads lengths before the vent.

DENTITION.—From three skulls in my collection. *Maxillary*. An edentulous space anteriorly that would take about 2 teeth, followed by from 6 to 8 teeth. *Palatine*. Edentulous? *Pterygoid*. A long edentulous space anteriorly, followed by from 4 to 6 teeth. *Mandibular*. An edentulous space anteriorly that would take two or three teeth, followed by from 9 to 10 teeth.

DISTRIBUTION.—Peculiar to Ceylon. A denizen of the low country ascending to 3,000 feet. South Prov. (Galle), West Prov. (Colombo, Matugama, Veyangoda), Sab'wa Prov. (Ratnapura and Yatiyantota Districts), Central Prov. (Peradeniya.)

NOTE.—There are two specimens in the Indian Museum labelled Nicobar Islands on the authority of de Roepstorff. This is to be distrusted. The same authority is responsible for two other equally startling records from the Nicobar Islands, viz., *Polyodontophis sagittarius* which does not occur south of Lower Bengal, and *Amblycephalus monticola* a hill-snake that does not occur south of Assam.

Oligodon templetoni Günther.

O. templetoni, Abercromby, *Sn. of Ceylon*, 1910, p. 73. Boulenger, *Cat. II*, 1894, pp. 241 and 359. Sarasin, *Zool. Jahr. Jena.*, 1910, p. 127. Wall, *Spol. Zeylan.* 1921, p. 400. *Oph. Tap.* 1921, p. 245. Willey, *Spol. Zeylan.* 1906, p. 238.

COLOUR.—Dorsally brown, with a more or less distinct lighter vertebral stripe from the nape to the tip of the tail. A series of darker brown or blackish brown transverse lateral bars, involving the 5th, 6th and 7th rows above the ventrals. There are about 18 of these on the body, and 5 on the tail. Ventrally dappled with squarish black and white (yellow or pinkish in life) spots. A fine peppering of dark brown on the

head, a dark oblique stripe below the eye. An oblique dark bar behind the gape nearly meeting its fellow over the nape.

FOOD AND BREEDING.—Nothing known.

GROWTH.—(a) *The Young*.—The smallest specimen I have seen was 126 mm. (5 inches) in length, but as the navel was completely effaced it is certain that the young are less than this when hatched or born.

(b) *Maximum Length*.—The longest I have measured was 290 mm. ($11\frac{1}{2}$ inches).

LEPIDOSIS.—(a) *Typical*.—As detailed in the synopsis.

(b) *Anomalies*. *Loreal*.—Sometimes touches the eye below the praocular. *Postoculars*. Sometimes only the upper touches the parietals. *Supralabials*. The 6th is sometimes cuneate and fails to reach the edge of the lip. *Sublinguals*. The posterior rarely touch the 3rd as well as the 4th infralabial.

DENTITION.—From one skull in my collection. *Maxillary*. An edentulous space that would carry 2 teeth followed by 7 teeth. *Palatine*. Edentulous. *Pterygoid*. Edentulous. *Mandibular*. An edentulous space anteriorly that would take 1 tooth, followed by 7 teeth.

DISTRIBUTION.—Peculiar to Ceylon. A low country species ascending to about 3,000 or 4,000 feet. South Prov. (Udugama), West Prov. (Hewissa, Matugama), Sab'wa Prov. (Ratnapura, Balangoda), Cent. Prov. (Peradeniya).

Oligodon melanurus Wall.

O. melanurus, Wall, *Bomb. N. H. J.*, XIX, p. 349.

TYPE.—In the British Museum. Co-type in the Bombay Natural History Society's collection. Both from Sukna, Darjiling District.

COLOUR.—Dorsally blackish. Under a lens the ground colour is seen to be blackish and finely powdered with lighter blue-grey speckling. Ventrally uniform deep bluish grey, the throat and chin rather paler. Head like the body above, and without marks.

FOOD.—Nothing known.

BREEDING.—The type was egg bound and contained four (five ?) eggs about 12 mm. ($\frac{1}{2}$ an inch) long. This and a male (the co-type) were dislodged from a heap of loose metal on the road side. Unfortunately no date of capture was recorded.

GROWTH.—*Maximum Length*.—333 mm. (1 foot, $1\frac{1}{8}$ inches).

LEPIDOSIS.—(a) *Typical*.—As detailed in the synopsis.

(b) *Anomalies*. *Anal*.—In the co-type the anal is entire.

DISTRIBUTION.—*Bengal*. (Sukna, Darjiling District, 540 feet).

NOTE.—Only two specimens are known.

Oligodon travancoricus Beddome.

O. travancoricus, Ferguson, *Bomb. N. H. J.*, X, p. 71. Sarasin, *Zool. Jahr. Jena* 1910, p. 138.

COLOUR.—Dorsally light brown, with a series of dark brown or blackish, light-edged bars, 25 to 33 on the body, and 5 to 6 on the tail. These bars are frequently indented in the middle line anteriorly and posteriorly. Belly marked with squarish black and white dapplings.

On the head there is a dark light-edged transverse praefrontal bar reappearing below the eye, a similar transverse bar over the frontal passing obliquely over the temporal region, and a similar bar behind the parietals passing to the side of the neck. These bars may be more or less confluent in the median line.

FOOD AND BREEDING.—Nothing known.

GROWTH.—*Maximum Length.*—465 mm. (1 foot, 6½ inches). Tail 65 mm. (2¾ inches).

LEPIDOSIS.—(a) *Typical.*—As detailed in the synopsis.

(b) *Anomalies. Supralabials.*—The 6th sometimes reaches the edge of the lip, the 5th and 6th are partially confluent on one side in one specimen.

DENTITION.—From one skull in my collection. *Maxillary.*—An edentulous space anteriorly that would take 2 to 3 teeth, followed by 7 teeth. *Palatine.* Edentulous. *Pterygoid.* An edentulous space anteriorly that would take 5 to 6 teeth, followed by 12 teeth. *Mandibular.* An edentulous space anteriorly that would take 1 tooth, followed by 8 teeth.

DISTRIBUTION.—*Western Ghats.*—South of the Palghat gap. (High Range, Travancore. Tinnevelly Hills).

Oligodon cruentatus (Günther).

Simoles cruentatus, Boulenger, Cat. II, 1894, p. 231. *Faun. Mal. Pen. Rept.* 1912, p. 150. Flower, P. Z. S., 1896, p. 885. Wall and Evans, *Bomb. N. H.* J., XIII, pp. 349 and 617.

COLOUR.—Dorsally brown, with the anterior inferior borders of many scales edged lighter and darker to form a sparse variegation. A light vertebral stripe from the nape to the tail tip involving the vertebral and the next two rows. An indistinct dark stripe from the neck to the vent on the contiguous halves of the 3rd and 4th rows above the ventrals where the rows are 17, and of the 2nd and 3rd where 15. A blackish band on the base of the tail, and a similar band subterminally. Ventrally dappled with squarish black and yellow (white in spirit) spots to the vent. Crimson beneath the tail. Head with a dark transverse bar on the edge of the praefrontals and frontal reappearing below the eye. An oblique dark temporal bar from behind the gape to the parietal, usually meeting its fellow on the frontal. A dark sagitta with its point on the frontal or detached, and the arms passing to the side of the neck. In old specimens these head marks are more or less disintegrate and obscure. Chin whitish with some fine mottling on the lips.

FOOD.—In one specimen I found four soft-shelled eggs in the stomach dubiously snake's or lizard's.

BREEDING.—Nothing known.

GROWTH.—(a) *The Young.* I have never had a specimen smaller than 245 mm. (9½ inches). The annual growth cannot be estimated from my records.

(b) *Maximum Length.* 387 mm. (1 foot, 3½ inches).

LEPIDOSIS.—(a) *Typical.* As detailed in the synopsis.

(b) *Anomalies. Loreal.* Rarely absent, being confluent with the praefrontal. *Præocular.* Rarely two. *Supralabials.* I have seen the

2nd and 3rd confluent on one side in one specimen. *Infralabials*. Sometimes only four, in which case the 4th only touches the posterior sublinguals. *Subcaudals*. Rarely one or more at the base of the tail or entire.

DENTITION.—From two skulls in my collection. *Maxilla*. 14 to 16, no edentulous space. *Palatine*. 8 to 10, no edentulous space anteriorly or posteriorly. *Pterygoid*. 14 to 18, no edentulous space anteriorly. *Mandibular*. 17 to 18, no edentulous space anteriorly.

DISTRIBUTION.—*Burma*. Pegu, Toungoo, Rangoon, Minglegon, Yuathit, Prome District, Mandalay. Bhamo.

NOTE.—Stoliczka's record of Penang seems to me to call for confirmation.

Oligodon violaceus (Cantor).

Simotes violaceus, Boulenger, Cat. II, 1894, p. 222; l.c. III, 1896, p. 640. Slater, List Sn. Ind. Mus. 1891, p. 23. Wall and Evans, Bomb. N. H. J., XIII, pp. 350 and 618. Wall, Bomb. N. H. J., XII, pp. 672 and 766; l.c. XIX, p. 831.

COLOUR.—Very variable. I would divide individuals into the following categories, all of which however are completely connected by intermediate forms.

Var A. *violaceus* Cantor (=Var *cincereus* Günther). Dorsally varying shades of brown with sometimes a ruddy or *violaceus* tinge. Rarely the colour of a boiled prawn. There are no cross bars or stripes. Belly whitish with or without dark, squarish, lateral spots.

Var B. *multifasciatus* Jan (= *semifasciatus* Anderson). Usually darker brown than the last, the anterior inferior margins of many scales edged darker and lighter to form a variegation. The dark streaks congregate to form cross bars, varying from 19 to 27 on the body, and 3 to 7 on the tail. In most specimens an intermediate series of less distinct bars are to be seen, but in a few examples the intermediate series are as distinct as the former, and then there may be from 39 to 55 bars on the body. In some specimens all the bars are obscure and the belly is unspotted. To such Günther applied the name *swinhonis*. Belly with dark, squarish, lateral spots usually disposed on alternate sides and alternate ventrals. Some specimens exhibit an ill-defined, darker, longitudinal median stripe from the neck to tail tip, involving the vertebral and the next two and a half rows and enclosing a narrow pale vertebral stripe. A similar narrower lateral stripe passes from the neck to the vent, and involves the contiguous edges of the 2nd and 3rd rows above the ventrals. The head in all varieties has a dark *prefronto-frontal* cross bar which reappears below the eye. A similar dark oblique temporal bar passes from behind the gape to the parietals. A dark sagitta on the nape is projected forwards to the frontal, or has its point detached. These head marks are very distinct in some examples, obscure in others, especially in old specimens.

The two varieties referred to are completely connected by specimens showing every gradation, from faint (*swinhonis*) to dark cross bars, and every degree from scanty and small, to numerous and dark, squarish spots on the belly.

FOOD AND BREEDING.—Nothing known.

GROWTH.—(a) *The Young.* I have seen examples 178 and 185 mm. long (7 and 7½ inches) which appeared to have been recently born or hatched. No dates unfortunately are on record.

(b) *Maximum length.*—760 mm. (2 feet, 6 inches). Tail 75 mm. (3 inches).

LEPIDOSIS.—(a) *Typical.* As detailed in the synopsis.

(b) *Anomalies.* *Internasals.*—Quite divided by the rostral in one example. *Parietals.* Sometimes a small portion is detached to form a spurious upper anterior temporal. *Supralabials.* Rarely the 2nd and 3rd, or the 4th and 5th are confluent. The 3rd may be cuneate, and not reach the edge of the lip. In some specimens the 3rd as well as the 4th is divided into an upper and a lower part. In some the 4th is not so divided. *Infralabials.* The 3rd and 4th are sometimes confluent. *Subcaudals.* It is not very unusual to find some of these shields entire at the base of the tail.

DENTITION.—From one skull in my collection. *Maxillary.* 10 to 11, no edentulous space anteriorly. *Palatine.* 8, no edentulous space anteriorly or posteriorly. *Pterygoid.* 11 to 13, no edentulous space anteriorly. *Mandibular.* 15 to 16, no edentulous space anteriorly.

DISTRIBUTION.—*Assam.* Chittagong. Burma. Indo-China. South China.

Var (A) *violaceus.* Uniform or with bars and stripes very faint.

Burma.—Tounggyi, S. Shan States (Brit. Mus.). Rangoon. Mingledon. Arakan Hills. (F.W.). Indo-China. Camboja (Brit. Mus.). S. China. Hongkong (F.W.).

Var (B) *multifasciatus.* (a) With 19 to 27 distinct bars on body, 3 to 7 on tail, and less distinct intermediate bars. *Assam.* Cherrapunji, Khasi Hills. Samaguting, Naga Hills. *Bengal.* Tinsukia (F.W.). Chittagong Hills (Ind. Mus.). *Burma.* Mansi, Katha District. Tounggyi, S. Shan States (Bombay colln.). Kunchoung (F.W.).

(b) With 39 to 55 bars on the body, 4 to 8 on the tail; all equally distinct. *Assam.*—Nazira (Ind. Mus.). Halem (F.W.). Naga Hills (Anderson). *Burma.* Kunchoung (F.W.).

NOTE.—Boulenger mentions Bengal as part of its habitat. I cannot trace the authority for this. There is a specimen from the Chittagong Hills, which are politically in Bengal, in the Indian Museum, but I have found more than one specimen from the Eastern Himalayas in museums, wrongly identified, which proved to be *albocinctus*. (One such is specimen No. 16523 in the Indian Museum).

Oligodon venustus Jerdon.

O. venustus, Boulenger, Cat. II, 1894, p. 235. Sarasin, Zool. Jahr. Jena, 1910, p. 138. *Bomb. N. H. J.*, XXIII, p. 169; l.c. XXVI, p. 567.

COLOUR.—Dorsally brown, with a series of from 23 to 31 spots on the body, and from 6 to 7 on the tail. These spots are large, and shaped like walnut kernels. They are demarcated by a blackish areola, outside which is a pale border. Sometimes some of the spots are bisected in the median line. In many specimens a similarly-coloured, smaller spot is seen laterally especially in the fore body. Ventrally dappled with squarish black and yellow spots. Sometimes the black is replaced by a few small brownish spots disposed laterally. Head with a dark pia-

frontal bar reappearing below the eye, a dark oblique temporal streak from behind the gape to the parietal shields, usually meeting its fellow. A dark sagitta on the nape with its point on the frontal and usually confluent with the temporal stripe in the median line, and sometimes with the praefrontal bar. In old specimens the head marks may be obscure, and more or less disintegrated. Chin yellow with black sutures.

FOOD.—I have found reptilian eggs (lizard's, possibly snake's), and frog's spawn, with occasionally a snail with more or less of the shell intact.

BREEDING.—The young appear in the month of June and preceding months.

GROWTH.—(a) *The Young.* The smallest specimen I have seen, which appeared to be newly hatched or born, measured 112 mm. ($4\frac{1}{2}$ inches).

(b) *Early life.* Specimens measuring from 240 to 264 mm. ($9\frac{1}{2}$ to $10\frac{1}{2}$ inches) in August and September seem to indicate that the young double their length in the first year of life.

(c) *Maximum length.* 495 mm. (1 foot, $7\frac{1}{2}$ inches). The tail 70 mm. ($2\frac{3}{4}$ inches).

LEPIDOSIS.—(a) *Typical.* As detailed in the synopsis. The 6th supralabial is almost as frequently excluded from the edge of the lip, as extending to the labial margin.

(b) *Anomalies.* *Loreal.*—Frequently absent, being either confluent with the nasal, or the praefrontal. *Supralabials.* Rarely the 2nd and 3rd, or the 5th and 6th are confluent. Rarely there is a supernumerary making eight. *Costals.* Rarely the 3rd and 4th rows above the ventrals blend, and re-divide several times before the 15 rows are finally established.

DENTITION.—From seven skulls in my collection. *Mucillary.* An edentulous space anteriorly that would take 1 or 2 teeth, followed by from 7 to 8 teeth. *Palatine.* An edentulous space anteriorly that would take 2 to 3 teeth, followed by from 1 to 3 teeth. An edentulous space posteriorly that would take about 5 or 6 teeth. *Pterygoid.* An edentulous space anteriorly that would take about 5 teeth, followed by from 4 to 8 teeth. *Mandibular.* No edentulous space anteriorly. 9 to 11 teeth.

DISTRIBUTION.—*Western Ghats.* South of the Goa gap. Wynad. Nilgiris. Palnis. Cochin. Travancore.

Oligodon melanozonatus Wall.

O. erythrorhachis, Annandale, Rec. Ind. Mus. 1912, p. 48. *O. melanozonatus*, Wall, Rec. Ind. Mus., XXIV, p. 29.

TYPES.—In the Indian Museum (Nos. 16798 and 16799).

COLOUR.—Dorsally light brown obscurely mottled with blackish. A series of twenty rather ill-defined, black cross bars on the body, and four on the tail. In the smaller and half grown specimen these are edged anteriorly and posteriorly with black as in *albocinctus*. Ventrally white with transverse, black, irregularly-disposed cross bars, many as broad as the ventral shields. The tail is similarly marked beneath. Head with an obscure, dark, praefronto-frontal bar, reappearing below the eye,

a black-edged chevron with its point on the frontal and the arms passing obliquely over the temporal region. A pale, black-edged sagitta on the nape with its point directed forwards.

FOOD AND BREEDING.—Nothing known.

GROWTH.—*Maximum Length.* 513 mm. (1 foot, 8½ inches). Tail 83 mm. (3⅔ inches).

LEPIDOSIS.—(a) *Typical.* As detailed in the synopsis. In one specimen there are seven supralabials on the left side. A cuneate 6th is suggested on both sides in one example, which appears to be confluent with the 5th. The same may be said of the right side in the other specimen.

DISTRIBUTION.—*Assam.* Upper Rotung Valley, Abor Hills, at about 2,000 feet elevation.

***Oligodon erythrogaster* Boulenger.**

Simotes octolineatus, Günther, P. Z. S. 1861, p. 216. *Rept. Brit. Ind.*, 1864, p. 206 (footnote).

Oligodon erythrogaster, Boulenger, Rec. Ind. Mus., 1907, p. 216. Wall, *Bomb. N. H. J.*, XIX, p. 1000; l.c. XXII, p. 639.

TYPE.—In the Indian Museum (No. 16108).

COLOUR.—Dorsally brown, lighter in the flanks. A light vertebral stripe with a median dark line passes from the nape on to the tail, involving the vertebral and half the next row of scales. A blackish stripe passes from the nape to the tail tip involving the contiguous halves of the seventh and eighth rows above the ventrals, confluent on the neck with its fellow and passing forwards to the frontal, and similarly confluent on the tail. Three light stripes pass down each side of the body to the vent which are separated by narrow black stripes. Belly coral red in the median line to the end of the tail, with white, black-edged spots laterally, the edges of the ventrals light brown. Head with a dark praefrontal band reappearing below the eye, a dark oblique temporal stripe from the supraocular to behind the gape. A dark sagitta on the nape, its point (sometimes detached) on the frontal, confluent posteriorly with the uppermost dark body stripe.

FOOD.—Nothing known.

BREEDING.—An egg-bound female was killed on the 11th of June 1911, which contained four elongate eggs 27×5 mm. ($1\frac{1}{16} \times \frac{1}{5}$ of an inch). This specimen measured 455 mm. (1 foot 5¾ inches). Tail 75 mm. (3 inches).

GROWTH.—*Maximum Length.* The female just referred to is the largest of four specimens examined by me.

LEPIDOSIS.—(a) *Typical.* As detailed in the synopsis. The sixth supralabial is cuneate, and fails to reach the edge of the lip. In one example on one side the sixth supralabial has a small detached fragment which comes to the edge of the lip. The costals sometimes reduce posteriorly to 13 rows.

DENTITION.—From one skull in my collection. *Maxillary.* An edentulous space anteriorly that would take 2 to 3 teeth, followed by 7 to 8 teeth. *Palatine.* An edentulous space anteriorly that

would take 2 to 3 teeth, followed by 2 to 3 teeth. An edentulous space posteriorly that would take 5 to 6 teeth. *Pterygoid*. An edentulous space anteriorly that would take about 6 teeth, followed by 4 teeth. *Mandibular*. An edentulous space anteriorly that would take 1 to 2 teeth, followed by 5 to 6 teeth. An edentulous space posteriorly that would take about 5 teeth.

DISTRIBUTION.—*Eastern Himalayas*. Nepal to Sikkim. Tindharia (F.W.).

Oligodon theobaldi (Günther.)

Sinotes beddomii, Boulenger, Cat. II, 1894, p. 229. Sarasin, Zool. Jahr. Jena, 1910, p. 138. Wall, Bomb. N. H. J. XXIII, p. 170.

Sinotes theobaldi, Boulenger, Cat. II, 1894, p. 230. Slater, List. Su. Ind. Mus. 1891, p. 24. Wall and Evans, Bomb. N. H. J. XIII, pp. 350 and 618. Wall, Bomb. N. H. J. XVIII, p. 783; l.c. XXIII, p. 170.

COLOUR.—Dorsally brown of various shades, the anterior inferior borders of many scales edged lighter and darker to form a variegation. A pale vertebral line from the nape on to the tail. A dark stripe from the nape to the tail tip involving the edge of the vertebral and the next three rows. A dark stripe passes from the neck to the vent on the confines of the 3rd and 4th rows above the ventrals where the scales are 17, and on the 2nd and 3rd where 15. Ventrally yellow, immaculate, or with a few scattered dark spots posteriorly. Head with a dark prefrontal bar reappearing below the eye, an oblique dark temporal stripe from behind the gape meeting its fellow on the frontal to form a chevron. A dark chevron on the nape, its point (sometimes detached) passing to the posterior angle of the frontal and its arms to the side of the neck. Chin and throat immaculate.

FOOD.—Nothing known.

BREEDING.—I have seen a gravid female with three eggs measuring 18 mm. ($\frac{3}{4}$ of an inch) in length. The dam was 295 mm. ($11\frac{1}{2}$ inches) long. The date of capture is not on record. Judging from these eggs it is probably oviparous.

GROWTH.—*The Young*. My smallest specimen was 115 mm. (4½ inches) long, but it is fairly certain the young when hatched or born are appreciably less than this.

(b) Maximum Length. 380 mm. (1 foot, 3 inches). Tail 30 mm. ($1\frac{1}{6}$ inches.)

LEPIDOSIS.—(a) Typical. As detailed in the synopsis.

(b) Anomalies. *Supralabials*. I have seen the 5th confluent with the lower postocular. *Infralabials*. In some specimens the 5th only touches the posterior sublinguals. *Subcaudals*. I have seen the last two entire in one example.

DENTITION.—From two skulls in my collection. *Maxillary*. 15 to 16 teeth; no edentulous space anteriorly. *Palatine*. 9 teeth; no edentulous space anteriorly or posteriorly. *Pterygoid*. 16 to 18 teeth; no edentulous space anteriorly. *Mandibular*. 16 to 17 teeth; no edentulous space anteriorly or posteriorly.

DISTRIBUTION.—*Assam.* Tura, Garo Hills (Bombay colln.). *Burma.* As far north as Myitkyina (Bombay colln.) and south to Tenasserim (Ind. Mus.).

NOTE.—The locality of specimen No. 11712 in the Indian Museum labelled Karachi is obviously a mistake. Its identity is correct. Wynad, on the authority of Beddome (types of *beddomei* in the British Museum), must also be discredited.

Oligodon affinis Günther.

O. affinis, Boulenger, Cat. II, 1894, p. 236. Ferguson, *Bomb. N. H. J.*, X, p. 70. Sarasin, *Zool. Jahr. Jena*, 1910, p. 138. Wall, *Bomb. N. H. J.* XXVI, p. 568.

COLOUR.—Brown dorsally, the anterior inferior borders of many scales edged darker to form a sparse variegation. A series of from 31 to 41 dark, light edged, linear cross bars over the body becoming indistinct or obsolescent on the tail. These bars involve about 5 to 7 scale rows in the breadth of the snake. Ventrally boldly dappled with squarish black and yellow spots, from throat to tail tip, many of which may be confluent across the belly. Head with a dark præfronto-frontal bar reappearing below the eye, a dark oblique temporal streak from the gape meeting its fellow across the parietals to form a chevron, and usually confluent with the præfrontal bar in the median line. A thin chevron on the nape, its point confluent with the preceding chevron, and the arms passing to the sides of the neck. Chin white with black sutures.

FOOD.—I have found the soft shelled eggs of a lizard (or snake?) in the stomach.

BREEDING.—A specimen 100 mm. (4 inches) in length captured in July indicates that the season for the appearance of the young is somewhat earlier in the year.

GROWTH.—(a) *The Young.* My smallest specimen was 100 mm. (4 inches), but from what we know of other species it is almost certain that the young are even smaller than this when hatched or born.

(b) *Maximum Length.* 342 mm. (1 foot 1½ inches.) Tail 44 mm. (1¾ inches).

LEPIDOSIS.—(a) *Typical.* As detailed in the synopsis.

(b) *Anomalies.* *Parietals.* Sometimes a detached fragment forms a spurious upper anterior temporal. *Loreal.* Rarely present. *Supralabials.* The 2nd in one specimen on one side was cuneate, and failed to reach the edge of the lip.

DENTITION.—From one skull in my collection. *Maxillary.* An edentulous space anteriorly that would take 2 teeth, followed by 7 teeth. *Palatine.* With a single tooth near its middle. *Pterygoid.* An edentulous space anteriorly that would take 5 to 6 teeth, followed by 4 teeth. *Mandibular.* An edentulous space anteriorly that would take 1 tooth, followed by 8 teeth. An edentulous space posteriorly that would take 3 teeth.

DISTRIBUTION.—*Western Ghats.* South of the Goa gap. Wynad to Travancore.

Oligodon arnensis (Shaw).

Simotes arnensis, Abercromby, *Sn. of Ceylon*, 1910, p. 72. Boulenger, *Cat. II*, 1894, pp. 229 and 359. Ferguson, *Bomb. N. H. J.*, X, p. 71. Sarasin, *Zool. Jahr. Jena*, 1910, p. 130. Sclator, *List Sn. Ind. Mus.*, 1891, p. 24. Wall, *Bomb. N. H. J.*, XVIII, p. 115; l.c. XIX, p. 532; l.c. XXII, p. 749. Willey, *Spol. Zeylan.* 1906, p. 233.

COLOUR.—Dorsally brown of various shades, with sometimes a purplish or ruddy tinge. The whole body and tail are crossed by black bars edged with yellow or buff. These bars vary considerably in number according to locality. In Ceylon specimens there are from 13 to 18 on the body and 3 to 6 on the tail. In Indian specimens south of the Nerbudda and the Ganges Valley, they usually number from 19 to 30 (15 and 16 in two Travancore examples) on the body, and 4 to 16 on the tail. In specimens north of the Nerbudda and in the Ganges Valley they are usually from 28 to 40 on the body (47 in a specimen from Bihar), and 7 to 20 on the tail. Where the number is small the bars involve from 4 to 5 scales vertebrally in the fore body, 2 to 3 behind; but where it is large they involve from 1 to 2 scales vertebrally. These bars usually break up in the flanks, and in some specimens there is a darker variegation to be seen in the flanks intermediate to the bars. In some specimens the bars are replaced by twin, or quadrimaculate transversely-placed spots as one sees in variety *fasciatus* of *tenuirostris*. The belly is partly white and usually immaculate, but some specimens show a series of dark spots placed laterally and with a tendency to be disposed on alternate ventrals, and alternate sides. The head has a dark prefronto-frontal bar, which reappears below the eye, a dark oblique temporal streak from behind the gape, which usually meets its fellow on the frontal to form a chevron. A short dark oblique streak is frequently seen in the suture between the 5th and 6th supralabials. A dark chevron is placed on the nape with its point in the interparietal suture, and its arms pass to the side of the neck. Where there are few cross bars on the body this chevron involves from 8 to 10 or more scales vertebrally, but where the cross bars are numerous it involves from 3 to 4 scales vertebrally. All the head markings are almost always discrete, very distinct, and usually edged with buff or yellow.

FOOD.—On two occasions I have found the soft shelled eggs of a lizard (or snake?) in the stomach. A plug of hair in the cloaca of one example indicates that it will eat small mammals.

BREEDING.—(a) *Method of Reproduction.* Eggs I have taken from the abdomen were of such a size and character as to leave little doubt that it is oviparous in habit.

(b) *Season.* I had a female in Fyzabad with eggs seemingly fit for discharge in August, and another with less advanced eggs in the same month.

(c) *The Eggs.* I have seen half a dozen or more egg-bound females and find that the clutch numbers from two to five. The most advanced eggs measured 36 by 10 mm. ($1\frac{4}{10}$ by $\frac{7}{10}$) of an inch. No embryo could be discovered within.

GROWTH.—(a) *The hatchling.* I have had three young which appeared to be hatchlings and which measured from 165 to 170 mm. ($6\frac{1}{2}$ to $6\frac{3}{4}$ inches).

(b) *Maturity.* The smallest egg-bound female I have seen measured 442 mm. (1 foot, $5\frac{1}{2}$ inches).

(c) *Maximum Length.* 635 mm. (2 feet, 1 inch) is the largest measurement I know.

LEPIDOSIS.—(a) *Typical.* As detailed in the synopsis.

(b) *Anomalies.* *Rostral.* I have once seen this entirely separating the nasals. *Parietals.* Sometimes a small fragment is detached to form a spurious upper anterior temporal. *Loreal.* Not infrequently absent, being confluent with either the praefrontal or the posterior nasal. *Præoculars.* Rarely two. *Supralabials.* Two are sometimes confluent, the series counting six in consequence. The 2nd is rarely divided and the series then counts eight. The 6th is sometimes cuneate, and fails to reach the edge of the lip. The 6th and 7th are rarely confluent. *Infralabials.* Occasionally there are five. *Anal.* I have twice seen this entire. *Subcaudals.* Rarely a few at the base of the tail are divided.

DENTITION.—From three skulls in my collection. *Maxillary.* An edentulous space anteriorly that would take about 3 teeth, followed by from 8 to 11 teeth. *Palatine.* An edentulous space anteriorly that would take about 2 teeth, followed by from 3 to 6 teeth. An edentulous space posteriorly that would take about 4 teeth. *Pterygoid.* An edentulous space anteriorly that would take about 4 to 6 teeth, followed by 10 to 17 (?18) teeth. *Mandibular.* No edentulous space anteriorly, 13 to 14 teeth.

DISTRIBUTION.—*Ceylon.* *Peninsular India.* To Sind and Baluchistan in the North-West. North-West Frontier, Western Himalayas to 5,500 feet. To Bihar and Lower Bengal in the North-East. (Calcutta. Kaliganj, Rangpur District.)

***Oligodon woodmasoni* (Sclater).**

Simotes woodmasoni, Annandale, J. A. S. Beng. 1905, pp. 173 and 175. Bouleenger, Cat. II, 1894, p. 223.

TYPES.—In the Indian Museum (Nos. 8459 and 12547).

COLOUR.—Dorsally brown with a ruddy tinge. A light vertebral stripe from the nape well on to the tail. Three other narrower light stripes pass down each side of the body from the neck to the vent. These are separated by blackish stripes. As age advances the blackish stripes tend to fade so that the larger of the two specimens seen by me is nearly uniform in colour. Belly dusky reddish beneath, the edges of the ventrals lighter. Head with a dark praefrontal bar reappearing below the eye. A dark streak on the first three supralabials to the anterior nasal. An oblique temporal streak from behind the gape to the parietals. A dark sagitta on the nape projected forwards to the frontal shield.

FOOD AND BREEDING.—Nothing known.

GROWTH.—(a) *Early Life.* A specimen measuring 177 mm. ($6\frac{1}{2}$ inches) appears to be a hatchling.

(b) *Maximum Length.* 600 mm. (1 foot, $11\frac{5}{8}$ inches). Tail slightly deficient.

LEPIDOSIS.—(a) *Typical.* As detailed in the synopsis.

(b) *Anomalies.* *Loreal.* Absent on one side in one specimen.

DISTRIBUTION.—*Andamans, Nicobars,*

Oligodon albocinctus (Cantor).

Simotes albocinctus, Annandale, Rec. Ind. Mus., 1912, pp. 37, 48 and 53, Boulenger, part. Cat. II, 1894, p. 220; Rec. Ind. Mus., 1913, p. 338. Evans, Bomb. N. H. J., XVI, p. 169. Slater, List Sn. Ind. Mus., 1891, p. 23. Venning, Bomb. N. H. J., XX, p. 338. Wall, Bomb. N. H. J., XIX, pp. 348, 757a, 830 and 898; i.e. XXII, p. 756.

COLOUR.—Variety (A) *O. albocinctus* (Cantor). Dorsally brown of various shades from the very lightest to the deepest hue, even to blackish brown. Many specimens exhibit a ruddy tinge which may be pronounced especially in the flanks. More rarely the prevailing hue is salmon pink. The back bears a series of whitish, greyish, buff, yellow or dusky brown cross-bars. These bars are well defined with black outside and have often a thin edging of buff or yellow. In ruddy specimens they are yellow. They are very well defined and conspicuous in light specimens, but in very dark specimens they may be very obscure. They number from 19 to 27 on the body and 4 to 8 on the tail. Frequently some intermediate scales in the flanks are edged with black and form a patch of variegation. In the space preceding the first bar there are usually two dark lines, parallel or curved so as to form an incomplete ellipse. In some specimens two more or less distinct, but ill-defined darker stripes pass down the body. The upper and broader passes from the neck well on to the tail and involves the edge of the vertebrals, and the next two rows. The lower passes from neck to vent, and involves the contiguous halves of the 3rd and 4th rows above the ventrals. The belly is whitish, yellowish or brownish (pinkish in the ruddy tinged specimens) with squarish, brown or blackish, lateral spots or mottling, the black in dark specimens often predominating. The head has a dark pre-fronto-frontal bar which reappears below the eye, and a dark oblique temporal bar from behind the gape to the parietals. The brown on the neck sends forwards a projection to the frontal shield, the tip of which is often detached. These head marks are usually well defined with blackish outside with a narrow edging of buff or yellow. In ruddy specimens they are yellow. The chin and throat are usually immaculate or there may be a dusky mark in the suture between the 4th and 5th infralabials.

Variety (B) *amabilis* (Günther). Differs from the preceding in having an intermediate and less conspicuous series of cross-bars. These specimens very closely resembles variety *multifasciatus* of *violaceus*, and I have found several in museums wrongly identified as *violaceus*.

FOOD.—I have only three records that reveal the nature of the diet. One contained a mouse, another a newly born mouse, and a third the tail of a mouse.

BREEDING.—My only record of a gravid female is one killed at Dibrugarh in July. Only three follicles were impregnated. I have records of well over one hundred specimens and the dearth of egg-bound females is therefore remarkable.

GROWTH.—(a) *The Young*.—The smallest I have seen and which appeared to be recently hatched or born was 200 mm. (8 inches) long.

(b) *Maturity*.—My only gravid female was 742 mm. (2 feet, 5½ inches) long.

(c) *Maximum Length*.—915 mm. (3 feet). Tail (imperfect) 100 mm. (4 inches).

LEPIDOSIS.—(a) *Typical*.—As detailed in the synopsis.

(b) *Anomalies*.—*Loreal*. Rarely absent, being confluent with the praefrontal. *Temporals*. The anterior is rarely confluent with the posterior superior shield. *Supralabials*. I have seen the 1st divided into an upper and a lower part in one example on both sides. The 2nd or 3rd are rarely divided vertically, thus making 8 shields in the series. The 5th and 6th, or the 6th and 7th are rarely confluent making the series 6. The 6th is rarely confluent with the anterior temporal. *Costals*. I have once seen 21 rows. The rows rarely reduce to 17 two heads-lengths before the vent. *Subcaudals*. Some at the base of the tail are rarely entire.

DENTITION.—From three skulls in my collection. *Maxillary*. 10 to 12, no edentulous space anteriorly. *Palatine*. No edentulous space anteriorly, 8 to 10 teeth, followed by an edentulous space that would take 1 to 2 teeth. *Pterygoid*. An edentulous space anteriorly that would take 1 tooth, followed by 18 to 23 teeth. *Mandibular*. 15 to 18 teeth. No edentulous space anteriorly or posteriorly.

DISTRIBUTION.—Variety (A) *Eastern Himalayas*. Buxa Dooars to Sikkim. *Assam*. Hills and plains as far north as the Abor Hills and south to Chittagong in Eastern Bengal. *Burma*. Kachin Hills. (Sadon. Lat. 25° 4'. Long. 98°. Sima. Lat. 25° 2'. Long. 97°.) Manipur (Assam). Chin Hills.

Variety (B) *Assam*. Plains and hills. *Burma*. Arakan Hills.

Oligodon juglandifer (Wall).

Simotes albocinctus, part. Boulenger, Cat. II, 1894, p. 220. Wall, part. *Bomb. N. H. J.*, XIX, p. 348.

Simotes juglandifer, Wall, *Bomb. N. H. J.*, XX, p. 1162.

TYPE.—In the Bombay Natural History Society's Collection (No. 210) from Tindharia, Darjiling District.

COLOUR.—Dorsally brown of various shades from deep brown to blackish brown. Many scales edged buff and black to form a variegation which is most pronounced in the flanks. A series of from 22 to 28 large dark brown or black spots on the body, and 6 to 8 on the tail. These spots are shaped like walnut kernels, having an indentation anteriorly and posteriorly, but are sometimes completely bisected. In some specimens a smaller spot is to be seen outside and close to the median spots. In some specimens there is an intermediate series of less distinct and smaller spots arranged transversely. In some specimens four dark stripes may be seen, the upper and broader passing from the neck to tail tip and involving the edge of the vertebrals and next two rows, sometimes confluent with that of the opposite side. A narrow dark stripe passes from the neck to the vent on the edges of the 3rd and 4th rows above the ventrals. Belly whitish, buff or brownish with a series of squarish, deep brown or black lateral spots, usually disposed on alternate ventral shields, those of one side usually alternating with those on the other. Head with a dark praefronto-frontal bar reappearing below the eye. A dark oblique black-edged bar from behind the gape to the parietals. A dark black-edged mark on the nape sends forward a projection to the frontal, the point of which is often detached. Chin and throat immaculate, or with a dark mark in the suture between

the 4th and 5th infralabials. I have seen specimens the colour of a boiled lobster.

FOOD AND BREEDING.—Nothing known.

GROWTH.—(a) *The Hatchling*.—My smallest specimen possibly a hatchling measured 233 mm. (9 $\frac{1}{4}$ inches).

(b) *Maximum Length*.—853 mm. (2 feet, 9 $\frac{3}{4}$ inches).

LEPIDOSIS.—(a) *Typical*.—As detailed in the synopsis.

(b) *Anomalies*.—*Supralabials*. The 5th and 6th are rarely confluent reducing the series to six. *Costals*. In one specimen the 4th or 5th row above the ventrals divided and re-united several times so that they counted 21 in places.

DENTITION.—From four skulls in my collection. *Maxillary*.—No edentulous space anteriorly, 10 to 12 teeth. *Palatine*. An edentulous space anteriorly that would take 1 or 2 teeth followed by 6 to 8 teeth. An edentulous space posteriorly that would take 2 or 3 teeth. *Pterygoid*. An edentulous space anteriorly that would take one tooth, followed by from 16 to 18 teeth. *Mandibular*. No edentulous space anteriorly or posteriorly, 13 to 17 teeth.

DISTRIBUTION.—*Eastern Himalayas*. Sikkim. Assam.

Oligodon purpurascens¹ (Schlegel.)

Simotes purpurascens, Boulenger, Cat. II, 1894, p. 219. *Faun. Mal. Pen. Rept.*, 1912, p. 148. Flower, P. Z. S., 1896, p. 884; l.c. 1899, p. 671. Malcolm-Smith, Journ. N. H. Siam, 1915, p. 213. Moequard, Rept. L'Indo-Chine, 1907, p. 46.

Simotes cyclurus, Boulenger, Cat. II, 1894, p. 219. *Faun. Mal. Pen. Rept.* 1912, p. 149. Flower, P. Z. S., 1896, p. 884; 1899, p. 671. Malcolm-Smith, Journ. N. H. Siam, 1914, p. 97; l.c. 1915, pp. 213 and 245. Moequard, Rept. L. Indo-Chine, 1907, p. 46. Wall and Evans, Bomb. N. H. J. XIII, pp. 350 and 617. Wall, P. Z. S., 1903, p. 92. Bomb. N. H. J., XVIII, p. 781; l.c. XIX, p. 348. Werner, Abh. Akad. der. Wiss., 1903, p. 365.

COLOUR.—This, like *laeniolatus*, is an extremely variable species. There are two main types of colouration. (A) *Oligodon purpurascens cyclurus* (Cantor). Dorsally the prevailing hue varies between a pale brown, and blackish brown. Usually some of the scales are narrowly edged with blackish, and others with buff thus producing a variegation. The darker variegation tends to form narrow cross-bars involving one or two scale rows in the length of the snake. In many specimens these cross-bars are conspicuous on the body and tail, and most of these exhibit a less conspicuous series of intermediate bars. Many specimens also exhibit four more or less conspicuous, darker, longitudinal stripes. The upper and broader of these passes from the neck to the tail tip, involving the edge of the vertebral and the next one and a half rows, leaving a pale linear stripe between them, but in many specimens these stripes are confluent vertebrally. The lower and narrower stripe passes from the

¹ I have for some years suspected that *purpurascens* (Schlegel) and *cyclurus* (Cantor) would prove to be one and the same species. My last visit to England this year (1922) gave me an opportunity of examining all the species labelled *purpurascens* and *cyclurus* in the British Museum. I examined the two supposed species side by side, and made most careful measurements of head shields and tried in vain to find some difference by which the two could be separated. The scale rows and ventrals and subcaudals agree in their ranges, and one is forced to the conclusion that they should be regarded as a single species. Schlegel's *purpurascens* (1837) has precedence over Cantor's *cyclurus* (1839). I have also examined equally carefully a large series of both *cyclurus* and *purpurascens* in the Indian Museum, and Bombay Natural History Society's collections.

neck to the vent, and involves the contiguous halves of the 3rd and 4th rows above the ventrals. These stripes, in the young especially, are sometimes well defined and conspicuous; in many specimens however they are ill-defined, inconspicuous, or even absent. The belly is whitish or yellowish, sometimes immaculate, sometimes with a few scattered brownish, squarish, lateral spots especially posteriorly, and sometimes so heavily spotted that blackish brown is the predominant colour. The head has a dark praefronto-frontal bar which reappears below the eye, and a dark oblique temporal stripe from behind the gape to the parietal shields. A dark sagitta on the nape sends forwards a process to the frontal shield, the point of which may be detached. In some specimens this is more or less confluent with the oblique temporal stripe. The chin and throat are usually immaculate or there may be some dusky marks in some of the sutures.

Specimens may be grouped as follows:—

A (a). Very pale brown. Variegations absent or faint. Stripes absent or faint. Belly immaculate.

A (b). Brown. Variegations (and consequently the cross-bars) more or less conspicuous. These number from 9 to 17 on the body, 3 to 4 on the tail, or twice this number if the narrower intermediate series are counted. Stripes absent or faintly suggested. Belly with a few dark spots mostly posteriorly.

A (c). Dark Brown. Variegations forming conspicuous narrow dark bars. Stripes conspicuous. Belly heavily spotted. These sub-varieties are all completely connected by intermediate forms.

(B) *O. purpurascens maculatus*. Differs from A in having a series of large dark spots across the back, involving four or more scale rows vertebrally in the length of the snake, and reminding one of the ornamentation seen in *venustus*, *splendidus* and variety *spilonotus* of *taeniatus*. There are from 9 to 13 (16 Malcolm-Smith) on the body, and 2 to 4 on the tail. Sometimes these are indented mesially or even completely bisected, especially in the fore body and on the tail. Sometimes they are more or less confluent with a smaller spot placed outside them. Some of these specimens exhibit some variegation as in variety A (b) and a series of cross-bars between the large dark spots. Some have stripes similar to variety A (c). The belly is immaculate. Head marks as in A. This variety is rare within Indian limits, but the common one in Siam.

Variety (C). *O. purpurascens purpurascens* (Schlegel). This differs from A in having the narrow cross-bars light instead of dark. In the British Museum specimen from Borneo there are 12 such bars on the body and 3 on the tail. A somewhat similar specimen in the same museum I would label *albocinctus* were it not for the locality, viz., Java. I count the ventrals 182 and subcaudals 41. This differs from all the other specimens (labelled *purpurascens*) in the British Museum in that the third supralabial is not divided into an upper and lower part, and does not touch the eye. In this respect as in others it agrees with *albocinctus*.

Variety (D). An unusual specimen in the British Museum from Borneo, donor Mr. Shortridge, has 11 broad cross-bars on the body and 3 on the tail. These cross-bars are not indented mesially, are bordered with black posteriorly, and by double black lines anteriorly enclosing a

whitish zone. There are also black bordered light annular spots dorso-laterally in the bars.

FOOD.—In one instance a toad had furnished the meal. In two specimens I have found the soft-shelled eggs of a lizard (or snake ?) in the stomach. On one occasion Mr. Jacob, I. F. S., sent me four soft shelled eggs, which he suspected had been laid by a *cyclurus* which he found in close proximity. On investigating the eggs which measured 27×16 mm. ($1\frac{1}{6} \times \frac{2}{3}$ of an inch) I extracted embryos that were about 62 mm. (2½ inches) long. Large ventral shields were evident, but there were no longitudinal series of costals, and the head shields were not of an ophidian character. I very much suspect these were lizards' eggs, and that the snake intended to make a meal of them.

BREEDING.—Nothing known.

GROWTH.—(a) *The Young.*—The smallest example of *cyclurus* I have examined, apparently newly hatched or born, measured 175 mm. (7 inches). A *purpurascens* in the British Museum is 207 mm. (8½ inches).

(b) *Maximum length.*—The largest variety *cyclurus* of which I have any knowledge is one in the Indian Museum [No 7163 type of *O. crassus* (Theobald)] which measures 946 mm. (3 feet, 1½ inches). Flower records a *purpurascens* 950 mm. (3 feet, 1½ inches).

LEPIDOSIS.—(a) *Typical.*—As detailed in the synopsis. The 4th supralabial is normally divided into an upper and a lower part both of which touch the eye.

(b) *Anomalies.*—*Supralabials.* Sometimes the 2nd and 3rd, 3rd and 4th, or 4th and 5th are confluent, thus reducing the series to 7. Rarely the 3rd or the 7th (in one specimen both) are cuneate, and fail to reach the edge of the lip. Sometimes the 4th is an entire shield. Sometimes both the 3rd and 4th are divided. Rarely the 3rd only is divided and the 3rd, 4th and 5th touch the eye. *Infralabials.* Sometimes the 3rd and 4th are confluent. *Costals.* Rarely the 4th or 5th row above the ventrals divides and may re-unite and redivide repeatedly so as to make 21 rows in places. Rarely the rows are 17 at a spot two head-lengths behind the head becoming 19 shortly after. Rarely again the 4th and 5th rows unite and redivide anteriorly so as to reduce the rows in places to 17.

DENTITION.—From four skulls of *cyclurus* and two of *purpurascens* in my collection. *Maxillary.* 9 to 10 teeth in *cyclurus*, 9 to 10 in *purpurascens*. No edentulous space anteriorly; 6 to 8 teeth in *cyclurus*, 9 in *purpurascens*, an edentulous space posteriorly that would take 1 or 2 teeth. *Pterygoid.* An edentulous space anteriorly that would take 1 or 2 teeth, followed by 11 to 16 teeth in *cyclurus*, 13 to 18 in *purpurascens*. *Mandibular.* 13 to 16 teeth in *cyclurus*, 15 to 16 in *purpurascens*. No edentulous space anteriorly or posteriorly.

DISTRIBUTION.—*Bengal, Eastern Himalayas, Assam, Burma, Malay Peninsula, Siam, Cochin-China, South China.*

Variety A (a). *Bengal.* Calcutta, Jessore, Purnea District. *Burma.* Pegu, Rangoon. *Siam.* Bangkok.

Variety A (b). *Bengal,* Calcutta, Rangpur District, Purnea District, Jalpaiguri District. *Eastern Himalayas.* Sikkim. *Assam.*

Garo Hills, Khasi Hills, North Cachar. *Burma*. Rangoon, Diamond Island, Amherst, Tavoy.

Variety A (c). *Bengal*. Jalpaiguri District. *Eastern Himalayas*. Sikkim. *Assam*. Garo Hills, Khasi Hills. *Burma*. *Siam*.

Variety B. *Burma*. Toungoo, Karen Hills, Kalaw, S. Shan States, Tavoy. *Siam*. (The common form.)

Variety C.—Borneo. Java?

Variety D.—Borneo.

Oligodon splendidus (Günther.)

Simotes splendidus, Boulenger, Cat. II, 1894, p. 217. Evans, *Bomb. N. H. J.*, XVI, p. 362. Sarasin, *Zool. Jahr. Jena*, 1910, p. 142. Venning, *Bomb. N. H. J.*, XXIII, p. 164. Wall and Evans, *Bomb. N. H. J.*, XIII, p. 537. Wall, *Bomb. N. H. J.*, XVIII, p. 781. *Rec. Ind. Mus.*, II, p. 105.

COLOUR.—Pale brown dorsally, each scale with a dark centre. A series of very large, dark, median spots shaped like walnut kernels, with smaller lateral spots below. These spots have an outer zone of dark brown which is edged with buff. There are from 14 to 17 of these spots on the body, and from 3 to 5 on the tail. Belly whitish or yellowish, with squarish, dark brown, lateral spots, usually on alternate ventrals and alternate sides. Head with an indistinct, dark, praefronto-frontal bar reappearing below the eye, and an indistinct dark oblique streak from behind the gape to the parietal. A large dark, mark on the nape, indented mesially behind, is projected forward to the frontal shield.

FOOD.—Nothing known.

BREEDING.—(a) *Method of Reproduction*.—I have seen eggs of such a size and character as to make it tolerably certain that this species is oviparous.

(b) *Season*.—Venning acquired a female in an advanced state of egg bearing on the 14th of February.

(c) *The Eggs*.—I have seen two egg-bound females which contained three and six eggs, respectively. The largest eggs measured 457 x 152 mm. ($1\frac{1}{2} \times \frac{1}{2}$ an inch). No trace of an embryo could be discovered.

GROWTH.—(a) *The Hatchling*.—The smallest I have seen and apparently a hatchling measured 155 mm. ($6\frac{1}{8}$ inches).

(b) *Maturity*.—The smaller of two egg-bound females measured 465 mm. (1 foot, 9 $\frac{3}{4}$ inches).

(c) *Maximum Length*.—730 mm. (2 feet, 4 $\frac{3}{4}$ inches). Tail 100 mm. (4 inches).

LEPIDOSIS.—(a) *Typical*.—As detailed in the synopsis.

(b) *Anomalies*.—*Praefrontals*. Rarely four in a transverse series. *Postoculars*. Rarely three. *Supralabials*. The 4th is sometimes divided into two upper and one lower part.

DENTITION.—From one skull in my collection. *Maxillary*. No edentulous space anteriorly; 11 teeth. *Palatine*. No edentulous space anteriorly or posteriorly; 9 teeth. *Pterygoid*. No edentulous space anteriorly; 14 to 15 teeth. *Mandibular*. No edentulous space anteriorly or posteriorly; 13 to 14 teeth.

DISTRIBUTION.—*Burma*.—Between Lat. 20° and 23° , and Long. 94° and 98° . Yamethin, Sagaing, Pyawbwe, Shwebo, Monywa, Mandalay, N. Shan States (Ruby Mines), S. Shan States (Kyaukse, Pakokku).

Synopsis of the salient features of lepidosis and dentition differentiating the species of Oligodon.

COSTALS.		VENTRALS.		Two heads-lengths before vent.		Two heads-lengths behind head.		Midbody.		Number.		Anal.		Subcaudals.		Rostral shields in contact.		Inter nasals.		NASALS.		SUPRALABIALS.		INFRA-LABIALS.		MAXILLA.		PALATINE.		PTERYGOID.		MANDIBLE.		Parietal contributing to orbital ring.	
13	13	13	132 to 142	22 to 27	8	2	2	22 to 27	37 to 40	6	0	+	+	Entire.	Postocular.	1	1	1	1	3rd & 4th	3rd & 4th	4 or 5	4	3rd	Touching the eye.	4th	No. of teeth.	No. of teeth.	Edentulous space anteriorly.	No. of teeth.	Edentulous space anteriorly.	planiceps.	
13	13	13	186 to 208	37 to 40	6	2	1	37 to 40	6	2	Semidivided.	Postocular.	1	1	1	1	4th & 5th	5th & 6th	6	4	4th	Touching ant temporals.	5th	15 to 16	0	8 to 10	0	Edentulous space anteriorly.	Edentulous space posteriorly.	herberti.
13	13	13	200	27 to 34	39	6	2	27 to 34	6	2	Divided.	Loreal.	0	1	1	1	5th & 6th	6th	7	4	4th	Touching post sublinguals.	6th	13 to 16	13 to 16	13 to 16	Edentulous space anteriorly.	Edentulous space anteriorly.	medougalli.	
15	15	13	144 to 159	46 to 64?	6	2	27 to 34	6	2	Postocular.	1	1	1	1	3rd & 4th	3rd & 4th	4	4	4th	Postocular.	7	15 to 16	0	8 to 10	0	Edentulous space anteriorly.	Edentulous space posteriorly.	torquatus.
15	15	13	154	46 to 64?	6	2	46 to 64?	6	2	Postocular.	0	1	1	1	5th & 6th	6th	7	4	4th	Postocular.	8	13 to 16	13 to 16	13 to 16	Edentulous space anteriorly.	Edentulous space anteriorly.	erythrorhachis.	
15	15	13	162 to 188	27 to 51	6	2	27 to 51	6	2	Postocular.	1	1	1	1	3rd & 4th	3rd & 4th	4	4	4th	Postocular.	7	6 to 7	6 to 7	6 to 7	Edentulous space anteriorly.	Edentulous space anteriorly.	dorsalis.	
15	15	15	158 to 173	25 to 29	6	0	29 to 56	6	2	Postocular.	1	1	1	1	5th & 6th	6th	7	4	4th	Postocular.	7	1 to 2	1 to 2	1 to 2	Edentulous space anteriorly.	Edentulous space anteriorly.	brevicauda.	
15	15	15	158 to 218	29 to 56	6	2	29 to 56	6	2	Postocular.	1	1	1	1	5th & 6th	6th	7	4	4th	Postocular.	7	5 to 6	5 to 6	5 to 6	Edentulous space anteriorly.	Edentulous space anteriorly.	taeniolatus.	

Synopsis of the salient features of lepidosis and dentition differentiating the species of Oligodon—contd.

COSTALS.		VENTRALS.		Subcaudals.		Rostral shields in contact.		NASALS.		SEPHALO- LABIALS.		INFRA- LABIALS.		MAXI- LLA.		PALATINE.		PTERY- GOID.		MANDIBLE.		Two heads-lengths behind Head.						
MIDBODY.	TWO HEADS-LENGTHS BEFORE VENT.	NUMBER.	ANGULATE.	ROUNDED.	ANAL.	INTERNASSAS.	ENTIRE.	SEMITDIVIDED.	DIVIDED.	LOREAL.	POSTOCULAR.	ANTERIOR TEMPORAL.	NUMBER.	TOUCHING THE EYE. TOUCHING ANT TEMPORAL, PARAL.	NUMBER.	TOUCHING POST SUBLINGUALS.	NO. OF TEETH.	EDENTATEOUS SPACE ANTERIORLY.	EDENTATEOUS SPACE POSTERIORLY.	NO. OF TEETH.	EDENTATEOUS SPACE ANTERIORLY.	NO. OF TEETH.	EDENTATEOUS SPACE POSTERIORLY.	NO. OF TEETH.	EDENTATEOUS SPACE ANTERIORLY.	NO. OF TEETH.	EDENTATEOUS SPACE POSTERIORLY.	INDIVIDUAL CONTRIBUTING TO ORBITAL RING.
17	17	17	180 to 190	+	:	1	46 to 57	6	2	6	3rd & 4th	5th	4th	10 to 12	0	8 to 10	0	18 to 23	+	15 to 18	0	0	No	woodmasoni.	
19	19	15	177 to 208	+	..	1	47 to 69	6	2	6	3rd & 4th	5th	4th	10 to 12	0	8 to 10	0	16 to 18	+	13 to 17	0	0	No	albocinctus.	
21	21	15	162 to 208	+	..	1	53 to 68	6	2	6	3rd & 4th	5th	4th	10 to 12	0	8 to 10	0	18 to 18	+	15 to 16	0	0	Yes	juglandifer.	
19	19	15	165 to 190	+	..	1	40 to 60	6	2	6	3rd & 4th	6th	4th & 5th	9 to 10	0	9	0	18	+	15 to 16	0	0	Yes	purpura- scens.	
21	21	17	189 to 195	+	..	1	29 to 50	6	2	8	4th & 5th	6th & 7th	4th & 5th	9 to 10	0	6 to 8	0	11 to 16	+	13 to 16	0	0	No	cyclurus.	
21	21	17	169 to 193	+	..	1	35 to 47	8	4	8	4th & 5th	6th & 7th	4th & 5th	11	0	9	0	0	14 to 15	0	13 to 14	0	0	Yes	splendidus.

CORRECTIONS AND AMPLIFICATIONS OF INDIAN LOCALITIES
IN DR. T. H. BECKER'S MONOGRAPH OF THE DOLICHO-
PODIDAE (DIPTERA).

By SYDNEY RIBEIRO, Entomological Assistant, Zoological Survey of India.

In a recent number of *Capita Zoologica* (Vol. I, pt. 4) Dr. T. H. Becker has published a monograph of the Indo-Australian Dolichopodidae in which a large number of specimens belonging to the Indian Museum are described or recorded. Many of the records of Indian species are wrongly transcribed from the printed labels, others incomplete, and Dr. Annandale has asked me to draw up a correct and amplified list. This I have done in the form of a table with four columns showing (1) the name of the species recorded, (2) the page of the monograph on which it is recorded, (3) the record as given, and (4) the record corrected and with such additions added as I have been able to find on reference to the specimens named by Dr. Becker. If corrections or additions are unnecessary the entry in the last column is left blank.

Name.	Page.	Record as given.	Record corrected, with additions.
<i>Dolichopus</i>			
<i>zickzack</i> Wied. ...	8	Bengalen : Monghyr.— Burma : Rangoon.— Assam : Sylhet.—Kalkutta, Ceylon.	Calcutta and Monghyr, Bengal ; Sylhet, Assam ; Rangoon, Burma ; Ceylon.
<i>angustinervis</i> Beck. ...	9	Assam : Shillong, Darjiling.	Darjiling, E. Himalayas ; Shillong, Assam.
<i>saphirus</i> Beck. ...	10	India orient : Matheran, 800 m hoch.	Matheran, (800 ft.), Thana dist., Bombay Pres.
<i>Paraclius</i>			
<i>adligatus</i> Beck. ...	13	Assam : Sylhet ...	Sylhet, Assam ; Mergui, Tenasserim.
<i>pilosellus</i> Beck. ...	14	Madha, Yenna-Tal, Satara-Distrik, 2200 Fuss hoch.	Medha, Yenna Valley, Sa- tara dist., (2200 ft.), Bombay Pres.
<i>nudus</i> Beck. ...	14	Helvak, Kogna-Tal, Satara distrik, 2000 Fuss hoch.	Helvak, Koyna Valley, Satara dist., (2000 ft.), Bombay Pres.
<i>abbreviatus</i> Beck. ...	15	Assam : Sylhet.—Nepal : Tharbani.	
<i>regularis</i> Beck. ...	15	Balighai bei Puri, Orissa. —Kuranda.	

Name.	Page.	Record as given.	Record corrected, with additions.
<i>Hercostomus</i>			
<i>neglectus</i> Beck.	22	Indien : Simla; 7000 Fuss hoch.	Simla, (7000 ft.), Simla Hills, W. Himalayas.
<i>elongatus</i> Beck.	22	Indien : Simla, 7000 Fuss hoch.	Simla, (7000 ft.), Simla Hills, W. Himalayas.
<i>promotus</i> Beck.	23	Indien : Dharampar, Simmla-Hügel, 5000 Fuss hoch.	Dharampur, Simla Hills, (5000 ft.), W. Himalayas.
<i>chetifer</i> Walk.	26	Naini-Tal, Kumaon, 6000 Fuss hoch.	Naini Tal, (6000 ft.), Kumaon, W. Himalayas.
<i>compositus</i> Beck.	27	Mahabaleshwar, Satara-Distrik, 4200 Fuss hoch.	Mahabaleshwar, Satara dist., (4200 ft.), Bombay Pres.
<i>gregalis</i> Beck.	29	Darjiling, (7000 ft.), E. Himalayas ; Dawna Hills, (2000—3000 ft.), Tenasserim.
<i>flavidinctus</i> Beck.	29	Thisgannyinaung, Dawna-Berge.	Thingannyinaung to Sukhi, Dawna Hills, (900—2100 ft.), Tenasserim.
<i>caecus</i> Beck.	30	N.—Indien : Darjiling, 7000 Fuss hoch.	Darjiling, (7000 ft.), E. Himalayas.
<i>bilinclus</i> Beck.	31	Assam, Mangaldal.	Mangaldai, N. E. Assam.
<i>geminatus</i> Beck.	31	Kalkutta ; Assam, Sylhet.	Calcutta, Bengal ; Sylhet, Assam.
<i>Tachytrechus</i>			
<i>salinarius</i> Beck.	32	Bombay ; Westbengalen, Parosnath.	Parosnath Hill, Chota Nagpur, Bihar ; Bombay.
<i>crassitarsis</i> Meij.	32	Nepal, Terai, Kumdhik.	Kumdhik, Nepal Terai.
<i>Thinophilus</i>			
<i>tesselatus</i> Beck.	35	Bengalen : Rajshai.	Calcutta and Rajshai.
<i>diminuatus</i> Beck.	36	Indien vom Gangesdelta, Channel Creek.	Channel Creek nr. Mud Point, Ganges delta, I. Bengal ; Balighai nr. Puri, Orissa.
<i>indigenus</i> Beck.	37	Sukhwani, Nepal ; Bijarani, Naini Tal dist. (Plains), U. Provs. ; Yaikam, Travancore, Madras Provs.; Balighai near Puri, Orissa ; Calcutta, Kushtia and Madhupur, Bengal ; Sylhet, Assam.

Name.	Page.	Record as given.	Record corrected, with additions.
<i>Hydrophorus</i>			
<i>praecox</i> Lehm. ...	41	Indien : Pankong-Tal und vom Suezkanal.	Pankong Valley, (Takhtze to Chagra); Pamirs [Yarkand Mission]; Suez Canal, on board ship at night. (N. Anandale Coll.).
<i>Paralleloneurum</i>			
<i>ciliifemoratum</i> Beck. ...	43	Umgegend von Kalkutta und aus Bengalen.	Calcutta and Madhupur, Bengal; Sylhet, Assam.
<i>Phalacrosoma</i>			
<i>imperfectum</i> Beck. ...	46	Indien : Kumaon, 5700 Fuss hoch, nahe bei Bhowali.	Near Bhowali, (5700 ft.), Kumaon, W. Himalayas.
<i>fulgidipes</i> Beck. ...	46	Baragh, Simla—Hügel.	Barogh, (5000 ft.), Simla Hills, W. Himalayas.
<i>Medetera</i>			
<i>adsumpta</i> Beck. ...	48	Umgebung von Kalkutta.	
<i>grisescens</i> Moij. ...	50	Indien : Trivandrum, Travancore. — Assam, Sylhet.	Almora, (5500 ft.), Kumaon, and Bijnor, U. Provs.; Shencottah, Madras Frontier, W. side of W. Ghats, and Trivandrum, Travancore, Madras Pres.; Puri, Orissa; Bhogaon, Purneah dist. and Chakradharpur, Chota Nagpur, Bihar; Bettiah, Champaran dist., Calcutta, Madhupur and Port Canning, Bengal; Sylhet, Assam; Dawna Hills, (2000—3000 ft.), Tenasserim; Mandalay, U. Burma; Bombay.
<i>platychira</i> Meij. ...	51	Calcutta, Bengal; Sylhet, Assam.
<i>nudicoxa</i> Beck. ...	52	Darjeling, Himalaya, 7000 Fuss hoch.	Darjiling, (7000 ft.), E. Himalayas. Paresnath Hill, (4300—4400 ft.), Chota Nagpur, Bihar.
<i>Syntormon</i>			
<i>cilitarsis</i> Beck. ...	56	Assam, Sylhet.	
<i>elongatus</i> Beck. ...	57	Kufri, 5000 Fuss hoch, Simia—Berge.	Kufri, (8000 ft), Simla Hills and Mundali, (9000 ft.), Jaunsa divn., Dehra Dun dist., W. Himalayas.

Name.	Page.	Record as given.	Record corrected, with additions.
<i>Syntormon</i>			
<i>frivolis</i> Beck. ...	58	Indien : Dharampur, 5000 Fuss hoch, Simia — Berge.	Dharampur, (5000 ft.), and Kufri, (8000 ft.), Simla Hills ; and Mundali, (9000 ft.), Jaunsa divn., Delra Dun dist., W. Himalayas.
<i>Neurogona</i>			
<i>angulata</i> Meij. ...	61	Calcutta, Bengal ; Sylhet, Assam.
<i>denudata</i> Beck. ...	62	Kalkutta ; Assam, Syl- hot ; Ceylon.	
<i>pectinata</i> Beck. ...	64	N.—Bengalen : Purneah und Kalkutta.	Purneah, Bihar ; Calcutta, Bengal.
<i>exemta</i> Beck. ...	64	Ceylon : Peradeniya.	Peradeniya, Ceylon.
<i>Diaphorus</i>			
<i>vagans</i> Beck. ...	71	Ceylon.	
<i>detectus</i> Beck. ...	72	Nordbengalen : Purneah, und aus Pallode.	Purneah, Bihar ; Pallode, 20 miles N. E. of Trivandrum, Travancore, Madras Pres.
<i>maurus</i> O.—Sack. ...	72	Umgebung von Kalkutta	Calcutta, (environs), Ben- gal ; Sukne, (500 ft.), E. Himalayas.
<i>aptatus</i> Beck. ...	73	Kalkutta.	
<i>slavipilus</i> Beck. ...	74	Bombay ; Annandale ; Shasthancottab.	Shasthaneottah, 12 miles N. N. E. of Quilon, Travancore, Madras Pres. (N. Annandale Coll.) ; Bombay.
<i>nigerrimus</i> Meij. ...	75	Annandale : Travancore.	Tenmalai, W. Ghats, (W. side), Travancore, Madras Pres. (N. Annandale Coll.)
<i>wonosobensis</i> Meij. ...	75	Kurseong, (5000 ft.), E. Himalayas.
<i>hirsutipes</i> Beck. ...	75	Himalaya : Darjiling, 5000 Fuss hoch.	Darjiling, (7000 ft.), E. Himalayas.
<i>protervus</i> Beck. ...	76	Kalkutta.	Calcutta, Bengal ; Dawna Hills, (2000—3000 ft.), Tenassorim.
<i>minor</i> Meij. ...	77	Himalaya : Darjiling, 7000 Fuss hoch.	Darjiling, (7000 ft.), and Kurseong, (4700 ft.), E. Himalayas.
<i>simulans</i> Beck. ...	77	Indien : Colombo.	Colombo, Ceylon.
<i>discrepans</i> Beck. ...	78	Umgebung von Kalkutta.	

Name.	Page.	Record as given.	Record corrected, with additions.
<i>Diaphorus.</i>			
<i>mandarinus</i> Wied. ...	78	Verschiedenen Indiens. Teilen	Sarah, Nepal ; Khurda Road, Puri dist. and Balighai nr. Puri, Orissa; Katihar, Purneah dist., Bihar ; Asansol, Calcutta, (environs), Kuchtea, Monghyr, Rajmahal and Tinpahar, Bengal ; Margherita, Mazbat, Mangaldai dist., Sadiya and Sylhet, Assam ; Mandalay and Rangoon, Burma ; Kawkareik, Amherst dist. and Dawna Hills, (at base), Tenasserim ; Parel Bombay.
<i>submixtus</i> Beck. ...	79	Assam : Margharita.	Margherita and Sadiya, Assam.
<i>habilis</i> Beck. ...	80	Bengalen : Monghyr.	Siripur, Saran, Bihar ; Monghyr, Bengal.
<i>intermixtus</i> Beck. ...	81	Assam : Margherita.	Naini Tal, (6000—7000 ft.), Kumaon, U. Provs. ; Calcutta, Bengal ; Margherita, Assam.
<i>Asyndetus</i>			
<i>latifrons</i> Lw. ...	86	Indian : Katihar, Distrikt Purneah.	Katihar, Purneah dist., Bihar ; Asansol, Barrackpore nr. Calcutta, and Rajshai, Bengal ; Sylhet, Assam.
<i>Chrysotus</i>			
<i>gratiosus</i> Beck. ...	86	Kalkutta.	
<i>degener</i> Frey ...	87	Umgebung von Kalkutta.	Calcutta, (environs), Madhupur and Monghyr, Bengal ; Sylhet, Assam ; Mandalay and Rangoon, Burma.
<i>alternus</i> Beck. ...	88	Indien : Burma, von den Dasona-Hügeln, 2—3000 Fuss hoch.	Shamdang, (c. 3000 ft.), Sikkim ; Calcutta and Barrackpore nr. Calcutta, Bengal ; Dawna Hills, (2000—3000 ft.), Tenasserim.
<i>discretus</i> Beck. ...	89	Mundeli ; Dehra Dun—Distrikt, 9000 Fuss hoch ; Umgebung von Kalkutta.	Darjiling, (7000 ft.), and Kurseong, (4700 ft.), E. Himalayas ; Mundali, (9000 ft.), Jaunsa divn., Dehra Dun dist., W. Himalayas ; Calcutta, Bengal ; Dawna Hills, (2000—3000 ft.), Tenasserim.

Name.	Page.	Record as given.	Record corrected, with additions.
<i>Trigonocera</i>			
<i>specialis</i> Beck. ...	92	Umgebung von Kalkutta.	
<i>obscura</i> Moij. ...	92	Mandalay, U. Burma.
<i>Sympycnus</i>			
<i>bisulcus</i> Beck. ...	94	Base of Dawna Hills, Tenasserim; Mahabaleshwar, (4200 ft.), Satara dist., Bombay Pres.
<i>gummigutti</i> Beck. ...	95	Nepal: Thamaspur, und von Burma, Dawna-Hügel; von Darjiling, 7000, Fuss hoch.	Darjiling, (7000 ft.), E. Himalayas; Thamaspur, Nepal; Dawna Hills, (at base and at 2000—3000 ft.), Tenasserim.
<i>apicalis</i> Moij. ...	96	Sylhet, Assam.
<i>scutatus</i> Moij. ...	97	Sylhet, Assam.
<i>simplex</i> Moij. ...	99	Indien: Ost-Himalaja, Sukna, 500 Fuss hoch.	Sukna, (500 ft.), and Darjiling, (7000 ft.), E. Himalayas; Barrackpore nr. Calcutta, Bengal; Margherita and Sadiya, Assam.
<i>vivus</i> Beck. ...	99	Burma, Downa-Hügel.	Dawna Hills, (at base), Tenasserim.
<i>strenuus</i> Beck. ...	101	Ceylon: Paradonyia.	Peradeniya, Ceylon.
<i>turbidus</i> Beck. ...	105	Östlichen Himalaja: Kurseong, 4700 Fuss hoch.	
<i>subjectus</i> Beck. ...	105	Burma: Daema-Hügel, 2—3000 Fuss hoch; Bergalen: Madhupur.	Madhupur, Bengal; Dawna Hills, (at base and at 2000—3000 ft.), Tenasserim.
<i>Chrysosoma</i>			
<i>bifiguratum</i> Beck. ...	140	Assam.	
<i>pulcherrimum</i> Beck. ...	145	Ceylon; Ceylon: Nilgiri hill [H. L. Andrews].	Maddathoray, W. base of W. Ghats, Travancore and Nilgiri Hills, (2500 ft.), Madras Pres. [H. L. Andrews Cole.]; Ceylon.
<i>vittatum</i> Wied. ...	146	Indien: Mazbat Mangaldai, Assam; Kalkutta; Ceylon.	Trivandrum, Travancore, Madras Pres.; Calcutta, Bengal; Ceylon; Mazbat, Mangaldai dist., Assam.

Name.	Page.	Record as given.	Record corrected, with additions.
<i>Chrysosoma</i>			
<i>stragulum</i> Beck. ...	150	Indien : Kalattupuzker, base of W. Ghats ; Trivandrum, Travancore ; Madduthoray, Travancore.	Kulattupuzha and Mad- dathoray, W. base of W. Ghats and Trivan- drum, Travancore, Madras Pres.
<i>infirme</i> Beck. ...	152	Ceylon : Paradeniya.	Peradeniya, Ceylon.
<i>sagax</i> Beck. ...	152	Indien : (Fundort ?), Koll. Winthem im Wiener Hof-Mus.	
<i>elongatum</i> Beck. ...	153	Himalajagebirgo, 5000 Fuss hoch, Kurseong.	Kurseong, (4700-5000 ft.), E. Himalayas ; Assam- Bhutan Frontier, Man- galdai dist., N. E. Assam.
<i>petulans</i> Beck. ...	154	Ceylon : Paradeniya.	Peradeniya, Ceylon.
<i>decoratum</i> Beck. ...	154	Nordlichen Indien, aus Assam, Sylhet.	
<i>ovale</i> Beck. ...	155	Ceylon : Hineratgoda ; S.—Ceylon; Walisama ; S.—Indien : Nilgiris, 6000 Fuss hoch, Coonor ; Ceylon : Rambodde.	Coonoor, (6000 ft.), Nil- giri Hills, Madras Pres.; Hineratgoda, Rambode and Walisama, Cey- lon.
<i>punctiforme</i> Beck. ...	156	Himalajagebirge : Sukna, 500 Fuss hoch.	Sukna, (500 ft.), E. Him- alayas.
<i>inscriptum</i> Beck. ...	157	Kalkutta ; Bengal : Rajmahal ; Assam : Sylhet.	Calcutta, Kushtea and Rajmahal, Bengal ; Syl- het, Assam.
<i>mirandum</i> Beck. ...	157	Himalajagebirge : Sukna, 500 Fuss hoch.	Sukna, (500 ft.), E. Him- alayas ; Siliguri, (base of E. Himalayas), N. Ben- gal.
<i>spengeli</i> Enderl. ...	158	Süd-Indien : Bangalore, 3000 Fuss hoch.	
<i>annotatum</i> Beck. ...	160	Ceylon : Pattipola, 2000 m. hoch ; Ceylon.	
<i>extractum</i> Beck. ...	161	Ceylon : Peradeniya.	
<i>duplicatum</i> Beck. ...	162	Ceylon : Peradeniya, Colombo, Victoria Gar- dens, Trivandrum, Travancore.	Trivandrum, Travancore, Madras Pres. Peradeniya and Colombo, Victoria Gardens, Ceylon.
<i>excisum</i> Beck. ...	162	Ceylon : Paradeniya.	Peradeniya, Ceylon.
<i>congruens</i> Beck. ...	163	Insel Ceylon.	

Name.	Page.	Record as given.	Record corrected, with additions.
<i>Chrysosoma</i>			
<i>foliatum</i> Beck. ...	164	Ost-Bengalen : Chittagony ; Chittagong, Mazbat und Sukna.	Sukna, (500 ft.), E. Himalayas ; Chittagong, E. Bengal ; Mazbat, Mangaldai dist., Assam.
<i>albidum</i> Beck. ...	164	Himalaja-Gebirge : Karlong, 5000 Fuss hoch ; Sukna, 500 Fuss hoch.	Ghumti, (4000 ft.), Darjiling dist., Kurseong, (5000 ft.), Sukna, (5000 ft.), and Gangtok, (6150 ft.), Sikkim, E. Himalayas.
<i>undulatum</i> Beck. ...	165	Assam : Dibrugarh (Abor Exped.) ; Sylhet.	Dibrugarh (Abor Exped.) and Sylhet, Assam.
<i>brunnipenne</i> Beck. ...	165	Trichinopoly, dem südlichen Indien.	
<i>faciatum</i> Beck. ...	166	Ceylon : Pattipola, 2000 m. hoch.	
<i>oculatum</i> Beck. ...	167	Kumaon bei Rhowali, 5700 Fuss hoch.	Near Rhowali, (5700 ft.), Kumaon, W. Himalayas.
<i>leucopogon</i> Wied. ...	168	Ceylon und Kalkutta.	Agisamtongee, S. end of Kayangulum, Bangalore, and Kulattupuzha, W. base of W. Ghats, Travancore ; Madras, (town), Marikuppam, (2500 ft.), and Oorgaum, (2500 ft.), Madras Pres. ; Puri, Orissa ; Adra, Manbhumi dist., Bihar ; Calcutta, Khargpur, Monghyr and Rajmahal, Bengal ; Moulmein, Tenasserim ; Rangoon, Burma ; Bombay and Parol, Bombay Pres. ; Ceylon.
<i>tenuipes</i> Beck. ...	174	bei Barogh, Simia hills, 5000 Fuss hoch ; Kurseong, Himalaja-Gebirge, 6000 Fuss hoch.	Kurseong, (6000 ft.), E. Himalayas ; Barogh, (5000 ft.), Simla Hills, W. Himalayas.
<i>triligatum</i> Beck. ...	177	Südlichen Indien : Perumal ; Trichinopol ; Coonoor, 6000 Fuss hoch, Nilgiris.	Perumal, Trichinopoly and Coonoor, (6000 ft.), Nilgiri Hills, Madras Pres.
<i>compressum</i> Beck. ...	178	Nordbengalen : Kushtea ; Bhogaon, Purneah-Distrikt.	Bhogaon, Purneah dist., Bihar ; Kushtea, Bengal.
<i>muticum</i> Thoms. ...	183	Moulmein, Indien.	Dibrugarh, N. E. Assam ; Moulmein, Tenasserim.

Name.	Page.	Record as given.	Record corrected, with additions.
<i>Chrysosoma</i>			
<i>armillatum</i> Big.	183	Colombo ; Ceylon : Kanthalay.	Colombo and Kanthalai, Ceylon.
<i>grossum</i> Beck.	186	Indien : Margherita.	Margherita, Assam
<i>posticum</i> Beck.	188	Bombay.	
<i>integrum</i> Beck.	189	Ceylon : Matale ; Südlichen Indien ; Trichinopol ; Kahtal, Purneah-Distrik [C. A. Paiva coll.] ; Colombo, Victoria Gardens.	Allahabad, U. Provs.; Purneah and Katihar, Purneah dist. (C. A. Paiva coll.) and Siripur, Saran, Bihar ; Calcutta and Rajmahal, Bengal ; 10 miles off Masulipatam, (on board ship), and Trichinopoly, Madras Pres.; Colombo, Victoria Gardens, Matale, Ceylon.
<i>perturbans</i> Beck.	192	Südlichen Indien : Trichinopoly.	Trichinopoly, Madras Pres.
<i>Sciopus</i>			
<i>tardus</i> Beck.	198	Rajmahal, Bengal.
<i>gravipes</i> Beck.	199	Ostbengalen : Chittagong.	
<i>interdictus</i> Beck.	201	Bombay.	
<i>turbidus</i> Beck.	202	Batighai, Orina.	Balighai nr. Puri, Orissa.
<i>pediformis</i> Beck.	204	Himalaja-Gebirge : Darjiling, 6—7000 Fuss hoch.	Darjiling, (6000—7000 ft.), E. Himalayas.
<i>penicillatus</i> Beck.	204	Shillong, Provinz Assam.
<i>infans</i> Beck.	205	Baighai, Orissa.	Balighai nr. Puri, Orissa.
<i>adhaerens</i> Beck.	207	Südlichen Indien : Trichinopoly.	Trichinopoly, Madras Pres.
* <i>insecans</i> Beck.	208	Kuranda, Indien.	?
<i>bifarius</i> Beck.	208	Indien : Cairuna, Kuranda.	?
* <i>triangulifer</i> Beck.	211	Indien : Kuranda.	?
<i>variabilis</i> Meij	211	Sylhet, Assam.
<i>fruticosus</i> Beck.	212	Sylhet, Assam.	

Name.	Page.	Record as given.	Record corrected, with additions.
<i>Condylostylus</i>			
<i>poecilus</i> Beck. ... 221	Indien; Madura, Shembaganur.	Shembaganur, Palni Hills and Madura, Madras Pres.	
<i>profundus</i> Beck. ... 222	Südindien: Kodaikanal; Trichinopoly.	Kodaikanal, Palni Hills and Trichinopoli, Madras Pres.; Sadiya, Assam.	
<i>imperfectus</i> Beck. ... 223	Indien, ohne speziellere Fundangabe.		
<i>conspicuus</i> Beck. ... 225	Unehagaon, Nainital-district, am Fusse des Himalaja und Kalkutta.	Kathgodam, (1200 ft.), Kumaon and Unehagaon, Naini Tal dist., base of W. Himalayas; Bangalore, (3000 ft.), Madras Pres.; Calcutta and Kushtia, Bengal; Mergui and Rangoon, Burma.	
<i>impar</i> Beck. ... 226	Colombo.	Colombo, Ceylon.	
<i>vigilans</i> Beck. ... 226	Himalajagebirge: Darjiling, 7000 Fuss hoch.	Darjiling, (7000 ft.), and Kurseong, (4700—5000 ft.), E. Himalayas; Margherita, Assam.	
<i>jacobsoni</i> Meij. ... 228	Dawna Hills, (2000—3000 ft.), Tenasserim.	
<i>Megistostylus</i>			
<i>longicornis</i> Fabr. ... 230	Ceylon; Indien.	Sikkim, E. Himalayas; Ceylon.	
<i>Mesorhaga</i>			
<i>obscura</i> Beck. ... 232	Ceylon: Pattipola, 2000 m. hoch.	Pattipola, (2000 ft.), Ceylon.	
<i>termialis</i> Beck. ... 233	Ceylon.		
<i>nigro-viridis</i> Beck. ... 235	Ceylon.		

A LIST OF THE INDIAN CICINDELIDAE WITH LOCALITIES.

By C. DOVER, F.E.S., and S. RIBEIRO, Assistant, Zoological Survey of India.

Canon Fowler's comparatively recent volume on the Cicindelidae in the "Fauna of British India" series renders the publication of a catalogue of this family unnecessary. We believe, however, that a revised list of the Indian tiger-beetles with special reference to the collection of the Zoological Survey of India will prove useful, and in submitting the following list hope that this may be so. In preparing it we have arranged the species alphabetically under the genera, partly to facilitate reference, and partly because the affinities of the various species are yet imperfectly known. We have endeavoured to incorporate all information as to geographical distribution and have also added footnotes to certain species where it has been thought necessary. As far as possible the precise situation of the various localities mentioned here has been given. It may be stated that when localities like "Sikhim" is mentioned this does not imply that the insect occurs all over Sikhim, but that the specimen has not been more accurately labelled than this. The localities of specimens in the Museum collection have been marked with an asterisk.

The life-histories and habits of the Cicindelidae are of great interest, but those of the Indian species are practically unknown, and there is a wide field for amateur observations here.

We are indebted both to Drs. N. Annandale and S. W. Kemp of the Zoological Survey of India for their valuable suggestions and help rendered to us in our study of this group. Our thanks to Dr. Walther Horn of the Deutsches Entomologisches Museum, Berlin, the leading authority on the Cicindelidae, for his kindness in looking through and checking this list, as well as for identifying many specimens.

References to Literature.

(References to works previous to Fowler's volume are not given.)

- 1912. FOWLER, W. W., Fauna of British India, General Introduction and Cicindelidae and Paussidae (London).
- 1912. GRAVELY, F. H., The Habits of some Tiger-Beetles from Orissa.—*Rec. Ind. Mus.*, VII, p. 207.
- 1913. HORN, W., 50 neue Cicindelidae.—*Arch. Nat. Berlin*, 79A, II.
- 1913. HORN, W., Matériaux pour servir à l'étude de la Faune Entomologique de L'Indo-Chine. Cicindelinae. No. I.—*Ann. Soc. Ent. Belgique*, LVII, pp. 362—366.
- 1914. HORN, W., Matériaux pour servir à l'étude de la Faune Entomologique de L'Indo-Chine. Cicindelinae. No. II.—*Ann. Soc. Ent. Belgique*, LVIII, pp. 137—140.
- 1914. FLETCHER, T. B., Note on Tiger-Beetles from Coorg.—*Journ. Bomb. Nat. Hist. Soc.*, XXIII, p. 329.

1915. HORN, W., Wytsman's *Genera Insectorum*, Cicindelidae.
1917. FLEUTIAUX, E., Enumeration des Cicindelidae récoltés en Indo-Chine française par M. Vitalis de Salvaza, de 1914 à 1916.—*Bull. Soc. Ent. France*, p. 48.
1917. FLEUTIAUX, E., Nouvelle liste de Cicindelidae de L'Indo-Chine. *Ibid.*, p. 368.
1919. FLEUTIAUX, E., Sur quelques Cicindelidae d'Indo-Chine (rectifications).—*Ibid.*, p. 252.
1919. GRAVELY, F. H., A note on the Marine Invertebrate Fauna of Chandipore, Orissa.—*Rec. Ind. Mus.*, XVI, p. 398.
1920. FLEUTIAUX, E., Tableau pour la détermination rapide des Tricondyla d'Indo-Chine.—*Bull. Soc. Ent. France*, p. 308.
1921. ANNANDALE, N. AND DOVER, C., The Cicindelid Beetles of Bar-kuda Island.—*Rec. Ind. Mus.*, XXII, pp. 335—337.
1921. DOVER, C. AND RIBEIRO, S., Records of some Indian Cicindelidae. —*Rec. Ind. Mus.*, XXII, pp. 721—727.

Family CICINDELIDAE.

Division ALLOCOSTERNALIAE.

Subfamily COLLYRINAE.

Genus **Collyris** Fabr. (1801).

brevipennis Horn. Bombay, Talewadi* and Castle Rock,* N. Kanara dist., Bombay Pres.

brevipennis var. *elongato-subtilis* Horn.¹ Anaimalai Hills, Madras Pres.

brevipennis var. *subtilesculpta* Horn. Coonoor and Ouchterlony Valley* (3,000—3,500 ft.), Nilgiri Hills and Dindigul, Madras Pres. ; Coorg, S. India.

dohrni Chaud. Colombo, Ceylon.

longicollis Fabr.² Sahibganj,* Burdwan dist., Bengal.

mniszechi Chaud. Naga Hills, Assam ; Karen Hills, Burma ; Laos, Indo-China ; Malacca ; Borneo ; (?) Java ; Momeit, Yunnan, W. China.

Genus **Neocollyris** Horn (1901).

acute-apicalis Horn.³ Chandkhira, Sylhet, Assam.

ampulicollis Horn.⁴ Wallardi, Travancore.

andrewesi Horn. Trivandrum* and Wallardi, Travancore ; Anaimalai Hills (3,000—4,000 ft.), and Nilgiri Hills,* Madras Pres. ; N. Kanara dist., Bombay Pres. ; Mahe, W. coast, India ; (?) Ceylon.

apicalis Chaud. Patkai Hills, Assam ; S. Burma ; Tonkin, Indo-China ; Singapore ; Sumatra ; Malacca.

¹ Cf. Horn (1913, p. 2). Not recorded in the "Fauna."

² Cf. Fowler (1912, p. 225).

³ Cf. Horn (1913, p. 4). Not recorded in the "Fauna."

⁴ Cf. Horn (1913, p. 5). Not recorded in the "Fauna."

aptera Lund. Naga Hills and Sylhet, Assam ; Tenasserim.

apteroides Horn. Manipur, Assam.

attenuata Redt. Soom* (4,500—5,000 ft.), Singla* (1,500 ft.), and Mungphu, Darjiling dist., E. Himalayas ; Nepal ; Simla, Punjab, W. Himalayas ; Calcutta, Bengal ; Khasi Hills, Assam.

auripennis Horn. Manipur, Assam ; Tonkin and Annam, Indo-China. *bipartita* Fleut. India ; Karen Mts., Burma.

bonelli Guér. Sikhim* ; Barkuda Id.,* Lake Chilka and Trichur, Cochin State, Madras Pres. ; Calcutta,* Khargpur,* and Rangamati,* (Chittagong Hill Tracts), Bengal ; Siripur,* Saran, Bihar ; Central Provs.* ; Sibsagar,* Khasi Hills,* Naga Hills, Patkai Hills and above Tura,* Garo Hills, Assam ; Maymyo, Tharawaddy and Pegu, Burma ; Tenasserim ; Hills near Taiping,* Perak, Malay Peninsula ; Java* ; S. Kashmir, the Himalayas and S. China to Sumbawa, Sumba, and the Banguey Is., N. Borneo.

bonelli var. *batesi* Horn. Khasi Hills, Assam ; Siam ; Cochin China.

bonelli var. *diversipes* Fowl. Pashok* (3,500 ft.), Darjiling dist., E. Himalayas ; Tura* (1,200—1,500 ft.), Garo Hills, Assam ; Burma ; Java ; Borneo.

bonelli var. *ortygia* Buq. Singla* (1,500 ft.), Darjiling dist., E. Himalayas ; Kashmir, W. Himalayas ; Nepal ; Calcutta* and Rajmahal,* Bengal ; Siripur,* Saran, Bihar ; N. Cachar and Tura* (1,200—1,500 ft.), Garo Hills, Assam ; Bhamo and Tharawaddy, Burma ; Mahe, W. coast, India ; S. China ; Tonkin and Annam, Indo-China ; Malacca ; Singapore ; Sumatra ; Java ; Bali Id., Sumbawa and Sumba, E. Indies ; Siam ; Banguey Is., N. Borneo.

bonelli subsp. *distincta* Chaud.¹ Darjiling and Pashok* (3,000 ft.), Darjiling dist., E. Himalayas ; Chatrapur,* Ganjam dist., Madras Pres. ; Balighai* near Puri, Orissa ; Calcutta,* Bengal ; Khasi Hills, Assam ; Castle Rock,* N. Kanara dist., Bombay Pres.

boysi Chaud.² Bhutan ; Above Tura* (3,900 ft.), Garo Hills, Assam. *brevilabris* Horn. Assam ; Karen Hills, Burma.

ceylonica Chaud. Ceylon.

compressicollis Horn.³ Assam.

crassicornis Chaud. Sikhim.

crassicornis Dej. Anaimalai Hills, Pondicherry and Gopkuda Id.,* Lake Chilka, Madras Pres. ; Travancore ; Chandipore* near Balasore, Orissa (sea-coast) ; Calcutta* (and environs)* and Maldah,* Bengal ; Chota Nagpur and Ranchi,* Bihar ; Sibsagar,* Assam ; N. Shan States,* Burma ; Ceylon ; Siam ; Malay Peninsula ; Java ; Sumatra ; Indo-China ; Celebes ; Moluccas.

orientata Schm.-Goeb. Sibsagar,* Assam ; Karen Hills, Tharawaddy, Taung-ngu, Rangoon and Pegu, Burma ; Tenasserim ; Siam ; Malacca ; Sumatra ; Borneo.

¹ Cf. Dover and Ribeiro (1921, p. 722). Recorded as a distinct species in the "Fauna."

² Fowler (1912, p. 257) treats this as synonymous with *N. saphyrina* but Horn (1908, p. 104) regards it as a distinct species. The Museum possesses a single example of *N. boyisi*, named by Horn.

³ Cf. Horn (1908, p. 106). Not recorded in the "Fauna."

cylindrica Schm.-Goeb. Burma.

cylindripennis Chaud. Karen Hills, Burma ; Siam.

feae Horn. Rangamati,* Chittagong Hill Tracts, Bengal ; Karen Hills, Burma.

foveifrons Horn.¹ Khasi Hills, Assam ; Tonkin, Indo-China.

fuscitarsis Schm.-Goeb. Sikhim* ; Singla* (1,500 ft.), Darjiling dist., E. Himalayas ; Assam ; Rangoon and Tharawaddy, Burma ; Moulmein, Tenasserim ; Tonkin, Indo-China ; Malacca ; Java ; Sumatra.

insignis Chaud. Sikhim* ; Darjiling, Mungphu,* Pankabari,* Pashok* (2,500 ft.), and Singla* (1,500 ft.), Darjiling dist., E. Himalayas ; Bhutan ; Khasi Hills and above Tura* (3,900 ft.), Garo Hills, Assam ; Burma.

kollari Horn. (?) S. India ; Centr. India.

lesnei Horn. Karen Hills, Burma.

linearis Schm.-Goeb. Assam ; Maymyo and Pegu,* Burma ; Cambodia, Saigon and Tonkin, Indo-China ; Siam.

linearis var. *srnkae* Horn. Pegu dist. and Ruby Mines, Burma ; Cambodia, Saigon and Tonkin, Indo-China.

maindroni Horn. Wallardi, Travancore ; Nilgiri Hills, Madras Pres.

moesta Schm.-Goeb. Burma ; Siam ; Perak,* Malay Peninsula ; Cambodia and Tonkin, Indo-China ; Malacca.

nilgirica Fowl. Nilgiri Hills, Madras Pres.

orichalcina Horn. Nilgiri Hills, Madras Pres. ; Naga Hills (3,500—5,000 ft.), Assam ; Yunnan, W. China.

parvula Chaud. Coonoor and Anaimalai Hills (in neighbourhood), and Yercaud,* Shevaroy Hills (3,000—5,000 ft.), Salem dist., Madras Pres. ; Sidapur,* Coorg, S. India ; Castle Rock,* N. Kanara dist. and Belgaum, Bombay Pres. ; Mahé, W. coast, India.

parvula var. *amoena* Chaud. Castle Rock,* N. Kanara dist. and Belgaum, Bombay Pres.

planifrons Horn. Ceylon.

plicaticollis Chaud. Ceylon.

plicicollis Horn. Nilgiri Hills, Madras Pres.

plicicollis var. *subtile-slavescens* Horn.² Shembaganur, Madura dist., Madras Pres.

punctatella Chaud. Balangoda, Ceylon.

redtenbacheri Horn.³ Mungphu, Singla* (1,500 ft.), Ghumti (1,500—5,000 ft.), and Pashok* (1,000—3,500 ft.), Darjiling dist., E. Himalayas ; Simla and Kousanie* (6,075 ft.), Kumaon, W. Himalayas ; Nepal ; Bhutan ; Pinjore,* Patiala State, Punjab ; Khasi Hills, Naga Hills,* Patkai Hills, Tura* (200—1,500 ft.), and above Tura* (3,000 ft.), Garo Hills, Assam ; Arrakan,* Burma ; Tenasserim.

roeschkei Horn. Calcutta, Bengal.

rubens Bates. Assam (plains).

¹ Towler (1912, p. 268) apparently doubts the validity of this species. It is included in his volume on Dr. Horn's authority.

² Cf. Horn (1913, p. 4). Not recorded in the "Fauna."

³ Fowler (1912, p. 240) remarks that he is doubtful whether this is a distinct species, or only a variety of *N. attenuata*.

- rufipalpis* Chaud. Assam* ; Karen Hills, Burma ; Cambodia and Tonkin, Indo-China ; Sumatra ; Java.
- saphyrina* Chaud. Pashok* (2,500 ft.), Mungphu,* Darjiling dist., E. Himalayas ; Nepal Valley* (4,500—6,500 ft.), E. Himalayas ; Pedong, Bhutan ; Bengal ; Assam ; N. India.
- sarawakensis* Thoms. Sylhet,* Assam ; Burma ; Sumatra ; Borneo ; Malacca.
- sarawakensis* var. *dohertyi* Horn.¹ Burma ; Sumatra ; Siam ; Malacca.
- saundersi* Chaud. Kandy and Bandarawela and Kandy, Ceylon.
- saundersi* var. *continentalis* Horn. Wallardi, Travancore.
- saundersi* var. *lactior* Horn. Colombo, Kandy,* Korale and Morawak, Ceylon.
- schaumi* Horn. Andaman Is.*
- similis* Lesne. Nilgiri Hills, Madras Pres. ; Assam ; Laos, Indo-China ; Perak, Malay Peninsula.
- smaragdina* Horn. Soom* (4,000—5,000 ft.), Mungphu* and Kurseong,* Darjiling dist., E. Himalayas ; Bhutan.
- smithi* Chaud. (?) Tibet ; Dacca, E. Bengal ; Naga Hills* and Cachar, Assam ; Pegu,* Burma.
- subclavata* Chaud. (?) Kollar dist., Mysore and Nilgiri Hills, Madras Pres. ; Bengal ; Andaman Is.* ; Indo-China.
- subclavata* var. *andamana* Bates. Andaman Is.
- subtilis* Chaud. Palni Hills and Kodaikanal, S. India ; Karen Hills, Burma ; Siam ; Sumatra ; Java.
- subtilis* var. *brachycephala* Horn. Karen Hills, Burma.
- variicornis* Chaud. Upper Teesta Valley* (4,000 ft.), Sitong* (3,800—4,000 ft.), and Singla* (1,500 ft.), Darjiling dist., E. Himalayas ; Tura* (1,200—1,500 ft.), Garo Hills, Sylhet, Khasi Hills, Patkai Hills and Naga Hills, Assam ; Tharawaddy, Burma ; Bukit Besar,* Nawngchik (2,500 ft.), Siamese Malay States.
- variicornis* var. *flavolabiata* Horn. Andaman* and Nicobar Is.*
- variicornis* var. *gestroi* Horn. Karen Hills, Burma ; Tenasserim.
- variitarsis* Chaud. Darjiling, Singla* (1,500 ft.), and Sukna* (500 ft.), Darjiling dist., E. Himalayas ; Nepal ; Bengal ; Tura* (1,200—1,500 ft.), Garo Hills and Dunsiri Valley,* Assam ; Karen Hills and Teinzo, Burma ; Tenasserim ; Tonkin, Indo-China ; Penang.
- variitarsis* var. *brachycephala* Horn. Bhamo, Rangoon and Karen Hills, Burma.

Genus *Tricondyla* Latr. (1822).

- coriacea* Chev. Chilaw,* Kanthalai, Kekirawa, Palatupana and Trincomali, Ceylon.
- cyanea* var. *annulicornis* Schm.-Goeb. Pegu,* Burma ; Tenasserim ; Siam ; Cambodia.
- gestroi* Fleut. Sylhet, Assam ; Kareni (? Karen Hills), Burma ; Tonkin, and Cambodia, Indo-China.
- gounelli* Horn. Ramnad, Madura dist., and Anaimalai Hills (3,500—4,000 ft.), Madras Pres. ; Trivandrum, Travancore.

¹ According to Fowler (1912, p. 269) an examination of further specimens might necessitate the separation of this form as a distinct species.

gounelli var. *horni* Mndr. Travancore (mountains) and Trivandrum,* Travancore.

granulifera Mots. Haragaru and Nalanda, Ceylon.

granulifera var. *rugosa* Chaud. Ceylon.

macrodera Chaud. Sikkim*; Darjiling,* Mungphu,* Ghumta* (4,000 ft.), Kalimpang* (600—1,500 ft.), Pashok* (2,000—3,000 ft.), Singla* (1,500 ft.), and Gopaldhara,* Rungbong Valley, Darjiling dist., E. Himalayas; Bhutan*; Tura* (1,500 ft.), and above Tura* (3,500—3,900 ft.), Garo Hills, Pasighat,* Sadiya dist., Sibsagar,* and Cachar, Assam; Tonkin, Indo-China.

mellyi Chaud. Maini Mukh,* Chittagong Hill Tracts, Bengal; above Tura* (3,500—3,900 ft.), Garo Hills and Silcuri, Cachar* dist., Assam; Laos and Tonkin, Indo-China.

nigripalpis Horn. Centr. Ceylon.

tuberculata Chaud. Sylhet, Silcuri,* Cachar dist., Assam.

Genus *Deroecrania* Chaud. (1860).

agnes Horn. Centr. Ceylon.

brevicollis Horn. Trichinopoly, Madras Pres.

concinna Chaud. Kandy,* Ceylon.

flavicornis Horn. Ceylon.

fusiformis Horn. Centr. Ceylon.

gibbiceps Chaud. Centr. Ceylon.

halyi Horn. Ceylon.

honorei Fleut. Rammad, Madura dist., Trichinopoly and Trichur, Cochin State, Madras Pres.; Kodaikanal, Palni Hills, S. India; Bombay.

longesulcata Horn. Wallardi, Travancore; Coonoor, Anaimalai Hills and Nilgiri Hills (4,500—6,000 ft.), Madras Pres.; Castle Rock,* N. Kanara dist., Bombay Pres.

nematodes Schaum. Centr. Ceylon.

nietneri Mots. Balangoda Ridge, Kandy,* Ceylon.

schaumi Horn. Ceylon.

scitiscabra Walk. Kandy,* Ceylon.

Division *PLATYSTERNALIAE*.

Subfamily *THERATINAE*.

Genus *Therates* Latr. (1817).

chenelli Bates. Naga Hills (2,000 ft.), Assam; Karen Hills and N. Shan States,* Burma.

chenelli var. *concinnus* Gestro. Ruby Mines, U. Burma.

dohertyi Horn. Pashok* (5,000 ft.), Darjiling dist., E. Himalayas; Patkai Hills, Assam; Pegu, Burma; Tenasserim.

dormeri Horn. Patkai Hills, Assam; Borneo.

gestroi Horn. Khasi Hills and above Tura* (3,500—3,900 ft.), Garo Hills, Assam; Lakhon, Lower Siam.

gestroi var. *annandalei* Horn. Kurseong* (4,700—5,000 ft.), Ghumti* (4,000 ft.), Pashok* (2,000, 4,000 and 5,000 ft.), Mahanadi (4,120 ft.),* and Kalimpong* (600—4,500 ft.), Darjiling dist., E. Himalayas.

hennigi Horn. Khasi Hills and above Tura* (3,900 ft.), Garo Hills, Assam.

obliquus Fleut. Pashok* (5,000 ft.), Darjiling dist., E. Himalayas; Ruby Mines, U. Burma; Momeit, Yunnan, W. China.

waagenorum Horn. Darjiling, E. Himalayas; Pegu,* Burma; Tenasserim.*

Subfamily CICINDELINAE.

Genus **Prothyma** Hope (1838).

belloides Horn. N. Kanara, Bombay.

bouvieri Horn. Maymyo, Burma; Lakhon, Lower Siam.

exornata Schm.-Goeb. Tharawaddy, Pegu,* N. Chin Hills and Karen Hills, Burma; Annam, Cambodia and Laos, Indo-China.

feae Gestro. Bhamo and Teinzo, Burma.

hennigi Horn.¹ Khasi Hills, Assam.

inornata Horn.² Kulu, Punjab, W. Himalayas; Assam.

limbata Wied. Simla, Punjab and Jhelum Valley,* Kashmir, W. Himalayas; Calcutta, Bengal.

paradoxa Horn. Mahe, W. coast, India; Pondicherry and Kanara, Madras Pres.; Trivandrum* and Karunagapalli,* Travancore; Ceylon.*

proxima Chaud. Kurseong, Darjiling dist., E. Himalayas; Dharwar, U. Provs.; Ramnad, Kanara, Cocanada and Coimbatore,* Madras Pres.; Balugaon,* Puri dist., and Cuttack,* Orissa; Calcutta and Birbhum,* Bengal; (?) Kathiawar, Bombay Pres.; Nagpur, Central Provs.

reconciliatrix Horn. Dacca,* E. Bengal; above Tura* (3,500 ft.), Garo Hills, Assam; Tenasserim.

schmidt-goebeli Horn. (?) Bengal; Gokteik, U. Burma; Cambodia and Laos, Indo-China.

scrobiculata Wied. Maldah, Bengal; Chota Nagpur, Bihar; Balasore,* Orissa (sea-coast).

Genus **Heptodonta** Hope (1838).

arrowi Horn. Tavoy and N. Chin Hills, Burma.

eugenia Chaud. Bet. Yenang-Yaung and Mandalay, Bhamo, Teinzo and Tharawaddy, Burma; Cochin China; Siam; Penang.

ferrarii Gestro.³ N. Shan States* and Karen Hills, Burma; Cambodia, Laos and Tonkin, Indo-China.

¹ To quote Fowler (1912, p. 309) "this is a somewhat abnormal species and ought, perhaps, to be referred to *Heptodonta*." Further material might necessitate the erection of a separate genus for this species.

² Fowler (1912, p. 305) remarks that *P. inornata* and *P. feae* are in all probability subspecies of *P. limbata*.

³ Fowler (1912, p. 312) treats this species as a synonym of *H. pulchella*. See Dover and Ribeiro (1921, p. 723).

kraatzi Horn. Darjiling, Mungphu* (3,000—4,000 ft.), Ghumti* (1,500—5,000 ft.), Pashok* (2,000 ft.), and Singla* (1,500 ft.), Darjiling dist., E. Himalayas; Khasi Hills and above Tura* (3,000—3,500 ft.), Garo Hills, Assam.

nodicollis Bates. Darjiling, Mungphu,* Pashok* (5,000 ft.), Sureil* (5,000 ft.), above Sureil* (5,500 ft.), Lebong* (6,000 ft.), and Kurseong* (5,000 ft.), Darjiling dist., E. Himalayas; Shillong* (5,500—6,400 ft.), Khasi Hills, Assam; Tonkin, Indo-China.

pulchella Hope. Darjiling, Mungphu,* Rungeet Valley* (700—800 ft.), and Gopaldhara,* Rungbong Valley, Darjiling dist., E. Himalayas; Simla, Punjab, W. Himalayas; Nepal; Tura* (1,000—1,200 ft.), Garo Hills, Assam; Karen Hills, Burma; Yunnan, W. China.

Genus **Cicindela** Linn. (1735).

agnata Fleut. Kurseong, Darjiling dist., E. Himalayas; Malabar, Ramnad, Madura dist. and Chounambar and Killinour, Pondicherry; (environs), Madras Pres.; Cuttack* (bank of R. Mahanadi), Sur Lake* and Balighai* near Puri, Orissa; Asansol and Berhampur, Murshidabad dist., Bengal; Chakardharpur,* Chota Nagpur, Bihar.

albina Wied. Darjiling, E. Himalayas; Balighai* near Puri, Orissa; Asansol, Bengal; Lohardaga,* Chota Nagpur, Bihar; Rawalpindi* (bank of R. Sohan), Punjab.

albopunctata Chaud. Sikkim*; Darjiling, Mungphu, Ghoom* (7,500 ft.), and Kurseong, Darjiling dist., E. Himalayas; Simla* and Kangra Valley* (4,500 ft.), Punjab, W. Himalayas; Soondrijal,* Nepal; Bhutan; Chamusuri, Moradababu and Mussoorie,* U. Provs.; Doiphang Valley,* Darrang dist., Assam-Bhutan Frontier.

andersoni Gestro. Karen Hills, N. Chin Hills, and Teinzo, Burma; Cochin China.

andrewsi Horn. N. Kanara, Bombay Pres.

andrewsi var. *mauritii* Horn.¹ Sidapur,* Coorg, S. India; N. W. Frontier Province.

andrewesi var. *unica* Fleut. "Inde" (Fleutiaux); "Ex occidentali meridionali Indiae Orientalis Anticae ora" (Horn); S. W. India.

angulata Fabr. Sikkim*; Mungphu, Pankabari* and Sitong,* Darjiling dist., E. Himalayas; Coromandel, Pondicherry (environs), and Ponear, Madras Pres.; Cuttack* (bank of R. Mahanadi), Orissa; Asansol, Calcutta,* Maldah,* Tetara, Damukdia,* Dacca, Bengal; Chota Nagpur, Bihar; Tura* (1,000 ft.), Garo Hills, Assam; Pegu* and Sittaung R.,* Burma; Karachi, Sind; Hainan; Annam and Cambodia, Indo-China.

angulata var. *plumigera* Horn. Mysore, Nilgiri Hills and Trichinopoly, Madras Pres.; Cambodia, Indo-China; Sumatra.

angulicollis Horn. Madura, Madras Pres.; Kodaikanal and Tope,* foot of Palni Hills, S. India; Dacca, E. Bengal.

¹ Referred to as a distinct species by Fowler, but we prefer to follow Horn (1915, p. 302) in regarding it as a variety of *andrewesi*.

assamensis Parry. Sikhim* ; Darjiling, Mungphu,* Gopaldhara,* Rungbong Valley, Pashok* (2,000 ft.) and Sukna* (500 ft.), Darjiling dist., E. Himalayas ; Tura* (1,200—1,500 ft.), and above Tura* (3,000 ft.), Garo Hills, Sylhet, Sibsagar, Nemotha* and Silcuri,* Cachar, Khasi Hills, Naga Hills* and Patkai Hills, Assam ; Arrakan,* Burma ; Penang.

atkinsoni Gestro. Rangoon and Karen Hills, Burma.

atelesta Chaud.¹ Mungphu and Kurseong, Darjiling dist., E. Himalayas ; Simla, Punjab, W. Himalayas ; Mysore, Madras Pres. ; Asansol and Calcutta, Bengal ; Chota Nagpur, Nowatoli and Ranchi, Bihar ; Sylhet, Assam ; Surat, Bombay Pres. ; Nagpur, Central Provs.

aulica Dej. Karachi,* Sind ; Persia ; Arabia ; Bagdad, Mesopotamia ; Brit. Somaliland ; Benguela ; Senegal ; Abyssinia ; Egypt ; Cape Verde Is. ; Angola ; Tunis ; St. Vincent ; Greece.

aulica subsp. *diania* Tsch. Baluchistan ; Sarhad and Bushire,* Persia.

aurofasciata Dej. Darjiling, E. Himalayas ; Travancore ; Mysore, Cochin, Ootacamund* (5,000—7,000 ft.), and Coonoor* (6,000 ft.), Nilgiri Hills,* Shimoga and Pondicherry, Madras Pres. ; S. India* ; N. Kanara, Bombay Pres.

aurofasciata var. *levida* Gory. Nilgiri Hills (2,500 ft.), Mysore, Bangalore* and Shimoga, Madras Pres. ; Belgaum and N. Kanara, Bombay Pres.

aurofasciata var. *seminigra* Fowl.² India.

aurovittata Brullé. Pondicherry and Barkuda Id.,* Madras Pres. ; Calcutta and environs,* Bengal ; Rangoon, Burma ; Ceylon ; Andaman Is.,* Nicobar Is. ; Philippine Is. ; (?) Centr. Japan.

aurulenta Fabr. Sikhim ; Mungphu* (3,000—4,000 ft.), Darjiling dist., E. Himalayas ; Chota Nagpur, Bihar ; Karen Hills, Tavoy,* Pegu* and N. Shan States,* Burma ; Tenasserim* ; Ceylon ; Singapore and Hills near Taiping,* Perak, Malay Peninsula ; Sarawak,* Sandakan* and Banguey Id., Borneo ; Java ; Tamansari* (1,600 ft.), Idjen Massif, E. Java ; Tonkin, Cambodia and Macao, Indo-China ; Siam ; Hong Kong ; Sinkep Id.,* off E. coast and Nias Id., Sumatra ; Formosa ; Lombok, E. Indies.

aurulenta var. *batesi* Fleut.³ Doiphang Valley,* Darrang dist., Assam-Bhutan Frontier ; Pegu, Burma ; Tonkin, Indo-China ; Formosa.

aurulenta var. *flavomaculata* Chevr. Mungphu, Darjiling dist., E. Himalayas ; Pegu, Burma ; Tenasserim ; Tonkin and Macao, Indo-China ; Hong Kong.*

aurulenta var. *virgula* Fleut. Sikhim* ; Kurseong, Ghumti* (4,000 ft.), Soom* (2,500—4,000 ft.), Pashok* (2,000 ft.), Singla* (1,500 ft.), Rungpo* (1,400 ft.), Rungeet Valley (100—800 ft.), Mungphu,* and Gopaldhara,* Rungbong Valley, Darjiling dist., E. Himalayas ;

¹ Cf. Horn (1915, p. 294). Fowler (1912, p. 358) treats this as a variety of *imperfecta*.

² Horn, *Genera Insectorum*, 1915, p. 304, seems to give this and the preceding form as synonyms of *aurofasciata*, but we prefer to follow Fowler in regarding them as varieties.

³ Not recorded in the "Fauna." Cf. Fleutiaux, *Ann. Soc. Ent. France*, p. 491, 1893.

Monda,* Nepal Terai; Chitlong,* little Nepal Valley; Bhutan; Buxa,* frontier of E. Bengal; Tura* (1,000—1,400 ft.), and Garobadha,* Garo Hills, Naga Hills,* Sylhet, Sibsagar,* Silonibari,* and Dejoo,* N. Lakhimpur and Shillong* (5,500—6,400 ft.), Khasi Hills,* Assam; Teinzo, Burma; Tonkin and Cambodia, Indo-China; Hong Kong and Shanghai, China; Singapore, Malay Peninsula.

azureocincta Bates. N. Kanara, Bombay Pres.

bellana Horn. Karachi,* Sind; Fao, Persia.

bellana var. *nuda* Horn. Karachi, Sind.

belli Horn. Anaimalai Hills, Madras Pres.; Mahe, W. coast, India; Travancore; N. Kanara and Belgaum, Bombay Pres.

belli var. *umbropolita* Horn.¹ Teppukadu* (2,500—3,000 ft.), Karkur Ghat* (1,500 ft.), Coonoor and Ouchterlony Valley (2,500—5,000 ft.), Nilgiri Hills, Madras Pres.; Sidapur,* Margalli* and Pollibetta,* Coorg, S. India.

bicolor Fabr. Simla, Punjab and Dehra Dun, base of W. Himalayas; Calcutta (environs),* Maldah,* Pubna,* Murshidabad, Berhampur,* Birbhum,* Asansol and Sabibganj, Bengal; Khasi Hills, Assam; Burma.

bicolor var. *atavus* Horn.² N. India (Lashuwala).

bigemina Klug. Darjiling, Pankabari* and Kurseong,* Darjiling dist., E. Himalayas; Siliguri,* base of E. Himalayas; Punjab; Calcutta,* Maldah, Berhampur,* Murshidabad and Sara Ghat,* Bengal; Purneah dist.,* Chakardharpur,* Chota Nagpur, and Hazara,* Sarai Sateh, Bihar; Gwalior, Centr. India; N. Kanara, Bombay Pres.

bigemina var. *brevis* Horn. "India or." (Horn); Rawalpindi* (bank of R. Sohan), Punjab.

bigemina var. *iravaddica* Gestro. Mandalay, Katha, Teinzo and Tharawaddy, Burma.

bigemina var. *procera* Horn. "India or." (Boucard).

biramosa Fabr. Madras, Waltair,* Mandapam,* Madura dist., Chounambar, Pondicherry, Kanara,* Trichinopoly dist., and Pamben,* Madras Pres.; Tuticorin,* Sravikad* and Trivandrum, Travancore; Mahe, W. coast, India; Balighai* near Puri, Chandipore* near Balasore and Puri,* Orissa; Hooghly River, Sunderbunds* and Chittagong, Bengal; (?) Assam; Arrakan,* Burma; Tenasserim; Mormugao,* Portuguese India; Trincomali,* Ceylon; Andaman Is.*; Nicobar Is.; Java; China; Malacca.

biramosa var. *contracta* Fleut.³ Rangoon, Burma; Tenasserim.

burmeisteri var. *stoliczkanai* Bates. Jhelum,* Kashmir, N. W. Himalayas.

calligramma Schaum. Shimoga, Coimbatore, and Kanara,* Madras Pres.; Trivandrum,* Travancore; Tope,* foot of Palni Hills, S. India; Ceylon.

¹ Fowler in the Fauna gives this variety specific rank, but we prefer to follow Horn (1915, p. 291) in regarding it as a variety of *belli*.

² Not recorded in the "Fauna." Inserted here on the authority of Dr. Horn.

³ Described by Fleutiaux, Ann. Soc. Ent. France, 1893, p. 488, and regarded by Horn as a variety. Cf. Genera Insectorum, 1915, p. 308.

cancellata Dej. Kurseong and Sukna* (500 ft.), Darjiling dist., E. Himalayas ; Siliguri,* base of E. Himalayas ; Mysore, Trichinopoly, Malabar,* Killinour and Chounambar, Pondicherry and Cheria Id.,* Chilka Lake, Madras Pres. ; Quilon* and Tuticorin, Travancore ; Puri,* Sur Lake* and Balighai* near Puri, Orissa ; Karachi, Sind ; Berhampur,* Murshidabad, Asansol and Kunbir, Bengal ; Nowatoli, Bihar ; N. Kanara and Bombay,* Bombay Pres. ; Mahe, W. coast, India ; Garobadha,* Garo Hills, Assam ; Tharawaddy, Leo* (1,170 ft.), Arrakan* and N. Chin Hills, Burma ; Cambodia and Tonkin, Indo-China ; Java.

cancellata var. *candei* Chevr. Pondicherry, and Trichinopoly, Madras Pres. ; Mahe, W. coast, India ; Hong Kong.

cardoni Fleut. Sikhim ; Mysore, Chounambar and Pondicherry, Madras Pres. ; Asansol, Bengal ; Purulia* and Ghatkuri* (1,200 ft.), Manharpur, Singbhumi dist., and Nowatoli,* Chota Nagpur, Bihar ; Taloshi* (2,000 ft.), and Medha* (2,200 ft.), Yenna Valley, Satara dist. and N. Kanara, Bombay Pres. ; Burma ; Chilaw,* Ceylon.

cariana Gestro. N. Shan States,* U. Burma.

catena Fabr. Darjiling, E. Himalayas ; Chatrapur,* Tindivanam, Villapouram, Coromandel, Pondicherry, Barkuda Id.* and Cheria Id.,* Chilka Lake, and Trichur, Cochin State, Madras Pres. ; Quilon,* Travancore ; Tope,* foot of Palni Hills, S. India ; Calcutta,* Bengal ; Chota Nagpur and Ranchi,* Bihar ; Teinzo, Burma ; Bombay ; Mahe, W. coast, India ; Karachi, Sind ; Colombo,* Peradeniya,* Kandy and Matale, Ceylon.

ceylonensis Horn. Trincomali and Wellawaya,* Ceylon.

ceylonensis var. *diversa* Horn. Damboolla, Ceylon.

chinensis De Geer. Simla,* Punjab, W. Himalayas ; Tonkin, Indo-China ; Korea ; Japan ; China.*

chlorida Chaud. Malabar Coast, Podanur near Coimbatore, Madras Pres. ; Tope,* foot of Palni Hills, S. India.

chloris Hope. Rungpo* (1,400 ft.), and Rungeet Valley* (700—800 ft.), Darjiling dist., E. Himalayas ; Siliguri,* base of E. Himalayas ; Gilgit* (4,400 ft.), and Srinagar* (6,000 ft.), Kashmir, W. Himalayas ; Soondrijal,* Benikhola* and Kumdhik,* Nepal Terai ; Kathgodam* (1,200 ft.), Kumaon, Hathikund* and Jumnagwar,* Ghurwal dist., Amangarh,* Patair,* Boxar,* Kaladhungi, Rudpur,* Bindukhera,* Bijnari,* Horai,* Jalaban* and Kiari,* Naini Tal dist., and Bhogpur,* Bijnor dist., U. Provs. ; Rawalpindi* (bank of R. Sohan), Punjab ; Mangaldai dist., N. E.,* Assam-Bhutan Frontier.

chloropleura Chaud. Pashok* (2,500—3,000 ft.), Kalimpong* (600—4,500 ft.), Rungeet Valley* (700—800 ft.), and Pashok Spur,* Darjiling dist., E. Himalayas ; Simla, Punjab, W. Himalayas ; Kashmir, N. W. Himalayas ; Sylhet, Assam.

cognata Wied. Darjiling, E. Himalayas ; Madras ; Cuttack* (bank of R. Mahanadi), Orissa ; Asansol, Berhampur and Goalbathan,* Bengal ; Chota Nagpur, Bihar ; (?) Cambodia, Indo-China.

copulata Schm.-Goeb. Karachi,* Sind.

corbetti Horn. Tharawaddy, Burma.

corticata Putz. Ramnad, Pondicherry, Mysore, Nilgiri Hills (1,250—3,500 ft.), and Trichinopoly, Madras Pres.; Trivandrum,* Travancore; Mahe, W. coast, India; Tope,* foot of Pahni Hills, S. India; Ceylon.

corticata var. *laeticolor* Horn. Ceylon.

crassipalpis Horn. Podanur near Coimbatore (1,000 ft.), Madras Pres.

cyanea Fabr. Sahibganj, Bengal; Palkot,* Chota Nagpur, Bihar.

cyanea var. *dejeani* Hope. Berhampur* and Murshidabad, Bengal; Chota Nagpur, Bihar.

davisoni Gestro. Thagata, Tenasserim.

decempunctata Dej. Murshidabad,* Rajmahal* and Birbhum, Bengal; Assam; Rangoon and Palon, Burma; Cambodia and Tonkin, Indo-China.

decempunctata var. *obscure-dilutata* Horn.¹ Delhi and Lahore* (river-side), Punjab.

delavayi Fairm.² Mungphu, Darjiling dist., E. Himalayas; Yunnan, W. China.

delavayi var. *funebris* Schm.-Goeb. Sikhim*; Mungphu* and Kurseong,* Darjiling dist., E. Himalayas; Punjab; N. E. Frontier, Assam; Naga Hills (4,000 ft.), Assam; Karen Hills, Burma.

despectata Horn.³ Tavoy,* Burma; Philippine Is.; Perak,* Malay Peninsula.

discrepans Walk. Colombo, Kandy and Nalanda, Ceylon.

discrepans var. *lacrymans* Schaum. Kandy,* Ceylon; (?) Nias.

discreta Schaum.⁴ Assam; Cambodia; Simalur Id., and Nias Id., Sumatra; Java; 10 miles S. of Kuching* and Kapit,* Sarawak, Borneo; Celebes; Toe-Kang-Besi Is.; Buru, Siberia; Philippine Is.; New Guinea; N. Queensland.

discreta var. *reductula* Horn.⁵ Tura* (1,000—1,500 ft.), Garo Hills, Assam; N. Chin Hills, Burma; Sarawak, Borneo; Sumatra; Amboyna, Malay Peninsula.

dissimilis var. *singalensis* Horn.⁶ Ceylon.

distinguenda Dej. Barkuda Id.,* Lake Chilka, Madras, and Pondicherry, Madras Pres.; Mahe, W. coast, India; Ceylon.

distinguenda var. *lunulata* Horn. Madras.

dives Gory. Sikhim; Mysore, Madras Pres.; Bengal; Belgaum and N. Kanara, Bombay Pres.; Mhow, Centr. India.

dormeri Horn. Kandy,* Ceylon.

dromicoides Chaud. Darjiling, Ghumti* (4,000 ft.), Mungphu,* Soom to Darjiling (4,500—7,000 ft.), below Sureil* (4,800 ft.), and Kurseong,* Darjiling dist., E. Himalayas; Nepal; Simla, Punjab, Kumaon* and Kousanie* (6,075 ft.), U. Provs., W. Himalayas; Nowatoli and Chota Nagpur, Bihar; Khasi Hills, Assam; Bombay.

¹ Not recorded in the "Fauna." Cf. Horn, 1913, p. 28.

² Cf. Horn, 1915, p. 289 for this species and its variety.

³ Not recorded in the "Fauna." Horn (1915, p. 290) records it doubtfully from Perak, but we have specimens both from Tavoy in Burma and Perak in the Malay Peninsula.

⁴ This species is included here on the authority of Dr. Horn. Cf. Horn, 1915, p. 298.

⁵ Previously known as *reducta*. Cf. Horn, 1915, p. 298.

⁶ Not recorded in the "Fauna." Cf. Horn, 1915, p. 293.

duponti Dej. Merchiston* State near Trivandrum, Travancore; Trichur, Cochin State, Nilgiri Hills, Mysore and Shimoga, Madras Pres.; Coorg (2,000 ft.), S. India; Bengal; Chota Nagpur, Bihar; Maflong,* Shillong, Cheerapunji,* Nongpriang,* and Maupun,* Khasi Hills, Tura* (1,000—1,400 ft.), Garo Hills, Sylhet and Patkai Hills,* Assam; Arrakan, Pegu, N. Chin Hills, N. Shan States, Karen Hills and Tharawaddy, Burma; Talewadi* near Castle Rock, N. Kanara dist., Bombay Pres.; Mahe, W. coast, India; Annam and Tonkin, Indo-China; Penang; Yunnan, W. China.

duponti var. *barmanica* Gestro. Nilgiri Hills,* Madras Pres.; Coorg* (2,000 ft.), S. India; Sylhet and N. Manipur, Assam; N. Shan States,* Karen Hills, Tharawaddy and Pegu,* Burma.

eruditia Wied. (?) Tenasserim; Kashmir and Kulu, W. Himalayas; S. India*; Maldah, Kierpur and Katihar, Purneah dist., Chota Nagpur and Pusa, Bihar; Agra and Allahabad, U. Provs.

fabriciana Horn.¹ N. Kanara, Bombay Pres.

fastidiosa Dej. Kurseong, Darjiling dist., E. Himalayas; Kashmir and W. Himalayas; Dehra Dun, base of W. Himalayas; Rawalpindi, Punjab; Barkuda Id.,* Lake Chilka and Rambha,* Ganjam dist. and Mysore, Madras Pres.; Nowatoli and Chota Nagpur, Bihar; Assam*; Pegu,* Burma; Mhow, Centr. India; Trincomali* and Anuradhapura (low country),* Ceylon.

fleutiauxi Horn.² Sikkim*; Singla* (1,500 ft.), Ghumti* (1,500—5,000 ft.), Mungphu,* Gopaldhara,* Rungbong Valley and Kurseong, Darjiling dist., E. Himalayas; Bengal.

foveolata Schaum. Malabar and Teppukadu* (3,000 ft.), Nilgiri Hills, Madras Pres.; Dacca, E. Bengal; Tura* (1,200 ft.), Garo Hills, Assam; Karen Hills, Maymyo,* Teinzo and Tharawaddy, Burma; Tonkin, Indo-China; Celebes; Sumatra; Philippine Is.

fuliginosa Dej. Bhamo, Pegu* and Teinzo, Burma; Ceylon; Malay States; Java; Shanghai, China; Borneo*; (?) Banka, E. Indies; Sumatra; Laos and Cambodia, Indo-China; Malacca.

funerea McLeay. Sikkim; Darjiling, Gopaldhara,* Rungbong Valley, Mungphu, Pankabari, Kalimpang* (600—4,500 ft.), Singla* (1,500 ft.), Kurseong, Rungeet Valley* (700—800 ft.), Rungpo* (900 ft.), and Sukna,* Darjiling dist., E. Himalayas; Soondrijal,* Nepal; Simla, Punjab, W. Himalayas; Mysore, Madras Pres.; Barway, W. Bengal; Sibsagar, Sylhet and Tura* (1,000 ft.), Garo Hills, Assam; Doiphang Valley,* Darrang dist., Assam-Bhutan Frontier; Allanmyo, N. Shan States,* Pegu,* Sadon* (5,000 ft.), Teinzo and Tharawaddy, Burma; Tenasserim; Poona, Bombay Pres.; Ghatkuri* (1,200 ft.), Manharpur, Singbhum dist., Bihar*; Hainan; Lampam,* Patalung, Siam; Perak,* Malay Peninsula; Java*; Sumatra; Celebes; Annam, Indo-China.

ganglbaueri Horn. Ceylon.

¹ Horn (1915, p. 303) gives this as a new name for *fabricii* Horn. Cf. Fowler, 1912, p. 403.

² Horn (1915, p. 303) gives this as a new name for *oberthuri* Fleutiaux. Cf. Fowler, 1912, p. 380.

germanica var. *kirilovi* Fisch.¹ Gilgit* (5,000 ft.), Kashmir, N. W. Himalayas; Persia (up to Sarhad and Seistan in the south-east); Turkestan; Kolyvan dist., Siberia; Altai Mts., Mongolia; Transcaspia.

grammophora Chaud. Mahanadi,* Darjiling dist., E. Himalayas; Calcutta and environs,* Maldah,* Sara Ghat,* Damukdia,* Goalbathan* and Asansol, Bengal; Katihar,* Purneah dist., Chota Nagpur, Pusa,* and Siripur,* Saran, Bihar; Kaladhungi,* Naini Tal dist., U. Provs.

guttata Wied. Dehra Dun, base of W. Himalayas; Shimoga and Ganjam, Madras Pres.; Chapra,* Ranchi* and Lohurdaga,* Bihar.

gyllenhali Dej. Bandra and Bassein, Bombay Pres.; Karachi,* Sind; E. Indies* (British Museum).

haemorrhoidalis Wied. Trivandrum, Travancore; Nilgiri Hills, Mysore, Karkur Ghat, Malabar, Barkuda Id.* and Gantesila,* Lake Chilka, Madras Pres.; Coorg, S. India; Angul* and Barkul* (0--1,000 ft.), Orissa; Calcutta, Bengal; Ranchi and Lohurdaga,* Bihar; Sibsagar,* N. E. Assam; N. Kanara, Bombay Pres.; Jubbulpore,* Centr. Provs.; Rawalpindi,* Punjab; Wellawaya,* Ceylon; Canton,* China.

haemorrhoidalis var. *xanthospilota* Hope, Madras.

hamiltoniana Thoms. Nadgani,* Malabar, Mysore and Nilgiri Hills,* Madras Pres.; Merchiston State* near Trivandrum, Travancore; Pollibetta,* Coorg, S. India.

helferi Schaum.² (?) Burma.

histrio Tsch. Karachi and Manora, Sind; Khorassan, Seistan and Bamfur, Persia.

holosericea Fabr. Asansol, Bengal; Nowatoli, Chota Nagpur, Bihar; Khasi Hills, Assam; Karen Hills, Burma; Java; Sumatra; Philippines.

humillima Gestro. Bhamo, Katha, Rangoon,* Shwego, Teinzo and Tharawaddy, Burma.

indica Fleut. Bombay.

intermedia Chaud. Jhelum Valley,* Kashmir, N. Punjab to Simla and Kumaon, W. Himalayas; Kathgodam* (1,200 ft.), and Kousanie* (6,075 ft.), U. Provs.; Assam.

interrupto-fasciata Schm.-Goeb. Teinzo, Ruby Mines and Maymyo, Burma; Tonkin, Laos and Cambodia, Indo-China; Hong Kong.

kaleea Bates.³ (?) Sikkim; Assam; Kiachou and Yunnan, China; Formosa*; Tonkin, Indo-China.

lacunosa Putz. Habarane* and Puttalam, Ceylon.

lefroyi Horn. Chapra and Pusa, Bihar.

¹ Horn (1915, p. 346) records this variety from Barkul. Now, as Barkul is in the Puri district of Orissa and is situated in south-east India we doubt this distribution, since the variety has been taken only in the north-west of India and is more especially known from central and south-western Asia.

² Horn, 1915, p. 309 regards *C. limbata* as a synonym of *C. helferi* and doubtfully records its locality.

³ Included on the authority of Dr. Horn. Cf. Horn, 1915, p. 298.

leucoloma Chaud. Simla, Punjab, W. Himalayas ; (?) Ceylon ; "India" or "East India" ; N. India.

limosa Saund. Mungphu,* Darjiling dist., E. Himalayas ; Chandipore* near Balasore, Orissa ; Durgapur* (salt lakes) near Calcutta, Bengal ; Pegu,* Burma ; Ceylon ; Andaman* and Nicobar Is., Chusan Is.

lunulata Fabr. Nushki dist.* W. Baluchistan ; Bagdad, Mesopotamia ; Caucasus ; S. Russia ; Lybia ; Abyssinia ; Morocco ; Corsica and Lampedusa Is., Italy ; Peking, China.

lunulata var. *nemoralis* Oliv. Perso-Baluch Frontier* ; Pamir ; Centr. Mongolia ; Seistan ; Turkestan ; Persia ; Syria ; Sinai ; Palestine ; Suez ; Islands of Rhodes, Crete, Cyprus, Malta, Elba, Sicily and (?) Sardinia ; Italy ; S. Europe.

maindroni Horn. Kiamari Id., Bombay Pres. ; Karachi,* Sind ; Baluchistan.

malabarica Maind. and Fleut. Mahe, W. coast, India.

mariae Gestro. N. Chin Hills, Tharawaddy, Taung-ngu, Pegu,* Rangoon, and Karen Hills, Burma ; Tenasserim.

melancholica Fabr. Kalka* (2,400 ft.), base of Simla Hills, W. Himalayas ; Kaladhungi,* Naini Tal dist., U. Provs. ; Ambala,* Punjab ; Bengal ; Pusa,* Bihar ; Nagpur, Central Provs. ; Bandra, Bombay Pres. ; Karachi, Sind ; Bushire,* Persia ; Anatolia, Rhodes, Syria to Arabia, Caucasus to Turkestan ; the Islands of Cape Verde, Crete, Sicily, Sardinia, Fernando Po, Principe, St. Thomas, Annabon, (?) Comoro and Cephalonia, Greece ; the whole continent of Africa.

minuta Oliv. Darjiling and Pashok* (1,000 ft.), Darjiling dist., E. Himalayas ; Trichur, Cochin State and Pondicherry, Madras Pres. ; Cuttack (bank of R. Mahanadi), and Dhauli,* Puri dist.,* Orissa ; Calcutta,* Dacca, Berhampore and Maldah,* Bengal ; Chakardharpur,* Chota Nagpur, Chapra, and Pusa,* Bihar ; Garobadha* and Tura* (1,000 ft.), Garo Hills, Assam ; Pegu* and Tharawaddy, Burma ; Myawadi* (900 ft.), Burmo-Siamese Frontier, Amherst dist. ; Patan* (2,050 ft.), Koyna Valley, Satara dist., Bombay Pres. ; Mahe, W. coast, India ; Delhi,* Punjab ; Sumbawa, Sumatra ; Perak,* Malay Peninsula ; Tonkin, Indo-China ; Borneo ; Java ; China ; Philippines.

motschulskyana Horn.¹ N. Kanara, Bombay Pres.

mouhoti var. *anometallescens* Horn. Maymyo and Ruby Mines, Burma ; Momeit, Yunnan, W. China.

mouhoti var. *bramani* Dokht. Pegu, Burma ; Lukhon, Siam ; Laos and Tonkin, Indo-China ; China (proper).

mouhoti var. *cariana* Gestro. N. Shan States and Karen Hills, Burma ; Tenasserim.

mouhoti var. *goebeli* Horn. Bhamo, Rangoon and Teinzo, Burma.

multiguttata Dej. Nepal ; Calcutta,* Asansol and Sara Ghat,* Bengal ; Patkai Hills, Assam.

mutata Fleut. Bhamo, Pegu, Teinzo and Tharawaddy, Burma ; Tonkin, Indo-China.

¹ Horn (1915, p. 289) gives this as a new name for *motschulskyi*. Cf. Fowler, 1912, p. 342.

nietneri Horn. Ceylon.

nitida Wied. Darjiling and Kurseong, Darjiling dist., E. Himalayas ; Bhutan ; Pondicherry and Ramnad, Madras Pres. ; Cuttack* (bank of R. Mahanadi), Orissa ; Berhampur* (bank of R. Bhagarati), Calcutta,* Maldah,* Sara Ghat,* Rajmahal,* Asansol and Jugge-nathgunge* (R. Brahmaputra), Bengal ; Kaladhungi,* Naini Tal dist., U. Provs. ; Chota Nagpur, Bihar ; Pegu,* Burma ; Karachi, Sind ; Cambodia, Indo-China.

octogramma Chaud. Dharhar, Bengal ; Dinapur, Bihar ; Kangra Valley, Punjab ; Yunnan, W. China.

octonotata Wied. Darjiling,* Pankabari* and Sukna* (500 ft.), Darjiling dist., E. Himalayas ; Simla, Punjab, W. Himalayas ; Agra, U. Provs. ; Asansol, Dacca, Maldah,* Murshidabad and Miani Mukh* (at junction of Miani and Kassalong Rivers), Chittagong Hill Tracts, Bengal ; Pusa,* Bihar ; Tura* (1,200 ft.), and Garobadha,* Garo Hills, Khasi Hills, Lushai Hills,* Sylhet and Sibsagar,* Assam ; Jamayi, Pegu* and Tharawaddy, Burma.

ornata Fleut. Karachi,* Sind.

phalangoides Schm.-Goeb. N. Chin Hills, Karen Hills, Pegu* and Teinzo, Burma.

princeps Vig. "Indien," Surada (Ganjam) [Horn] ; (?) Madras ; Centr. India.

princeps var. *ducalis* Horn. Palkot,* Chota Nagpur, Bihar.

prothymoides Horn. Karkur Ghat, Malabar, Madras Pres.

quadrilineata Fabr. Chounambar, Madras Pres. ; Chandipore* near Balasore, Orissa sea-coast ; Sunderbunds, Gangetic delta and Channel Creek* near Mud Point, Bengal ; Arrakan, Burma ; Tenasserim ; Dabhol,* Koltharen* and Harni,* Ratnagiri dist., Bombay Pres. ; Cacara Bay,* Portuguese India ; Baluchistan ; Manora, Karachi, Sind ; Ceylon to North India.

quadrilineata var. *renei* Horn. Trichinopoly and Pamben,* Madras Pres. ; Karachi, Sind ; Baluchistan ; Ceylon.

rugosiceps Chaud. Nilgiri Hills, Ramnad, Pillar, Mysore and Coonoor (2,500—3,500 ft.), Madras Pres.

seriepunctata Horn. Kalimpong* (600—4,500 ft.), Pashok* (2,000 ft.), Singla* (1,500 ft.), Rungeet Valley* (700—800 ft.), and Kurseong, Darjiling dist., E. Himalayas ; N. Bengal.

sexpunctata Fabr. Sikhim* ; Mungphu, Darjiling dist., E. Himalayas ; Raxaul,* Nepal Frontier ; Gilgit,* Kashmir, N. W. Himalayas ; Bangalore* and Shimoga, Madras Pres. ; Coorg, S. India* ; Calcutta* and environs,* Sunderbunds, Diamond Harbour,* Bosondhar, Berhampur, Birbhum, Raniganj,* Damukdia,* Maldah,* Alipur Duars,* Sara Ghat,* Khargpur* and Basanti,* Bengal ; Purneah, Ranchi,* Champaran,* and Chota Nagpur, Bihar ; Sadiya,* and Shillong,* Assam ; Rangoon,* Pegu, Sittaung River, Arrakan,* delta of the Irrawaddy,* Kawkareik* and Farm Caves* near Moulmein, Burma ; Bombay ; Mahe, W. coast, India ; Wellawaya* and Peradeniya,* Ceylon ; Andamans ; Siam ; (?) Java ; Annam to Hong Kong ; E. Indies* ; Cambodia, Indo-China ; Formosa ; Philippines ; Senegal and Cameroon, Africa.

shivah Parry. Nepal ; N. Kanara, Bombay Pres. ; (?) Assam.

sinica Fleut. Taik-gyi, Pegu, Burma ; Saigon and Chiem Hoa, Tonkin, Indo-China ; Bangkok, Siam ; Java ; Hong Kong.

spinolae Gestro. Darjiling, Pashok* (1,000, 2,000 and 5,000 ft.), Singla* (1,500 ft.), Rungeet Valley* (700—800 ft.), and Sukna* (500 ft.), Darjiling dist., E. Himalayas ; Dehra Dun, W. Himalayas ; Asansol, Dacca and Rangamati,* Chittagong Hill Tracts, Bengal ; Sylhet, Patkai Hills and Tura* (1,000—1,500 ft.), Garo Hills, Assam ; N. Chin Hills, Bhamo, Teinzo, Tharawaddy, Karen Hills, Ruby Mines, and Rangoon, Burma ; Tenasserim ; Chota Nagpur, Bihar ; Tonkin, Indo-China ; Momeit, Yunnan, W. China.

striatifrons Chaud. Murshidabad * and Berhampur,* Bengal ; N. Kanara, Bombay Pres.

striolata Ill. Munghpu* Soom* (3,000—3,500 ft.), and Pashok* (3,000 ft.), Darjiling dist., E. Himalayas ; Dehra Dun, base of W. Himalayas ; Trivandrum,* Travancore ; Nilgiri Hills and Kanara,* Madras Pres. ; Pollibetta* and Coorg, S. India ; Calcutta, Maldah * and Rangamati,* Chittagong Hill Tracts, Bengal ; Khasi Hills, Assam ; Pegu,* Teinzo and Tharawaddy, Burma ; Tenasserim ; Chota Nagpur, Bihar ; Mahe, W. coast, India ; Tonkin, Indo-China ; Singtau, China ; Java ; Borneo ; Sumatra ; Celebes ; Formosa ; Philippines.

striolata var. *lineifrons* Chaud. Dehra Dun, base of W. Himalayas ; Nilgiri Hills, Madras Pres. ; above Tura* (2,500 ft.), Garo Hills, Assam ; N. Chin Hills, Karen Hills, Pegu,* Teinzo and Tharawaddy, Burma ; Tenasserim ; Perak, Malay Peninsula ; Cambodia and Tonkin, Indo-China.

striolata var. *parvimaculata* Fowl.¹ (?) Darjiling, E. Himalayas.

sublacerata var. *balucha* Bates. Skardo, Kashmir, N. W. Himalayas ; Baluchistan ; Perso-Baluch Frontier * ; Caucasus ; Armenia ; Transcasplia ; Turkestan (to Ferghana and the Pamirs).

sumatrensis Hbst. Siliguri,* base of E. Himalayas ; Kumdhik, Maho,* Nepal Terai ; Kashmir, N. W. Himalayas ; Trivandrum, Travancore,* Telicherry, Pondicherry, Barkuda Id.* and Cheria Id.,* Lake Chilka, Madras Pres. ; Cuttack,* Orissa ; Calcutta,* Damukdia,* Maldah and River Ganges, Bengal ; Bhamo, N. Chin Hills, Pegu, Tavoy, Teinzo and Tharawaddy, Burma ; Tura* (1,000 ft.), Garo Hills, Assam ; Chakardharpur,* Singbhumi dist. (base of hills), Chota Nagpur, Ranchi and Purulia,* Bihar ; Patan* (2,050 ft.), Koyna Valley, Satara dist., and Khandesh, Bombay Pres. ; Mahe, W. coast, India ; Cacara Bay,* Portuguese India ; Bhogpur,* Bijnor dist., U. Provs. ; Karachi, Sind ; Ceylon ; Hainan, Quang-Tang and Yunnan, China ; Central China * ; Perak, Malay Peninsula ; (?) Java ; Borneo ; Japan ; Formosa ; Sumbawa, E. Indies ; Sumatra ; Philippines ; Sinkep Id.*

sumatrensis var. *imperfecta* Horn. Mussoorie,* W. Himalayas ; Forest Tramway, mile 10—14* (0—300 ft.), Cochin State, Shimoga and

¹ Horn (1915, p. 305) classes this as a separate species but we prefer for the present to follow Fowler in regarding it as a variety.

Mysore, Madras Pres. ; Ranchi,* Bihar ; Castle Rock,* N. Kanara dist., Pimpli,* Vashishti Valley (100 ft.), Ratnagiri dist., and Medha* (2,200 ft.), Yenna Valley, Helvak* (2,000 ft.), Taloshi* (2,000 ft.), Patari* (2,050 ft.), Koyna Valley, Satara dist., and Surat,* Bombay Pres. ; Jubbulpore,* Centr. Provs.

tetragrammica Chaud. Anaimalai Hills* (3,000—4,000 ft.), Malabar Coast, Nilgiri Hills, Ramnad, Shembaganur and Trichinopoly, Madras Pres. ; Pykara (6,000 ft.), Coorg, S. India ; S. Bombay. *tetrasticta* Wied. Ganjam, Gantasila* and Barkuda Id.,* Lake Chilka, Madras Pres. ; Asansol, Birbhum,* Calcutta,* River Ganges,* Bengal ; Nowatoli and Chota Nagpur,* Bihar ; Dharwar, Bombay Pres.

triguttata Hbst. Darjiling dist. (600—4,500 ft.), and Kalimpong* (600—4,500 ft.), E. Himalayas ; Siliguri,* base of E. Himalayas ; Dehra Dun, base of W. Himalayas ; Calcutta * and Rangamati,* Chittagong Hill Tracts, Bengal ; Tura* (1,300 ft.), Garo Hills, Assam ; Bhamo, Karen Hills, Pegu,* Taung-ngu, Teinzo and Tharawaddy, Burma ; Tenasserim ; Hong Kong and Shanghai, China ; Sumbawa, E. Indies ; Borneo ; Sumatra ; Tonkin, Indo-China ; Singapore ; Malay Archipelago ; Celebes ; Philippines.

tritoma Schm.-Goeb. Karen Hills and Pegu, Burma.

undulata Dej.¹ Barkuda Id.,* Gopkuda Id.* and Samal Id.,* Lake Chilka, Mysore and Ramnad, Madras Pres. ; Balugaon,* Puri dist., Orissa ; Asansol, Calcutta,* Dacca, and Maldah * Bengal ; Nowatoli and Lohardaga,* Bihar ; Sylhet, Assam ; Gulistan,* Baluchistan ; Mahe, W. coast, India ; Gwalior, Centr. India ; Allahabad, U. Provs. ; Ceylon ; Hong Kong ; Perak, Malay Peninsula.

undulata var. *dubia* Horn. Singla* (1,500 ft.), and Darjiling, Darjiling dist., E. Himalayas ; N. Assam ; Pakokku* (180 ft.), Mandalay, and Myingyan,* Burma.

venosa Koll. Kurseong and Darjiling, E. Himalayas ; Kashmir, N. W. Himalayas ; Bhutan ; Pondicherry and Ramnad, Madras Pres. ; Cuttack, Orissa ; Berhampur, Calcutta, Maldah, Rajmahal and Sara Ghat, Bengal ; Pegu, and Hmawbi,* Burma ; Patna dist. and Chota Nagpur, Bihar ; Karachi, Sind ; Kaladhungi, U. Provs. ; Cambodia, Indo-China.

venus Horn. Nadgani, Malabar and Nilgiri Hills, Madras Pres.

venus var. *stuprata* Horn.² Travancore.

vigintiguttata Hbst. Sikhim ; Kurseong* (5,000 ft.), Mungphu* (3,000—4,000 ft.), and Pankabari,* Darjiling dist., E. Himalayas ; Bhutan * ; Barkul,* Puri dist., Orissa ; Berhampur,* Birbhum,* Dacca, Damukdia Ghat,* Murshidabad,* Rajmahal * and Sara Ghat,* Bengal.

viridicincta Horn. Dehra Dun, base of W. Himalayas ; Mysore, Kanara and Nilgiri Hills (3,000 ft.), Madras Pres. ; Pollibetta,* Coorg, S. India ; Chota Nagpur, Bihar.

¹ Cf. Annandale and Dover (1921, p. 336) and Dover and Ribeiro (1921, p. 724),

² Not recorded in the Fauna. Cf. Horn (1915, p. 287).

- viridilabris* Chaud. Kalka* (2,400 ft.), and Simla, Punjab, W. Himalayas ; Bengal ; Pusa,* Bihar ; East Indies.
- viridilabris* var. *fallaciosa* Horn. Karen Hills and Teinzo, Burma.
- viridilabris* var. *fusco-cuprascens* Horn. Nilgiri Hills (1,250 ft.), and Trichinopoly, Madras Pres.
- viridilabris* var. *labioaenea* Horn. Colombo, Kandy,* Negombo, Patta-lam and Peradeniya, Ceylon.
- viridilabris* var. *severini* Horn. Travancore and Nilgiri Hills* (3,000 ft.) ; Madras Pres. ; Pollibetta,* Coorg, S. India ; Bengal ; Teinzo, Burma ; Mahe, W. coast, India ; Chota Nagpur, Bihar ; Tonkin, Indo-China.
- vittigera* Dej. Berhampur, Calcutta, Dacca, Damukdia Ghat,* Goal-bathan* and Maldah,* Bengal.
- waterhousei* Horn. Ceylon.
- westermanni* Schaum. Coromandel, Madras and Tranquebar, Madras Pres.
- withilli* Hope. Trichur, Cochin State, Anaimalai Hills (3,000—4,000 ft.), Coonoor (environs), N. Malabar,* Nilgiri Hills,* Ootacamund, and Wyanaad, Madras Pres. ; Wallardi, Travancore ; Mahe, W. coast, India ; Talewadi* near Castle Rock, N. Kanara dist., Belgaum, Khanapur and Poona,* Bombay Pres. ; Tharawaddy, Burma.
- willeyi* Horn. Central Provinces, Ceylon.

Genus *Apteroessa* Hope (1838).

grossa Fabr. Tranquebar, Coromandel Coast, Madras Pres.

Subfamily MEGACEPHALINAE.

Genus *Megacephala* Latr. (1802).

- euphratica* Dej. Karachi, Sind ; Persia ; R. Euphrates, Mesopotamia ; Sinai ; (?) Syria ; Caucasus ; Andalusia ; Gulf of Tadjura ; Cyprus ; Rhodes ; Cartagena ; Egypt and Algeria to Tripoli, Africa.
- euphratica* var. *armeniaca* Cast.¹ Agra, U. Provs.* ; Afghan-Baluch Frontier* ; Armenia and Transcaspia to Amu Daria ; Seistan,* Persia.

¹ It is also found in the same localities as the above species.