

Poster presentation

Open Access

The amyloid precursor protein potentiates CHOP induction and cell death in response to ER Ca²⁺ depletion

Ekaterini Copanaki*, Tina Schürmann, Anne Eckert, Kristina Leuner, Walter E Müller, Jochen HM Prehn and Donat Kögel

Address: Department of Neurosurgery, Johann Wolfgang Goethe University Clinics, Frankfurt am Main, Germany

* Corresponding author

from Annual Meeting of the Study Group Neurochemistry. International Conference of the Gesellschaft für Biochemie und Molekularbiologie 2006 (GBM 2006): Molecular pathways in health and disease of the nervous system Witten, Germany. 28–30 September 2006

Published: 23 March 2007

BMC Neuroscience 2007, 8(Suppl 1):P26 doi:10.1186/1471-2202-8-S1-P26

© 2007 Copanaki et al; licensee BioMed Central Ltd.

Here we investigated the role of the amyloid precursor protein (APP) in regulation of Ca²⁺ store depletion-induced neural cell death. Ca²⁺ store depletion from the endoplasmic reticulum (ER) was induced by the SERCA (Sarco/Endoplasmic Reticulum Calcium ATPase) inhibitor thapsigargin which led to a rapid induction of the unfolded protein response (UPR) and a delayed activation of executioner caspases in the cultures. Overexpression of APP potently enhanced cytosolic Ca²⁺ levels and cell death after ER Ca²⁺ store depletion in comparison to vector-transfected controls. GeneChipR and RT-PCR analysis revealed that the expression of classical UPR chaperone genes was not altered by overexpression of APP. Interestingly, the induction of the ER stress-responsive pro-apoptotic transcription factor CHOP was significantly upregulated in APP-overexpressing cells in comparison to vectortransfected controls. Chelation of intracellular Ca²⁺ with BAPTA-AM revealed that enhanced CHOP expression after store depletion occurred in a Ca²⁺-dependent manner in APPoverexpressing cells. Prevention of CHOP induction by BAPTA-AM and by RNA interference was also able to abrogate the potentiating effect of APP on thapsigargin-induced apoptosis. Application of the store-operated channel (SOC)-inhibitors SK F96365 and 2-APB downmodulated APP-triggered potentiation of cytosolic Ca²⁺ levels and apoptosis after treatment with thapsigargin. Our data demonstrate that APP-mediated regulation of ER Ca²⁺ homeostasis significantly modulates Ca²⁺ store depletion-induced cell death in a SOC- and CHOP-dependent manner, but independent of the UPR.