



Estimating N_2O_5 uptake coefficients using ambient measurements of NO_3 , N_2O_5 , ClNO_2 and particle-phase nitrate.

G. J. Phillips¹, J. Thieser¹, M. J. Tang¹, N. Sobanski¹, G. Schuster¹, J. Fachinger², F. Drewnick², S. Borrmann², H. Bingemer³, J. Lelieveld¹, and J. N. Crowley¹

5 ¹Department of Atmospheric Chemistry, Max Planck Institute for Chemistry, Germany

²Particle Chemistry Department, Max Planck Institute for Chemistry, Germany

³Institute for Atmospheric and Environmental Sciences, Goethe University, Frankfurt, Germany

Correspondence to: John Crowley (john.crowley@mpic.de)

Abstract. We present an estimation of the uptake coefficient (γ) and yield of nitryl chloride (ClNO_2) (f) for the
10 heterogeneous processing of dinitrogen pentoxide (N_2O_5) using simultaneous measurements of particle and trace gas
composition at a semi-rural, non-coastal, mountain site in the summer of 2011. The yield of ClNO_2 varied between ($0.035 \pm$
 0.027) and (1.38 ± 0.60) with a campaign average of (0.49 ± 0.35). The large variability in f reflects the highly variable
chloride content of particles at the site. Uptake coefficients were also highly variable with minimum, maximum and average
 γ values of 0.004, 0.11 and 0.028 ± 0.029 , respectively, with no significant correlation with particle composition, but a weak
15 dependence on relative humidity. The uptake coefficients obtained are compared to existing parameterisations based on
laboratory datasets and with other values obtained by analysis of field data.

1 Introduction

The reaction of N_2O_5 with atmospheric aerosol represents an important nocturnal control on the atmospheric lifetime of NO_x
and can impact the production of atmospheric oxidants, such as ozone and hydroxyl radicals (Dentener and Crutzen, 1993;
20 Riemer et al., 2003; Brown et al., 2006; Macintyre and Evans, 2010). The reaction, in addition to loss of NO_x , results in the
formation of particulate nitrate and can result in the release of ClNO_2 in both marine and continental environments (Osthoff
et al., 2008; Kercher et al., 2009; Thornton et al., 2010; Mielke et al., 2011; Phillips et al., 2012; Riedel et al., 2012a; Bannan
et al., 2015).

Neglecting gas-phase diffusive effects, which are insignificant for transport of N_2O_5 to the sub-micron diameter particles
25 dealt with in this study, the rate of loss of N_2O_5 on a particle surface can be described by the following expression:




$$\frac{d[\text{N}_2\text{O}_5]}{dt} = -0.25\bar{c}\gamma A[\text{N}_2\text{O}_5] \quad (1)$$

where \bar{c} is the mean molecule speed of N_2O_5 , γ is the uptake coefficient, and A is the aerosol surface area density (i.e. the
particle surface area per volume of air). This expression is the basis of the numerous laboratory studies designed to derive γ



for tropospheric and stratospheric particles and its dependence on environmental variables such as temperature and relative humidity (Ammann et al., 2013).

The reaction of N_2O_5 with aqueous particles is complex. Following bulk accommodation, the uptake proceeds via disproportionation of N_2O_5 in the aqueous phase (Mozurkewich and Calvert, 1988) to form HNO_3 (and NO_3^-):




In the presence of chloride, $ClNO_2$ can also be formed (Finlayson-Pitts et al., 1989; George et al., 1994; Behnke et al., 1997).



10 The relative rate of formation of the NO_3^- (or HNO_3) and $ClNO_2$ products of R2 and R4 depends on the fate of $H_2NO_3^+$ (hydrated nitronium ion) and thus on the concentration of chloride available (Behnke et al., 1997; Schweitzer et al., 1998; Thornton and Abbatt, 2005; Bertram and Thornton, 2009; Roberts et al., 2009) and on the rate coefficients for its aqueous phase reaction with Cl^- (k_4) or H_2O (k_2).

We define the branching ratio to $ClNO_2$ formation as f so that the net yield of NO_3^- and $ClNO_2$ formed (per N_2O_5 taken up)
 15 can be written as:



f can be calculated from knowledge of the relative chloride and water content of a particle and the rate coefficients k_2 and k_4 :

$$f = \frac{k_4 [Cl^-]}{k_4 [Cl^-] + k_2 [H_2O]} \quad (2)$$

A value of k_4 / k_2 of 450 is presently recommended (Ammann et al., 2013), but may be modified for systems with a
 20 significant concentration of aromatics which can also react with hydrated nitronium ions (Ryder et al., 2015).

Nitrate formed in the uptake of N_2O_5 can partition to the gas phase as nitric acid, or remain in the particle depending on the pH, available ammonia/ammonium and temperature. Apart from highly acidic aerosol (Roberts et al., 2009) $ClNO_2$ is only weakly soluble / reactive and degasses completely from the particle. In the absence of appreciable chemical losses at night,
 25 $ClNO_2$ remains in the atmosphere until dawn when it is removed by photolysis over a period of a few hours (Ghosh et al., 2012).



The photochemical destruction of $ClNO_2$ recycles NO_x and activates Cl atoms for possible oxidation of VOCs, which may lead (via peroxy radical formation) to enhanced NO to NO_2 oxidation and ozone production (Simon et al., 2009; Sarwar et al., 2012). The chemical processes involved in the nighttime formation of N_2O_5 , its heterogeneous processing and the
 30 subsequent daytime photochemical reactions involving Cl atoms (and their impact on VOC oxidation and ozone formation) are illustrated in Fig. 1.

Laboratory studies (using synthetic surrogates for atmospheric aerosol) indicate that N_2O_5 uptake coefficients on aqueous, tropospheric aerosol are large (e.g. $1-3 \times 10^{-2}$ for ammonium sulphate) and show a complex dependence on environmental



variables such as temperature and relative humidity and also on the nitrate and chloride content of the particle. A detailed summary of the results of the laboratory studies is given by Amman et al. (2013).

Briefly, the presence of particulate nitrate has been seen to reduce γ , which is understood in terms of enhancing the rate of (R-1) compared to R2 and R4 (Wahner et al., 1998a; Wahner et al., 1998b; Mentel et al., 1999; Hallquist et al., 2003; Bertram and Thornton, 2009). In contrast, the presence of chloride increases γ through the competitive removal of H_2NO_3^+ in R4 (Bertram and Thornton, 2009). Laboratory experiments have also documented the reduction of γ caused by particle phase organics, including organic coatings, which lower the water activity and the hydrolysis rate of N_2O_5 and possibly the rate of accommodation of N_2O_5 at the gas-particle interface (Anttila et al., 2006; Badger et al., 2006; McNeill et al., 2006; Park et al., 2007; Cosman and Bertram, 2008; Cosman et al., 2008; Gaston et al., 2014). The uptake coefficient to dry particles is reduced for the same potential reasons (Hu and Abbatt, 1997; Thornton et al., 2003).

The loss of N_2O_5 to ambient aerosol samples (as opposed to chemically “simpler” aerosol surrogates prepared in the laboratory) was investigated by Bertram and colleagues (Bertram et al., 2009a; Bertram et al., 2009b; Riedel et al., 2012b) who generated gas-phase N_2O_5 and monitored its loss via uptake to atmospheric aerosol sampled into a flow-reactor. The results indicated that the uptake coefficient was highly variable, sometimes approaching a factor of 10 lower than derived from laboratory studies using synthetic aerosol and strongly dependent on aerosol composition, especially the water / organic content. Other, less direct measurements of N_2O_5 reactivity on ambient particles (discussed in detail later), indicate high variability and occasional low values that are inconsistent with laboratory investigations on pure samples.

In this paper we address the need for measurements of the N_2O_5 uptake coefficient to particles in the real atmospheric environment. We present simultaneous measurements of particle composition, N_2O_5 and ClNO_2 to derive both the uptake coefficient (γ) and the efficiency (f) of ClNO_2 production in number of different air masses and discuss the limitations and validity of this and similar approaches.

2 Methods

2.1 PARADE field site

The PARADE measurement intensive (Particles And RADicals: diel observations of the impact of urban and biogenic Emissions) took place between mid-August to mid-September 2011 at the Taunus Observatory (TO), Kleiner Feldberg, Germany. The observatory is operated by the Institute for Atmospheric and Environmental Science of the Goethe University, Frankfurt. The Kleiner Feldberg (825 m above sea-level) sits within the Taunus range approximately 30 km NW of Frankfurt am Main, Germany. The site, which may be described as rural with anthropogenic impact from local industrial / population centres, is described in more detail by, Wobrock et al. (1994), Handisides (2001), Crowley et al. (2010) and Sobanski et al. (2016). It is \approx 400 km distant from the nearest (North Sea) coastline.



2.2 Instrumentation

N_2O_5 and NO_3 were measured using cavity-ring-down spectroscopy (CRDS) with instruments described by Schuster et al. (2009) and Crowley et al. (2010). NO_3 was measured directly in a cavity at ambient temperature whereas the sum $\text{N}_2\text{O}_5 + \text{NO}_3$ was measured in a separate cavity at $\sim 100^\circ\text{C}$ after thermal decomposition of N_2O_5 to NO_3 in a heated section of the inlet, also at 100°C . Following corrections for the transmission of NO_3 and N_2O_5 through the inlets, filter and cavities, the difference signal is used to calculate N_2O_5 mixing ratios.

The CRDS was situated on a platform on the roof of the TO-laboratory at a height of 10 meters from the ground. Air was drawn through a 1 m length of $\frac{1}{2}$ " (OD) Teflon (perfluoroalkoxy, PFA) tubing at 50 (std) Lmin^{-1} (SLM) and sampled (18 SLM) from the centre of the flow via $\frac{1}{4}$ " PFA tubing and a Teflon filter (exchanged hourly using an automatic filter changer) into the two cavities of the CRDS. This setup keeps inlet residence times short (~ 0.1 s) and also reduces sampling of coarse particles and droplets. The total uncertainty associated with the N_2O_5 measurements was 15 % with the largest contribution the uncertainty in the NO_3 cross section and NO_3 losses.

The non-refractory composition of particulate matter with an aerodynamic diameter less than $1\ \mu\text{m}$ (PM₁) was measured with an Aerodyne HR-ToF aerosol mass spectrometer (AMS) (Jayne et al., 2000; DeCarlo et al., 2006). The AMS was deployed on board the MPIC mobile laboratory (MoLa) (Drewnick et al., 2012) located approximately 15 meters from the measurement station with air sampled from ~ 7 m above the ground via wide bore metal tubing at a flow rate of 90 (STD) L min^{-1} . Total organic, nitrate, sulphate and chloride in non-refractory particles of $< 1\ \mu\text{m}$ diameter (NR-PM₁) are reported here. As the AMS detects marine chloride (i.e. refractory chloride) with one to two orders of magnitude lower sensitivity than non-refractory chloride (Zorn et al., 2008; Ovadnevaite et al., 2012; Schmale et al., 2013; Drewnick et al., 2015) it is reasonable to assume that the majority of the Cl^- reported by the AMS is due to NH_4Cl arising from uptake of gas-phase HCl which may have marine or anthropogenic origin. The AMS was calibrated using the measurement of standard ammonium nitrate particles of known particle size via a DMA system.

Particle size spectra were measured using an optical particle counter (OPC, Model 1.109, Grimm), an aerodynamic particle sizer (APS, Model 3321, TSI) and a fast mobility particle size spectrometer (FMPS, Model 3091, TSI). All particles were sampled at ambient RH so corrections for hygroscopic particle growth were not necessary. The uncertainties associated with parameters required for calculation of the uptake coefficient are 25 % for the nitrate measurement and 30 % for the particle diameter measurement (Wiedensohler et al., 2012). Note that an uncertainty in the particle diameter of ~ 30 % implies an uncertainty in the particle surface area of ~ 70 %. This uncertainty applies not only to this study, but also to all previous laboratory and field studies that use similar instrumentation and thus a certain cancelling of errors occurs when comparing to other datasets.

NO_2 was measured during PARADE using a variety of techniques which all showed good agreement. The data used in this analysis were obtained using the MPIC NO_2 TD-CRDS system (Thieser et al., 2016). The instrument is a two-cavity system which also measured total peroxy nitrates and total alkyl nitrates (ΣPNs and ΣANs). The atmosphere was sampled by



drawing air at ~20 SLM through a 3/8" outer-diameter PFA tube 8 m in length and sub sampling approximately 4 SLM into the cavity. The main inlet was shared by the iodide CIMS system which was used to measure ClNO₂, speciated peroxy nitrates and peroxy acids (Phillips et al., 2012; Phillips et al., 2013b).

The ClNO₂ dataset and method was described by Phillips et al. (2012) and follows prior measurements of ClNO₂ using the same technique (Osthoff et al., 2008; Thornton et al., 2010; Mielke et al., 2011). The CIMS instrument was constructed by THS Instruments, Georgia, USA and is based on the CIMS technique described by Slusher et al. (2004) and Zheng et al. (2011). No indication of the production of ClNO₂ on the inlet walls was observed. ClNO₂ was monitored at I³⁷Cl⁻ ($m/z = 163.9$, with a LOD (2σ) of 12 pptv) for the entire period, however there are clear indications (i.e. non-zero daytime concentrations) that the signal at this mass is not solely due to ClNO₂ and consequently ICINO₂⁻ ($m/z = 207.9$) was monitored with a LOD (2σ) of 3 pptv, from the 02.09.2011. The impurity at $m/z = 163.9$ contains one chlorine atom ($m/z = 161.9$ was also observed at the correct isotope ratio) and has a diel cycle that strongly resembles that of HCl, which was measured using ion-chromatography during a subsequent campaign (Phillips et al., 2013a) at the same site and time of year. The mechanism of HCl detection at $m/z = 161.9$ and $m/z = 163.9$ remains unclear as the reaction $I + HCl \rightarrow ICl + H$ is endothermic by at least 200 kJ/mol.

The instrument sensitivity to ClNO₂ was determined by the measurement of a ClNO₂ standard synthesized by passing Cl₂ over a mixture of NaNO₂ and NaCl crystals in a flow of humidified N₂. The concentration of the standard was determined using thermal-dissociation cavity ring-down absorption spectroscopy (TD-CRDS) using the instrument described by Thieser et al. (2016). A zero measurement, using a bypass with 25 cm of metal wool heated to 473 K, was made once an hour and the accuracy of the ClNO₂ measurement is 25 % (Phillips et al., 2012).

Meteorological data were obtained from the public data depository of the Hessisches Landesamt für Umwelt und Geologie (HLUG) monitoring station situated at the peak of the Kleiner Feldberg approximately 10 meters from the main sample inlet location. Data is available from <http://www.hlug.de/start/luft/luftmessnetz.html>. Back trajectories were calculated using HYSPLIT (Draxler and Rolph, 2011; Stein et al., 2015).

3 Meteorological / chemical conditions during PARADE

The time series of a selection of particle and trace-gas concentrations measured during PARADE is shown in Figs. 2 and 3 along with temperature and relative humidity.

3.1 Particle characteristics

Fig. 2 indicates that the aerosol surface area (A) available for uptake of N₂O₅ was highly variable and was predominantly (> 75 % on average) associated with particles less than 550 nm in diameter (FMPS data). Relative humidity, an environmental parameter which influences the water content of the particles and is thus expected to influence the uptake of N₂O₅ significantly, varied between 25 and 100 %. On average, the sub-micron non-refractory aerosol was (by mass) 55 % organic,



26 % sulphate with nitrate and ammonium both 9.5 %, which may be considered typical for an anthropogenically influenced, rural region. Particulate sulphate and organic content were correlated, with the particulate sulphate to organic mass ratio, a parameter that potentially impacts on the N_2O_5 uptake to particles (Bertram et al., 2009b), varying between ~ 0.1 and 2.4 with a mean value of ~ 0.6 . The campaign averaged nitrate particle mass concentration was $0.56 \mu\text{g m}^{-3}$ during the day and $0.89 \mu\text{g m}^{-3}$ during the night at this site. The larger nighttime values reflect high rates of nitrate production from N_2O_5 uptake at nighttime rather than temperature dependent partitioning of $\text{HNO}_3 / \text{NH}_3 /$ ammonium nitrate (Phillips et al., 2013a), which we discuss below.

3.2 NO_x , O_3 and N_2O_5 formation

As reported previously for this site (Crowley et al., 2010; Phillips et al., 2012; Sobanski et al., 2016) local emissions result in high variability in NO_x with NO_2 usually between ~ 0.5 and 10 ppbv but with excursions up to ~ 20 ppbv. Daytime maxima for NO were between ~ 0.5 and 2 ppb and ozone levels were between 20 and 70 ppbv. The high variability in NO_2 and O_3 result in a highly variable NO_3 production term between <0.05 and >0.5 pptv s^{-1} (Sobanski et al., 2016). The NO_3 and N_2O_5 lifetimes were also highly variable and on some nights (notably 20-21.08, 30-31.08, 31.08-01.09, 01-02.09) on which extended NO_3 (and N_2O_5) lifetimes were observed we have compelling evidence that the inlets were sampling from a relatively low altitude residual layer (Sobanski et al., 2016).

3.3 Meteorology, ClNO_2 and particulate nitrate

The PARADE measurement period began on 15.08 and ran until 17.09. As described by Phillips et al. (Phillips et al., 2012), this period may be separated into three meteorologically distinct parts.

Period 1 (17.08 – 26.08): The period beginning on the 17.08 and ending on 26.08 was changeable with calculated air mass 24 hour back-trajectories suggesting the air was largely continental in origin, arriving from the west-to-south wind sector. The exceptions to the continental origin occurred up to sunrise on the 17.08 and on the 19-20.08 with air arriving at the site with relatively high humidity from the NW with 48-hour back-trajectories indicating an approximate UK/English channel origin. On the nights of the 17-18.08 and 19-20.08 concurrent increases in ClNO_2 and NR PM1 NO_3^- were observed. There is little indication of nocturnal production of ClNO_2 on the remaining nights of this initial period of the measurements. In fact, on the night of the 20-21.08 high concentrations of N_2O_5 were measured, $[\text{N}_2\text{O}_5]_{\text{max}} \approx 800$ pptv, during a period of low RH of 25 % (lowest observed during the measurement period). No concurrent increase in the concentration of submicron particulate nitrate or ClNO_2 was measured.

Period 2 (26.08 – 29.08): On the evening of 26.08 a front passed across the measurement site associated with heavy cloud cover and rain. Before the passage of the front, large, variable concentrations of NO_x were measured coinciding with an increase in PM1 particle mass. The rain was sustained following the passing of the front and a RH of 100% was measured into the morning of 28.08. After the front, concentrations of fine particles, as measured by the FMPS and AMS, and NO_x



were relatively low and remained suppressed until 29.08, with the exception of NR PM1 Cl⁻ which peaked on the nights of 27-28.08 and 28-29.08 possibly due to the marine influence of the post-frontal air and the increased supply of chloride from marine particles. Following 29.08 large concentrations of ClNO₂ were observed on a series of nights up to and including 3.09. Back-trajectories calculated with the HYSPLIT model suggest that the westerly flow following the front gradually weakened and a surface high pressure set in leading to high concentrations of NO_x and O₃ probably due to the influence of the near conurbation of Frankfurt-Wiesbaden-Mainz.

Period 3 (29.08- 09.09): The remainder of the measurement period was influenced by a mainly westerly flow and the observatory was frequently shrouded with cloud. Nocturnal production of modest concentrations of ClNO₂ occurred between 6.09 and 9.09 without a concomitant increase in NR PM1 NO₃⁻. Each of these overnight periods was impacted by cloud at the measurement site with the exception of the 6.09 where the RH was nevertheless above 90%. Measurements of particle composition during PARADE were only available at the submicron size; it may be possible that during misty and cloudy conditions the nitrate formation occurred on surfaces which were not measured, i.e. larger mist or fog droplets or, in some cases nitrate formed was scavenged within the clouds. Photolysis frequencies were attenuated by the cloud cover and rain at the site during this period, allowing measurable quantities of ClNO₂ to persist well past noon, see for example 8.09 to 9.09. N₂O₅ was not measured beyond 9.9; however, the little data available during this period suggest that relative productivity with respect to ClNO₂ was high.

4 Estimation of f and γ for N₂O₅ uptake to ambient aerosol

Apart from the measurement of the loss rate of synthetic N₂O₅ samples to ambient aerosol as carried out by Bertram and co-workers (see section 1), there are different methods by which ambient measurements can be analysed to derive an uptake coefficient (γ) for N₂O₅ interaction with atmospheric particles, which depend on the data available. These include 1) the analysis of product formation (gas and/or particle phase) resulting from the uptake of N₂O₅ (Mielke et al., 2013), 2) analysis of the steady-state N₂O₅ (or NO₃) lifetime (Brown et al., 2006; Brown et al., 2009; Brown et al., 2016a) and 3) box modelling of N₂O₅ chemistry with various observational constraints (Wagner et al., 2013). All three methods have their own strengths and weaknesses and all contain assumptions that are evidently not applicable in all cases.

We first discuss the methods which we have applied (1 and 2) to derive f and γ during the PARADE measurement campaign and compare the results to similar analyses in the literature.

4.1 Method 1: Using product formation rates to derive f and γ

For the PARADE campaign, periods of data were identified where clear correlations between particulate nitrate NO₃ and ClNO₂ were observed, as illustrated, for example, in Fig. 4. On these nights, there is no particulate nitrate formation without



concurrent ClNO₂ formation and the covariance of NO₃⁻ and ClNO₂ is taken as evidence that, to a good approximation, both ClNO₂ and particulate nitrate are produced only by the uptake of N₂O₅.

From Eq.1 and the definition of the branching ratio (R5), the rate of production of ClNO₂ (*p*ClNO₂) from the reaction of N₂O₅ with aerosol (surface area *A*) can be written as:

$$5 \quad p\text{ClNO}_2 = \frac{d[\text{ClNO}_2]}{dt} = f(0.25\gamma\bar{c}A[\text{N}_2\text{O}_5]) \quad (3)$$

The rate of production of particulate nitrate (*p*NO₃⁻) is:

$$p\text{NO}_3^- = \frac{d[\text{NO}_3^-]}{dt} = 2(1-f)(0.25\gamma\bar{c}A[\text{N}_2\text{O}_5]) + f(0.25\gamma\bar{c}A[\text{N}_2\text{O}_5]) \quad (4)$$

Combining and rearranging Eq.3 and Eq.4 we get:

$$f = 2 \left(\frac{p\text{NO}_3^-}{p\text{ClNO}_2} + 1 \right)^{-1} \quad (5)$$

10 and

$$\gamma = \frac{2(p\text{ClNO}_2 + p\text{NO}_3^-)}{\bar{c}A[\text{N}_2\text{O}_5]} \quad (6)$$

Note that the fractional formation of ClNO₂ (*f*) is the amount of ClNO₂ formed per N₂O₅ taken up to a particle and should not be confused with the yields of ClNO₂ per NO₃⁻ formed that have occasionally been reported (Osthoff et al., 2008; Mielke et al., 2013) and which represent lower limits to *f* as NO₃⁻ is not necessarily converted stoichiometrically to N₂O₅ in the atmosphere.

Eq.5 illustrates that measurement of the production rates of ClNO₂ and particulate nitrate is sufficient to derive the fractional yield of ClNO₂ (*f*) and that, by also measuring the concentrations of N₂O₅ and aerosol surface area (*A*), we can derive the uptake coefficient, γ (Eq.6). Note that without measurement of both particulate nitrate and ClNO₂ production, it is not possible to derive both the yield of ClNO₂ and the uptake coefficient. By analysing the ClNO₂ product alone, Eq. 3 can be used to derive a composite term ($\gamma \times f$) as described by Mielke et al. (2013).

The analysis assumes that, during the period over which data are analysed, the relevant properties of the air mass are conserved and the losses of either measured species are not significant. It also assumes that the efficiency of N₂O₅ uptake and ClNO₂ / NO₃⁻ production is independent of particle size. Later we examine the effect of considering uptake to coarse and fine aerosol particles separately as previously done by Mielke et al. (2013). Two further assumptions are 1) that measurement of *p*NO₃⁻ accounts for the total production of nitrate by R3, i.e. that NO₃⁻ formed from uptake of N₂O₅ does not significantly degas from the particle as HNO₃ and 2) that the formation of particulate nitrate via the net uptake of HNO₃ to aerosol as the temperature drops during the night is insignificant compared to that formed in N₂O₅ uptake. Note that the degassing of NO₃⁻ as HNO₃ will result in an underestimation of γ , and overestimation of *f*, whereas the net uptake of HNO₃ (forming particulate nitrate) will have the opposite effect.

30 Regarding assumption 1) we note that the increase of particle nitrate during night is accompanied by an equivalent increase in ammonium (see Fig. 4) as gas-phase ammonia repartitions to form ammonium nitrate and buffers the release of HNO₃. For a given ammonia concentration, as the temperatures during nighttime fall, the partitioning increasingly favours the



retention of particulate nitrate over release of HNO_3 . During a subsequent campaign at this site, we showed that the nighttime formation of particulate nitrate is dominated by N_2O_5 uptake (Phillips et al., 2013a). We also showed that, once corrected for the contribution of N_2O_5 , the measurements of gas-phase HNO_3 did not reveal nighttime increases coincident with those of particle nitrate, suggesting that N_2O_5 uptake is not an important source of HNO_3 at this site. This is likely to be related to the cool nights at altitude of > 800 m and the abundance of ammonia in this mixed rural / industrialised region of Germany. Additionally, the strong temperature dependence of partitioning between ammonia, HNO_3 and ammonium nitrate would result in a bias to high values of γ at low temperatures which prevent release of nitrate from the particles. The uptake coefficients obtained in this study showed no significant dependence on temperature (see later).

Regarding assumption 2) we note that, the strong correlation observed between ClNO_2 and NO_3^- concentrations is a useful indicator that the source of particulate nitrate is dominated by N_2O_5 uptake and not HNO_3 uptake as N_2O_5 and HNO_3 was completely different diel profiles and atmospheric lifetimes.

Three different types of analysis were used to derive uptake coefficients and the fractional formation of ClNO_2 . In the simplest method (1a) to derive f we use longer time periods (several hours or the whole night) where plots of $[\text{ClNO}_2]$ versus $[\text{NO}_3^-]$ are approximately linear and values of f are estimated from a linear fit of the data as exemplified in Fig. 5. On some nights (four in total) a linear relation is observed (Fig. 5a) whereas air mass changes can result in two distinct slopes (Fig. 5b, two nights in total) or such variability that analysis over a prolonged period is impossible (Fig. 5c, all other nights). For the example displayed in Fig. 5a a slope of 56.3 ± 2.2 pptv $\mu\text{g}^{-1} \text{m}^3$ (2σ statistical error) results in a value of $f = 0.24 \pm 0.07$, where the error in f includes overall uncertainty in the measurement of ClNO_2 and NO_3^- . Using this method, a total of 8 values of f were obtained during the campaign.

In order to derive γ , absolute production rates of NO_3^- and ClNO_2 are required and in method 1b shorter periods of data, usually between 1 and 3 hours, were chosen by inspection of the time series such that NO_3^- and ClNO_2 concentrations both increase during a period of relatively constant composition and environmental variables such as temperature and RH. It is more likely that the assumptions hold during the shorter time periods chosen for 1b than the longer sections of data used in 1a. In this case, values of $p\text{ClNO}_2$ and $p\text{NO}_3^-$ and average values of A and $[\text{N}_2\text{O}_5]$ are calculated for the same period and inserted into Eqs. 5 and 6 to derive f and γ . An example of this analysis is shown in Fig. 6, which indicates (grey shaded area) time periods in which ClNO_2 and NO_3^- concentrations increased whilst other parameters (e.g. N_2O_5 , A and RH) were relatively constant. In this example between $\sim 21:00$ and $23:20$ on the night 29.08-30.08, values of $p\text{ClNO}_2$ (14.6 ± 3.0 pptv/hr), $p\text{NO}_3^-$ (21.0 ± 5.1 pptv/hr), \bar{A} (7.6 ± 2.2) $\times 10^{-7}$ $\text{cm}^2 \text{cm}^{-3}$, $[\text{N}_2\text{O}_5]$ (153 ± 43 pptv) were obtained. The uncertainties quoted were derived by propagating statistical uncertainty (e.g. in the production rate of ClNO_2 from fitting to the data) and absolute uncertainty in the measurements of the concentrations of ClNO_2 , N_2O_5 , particulate nitrate and aerosol surface area as listed in section 2.2. Generally, the absolute error in the concentration measurements dominates the overall uncertainty for each parameter, the major exception being N_2O_5 which could be sufficiently variable over the averaging period for this to also contribute significantly. Inserting this set of parameters into Eqs. 5 and 6 results in $f = (0.82 \pm 0.26)$ and $\gamma = (7.3 \pm 3.1) \times$



10^{-3} at $\text{RH} = 77 \pm 2 \%$ and a temperature of $8 \pm 1^\circ\text{C}$. Using this method a total of 12 values of γ were obtained during the campaign.

A more rigorous analysis, which avoids use of average values of A and $[\text{N}_2\text{O}_5]$ was also carried out. In this case $p\text{NO}_3^-$ and $p\text{ClNO}_2$ are calculated from 10 min averaged datasets using the right-hand sides of Eq. 3 and 4 respectively, taking measured values of N_2O_5 and A at each time step and using an initial estimate for f and γ . The predicted concentrations of ClNO_2 and NO_3^- were then calculated for each time step by integration over the analysis time period. Values of f and γ were then varied and the integration repeated until good agreement between observed and calculated $[\text{ClNO}_2]$ and $[\text{NO}_3^-]$ was obtained. An example of this type of analysis (for the night 19.08-20.08) is displayed in Fig. 7. The red lines are the results of the analysis in which the total particle surface area was used. Later, we discuss the effects of separately calculating the formation of NO_3^- and ClNO_2 from the fine and coarse aerosol fractions. Unlike method 1b, a limitation of this procedure is that the analysis is limited to the first period of the night. Similar to method 1b it cannot predict negative changes in concentrations of ClNO_2 or NO_3^- which are a result of changes in air-mass origin / age.

The values of f and γ derived from method 1a-1c are plotted versus RH in Fig. 8. The data are colour coded according to the method used.

The lower panel of Fig. 8 reveals high variability in the values of f obtained which range from 0.035 ± 0.027 to 1.38 ± 0.60 . The errors on each individual determination vary from ~ 35 to $\sim 100 \%$, which is the result of scatter in the data as well as uncertainty associated with measurement of ClNO_2 and particle- NO_3^- and can result in non-physical values larger than unity. A further possible reason for values of f that exceed unity is the degassing of NO_3^- as HNO_3 from the particles, though as described above, this effect is expected to be small.

The average value of γ obtained for the campaign considering both methods 1b and 1c was 0.027 with a standard deviation of 0.03. Minimum and maximum values were 0.004 and 0.11, respectively. The large standard deviation reflects the great variability in the values of γ obtained, with up to a factor of 10 difference in γ at the same relative humidity. Considering the large variability, the values of γ obtained using methods 1b and 1c over the same time period are in reasonable agreement with an average ratio of 1.7. We compare the values of γ obtained during PARADE with other ambient datasets below.

25 4.2 Method 2. NO_3^- steady state lifetime analysis

Atmospheric N_2O_5 is formed in a series of oxidation steps initiated mainly by the reaction of NO_2 with O_3 (R7).



30 Ambient concentrations of NO_3^- and N_2O_5 are thus coupled via the gas-phase, thermochemical equilibrium that exists due to R8 and R9, so that the relative amounts of NO_3^- and N_2O_5 are determined by temperature and NO_2 levels.



The so called “steady state” determination method for γ is based on the assumption that, after a certain period of time following sunset (often on the order of hours) the direct and indirect losses of NO_3 and N_2O_5 balance their production. The steady-state lifetimes can then be calculated from observations of the NO_3 and N_2O_5 concentrations and the production term $k_7[\text{NO}_2][\text{O}_3]$, where k_7 is the rate constant for reaction R7. This method was first used by Platt and co-workers to assess the heterogeneous loss of N_2O_5 (Platt and Heintz, 1994; Platt and Janssen, 1995; Heintz et al., 1996), and has been extended by Brown and co-workers to derive γ in regions distant from NO_x sources such as the marine environment (Aldener et al., 2006) and aloft (Brown et al., 2006; Brown et al., 2009) and most recently for the continental boundary layer (Brown et al., 2016a). $[\text{NO}_2]$ and $[\text{O}_3]$ measurements are required to calculate the rate of NO_3 production, which is generally assumed to be via R7 only, though a contribution of NO_2 oxidation by stabilised Criegee intermediates has recently been hypothesised to represent a potential bias to this calculation (Sobanski et al., 2016). The steady-state analysis does not require any information about products formed by N_2O_5 heterogeneous reactions.

The inverse steady-state lifetimes of NO_3 (τ_{NO_3}) and N_2O_5 ($\tau_{\text{N}_2\text{O}_5}$) are given by expressions Eq.7 and Eq.8, respectively:

$$(\tau_{\text{NO}_3})^{-1} \approx \gamma(0.25\bar{c}AK_{\text{eq}}[\text{NO}_2]) + k_g \quad (7)$$

$$(\tau_{\text{N}_2\text{O}_5})^{-1} \approx k_g(K_{\text{eq}}[\text{NO}_2])^{-1} + 0.25\bar{c}A\gamma \quad (8)$$

Where K_{eq} is the temperature dependent equilibrium constant describing the relative concentrations of NO_2 , NO_3 and N_2O_5 (R8 and R9), $[\text{NO}_2]$ is the concentration of NO_2 , and k_g is the pseudo first-order loss constant for NO_3 loss in gas-phase reactions (e.g. with NO or with hydrocarbons). A plot of $(\tau_{\text{NO}_3})^{-1}$ or $(\tau_{\text{N}_2\text{O}_5})^{-1}K_{\text{eq}}[\text{NO}_2]$ against $0.25\bar{c}AK_{\text{eq}}[\text{NO}_2]$ should be a straight line with γ as slope and k_g as intercept. This method thus relies on the fact that the relative concentrations of N_2O_5 and NO_3 vary with $K_{\text{eq}}[\text{NO}_2]$ and thus the contribution of their individual losses to the overall lifetime of both NO_3 and N_2O_5 also varies with $[\text{NO}_2]$ once changes in temperature (and thus K_{eq}) are accounted for. The method therefore assumes that, for a given analysis period in which NO_2 is changing sufficiently to change the relative loss rates of NO_3 and N_2O_5 , both γ and k_g are constant (i.e. do not depend on $[\text{NO}_2]$). This will not always be the case and we have often observed that the relation between $(\tau_{\text{NO}_3})^{-1}$ and $K_{\text{eq}}[\text{NO}_2]$ is non-linear. In environments where NO_3 losses are dominated by gas-phase reactions of NO_3 , the uncertainty associated with the derivation of γ via a steady-state analysis is clearly greatly enhanced.

Figure 9 illustrates some of these issues for data obtained over a period of several hours on the nights 1-2.09, 2-3.09 and 5-6.09. The first two nights (1-2.09 and 2-3.09) had long NO_3 lifetimes and were selected for analysis as long NO_3 lifetimes represent periods with low rates of NO_3 loss by gas-phase processes (i.e. k_g is small). We have previously shown (Sobanski et al., 2016) that the long NO_3 -lifetimes on these nights result from sampling from a low altitude residual layer. A large variability in the nighttime NO_2 mixing ratio on these nights (~ 1 to 13 ppbv, see Fig. 9a and 9c) should result in a significant shift in the NO_2 -to- N_2O_5 ratio and thus sensitivity in the NO_3 lifetime to the uptake of N_2O_5 to aerosol. On the night 1-2nd Sept., plume like features in the NO_2 mixing ratio at $\sim 22:00$, $01:30$ and $04:00$ were accompanied by decreases in the steady-state NO_3 lifetime. The inverse NO_3 lifetime is plotted against $0.25\bar{c}AK_{\text{eq}}[\text{NO}_2]$ in Fig. 9b. Here we have selected data that



was obtained from about 2 hr after sunset to the next dawn when NO_3 started to decrease as represented by the red data points in Fig. 9a. The slope of the plot results in values of $\gamma = (9.6 \pm 0.7) \times 10^{-3}$ and $k_g = (2.0 \pm 0.6) \times 10^{-4} \text{ s}^{-1}$, where the errors are statistical only. Over the period analysed, the relative humidity (blue line in Fig. 9a) was $60 \pm 2 \%$.

On the night 2-3.09 at about 19:30 UTC (≈ 1 hour after sunset), the NO_3 lifetime increases gradually to a value of ~ 900 s until 21:30 as NO_2 decreases from ~ 7 to 3 ppbv. A rapid increase in NO_2 at 21:30 is then accompanied by a much shorter NO_3 lifetime. After $\sim 22:00$, NO_2 slowly decreases and the NO_3 lifetime recovers to about 500 s. Thus, also in this night there is a clear anti-correlation between NO_2 and the NO_3 lifetime. Figure 9d plots the inverse of the NO_3 lifetime versus $0.25\bar{c}AK_{eq}[\text{NO}_2]$ with the data point colour coded according to relative humidity. The first period of the night (orange, yellow and green data-points) are best described (black line) by an uptake coefficient of $(2.7 \pm 0.2) \times 10^{-2}$ and $k_g = (1.0 \pm 0.4) \times 10^{-4} \text{ s}^{-1}$. The second period (starting ~ 5 hours after sunset) is best characterised by larger values of $\gamma = (4.5 \pm 0.3) \times 10^{-2}$ and $k_g = (1.5 \pm 0.1) \times 10^{-3} \text{ s}^{-1}$ (errors are statistical only). We note that prior to the peak in NO_2 at 21:30 the relative humidity of the air was stable at $\approx 70 \pm 3 \%$ whereas after 22:00 it remained at $80 \pm 3 \%$. A shift in air mass to one with larger water vapour content could help explain the larger values of γ obtained in the 2nd period of this night and also may also be the reason for the larger gas-phase losses of NO_3 if the more humidified air mass is more impacted by boundary-layer emissions. On two other nights when the NO_3 lifetimes were long (30-31.08 and 31.08-1.09) there was very little variation in NO_2 so that the steady-state lifetime of NO_3 was insensitive to the uptake coefficient.

In Fig. 9e and 9f we present data that were obtained with sufficient variation in NO_2 , but with relatively short NO_3 lifetimes. As in the datasets discussed above, NO_3 lifetimes are anti-correlated with NO_2 . However, the plot of the inverse of the NO_3 lifetime versus $0.25\bar{c}AK_{eq}[\text{NO}_2]$ in Fig. 9f has a very large slope, resulting in $\gamma \approx 0.3$ and also a negative intercept. Unrealistically large values of γ and negative intercepts can be caused by a covariance between NO_2 concentrations and gas-phase losses of NO_3 , which makes this type of analysis unviable. Similar observations of apparently negative gas-phase reactivity indicating breakdown of the steady-state method have previously been made (Morgan et al., 2015; Brown et al., 2016a). In addition to co-variance between NO_2 and other trace-gases that are reactive to NO_3 , negative intercepts can also result from analysis of time periods in which steady state was not acquired. During the PARADE campaign there were only three clear examples of the expected relationship between the NO_3 lifetime and NO_2 mixing ratios as defined by Eq. 7 or 8. This indicates that, over periods of a few hours, the changes in the NO_3 lifetime observed cannot be attributed solely to different rates of heterogeneous processing but that other air mass characteristics (e.g. influencing the gas-phase losses of NO_3) are also variable, which may be expected in a region that is impacted by fresh emissions of reactive gases. We also note that the average aerosol surface area during the PARADE campaign was about $70 \mu\text{m}^2 \text{ cm}^{-3}$, which is low compared to the ones reported e.g. by Aldener et al. (2006), Brown et al. (2006) and Brown et al. (2009) which range from 200 to $600 \mu\text{m}^2 \text{ cm}^{-3}$ and where plots of inverse NO_3 lifetime versus $0.25\bar{c}AK_{eq}[\text{NO}_2]$ resulted in straight lines and returned reasonable values of γ . We conclude that the steady-state approach to derive γ is best suited to air masses with high aerosol loading but remote from fresh emissions and only worked on a few occasions during the PARADE campaign when NO_3 was sampled



from the residual layer. The overall uncertainty associated with the determination of γ via the steady-state method is derived from uncertainty in the mixing ratios of O_3 (5 %) and NO_3 (25 %), the aerosol surface area (~70 %), the rate constant for reaction between O_3 and NO_2 (15 % at 298 K (Atkinson et al., 2004)) and the equilibrium constant, k_{eq} (20 % at 298 K, (Burkholder et al., 2016)) and when propagated in quadrature is equal to ± 75 %. The three values of γ obtained by the steady-state method (with total uncertainty) are also plotted in Fig. 8.

4.3 Method 3. Iterative box model of N_2O_5 formation and loss.

A further method to derive N_2O_5 uptake coefficients is to use a box model of NO_3 / N_2O_5 formation and loss (both gas-phase and heterogeneous) with γ treated as a variable to match observed N_2O_5 concentrations. Such an analysis was conducted by Wagner et al. (2013) who used an iterative box model of N_2O_5 chemistry to estimate the uptake coefficient that would be needed to account from the N_2O_5 concentration measured at the observation point at time “t” after sunset. The calculations were constrained by on-site measurements of N_2O_5 precursor gases (NO_2 and O_3) which were used to estimate the original NO_2 and O_3 concentrations (i.e. at sunset) for the same air mass, and measurements of the aerosol surface area. As the concentration of N_2O_5 at any time after sunset depends not only on its heterogeneous loss processes but also on gas-phase losses of NO_3 , the total NO_3 reactivity must also be known. As the authors point out, this can be a poorly constrained parameter as potentially not all reacting VOCs are measured and the reactions of NO_3 with other trace gases including NO and RO_2 during transport to the measurement site cannot be accurately assessed. This method is therefore expected to be least accurate when gas-phase losses of NO_3 are a substantial fraction of the overall loss rate of NO_3 and N_2O_5 . During much of the PARADE campaign (and also for previous measurements at this site (Crowley et al., 2010) NO_3 lifetimes were dominated by gas-phase losses and on the few occasions when NO_3 was long-lived this was due to sampling from a residual layer that was decoupled from boundary layer emissions. As we have recently shown (Sobanski et al., 2016) when sampling from the residual layer, measured VOC mixing ratios at the site are incompatible with the high NO_3 and N_2O_5 levels observed and a box-model approach that requires a constraint on gas-phase losses of NO_3 cannot provide reliable results. For this reason, we have not attempted to analyse our data in a similar fashion to Wagner et al. (Wagner et al., 2013).

4.4 Comparison of γ and f with literature values derived from ambient datasets

Previous determinations of γ from ambient datasets are summarised along with the results of this work in Table 1. The reported values of γ vary over more than two-orders of magnitude, between 0.0005 to ~0.1. The directly measured loss rates of N_2O_5 to ambient aerosol (Bertram et al., 2009b; Riedel et al., 2012b) convert to variable values of γ (~0.005 – 0.035) in Seattle (RH ~ 74 ± 13 %) that are comparable to those reported in this work. The largest values of γ were obtained at low organic to sulphate ratios, whereas their largest values at a coastal location impacted by chloride containing aerosol were



obtained when the molar $\text{H}_2\text{O} / \text{NO}_3^-$ ratio was high. In drier conditions in Boulder ($\text{RH} < 30\%$) γ was much lower (< 0.005) and independent of the organic to sulphate ratio.

Apart from the present study, the steady state method using NO_3 lifetimes has been successfully used for analysis of aircraft data (Brown et al., 2006; Brown et al., 2009; Morgan et al., 2015), ship data (Aldener et al., 2006) and also in ground-based studies in which residual layer air was sampled (Brown et al., 2016a). In their airborne studies over the NE U.S., Brown et al. (2006) report regional differences in γ which they assign to changes in the sulphate content or sulphate-to-organic ratio of the particles. In a further airborne study over Texas, Brown et al. (2009) analysed several residual layer plumes to derive ~30 values of γ between 5×10^{-4} and 0.006 during three flights. They found no dependence of γ on RH or aerosol composition, which was typically ammonium sulphate with an organic fraction of $> 50\%$. In contrast, airborne measurements over Europe (Morgan et al., 2015) suggest that γ is larger and dependent on the particulate nitrate content. Measurements of NO_3 and N_2O_5 on a polluted mountain site in Hong Kong (Brown et al., 2016b), enabled 10 derivations of γ which varied between 0.004 and 0.029. Measurements of ClNO_2 at the same location indicate that N_2O_5 uptake was, at least partially, to chloride containing particles. Values of (0.03 ± 0.02) for γ obtained over the ocean (Aldener et al., 2006) are, on average, at the higher range of ambient measurements, potentially reflecting the role of particulate chloride.

In their analysis of N_2O_5 and ClNO_2 datasets obtained during CalNex-LA at a coastal location, Mielke et al. (2013) did not derive separate values of f and γ but report a composite term, γf . For submicron particles their values of γf vary ~ two orders of magnitude during a single night with campaign minimum values close to 10^{-4} and maximum values of 0.05. In comparison, γf values from the present study vary between ~0.001 and 0.09. Mielke et al. (2013) report a campaign average value of $\gamma f = 0.0084$, which is similar to the average value of $\gamma f = 0.014$ obtained in the present study. Similar values may indeed be expected despite the different locations (semi-rural mountain site in PARADE versus polluted coastal in CalNex-LA) as most of the γ values reported here were derived using ClNO_2 observations, i.e. for particles with chloride content as would certainly be expected at the coastal location.

Wagner et al. (2013) used a box-model and observations of N_2O_5 to derive highly scattered values of γ with two peaks in the frequency distribution at 0.015 and 0.04, lower values of γ being associated with higher particulate nitrate content. As discussed above, this method to derive γ requires knowledge of the NO_3 lifetime with respect to gas-phase losses, and the authors suggest that uncertainty in this parameter may result in values of γ that are too high. The source of the very high scatter in γ derived by this method is unlikely to be related to aerosol composition, but probably results from the large variability of N_2O_5 (and NO_3) frequently observed in ground based measurements, which is the result of sampling advected air masses with strong vertical gradients in NO_3 in a poorly mixed boundary layer at night.

In summary, the derivation of γ from ambient datasets reveals great variability in the values obtained, with occasional evidence for a suppressing role of particulate organics and nitrate, which is consistent with laboratory observations. Below we examine our dataset for evidence of such effects and compare the γ values obtained to parameterisations.



The large spread in f derived in the present study is consistent with previous analyses of field measurements in which values between < 0.01 and > 0.9 have been reported (Riedel et al., 2012b; Wagner et al., 2013; Young et al., 2013). The large spread in f is to be expected as this parameter is controlled largely by particulate chloride content, though, as mentioned already, some dissolved organic species may compete with Cl^- for reaction with NO_2^+ and thus reduce the ClNO_2 yield for a given chloride content. We thus expect f to be largest in polluted coastal regions and lowest (or zero) in continental regions with no marine influence or anthropogenic chlorine emissions.

4.5 Factors affecting the ClNO_2 formation efficiency, f

From the PARADE dataset, we derived values of f that ranged from 0.029 ± 0.027 to 1.38 ± 0.60 , with a mean value of 0.49. To put these numbers into context, we rearrange Eq. 2 and use a k_2 / k_4 ratio of 2.2×10^{-3} to calculate which particulate chloride concentrations would be required to explain the values of f observed (Eq. 7).

$$[\text{Cl}^-] = 2.2 \times 10^{-3} [\text{H}_2\text{O}] / (f^{-1} - 1) \quad (9)$$

Assuming that the fate of H_2NO_3^+ is reaction with Cl^- and H_2O and taking a water concentration of ~ 40 M, we obtain $f \sim 0.035$ (at the lower limit of values obtained) when N_2O_5 is taken up to particles with $[\text{Cl}^-] = 3.4 \times 10^{-3}$ M. For $[\text{Cl}^-] = 0.85$ M, $f \sim 0.9$. Chloride concentrations exceeding 1 M may be found in sea-salt particles, which have $[\text{Cl}^-]$ close to 5 M when freshly generated (Sander and Crutzen, 1996). While the sodium content of the particles is conserved during transport, there are mechanisms by which chloride may be lost to the gas-phase including acid-displacement of HCl following uptake of HNO_3 or H_2SO_4 and also reactive losses via uptake of N_2O_5 and other inorganic trace gases and radicals (Keene et al., 1990; Keene et al., 1999; von Glasow et al., 2001). The presence of chloride in non-marine aerosol is related to the uptake of HCl to particles and formation of ammonium chloride, where the HCl may be either of anthropogenic or marine origin. As discussed by Phillips et al. (2012), there is indirect evidence for the presence of aged, sea-salt aerosol at the Kleiner Feldberg site during the PARADE campaign, which is largely based on the formation of ClNO_2 on air mass origin as derived from back trajectories and wind-direction measurements. During a campaign (INUIT, (Phillips et al., 2013a)) at the same site and time of year in 2012, we measured the inorganic particle composition via ion-chromatography and showed that the site is regularly impacted by marine aerosol during periods of strong north-westerly winds. During INUIT, high concentrations of particulate inorganic chloride (up to $\sim 2 \mu\text{g m}^{-3}$) were strongly correlated with sodium (up to $\sim 1.6 \mu\text{g m}^{-3}$) with a chloride / sodium molar mixing ratio that was significantly lower than unity, indicating loss of chloride from sea-salt during transport from the coastal source regions (~ 400 km distant). For the purpose of illustration, 1000 deliquesced particles cm^{-3} with an average diameter of 300 nm and a 0.85 M chloride concentration (as derived from our measurement of f) would result in $\sim 0.4 \mu\text{g m}^{-3} [\text{Cl}^-]$ suggesting that the high values of f obtained are compatible with chloride particle concentrations at this site, albeit not measured simultaneously. A high efficiency of ClNO_2 formation f (> 0.5) was measured in the period between 29.08 and 03.09 for which two-day back trajectories indicate that the site was influenced by marine air near the UK (Phillips et al., 2012). Over this period, NR PM1 Cl^- increases on a number of occasions in concert with nocturnal ClNO_2 production.



The lower values of f derived (e.g. 0.035) are associated with mM [Cl⁻] particle concentrations and are likely the result of N₂O₅ uptake to non-marine particles to which HCl has partitioned during transport.

4.6 Factors affecting the N₂O₅ uptake efficiency, γ

The average value of γ derived from the PARADE dataset from methods 1b, 1c and steady-state analysis is 0.028 with a large standard deviation (0.029) reflecting the high variability in this parameter. No significant dependence on temperature was observed (Fig. 10).

The values of γ derived from the PARADE dataset are compared in Fig. 8 to those predicted from different parameterisations available in the literature, all derived from laboratory studies. The parameterisation listed by the IUPAC panel considers, via the resistor model, the dependence of γ on the bulk accommodation coefficient (α_b), the solubility (H) and diffusivity (D_1) of N₂O₅ and the rate coefficient, $k_{\text{H}_2\text{O}}$, for its hydrolysis (Eq. 8).

$$\gamma_{\text{IUPAC}} = \left\{ \frac{1}{\alpha_b} + \frac{\bar{c}}{4HRT(D_1k_{\text{H}_2\text{O}}[\text{H}_2\text{O}])^{0.5}} \right\}^{-1} \quad (10)$$

IUPAC preferred values are listed (for ammonium sulphate) as $\alpha_b = 0.03$, $k_{\text{H}_2\text{O}} = 1.0 \times 10^5 \text{ M}^{-1}\text{s}^{-1}$, $D_1 = 1 \times 10^{-5} \text{ cm}^2 \text{ s}^{-1}$ and $H = 2 \text{ M atm}^{-1}$.

The more complex parameterisation of Bertram and Thornton (2009) considers the concentrations of particulate nitrate, [NO₃⁻], chloride, [Cl⁻], and water [H₂O] (Eq. 9).

$$\gamma_{\text{BT}} = Ak_1 \left(1 - \frac{1}{\left(\frac{k_2[\text{H}_2\text{O}(l)]}{k_{-1}[\text{NO}_3^-]} \right) + 1 + \left(\frac{k_4[\text{Cl}^-]}{k_{-1}[\text{NO}_3^-]} \right)} \right) \quad (11)$$

where $A = 3.2 \times 10^{-8} \text{ s}$, $k_1 = 1.15 \times 10^6 - 1.15 \times 10^6 \exp(-0.13[\text{H}_2\text{O}(l)]) \text{ s}^{-1}$, $k_2/k_{-1} = 0.06$ and $k_4/k_{-1} = 29$ as empirically derived and listed in Bertram and Thornton (2009). We refer to the calculated values of γ as γ_{IUPAC} and γ_{BT} , respectively.

Other parameterisations have been developed which also deal with the effects of particle organics e.g. as a particle coating (Anttila et al., 2006) and have been reviewed by Chang et al (Chang et al., 2011). We examine the effects of assuming that a hydrophobic coating covers the particles below.

For the parameterisation of Bertram and Thornton (2009) (henceforth referred to as BT), particulate nitrate, sulphate and ammonium concentrations as measured by the AMS and the relative humidity were then used as input parameters to calculate the particle liquid water content [H₂O(l)] using the E-AIM model (<http://www.aim.env.uea.ac.uk/aim/model3/model3a.php>) (Clegg et al., 1998; Wexler and Clegg, 2002). The particulate chloride content was calculated using values of f derived as described above and in Eq. 2. In those cases where f exceeded unity, it was set to 1.0 for calculation of particulate chloride content (Eq. 7) to avoid generation of negative concentrations. For the γ values obtained using the steady-state method, the chloride content is unknown and was set to zero. Given insufficient information on the identity of



the condensed organics or particle hygroscopicity, the influence of particle organic content on the particle water was not considered.

γ_{BT} (red stars) and γ_{IUPAC} (blue stars) were computed for each observational data point in Fig. 8. On average, the predictions and the measurements are in reasonable agreement. The variability in the γ_{BT} predicted values (for a given RH) stems from different particulate chloride content. As an example of the sensitivity to chloride, the low value predicted for 70 % RH in Fig. 8 was obtained for a calculation of uptake to particles which had a high nitrate content and, as the chloride content was not known for this particular period (γ was derived using the steady-state method), it was set to zero. Adding ~ 0.02 M chloride to the calculation (equivalent to $f \sim 0.2$) would increase the calculated γ by ~ 40 % (from 0.007 to 0.01). The BT-predicted γ , averaged over the same time periods as the measured values, is (0.028 ± 0.008) . This is entirely consistent with the campaign averaged value derived from methods 1b and 1c of (0.025 ± 0.027) , as described above. However, the BT-parameterisation lies at the upper range of the measurements for relative humidities between 65 and 75 % and at the lower end for the highest relative humidities encountered. As the particles during PARADE have significant organic content (see Fig. 2) this may reflect the dependence of the organic suppression of γ on relative humidity as reported by (Gaston et al., 2014).

As already stated, the BT-parameterisation accounts for a reduction in γ due to particulate nitrate but does not take the effect of particle organic content into account. γ values derived from observation of N_2O_5 uptake to ambient particles in Seattle (Bertram et al., 2009b) have revealed that the uptake coefficient can be dependent on the sulphur-to-organic ratio, with reduced uptake at low ratios. In Fig 10. we plot the measured uptake coefficients against the molar H_2O / nitrate ratio of the particles and the organic-to-sulphate ratio. Despite significant variation in both nitrate content (~ 0.2 to $7 \mu\text{g m}^{-3}$) and the organic-to-sulphate ratio (0.3 - 1.2), no trend is seen. The PARADE γ s are nonetheless consistent with the Seattle observations reported by Bertram et al. (2009b) in which γ was close to 3×10^{-2} for organic/sulphate ratios up to 3. The large organic/sulphate ratios up to ~ 13 observed by Bertram et al. were not encountered during the PARADE campaign. As discussed by (Gaston et al., 2014), the effect of organic content in a particle will depend on the amount and oxidation state of the organics, with aged particles containing a high fraction of oxidised organics least impacting on γ , especially at high relative humidities.

The lack of a measureable dependence on particulate nitrate content may be contrasted with the interpretation of Morgan et al. (2015), who report that the uptake coefficients they derived in their steady-state analysis of airborne data ($0.03 > \gamma > 0.007$) reveal a dependence on the H_2O / nitrate ratio, as would be expected from Eq. 8, with the lower values of γ associated with high nitrate content. The larger scatter in uptake coefficients in the present study would most likely disguise this effect. In addition, the formation of ClNO_2 is an unambiguous confirmation of the presence of chloride in the particles, which will efficiently counter the reduction in γ caused by nitrate. We also note that the effect of particulate nitrate content is difficult to assess for the PARADE campaign. As we have already shown for this site (Phillips et al., 2013a) a major fraction of the



particulate nitrate is in any case the result of N_2O_5 uptake, so that efficient uptake (i.e. high values of γ) may be accompanied by large nitrate concentrations, essentially implying a time dependence to γ as the uptake proceeds (and particulate nitrate increases) or, in other words, an apparent dependence of γ on N_2O_5 levels. While such effects are reported in laboratory experiments they will be very difficult to observe in ambient datasets and we see no dependence of γ on the N_2O_5 levels.

- 5 We have also examined the possibility that the formation of $ClNO_2$ occurs only in coarse particles, i.e. with $> 0.5 \mu m$ diameter as measured by the APS. In this case, assuming the coarse particles to be sea-salt we fixed γ at 0.02. (Ammann et al., 2013). Similar to the conclusions of Mielke et al. (2013), we find that even when the $ClNO_2$ efficiency is optimised by setting and f to 1, only a fraction ($\sim 20\%$) of the $ClNO_2$ observed can be accounted for. This indicates that chloride is distributed across the particle size spectrum and, after 2-3 days of transport across polluted NW-Europe, is also in the form of gas-phase HCl in equilibrium with ammonium chloride as well as in coarse particles as NaCl.

10 The presence of an organic coating can reduce the uptake coefficient of N_2O_5 to an aqueous particle and a parameterisation of γ based on laboratory studies has been developed to account for this (Mentel et al., 1999; Anttila et al., 2006; Riemer et al., 2009). In this case, the uptake coefficient is given by:

$$\gamma_{\text{coating}} = \frac{4RT H_{\text{org}} D_{\text{org}} R_{\text{aq}}}{\bar{c} L R_{\text{part}}} \quad (12)$$

- 15 Where H_{org} and D_{org} are the solubility and diffusivity, respectively of N_2O_5 in the organic coating (of thickness L). R_{aq} and R_{part} are the radii of the aqueous core and the particle, respectively. In order to calculate the thickness of the coating, we assume that the entire organic content of the particle is hydrophobic, has a relative density of 1.27 and forms a coating on an aqueous, inorganic core (of density 1.28 as returned from the E-AIM calculation). We set $H_{\text{org}} D_{\text{org}}$ to $0.03 \times H_{\text{aq}} D_{\text{aq}}$ as derived by Anttila et al. (2006) in their laboratory studies with $H_{\text{aq}} = 5 \text{ mol L}^{-1} \text{ atm}^{-1}$ and $D_{\text{aq}} = 10^{-9} \text{ m}^2 \text{ s}^{-1}$. The net uptake to a particle with an aqueous core (containing nitrate and chloride) and an organic coating is then given by:

$$\frac{1}{\gamma_{\text{net}}} = \frac{1}{\gamma_{\text{BT}}} + \frac{1}{\gamma_{\text{coating}}} \quad (13)$$

- 25 As shown in Fig. 8, the effect of the organic coating can be substantial and values of γ_{net} obtained are significantly lower than those calculated from the BT-parameterisation and also the observations. This is not totally surprising, as a significant fraction of the organic content of an aged particle is likely to be soluble in water, which means that the assumed thickness of the organic coating is an upper limit and the effect of organic in suppressing γ is overestimated (Gaston et al., 2014). The scatter in our data and the missing information regarding the chemical nature of the organics in the particle (to get realistic values of H_{org} and D_{org}) do not allow us to further evaluate the role of organics in suppressing γ .

5 Conclusions

- We present estimates of γ using ambient measurements of gas and particle composition at the Kleiner Feldberg observatory, near Frankfurt, SW Germany, during the PARADE observational experiment in the summer of 2011. Values of γ were



derived using different methods and reveal high variability with ($0.004 < \gamma < 0.11$) and an average value of 0.028 ± 0.027 . The results are compared with different parameterisations based on laboratory data and are in reasonable agreement when we neglect the potential effect of an organic coating on the particle but account for inorganic composition and relative humidity (Bertram and Thornton, 2009). The assumption that the organic fraction of the particle is in the form of a hydrophobic coating (Anttila et al., 2006) results in predicted γ values that are inconsistent with our dataset, and is clearly inappropriate for the aerosol encountered during the PARADE campaign.

Acknowledgements

We would like to thank the staff and department of the Johann Wolfgang Goethe-University, Frankfurt am Main for logistical support and the use of the Taunus Observatory. We acknowledge the help and support of all PARADE participants and our colleagues in the Department of Atmospheric Chemistry, MPIC.

References

- Aldener, M., Brown, S. S., Stark, H., Williams, E. J., Lerner, B. M., Kuster, W. C., Goldan, P. D., Quinn, P. K., Bates, T. S., Fehsenfeld, F. C., and Ravishankara, A. R.: Reactivity and loss mechanisms of NO_3 and N_2O_5 in a polluted marine environment: Results from in situ measurements during New England Air Quality Study 2002, *J. Geophys. Res. -Atmos.*, 111, art. D23S73, doi 10.1029/2006JD007252, 2006.
- Ammann, M., Cox, R. A., Crowley, J. N., Jenkin, M. E., Mellouki, A., Rossi, M. J., Troe, J., and Wallington, T. J.: Evaluated kinetic and photochemical data for atmospheric chemistry: Volume VI – heterogeneous reactions with liquid substrates, *Atmos. Chem. Phys.*, 13, 8045-8228, doi:10.5194/acp-13-8045-2013, 2013.
- Anttila, T., Kiendler-Scharr, A., Tillmann, R., and Mentel, T. F.: On the reactive uptake of gaseous compounds by organic-coated aqueous aerosols: Theoretical analysis and application to the heterogeneous hydrolysis of N_2O_5 , *J. Phys. Chem. A*, 110, 10435-10443, 2006.
- Atkinson, R., Baulch, D. L., Cox, R. A., Crowley, J. N., Hampson, R. F., Hynes, R. G., Jenkin, M. E., Rossi, M. J., and Troe, J.: Evaluated kinetic and photochemical data for atmospheric chemistry: Volume I - gas phase reactions of Ox, HOx, NOx and SOx species, *Atmos. Chem. Phys.*, 4, 1461-1738, 2004.
- Badger, C. L., Griffiths, P. T., George, I., Abbatt, J. P. D., and Cox, R. A.: Reactive uptake of N_2O_5 by aerosol particles containing mixtures of humic acid and ammonium sulfate, *J. Phys. Chem. A*, 110, 6986-6994, 2006.
- Bannan, T. J., Booth, A. M., Bacak, A., Muller, J. B. A., Leather, K. E., Le Breton, M., Jones, B., Young, D., Coe, H., Allan, J., Visser, S., Slowik, J. G., Furger, M., Prevot, A. S. H., Lee, J., Dunmore, R. E., Hopkins, J. R., Hamilton, J. F., Lewis, A. C., Whalley, L. K., Sharp, T., Stone, D., Heard, D. E., Fleming, Z. L., Leigh, R., Shallcross, D. E., and Percival, C. J.: The first UK measurements of nitryl chloride using a chemical ionization mass spectrometer in central London in the summer of 2012, and an investigation of the role of Cl atom oxidation, *J. Geophys. Res. -Atmos.*, 120, 5638-5657, doi:10.1002/2014jd022629, 2015.
- Behnke, W., George, C., Scheer, V., and Zetzsch, C.: Production and decay of ClNO_2 from the reaction of gaseous N_2O_5 with NaCl solution: Bulk and aerosol experiments, *J. Geophys. Res. -Atmos.*, 102, 3795-3804, 1997.
- Bertram, T. H., and Thornton, J. A.: Toward a general parameterization of N_2O_5 reactivity on aqueous particles: the competing effects of particle liquid water, nitrate and chloride, *Atmos. Chem. Phys.*, 9, 8351-8363, 2009.
- Bertram, T. H., Thornton, J. A., and Riedel, T. P.: An experimental technique for the direct measurement of N_2O_5 reactivity on ambient aerosols, *Atmos. Meas. Tech.*, 2, 231-242, 2009a.



- Bertram, T. H., Thornton, J. A., Riedel, T. P., Middlebrook, A. M., Bahreini, R., Bates, T. S., Quinn, P. K., and Coffman, D. J.: Direct observations of N_2O_5 reactivity on ambient aerosol particles, *Geophys. Res. Lett.*, 36, L19803, doi:10.1029/2009GL040248, 2009b.
- 5 Brown, S. S., Ryerson, T. B., Wollny, A. G., Brock, C. A., Peltier, R., Sullivan, A. P., Weber, R. J., Dube, W. P., Trainer, M., Meagher, J. F., Fehsenfeld, F. C., and Ravishankara, A. R.: Variability in nocturnal nitrogen oxide processing and its role in regional air quality, *Science*, 311, 67-70, 2006.
- Brown, S. S., Dube, W. P., Fuchs, H., Ryerson, T. B., Wollny, A. G., Brock, C. A., Bahreini, R., Middlebrook, A. M., Neuman, J. A., Atlas, E., Roberts, J. M., Osthoff, H. D., Trainer, M., Fehsenfeld, F. C., and Ravishankara, A. R.: Reactive uptake coefficients for N_2O_5 determined from aircraft measurements during the Second Texas Air Quality Study: Comparison to current model parameterizations, *J. Geophys. Res. -Atmos.*, 114, art. D00F10, doi:10.1029/2008JD011679, 10 2009.
- Brown, S. S., Dube, W. P., Tham, Y. J., Zha, Q., Xue, L., Poon, S., Wang, Z., Blake, D. R., Tsui, W., Parrish, D. D., and Wang, T.: Nighttime chemistry at a high altitude site above Hong Kong, *J. Geophys. Res. -Atmos.*, 121, 2457-2475, doi:10.1002/2015jd024566, 2016a.
- 15 Brown, S. S., Dube, W. P., Tham, Y. J., Zha, Q. Z., Xue, L. K., Poon, S., Wang, Z., Blake, D. R., Tsui, W., Parrish, D. D., and Wang, T.: Nighttime chemistry at a high altitude site above Hong Kong, *J. Geophys. Res. -Atmos.*, 121, 2457-2475, doi:10.1002/2015jd024566, 2016b.
- Burkholder, J. B., Sander, S. P., Abbatt, J., Barker, J. R., Huie, R. E., Kolb, C. E., Kurylo, M. J., Orkin, V. L., Wilmouth, D. M., and Wine, P. H.: Chemical Kinetics and Photochemical Data for Use in Atmospheric Studies, Evaluation No. 18, "JPL Publication 15-10, Jet Propulsion Laboratory, Pasadena, <http://jpldataeval.jpl.nasa.gov>, 2016.
- 20 Chang, W. L., Bhave, P. V., Brown, S. S., Riemer, N., Stutz, J., and Dabdub, D.: Heterogeneous atmospheric chemistry, ambient measurements and model calculations of N_2O_5 : A review, *Aerosol. Sci. Tech*, 45, 655-685, 2011.
- Clegg, S. L., Brimblecombe, P., and Wexler, A. S.: Thermodynamic model of the system $\text{H}^+ - \text{NH}_4^+ - \text{Na}^+ - \text{SO}_4^{2-} - \text{NO}_3^- - \text{Cl}^- - \text{H}_2\text{O}$ at 298.15 K, *J. Phys. Chem. A*, 102, 2155-2171, doi:10.1021/jp973043j, 1998.
- 25 Cosman, L. M., and Bertram, A. K.: Reactive uptake of N_2O_5 on aqueous H_2SO_4 solutions coated with 1-component and 2-component monolayers, *J. Phys. Chem. A*, 112, 4625-4635, doi:10.1021/jp8005469, 2008.
- Cosman, L. M., Knopf, D. A., and Bertram, A. K.: N_2O_5 reactive uptake on aqueous sulfuric acid solutions coated with branched and straight-chain insoluble organic surfactants, *J. Phys. Chem. A*, 112, 2386-2396, doi:10.1021/jp710685r, 2008.
- 30 Crowley, J. N., Schuster, G., Pouvesle, N., Parchatka, U., Fischer, H., Bonn, B., Bingemer, H., and Lelieveld, J.: Nocturnal nitrogen oxides at a rural mountain site in south-western Germany, *Atmos. Chem. Phys.*, 10, 2795-2812, 2010.
- DeCarlo, P. F., Kimmel, J. R., Trimborn, A., Northway, M. J., Jayne, J. T., Aiken, A. C., Gonin, M., Fuhrer, K., Horvath, T., Docherty, K. S., Worsnop, D. R., and Jimenez, J. L.: Field-deployable, high-resolution, time-of-flight aerosol mass spectrometer, *Anal. Chem.*, 78, 8281-8289, doi:10.1021/ac061249n, 2006.
- Dentener, F. J., and Crutzen, P. J.: Reaction of N_2O_5 on tropospheric aerosols - Impact on the global distributions of NO_x , O_3 , and OH, *J. Geophys. Res. -Atmos.*, 98, 7149-7163, 1993.
- 35 Draxler, R. R., and Rolph, G. D.: HYSPLIT (HYbrid Single-Particle Lagrangian Integrated Trajectory) Model access via NOAA ARL READY Website (<http://ready.arl.noaa.gov/HYSPLIT.php>). NOAA Air Resources Laboratory, Silver Spring, MD., 2011.
- Drewnick, F., Boettger, T., von der Weiden-Reinmueller, S. L., Zorn, S. R., Klimach, T., Schneider, J., and Borrmann, S.: Design of a mobile aerosol research laboratory and data processing tools for effective stationary and mobile field measurements, *Atmos. Meas. Tech.*, 5, 1443-1457, doi:10.5194/amt-5-1443-2012, 2012.
- 40 Drewnick, F., Diesch, J. M., Faber, P., and Borrmann, S.: Aerosol mass spectrometry: particle-vaporizer interactions and their consequences for the measurements, *Atmos. Meas. Tech.*, 8, 3811-3830, doi:10.5194/amt-8-3811-2015, 2015.
- Finlayson-Pitts, B. J., Ezell, M. J., and Pitts, J. N. J.: Formation of chemically active chlorine compounds by reactions of atmospheric NaCl particles with gaseous N_2O_5 and ClONO_2 , *Nature*, 337, 241-244, 1989.
- 45 Gaston, C. J., Thornton, J. A., and Ng, N. L.: Reactive uptake of N_2O_5 to internally mixed inorganic and organic particles: the role of organic carbon oxidation state and inferred organic phase separations, *Atmos. Chem. Phys.*, 14, 5693-5707, doi:10.5194/acp-14-5693-2014, 2014.
- George, C., Ponche, J. L., Mirabel, P., Behnke, W., Scheer, V., and Zetzsch, C.: Study of the uptake of N_2O_5 by water and NaCl solutions, *J. Phys. Chem.*, 98, 8780-8784, 1994.
- 50



- Ghosh, B., Papanastasiou, D. K., Talukdar, R. K., Roberts, J. M., and Burkholder, J. B.: Nitryl Chloride (CINO₂): UV/Vis Absorption Spectrum between 210 and 296 K and O(P³) Quantum Yield at 193 and 248 nm, *J. Phys. Chem. A*, 116, 5796-5805, doi:10.1021/jp207389y, 2012.
- Hallquist, M., Stewart, D. J., Stephenson, S. K., and Cox, R. A.: Hydrolysis of N₂O₅ on sub-micron sulfate aerosols, *Phys. Chem. Chem. Phys.*, 5, 3453-3463, doi:10.1039/b301827j, 2003.
- Handisides, G. M.: The influence of peroxy radicals on ozone production, Fachbereich Geowissenschaften, Johann Wolfgang Goethe Universität, Frankfurt am Main, 2001.
- Heintz, F., Platt, U., Flentje, H., and Dubois, R.: Long-term observation of nitrate radicals at the tor station, Kap Arkona (Rügen), *J. Geophys. Res. -Atmos.*, 101, 22891-22910, 1996.
- Hu, J. H., and Abbatt, J. P. D.: Reaction probabilities for N₂O₅ hydrolysis on sulfuric acid and ammonium sulfate aerosols at room temperature, *J. Phys. Chem. A*, 101, 871-878, 1997.
- Jayne, J. T., Leard, D. C., Zhang, X. F., Davidovits, P., Smith, K. A., Kolb, C. E., and Worsnop, D. R.: Development of an aerosol mass spectrometer for size and composition analysis of submicron particles, *Aerosol Science and Technology*, 33, 49-70, doi:10.1080/027868200410840, 2000.
- Keene, W. C., Pszenny, A. A. P., Jacob, D. I., Duce, R. A., Galloway, J. N., Schultz-Tokos, J. J., Sievering, H., and Boatman, J. F.: The geochemical cycling of reactive chlorine through the marine troposphere *Glob. Biogeochem. Cycles*, 4, 407-430, doi:10.1029/GB004i004p00407, 1990.
- Keene, W. C., Khalil, M. A. K., Erickson, D. J., McCulloch, A., Graedel, T. E., Lobert, J. M., Aucott, M. L., Gong, S. L., Harper, D. B., Kleiman, G., Midgley, P., Moore, R. M., Seuzaret, C., Sturges, W. T., Benkovitz, C. M., Koropalov, V., Barrie, L. A., and Li, Y. F.: Composite global emissions of reactive chlorine from anthropogenic and natural sources: Reactive Chlorine Emissions Inventory, *J. Geophys. Res. -Atmos.*, 104, 8429-8440, doi:10.1029/1998jd100084, 1999.
- Kercher, J. P., Riedel, T. P., and Thornton, J. A.: Chlorine activation by N₂O₅: simultaneous, in situ detection of CINO₂ and N₂O₅ by chemical ionization mass spectrometry, *Atmos. Meas. Tech.*, 2, 193-204, 2009.
- Macintyre, H. L., and Evans, M. J.: Sensitivity of a global model to the uptake of N₂O₅ by tropospheric aerosol, *Atmos. Chem. Phys.*, 10, 7409-7414, doi:10.5194/acp-10-7409-2010, 2010.
- McNeill, V. F., Patterson, J., Wolfe, G. M., and Thornton, J. A.: The effect of varying levels of surfactant on the reactive uptake of N₂O₅ to aqueous aerosol, *Atmos. Chem. Phys.*, 6, 1635-1644, 2006.
- Mentel, T. F., Sohn, M., and Wahner, A.: Nitrate effect in the heterogeneous hydrolysis of dinitrogen pentoxide on aqueous aerosols, *Phys. Chem. Chem. Phys.*, 1, 5451-5457, 1999.
- Mielke, L. H., Furgeson, A., and Osthoff, H. D.: Observation of CINO₂ in a mid-continental urban environment, *Env. Sci. Tech.*, 45, 8889-8896, doi:10.1021/es201955u, 2011.
- Mielke, L. H., Stutz, J., Tsai, C., Hurlock, S. C., Roberts, J. M., Veres, P. R., Froyd, K. D., Hayes, P. L., Cubison, M. J., Jimenez, J. L., Washenfelder, R. A., Young, C. J., Gilman, J. B., de Gouw, J. A., Flynn, J. H., Grossberg, N., Lefer, B. L., Liu, J., Weber, R. J., and Osthoff, H. D.: Heterogeneous formation of nitryl chloride and its role as a nocturnal NO_x reservoir species during CalNex-LA 2010, *J. Geophys. Res. -Atmos.*, 118, 10638-10652, doi:10.1002/jgrd.50783, 2013.
- Morgan, W. T., Ouyang, B., Allan, J. D., Aruffo, E., Di Carlo, P., Kennedy, O. J., Lowe, D., Flynn, M. J., Rosenberg, P. D., Williams, P. I., Jones, R., McFiggans, G. B., and Coe, H.: Influence of aerosol chemical composition on N₂O₅ uptake: airborne regional measurements in northwestern Europe, *Atmos. Chem. Phys.*, 15, 973-990, doi:10.5194/acp-15-973-2015, 2015.
- Mozurkewich, M., and Calvert, J. G.: Reaction probability of N₂O₅ on aqueous aerosols, *J. Geophys. Res. -Atmos.*, 93, 15889-15896, 1988.
- Osthoff, H. D., Roberts, J. M., Ravishankara, A. R., Williams, E. J., Lerner, B. M., Sommariva, R., Bates, T. S., Coffman, D., Quinn, P. K., Dibb, J. E., Stark, H., Burkholder, J. B., Talukdar, R. K., Meagher, J., Fehsenfeld, F. C., and Brown, S. S.: High levels of nitryl chloride in the polluted subtropical marine boundary layer, *Nature Geoscience*, 1, 324-328, 2008.
- Ovadnevaite, J., Ceburnis, D., Canagaratna, M., Berresheim, H., Bialek, J., Martucci, G., Worsnop, D. R., and O'Dowd, C.: On the effect of wind speed on submicron sea salt mass concentrations and source fluxes, *J. Geophys. Res. -Atmos.*, 117, doi:10.1029/2011jd017379, 2012.
- Park, S. C., Burden, D. K., and Nathanson, G. M.: The inhibition of N₂O₅ hydrolysis in sulfuric acid by 1-butanol and 1-hexanol surfactant coatings, *J. Phys. Chem. A*, 111, 2921-2929, doi:10.1021/jp068228h, 2007.



- Phillips, G. J., Tang, M. J., Thieser, J., Brickwedde, B., Schuster, G., Bohn, B., Lelieveld, J., and Crowley, J. N.: Significant concentrations of nitryl chloride observed in rural continental Europe associated with the influence of sea salt chloride and anthropogenic emissions, *Geophys. Res. Lett.*, 39, L10811, doi:10.1029/2012GL051912, 2012.
- 5 Phillips, G. J., Makkonen, U., Schuster, G., Sobanski, N., Hakola, H., and Crowley, J. N.: The detection of nocturnal N_2O_5 as HNO_3 by alkali- and aqueous-denuder techniques, *Atmos. Meas. Tech.*, 6, 231-237, doi:10.5194/amt-6-231-2013, 2013a.
- Phillips, G. J., Pouvesle, N., Thieser, J., Schuster, G., Axinte, R., Fischer, H., Williams, J., Lelieveld, J., and Crowley, J. N.: Peroxyacetyl nitrate (PAN) and peroxyacetic acid (PAA) measurements by iodide chemical ionisation mass spectrometry: first analysis of results in the boreal forest and implications for the measurement of PAN fluxes, *Atmos. Chem. Phys.*, 13, 1129-1139, doi:10.5194/acp-13-1129-2013, 2013b.
- 10 Platt, U., and Heintz, F.: Nitrate radicals in tropospheric chemistry *Isr. J. Chem.*, 34, 289-300, 1994.
- Platt, U., and Janssen, C.: Observation and role of the free radicals NO_3 , ClO , BrO and IO in the troposphere, *Faraday Discussions*, 100, 175-198, doi:10.1039/fd9950000175, 1995.
- Riedel, T. P., Bertram, T. H., Crisp, T. A., Williams, E. J., Lerner, B. M., Vlasenko, A., Li, S. M., Gilman, J., de Gouw, J., Bon, D. M., Wagner, N. L., Brown, S. S., and Thornton, J. A.: Nitryl Chloride and Molecular Chlorine in the Coastal Marine
- 15 Boundary Layer, *Env. Sci. Tech.*, 46, 10463-10470, doi:10.1021/es204632r, 2012a.
- Riedel, T. P., Bertram, T. H., Ryder, O. S., Liu, S., Day, D. A., Russell, L. M., Gaston, C. J., Prather, K. A., and Thornton, J. A.: Direct N_2O_5 reactivity measurements at a polluted coastal site, *Atmos. Chem. Phys.*, 12, 2959-2968, doi:10.5194/acp-12-2959-2012, 2012b.
- Riemer, N., Vogel, H., Vogel, B., Schell, B., Ackermann, I., Kessler, C., and Hass, H.: Impact of the heterogeneous
- 20 hydrolysis of N_2O_5 on chemistry and nitrate aerosol formation in the lower troposphere under photochemical conditions, *J. Geophys. Res. -Atmos.*, 108, 2003.
- Riemer, N., Vogel, H., Vogel, B., Anttila, T., Kiendler-Scharr, A., and Mentel, T. F.: Relative importance of organic coatings for the heterogeneous hydrolysis of N_2O_5 during summer in Europe, *J. Geophys. Res. -Atmos.*, 114, art. Nr D17307, doi:10.1029/2008JD011369, 2009.
- 25 Roberts, J. M., Osthoff, H. D., Brown, S. S., Ravishankara, A. R., Coffman, D., Quinn, P., and Bates, T.: Laboratory studies of products of N_2O_5 uptake on Cl containing substrates, *Geophys. Res. Lett.*, 36, doi:10.1029/2009gl040448, 2009.
- Ryder, O. S., Campbell, N. R., Shalowski, M., Al-Mashat, H., Nathanson, G. M., and Bertram, T. H.: Role of organics in regulating ClNO_2 production at the air-sea interface, *J. Phys. Chem. A*, 119, 8519-8526, doi:10.1021/jp5129673, 2015.
- Sander, R., and Crutzen, P. J.: Model study indicating halogen activation and ozone destruction in polluted air masses
- 30 transported to the sea, *J. Geophys. Res. -Atmos.*, 101, 9121-9138, 1996.
- Sarwar, G., Simon, H., Bhawe, P., and Yarwood, G.: Examining the impact of heterogeneous nitryl chloride production on air quality across the United States, *Atmos. Chem. Phys.*, 12, 6455-6473, doi:10.5194/acp-12-6455-2012, 2012.
- Schmale, J., Schneider, J., Nemitz, E., Tang, Y. S., Dragosits, U., Blackall, T. D., Trathan, P. N., Phillips, G. J., Sutton, M., and Braban, C. F.: Sub-Antarctic marine aerosol: dominant contributions from biogenic sources, *Atmos. Chem. Phys.*, 13,
- 35 8669-8694, doi:10.5194/acp-13-8669-2013, 2013.
- Schuster, G., Labazan, I., and Crowley, J. N.: A cavity ring down / cavity enhanced absorption device for measurement of ambient NO_3 and N_2O_5 , *Atmos. Meas. Tech.*, 2, 1-13, 2009.
- Schweitzer, F., Mirabel, P., and George, C.: Multiphase chemistry of N_2O_5 , ClNO_2 , and BrNO_2 , *J. Phys. Chem. A*, 102, 3942-3952, 1998.
- 40 Simon, H., Kimura, Y., McGaughey, G., Allen, D. T., Brown, S. S., Osthoff, H. D., Roberts, J. M., Byun, D., and Lee, D.: Modeling the impact of ClNO_2 on ozone formation in the Houston area, *J. Geophys. Res. -Atmos.*, 114, 2009.
- Slusher, D. L., Huey, L. G., Tanner, D. J., Flocke, F. M., and Roberts, J. M.: A thermal dissociation-chemical ionization mass spectrometry (TD-CIMS) technique for the simultaneous measurement of peroxyacyl nitrates and dinitrogen pentoxide, *J. Geophys. Res. -Atmos.*, 109, Art Nr. D19315, doi:10.1029/2004JD004670, 2004.
- 45 Sobanski, N., Tang, M. J., Thieser, J., Schuster, G., Pöhler, D., Fischer, H., Song, W., Sauvage, C., Williams, J., Fachinger, J., Berkes, F., Hoor, P., Platt, U., Lelieveld, J., and Crowley, J. N.: Chemical and meteorological influences on the lifetime of NO_3 at a semi-rural mountain site during PARADE, *Atmos. Chem. Phys.*, 16, 4867-4883, doi:10.5194/acp-16-4867-2016, 2016.



- Stein, A. F., Draxler, R. R., Rolph, G. D., Stunder, B. J. B., Cohen, M. D., and Ngan, F.: NOAA'S HYSPLIT ATMOSPHERIC TRANSPORT AND DISPERSION MODELING SYSTEM, *Bulletin of the American Meteorological Society*, 96, 2059-2077, doi:10.1175/bams-d-14-00110.1, 2015.
- 5 Thieser, J., Schuster, G., Phillips, G. J., Reiffs, A., Parchatka, U., Pöhler, D., Lelieveld, J., and Crowley, J. N.: A two-channel, thermal dissociation cavity-ringdown spectrometer for the detection of ambient NO₂, RO₂NO₂ and RONO₂, *Atmos. Meas. Tech.*, 9, 553-576, 2016.
- Thornton, J. A., Braban, C. F., and Abbatt, J. P. D.: N₂O₅ hydrolysis on sub-micron organic aerosols: the effect of relative humidity, particle phase, and particle size, *Phys. Chem. Chem. Phys.*, 5, 4593-4603, 2003.
- 10 Thornton, J. A., and Abbatt, J. P. D.: N₂O₅ reaction on submicron sea salt aerosol: Kinetics, products, and the effect of surface active organics, *J. Phys. Chem. A*, 109, 10004-10012, 2005.
- Thornton, J. A., Kercher, J. P., Riedel, T. P., Wagner, N. L., Cozic, J., Holloway, J. S., Dube, W. P., Wolfe, G. M., Quinn, P. K., Middlebrook, A. M., Alexander, B., and Brown, S. S.: A large atomic chlorine source inferred from mid-continental reactive nitrogen chemistry, *Nature*, 464, 271-274, doi:10.1038/nature08905, 2010.
- von Glasow, R., Sander, R., Bott, A., and Crutzen, P. J.: Modeling halogen chemistry in the marine boundary layer. 1. Cloud-free MBL, *JGR*, 107, 9,1-9,16, 2001.
- 15 Wagner, N. L., Riedel, T. P., Young, C. J., Bahreini, R., Brock, C. A., Dube, W. P., Kim, S., Middlebrook, A. M., Ozturk, F., Roberts, J. M., Russo, R., Sive, B., Swarthout, R., Thornton, J. A., VandenBoer, T. C., Zhou, Y., and Brown, S. S.: N₂O₅ uptake coefficients and nocturnal NO₂ removal rates determined from ambient wintertime measurements, *J. Geophys. Res. - Atmos.*, 118, 9331-9350, doi:10.1002/jgrd.50653, 2013.
- 20 Wahner, A., Mentel, T. F., and Sohn, M.: Gas-phase reaction of N₂O₅ with water vapor: Importance of heterogeneous hydrolysis of N₂O₅ and surface desorption of HNO₃ in a large teflon chamber, *Geophys. Res. Lett.*, 25, 2169-2172, 1998a.
- Wahner, A., Mentel, T. F., Sohn, M., and Stier, J.: Heterogeneous reaction of N₂O₅ on sodium nitrate aerosol, *J. Geophys. Res. - Atmos.*, 103, 31103-31112, 1998b.
- 25 Wexler, A. S., and Clegg, S. L.: Atmospheric aerosol models for systems including the ions H⁺, NH₄⁺, Na⁺, SO₄²⁻, NO₃⁻, Cl⁻, Br⁻, and H₂O, *J. Geophys. Res. - Atmos.*, 107, 2002.
- Wiedensohler, A., Birmili, W., Nowak, A., Sonntag, A., Weinhold, K., Merkel, M., Wehner, B., Tuch, T., Pfeifer, S., Fiebig, M., Fjaraa, A. M., Asmi, E., Sellegri, K., Depuy, R., Venzac, H., Villani, P., Laj, P., Aalto, P., Ogren, J. A., Swietlicki, E., Williams, P., Roldin, P., Quincey, P., Hueglin, C., Fierz-Schmidhauser, R., Gysel, M., Weingartner, E., Riccobono, F., Santos, S., Gruening, C., Faloony, K., Beddows, D., Harrison, R. M., Monahan, C., Jennings, S. G., O'Dowd, C. D., Marinoni, A., Horn, H. G., Keck, L., Jiang, J., Scheckman, J., McMurry, P. H., Deng, Z., Zhao, C. S., Moerman, M., Henzing, B., de Leeuw, G., Loeschau, G., and Bastian, S.: Mobility particle size spectrometers: harmonization of technical standards and data structure to facilitate high quality long-term observations of atmospheric particle number size distributions, *Atmos. Meas. Tech.*, 5, 657-685, doi:10.5194/amt-5-657-2012, 2012.
- 30 Wobrock, W., Schell, D., Maser, R., Jaeschke, W., Georgii, H. W., Wieprecht, W., Arends, B. G., Mols, J. J., Kos, G. P. A., Fuzzi, S., Facchini, M. C., Orsi, G., Berner, A., Solly, I., Kruisz, C., Svenningsson, I. B., Wiedensohler, A., Hansson, H. C., Ogren, J. A., Noone, K. J., Hallberg, A., Pahl, S., Schneider, T., Winkler, P., Winiwarter, W., Colvile, R. N., Choulaton, T. W., Flossmann, A. I., and Borrmann, S.: The Kleiner-Feldberg Cloud Experiment 1990 - an Overview, *J. Atmos. Chem.*, 19, 3-35, 1994.
- 35 Young, A. H., Keene, W. C., Pszenny, A. A. P., Sander, R., Thornton, J. A., Riedel, T. P., and Maben, J. R.: Phase partitioning of soluble trace gases with size-resolved aerosols in near-surface continental air over northern Colorado, USA, during winter, *J. Geophys. Res. - Atmos.*, 118, 9414-9427, doi:10.1002/jgrd.50655, 2013.
- Zheng, W., Flocke, F. M., Tyndall, G. S., Swanson, A., Orlando, J. J., Roberts, J. M., Huey, L. G., and Tanner, D. J.: Characterization of a thermal decomposition chemical ionization mass spectrometer for the measurement of peroxy acyl nitrates (PANs) in the atmosphere, *Atmos. Chem. Phys.*, 11, 6529-6547, doi:doi:10.5194/acp-11-6529-2011, 2011.
- 45 Zorn, S. R., Drewnick, F., Schott, M., Hoffmann, T., and Borrmann, S.: Characterization of the South Atlantic marine boundary layer aerosol using an aerodyne aerosol mass spectrometer, *Atmos. Chem. Phys.*, 8, 4711-4728, doi:10.5194/acp-8-4711-2008, 2008.



Table 1: Values of γ derived from ambient datasets

Platform (location / height)	γ (method)	Notes	reference
Ship (East Coast U.S., 11 m)	0.03 ± 0.02 (ss)	Polluted marine	(Aldener et al., 2006)
Aircraft (NE-U.S. < 1.5 km)	< 0.0016 - 0.017 (ss)	Continental / coastal residual layer. γ dependence on organic / sulphate content.	(Brown et al., 2006)
Aircraft (Texas, U.S. < 1 km)	0.0005-0.006 (ss)	Continental / coastal residual layer. γ variable but independent of RH or aerosol composition	(Brown et al., 2009)
Ground (US, Seattle/Boulder 5-10 m)	< 0.01-0.04 (AFR)	Urban / sub-urban environment. γ dependence on organic / sulphate ratio and RH.	(Bertram et al., 2009b)
Ground (US, La Jolla, 15 m)	<0.001– 0.029 (AFR)	Polluted coastal environment. γ dependence on nitrate content	(Riedel et al., 2012b)
Ground (U.S, Boulder 10-300 m)	0.002–0.1 (Box model)	Continental, pollution impacted boundary layer / residual layer. γ dependence on nitrate content	(Wagner et al., 2013)
Aircraft (N.W. Europe 500-1000 m)	0.01–0.03 (ss)	Continental, pollution impacted residual layer / free troposphere. γ dependence on nitrate content	(Morgan et al., 2015)
Ground US, Pasadena (10 m)	$\gamma f = 0.008$ (av)	Coastal (CalNex-LA). γf was suppressed by particle organic content and enhanced by particulate chloride content.	(Mielke et al., 2013)
Ground China, Hong Kong, (957 m)	0.004–0.029 (ss)	Coastal, pollution impacted mountain site.	(Brown et al., 2016a)
Ground (S.W. Germany, 650 m)	0.004–0.11 (1b, 1c, ss)	Semi-rural mountain site with anthropogenic influence, mixed boundary layer / residual layer	This work

Notes: ss = steady-state analysis. AFR = aerosol flow reactor. Av = averaged over a campaign.




Fig. 1. Gas- and aqueous-phase chemical processes forming ClNO_2 and particulate nitrate indicating the role of ClNO_2 in modifying the effect of N_2O_5 uptake on the chemical lifetime of NO_x and photochemical RO_2 generation.




Fig. 2. Time series of temperature, relative humidity and particle properties during PARADE. A = aerosol surface area. Non-refractory, PM1 organic, nitrate, sulphate and ammonium were measured by the AMS. Aerosol surface area was measured by an FMPS (20-500 nm) and by APS ($>0.5 \mu\text{m}$) and OPC ($>0.25 \mu\text{m}$).




Fig. 3. Time series of ClNO_2 and its precursor, N_2O_5 . Global radiation (yellow) is plotted to separate day-night periods, and to indicate the day-to-day, relative photochemical activity. The precursors to N_2O_5 (NO_2 and O_3) are shown as are the AMS measurement of particulate nitrate (as in Fig. 2) and chloride.




Fig. 4. Zoom-in on particle-nitrate and CINO₂ formation on 4 nights. Global radiation (yellow) is plotted to separate day-night periods and indicate the day-to-day, relative photochemical activity. Note that the CINO₂ plotted for these days was measured at $m/z = 161.9$ and therefore contains a contribution from HCl, hence the apparent daytime production.




Fig. 5. Plots of CINO_2 versus NO_3^- for three different campaign nights. The slopes are converted to a fractional formation of CINO_2 (f) via Eq 5.




Fig 6. Plot demonstrating the derivation of f and γ from ClNO_2 and NO_3^- measurements using analysis method 1b. The grey area represents the time period in which $p\text{ClNO}_2$ and $p\text{NO}_3^-$ were measured. Average values of RH, $[\text{N}_2\text{O}_5]$ and A over the same period are also indicated.




Fig. 7. Plot demonstrating the derivation of f and γ from ClONO_2 and NO_3^- measurements using analysis method 1c. The red lines are the predicted, integrated concentrations of NO_3^- and ClONO_2 calculated using the summed particle surface area (A) from both coarse and fine particles. In this particular case the analysis returns a value of $\gamma = 0.085$ and $f = 0.33$. The relative humidity over this period was $82 \pm 2\%$.




Fig. 8. Values of γ and f derived via methods 1a-1c and the steady-state analysis during PARADE. The vertical error bars represent total uncertainty including errors associated with the measurement of concentrations of trace gases and particle surface areas. The horizontal error bars are the standard deviation of the relative humidity over the measurement period (1-3 hours for methods 1b and 1c and up to 8 hours for method 1a.). The red stars uptake coefficients calculated using the Bertram and Thornton (2009) parametrisation, the blue stars include the impact of an organic coating (Anttila et al, 2006).




Fig. 9. Derivation of γ via analysis of NO_3 steady-state lifetime variation with NO_2 . The grey $\tau(\text{NO}_3)$ data points were not considered in the analysis as NO_3 will not always be in steady state in the first hours after sunset or during sunrise.




Fig. 10. Plot of γ versus temperature, molar ratio of particle H_2O to particulate nitrate and the particle organic to SO_4^{2-} mass-ratio. Error bars are total uncertainty, calculated as described in the text.