

# IDF



## **International Dragonfly Fund - Report**

### **Journal of the International Dragonfly Fund**

1- 21

#### **Prosejit Dawn**

DragonflyIndia Meet 2016.

*Published: 15.02.2017*

# 102

ISSN 1435-3393

The International Dragonfly Fund (IDF) is a scientific society founded in 1996 for the improvement of odonatological knowledge and the protection of species.  
Internet: <http://www.dragonflyfund.org/>

This series intends to publish studies promoted by IDF and to facilitate cost-efficient and rapid dissemination of odonatological data..

Editorial Work:	Milen Marinov and Martin Schorr
Layout:	Martin Schorr
IDF-home page:	Holger Hunger
Printing:	Colour Connection GmbH, Frankfurt
Impressum:	Publisher: International Dragonfly Fund e.V., Schulstr. 7B, 54314 Zerf, Germany. E-mail: <a href="mailto:oestlap@online.de">oestlap@online.de</a>
Responsible editor:	Martin Schorr

Cover picture:	<b><i>Amphithemis vacillians</i></b> , India, Stream near Mohakal, 2016
Photographer:	Prosenjit Dawn

## DragonflyIndia Meet 2016

Prosenjit Dawn<sup>1,2</sup>

<sup>1</sup>Assistant Professor, Dept. of Zoology, Shyampur Siddheswari Mahavidyalaya, Howrah.

<sup>2</sup>Zoological Survey of India, New Alipore, Kolkata – 700053, West Bengal, India  
Email: prosenjit.dawn@gmail.com

### Abstract

The third meeting of DragonflyIndia social group was organized this time in Tiyabon Ecoresort, Gorumara National Park, Jalpaiguri district of West Bengal, India. Activities in the framework of the meeting are briefly outlined. A list of 66 Odonata taxa recorded during the field work is added.

**Key words:** DragonflyIndia, India, West Bengal, odonatology, meeting, *Agrionoptera insignis* (Rambur, 1842), *Amphithemis vacillans* Selys, 1891, *Coeliccia bimaculata* Laidlaw, 1914, *Ischnura aurora* (Brauer, 1865), *Ischnura rubilio* (Selys, 1876)

### Introduction

**Brief history of DragonflyIndia:** DragonflyIndia as an e-community was created in 2005 on Yahoo groups. Due to change in the social media popularity the community was moved to Facebook later in 2009. Since then the Facebook group has grown to reach 6000+ members worldwide. DragonflyIndia meet is the annual gathering of the willing members and is being organized since 2014. This year the meeting was organized in Gorumara National Park, Jalpaiguri, West Bengal (IFig. 1) on 26<sup>th</sup> to 29<sup>th</sup> August, 2016.

**The 2016 meeting:** 26 participants from eight states of India and one from Sri Lanka participated in this meet (IFig. 2). The meet was held in Tiyabon Ecoresort adjacent to the Gorumara National Park in Jalpaiguri district of West Bengal. The four days meet featured extensive field visits to forest streams, Perennial Lakes, Flooded grassland and crop-fields and Rivers in and around Gorumara National Park (Fig. 3-4). On 28<sup>th</sup> August there was a trip to the outskirts of Neora Valley National Park in Himalayan foothills where we visited a hilly torrential stream (Fig. 5). Weather was overall sunny with an average temperature range 20-35°C except two hours of heavy rain in Samsing. Apart from the field trips there were presentations and workshop sessions. Two presentations about general biology, field identification and odonates of West Bengal were from the resource person's end associated with hand on training on specimen handling, collection, preservation and identification of odonate specimens both in field and lab (IFig. 6-7). A special session was organized on discussion about larval identification, larval sampling and rearing. A workshop was organized by Mr. A. Pendharkar on popularizing Odonatology study and their conservation among

commons through outreach; and the participants were asked to design insectarium, museum, dragonfly postcards, books, cartoons etc. as part of the workshop.




Figure 1: Map showing the sites visited during the meet.



Figure 2: Top from Left: Prosenjit Dawn, Parag Rangnekar, Nitin Kumbhar, Megha Sethi, Devsena Roychoudhury. Bottom from left: Kumaran Sathasivam, Amila Prasanna Sumanapala, Shuvendu Das, Sachin Kamble, Sagar Adhurya, Ananya Sarkar, Partha Sarathi Mondal, S. Nazneen, Shubhankar Patra, Dipti Thakuria, Neha Mujumdar, Omkar Dharwarkar, S. Shivakumar, Jignasa Patel, Prasenjit Shome, Monika Devi, Anand Pendharkar, Harkiran Kaur, Viswajit Bansode, Neha Katnoria and behind the camera Arjan Basu Roy. (Photo: Arjan Basu Roy)





**Figure 3: Perennial Pond with plenty of aquatic vegetation at Gorumara National Park. (Photo: Prosenjit Dawn)**



**Figure 4: Field visit to the forest in Gorumara National Park. (Photo: Prosenjit Dawn)**



**Figure 5: Participants photographing the odonates in the hilly stream at Samsing. (Photo: Neha Mujumdar)**



**Figure 6: Presentation - Dragonflies of Gorumara National Park, West Bengal. (Photo: Neha Mujumdar)**



**Figure 7: Hands on training on Odonata specimen preservation and identification by Mr. Parag Rangnekar. (Photo: Neha Mujumdar)**

#### **Sampling localities:**

Gorumara National Park and Samsing: Being situated in the Indomalayan Ecozone Gorumara National Park hosts Lower Gangetic plains moist deciduous forests mainly dominated by sal, teak, rain tree, silk cotton tree etc. and Terai-Duar savanna and grasslands. This small park covers only 80 km<sup>2</sup> area and mostly famous for its healthy population of Indian one horned Rhinoceros and Indian Elephant. Other remarkable animals are Gaur, Leopard, Peafowl, Hornbills, King Cobra etc. Samsing on the other hand as a part of the lower Neora Valley National Park has Eastern Himalayan hilly moist deciduous forest.

#### **Status of the regional odonate fauna:**

The Odonata fauna of West Bengal is well explored since the British period. This state has a interesting species composition because northern districts have Indo-Malayan elements, Himalayan species; on the other hand the central part of the state share species composition with dry deciduous central Indian forests and the southernmost region comprises the peninsular Indian and coastal species. Fauna of British India by Fraser (1933, 1934 and 1936) included taxonomic descriptions and distribution records of several odonate species from northern part of Bengal. Fraser (1934, 1935a, 1935b, 1935c, 1935d and 1940) separately published dragonfly fauna of Darjeeling and Jalpaiguri districts later. The first compilation of Odonata fauna from the state was done by Srivastava & Sinha (1993) reporting 185 species from the state in which several species were reported from Darjeeling district and Dooars of Bengal. Recent



compilation work by Dawn (in preparation) estimates a list of 230+ species from the state including several new distributional records from the Dooars region.

## Material and methods

Sampling in Gorumara National Park and environs took place between 26<sup>th</sup> to 29<sup>th</sup> August, 2016: Tiyabon resort and forest (26°51'32.10"N 88°48'14.29"E), Stream near Mohakal (26°46'12.63"N 88°47'35.64"E), Chukchuki lake (26°45'17.45"N 88°50'47.09"E), ponds and grasslands at Ramsai village (26°43'33.19"N 88°51'14.05"E) and Murti river near Murti forest camp (26°50'27.99"N 88°49'41.44"E).

On 28<sup>th</sup> August there was a trip to the outskirts of Neora Valley National Park in Himalayan foothills to Samsing: Hilly stream in Samsing (27°00'43.75"N 88°47'09.11"E).

## Results

The three days field trip during the meet resulted in a list of 66 taxa belonging to 44 genera and 9 families of odonates from the area (Table 1 in appendix). Two of them were unidentified as we couldn't catch them or manage proper photograph. Some interesting behavioural observations were also noted in field (see Fig. 19-20).

## Discussion

Some of the records need a brief comment.

### Note on imagines:

*Agrionoptera insignis* (Rambur, 1842) – This widespread species is not very common and only found in West Bengal within Indian limits. This excellent insect have many subspecies throughout the World, the one available from West Bengal are sometime believed to be *A. insignis dorothea* Fraser, 1927 (Fig. 8).

*Amphithemis vacillans* Selys, 1891 – After a gap of almost seventy years, recently this species is recorded from West Bengal and Assam. During the meet we came across more than 10 specimens of this species, and also identified the breeding place for this species as newly emerged tenerals were spotted (Fig. 9-10).

*Coelliccia bimaculata* Laidlaw, 1914 – This species is known to be endemic to North Eastern Himalayas. During the meet we recorded the species second time from West Bengal after Dawn (in preparation) from the same area (Image 11).

*Coelliccia renifera* (Selys, 1886) – Widespread species throughout the Himalayas. Being a habitant of forest streams, they are not very commonly seen damselflies (Fig. 12).

*Aristocypha cuneata* (Selys, 1853) – This is one of the cryptic species we encountered during the meet. They are typically found near the fast forest streams of Samsing, associated with *Aristocypha quadrimaculata*. Distinguishable by slightly bigger size, palest blue elongated triangular spot of thorax and typical marking patterns of wing (Fig. 13).

*Aristocypha quadrimaculata* Selys, 1853 – This species is locally common in the Eastern Himalayas. Smaller in size than *A. cuneata* and have distinct wing markings (Fig. 14).



Figure 8: *Agrionoptera insignis* (Rambur, 1842) male.  
(Photo: Prosenjit Dawn)



Figure 9: Teneral male of *Amphithemis vacillans* Selys, 1891.  
(Photo: Prosenjit Dawn)





Figure 10: *Amphithemis vacillans* Selys, 1891 mature male. (Photo: Prosenjit Dawn)



Figure 11: *Coeliccia bimaculata* Laidlaw, 1914 male. (Photo: Prosenjit Dawn)



Figure 12: *Coeliccia renifera* (Selys, 1886) male. (Photo: Prosenjit Dawn)



Figure 13: *Aristocypha cuneata* (Selys, 1853) male. (Photo: Prosenjit Dawn)



Figure 14: *Aristocypha quadrimaculata* Selys, 1853 male. (Photo: Prosenjit Dawn)



Figure 15: Exuvia of *Vestalis* sp. (Photo: Prosenjit Dawn)



Figure 16: Emerging individual of *Agriocnemis femina* (Brauer, 1868) female. (Photo: Prosenjit Dawn)



Figure 17: Exuvia of *Hydrobasileus croceus* Brauer, 1867. (Photo: Prosenjit Dawn)

*Aciagrion hisopa* sensu Fraser nec (Selys, 1876) – Taxonomic note: Oleg Kosterin (in lit., 09-vi-2015) stated that the specimens from India are not *A. hisopa* (see Fig. 23). "Fraser had a misconception of *A. hisopa*. The true *hisopa* is missing from India. However, Indian *Aciagrion* are not revised."

### Note on the Exuviae studied during the meet:

Generally, knowledge on Indian odonate larvae and exuviae is poor. We therefore decided to collect exuviae to start a more close insight in larval morphology of Indian Odonata which must be intensified at each opportunity.

1. *Vestalis* sp. – An exuvia was found in the vegetation beside a forest stream.

Diagnosis: Comparatively longer damselfly exuvia; antennae seven segmented with first segment longer than the rest six segments all together (Fig. 15).

2. *Agriocnemis* sp. – Female individuals found to emerge during daytime from pond full of aquatic vegetation (Fig. 16).

Diagnosis: Small sized larvae, leaflike caudal gills pointed at tip and with black transverse markings.

3. *Pseudagrion* sp. – Exuviae were found over the submerged aquatic vegetation.

Diagnosis: Medium sized larva with caudal gills rounded at tip and with a node in the middle.

4. *Hydrobasileus croceus* (Brauer, 1867) – Found hanging from low water side vegetation beside the perennial pond.

Diagnosis: Strong backwardly directed dorsal spines, lateral spines of 9th and 10th segment very long. Eyes protruding backwards and distinctly pointed (Fig.17).



**Figure 18: Exuvia of *Epophthalmia vittata* Burmeister, 1839. (Photo: Prosenjit Dawn)**

5. *Epophthalmia vittata* Burmeister, 1839 – Found hanging from vegetation and wooden bridge beside a perennial pond.

Diagnosis: Large exuviae with long spidery legs, antennae seven segmented, distinct dorsal spines present, labium spoon shaped and inner margin of Labial palps armed with strong numerous teeth (Fig. 18).



### Note on *Ischnura aurora* (Brauer, 1865) and *Ischnura rubilio* (Selys, 1876):

A confusion issue about the identification of *Ischnura aurora* and *Ischnura rubilio* has aroused. Before the primary review of this manuscript by Martin Schorr we overlooked these two species and identified every single individual as *Ischnura aurora* from the study area. Now after re-examining our old specimens we realize that following the discussion by Papazian et al., 2007 most individuals present in North Bengal are actually matching with *I. rubilio* rather than *I. aurora*. Result with the specimens in my personal collection from Central India also the same. So, we need a thorough study of the *Ischnura aurora* specimens from India if they are all misidentified and actually are *I. rubilio*.

### Acknowledgements

The organizers are thankful to all the active members of DiversityIndia, Dragonfly-India and participants of the Meet 2016 for making the meet a big success. Thanks are also due to the Nature Mates – Nature Club for their support. Special mention for Martin Schorr for his valuable comment on the manuscript; the International Dragonfly Fund and Worldwide Dragonfly Association for their financial support.

### References

- Asahina, S. 1984. A list of the Odonata from Thailand. Part VI. Platycnemididae – genus *Coeliccia*. Tombo 27: 2-20.
- Fraser F.C. 1933. The Fauna of British India Odonata 1. Taylor and Francis Ltd. London.
- Fraser F.C. 1934a. Fauna of British India Odonata 2. Taylor and Francis Ltd. London.
- Fraser F.C. 1934b. The dragonfly fauna of Darjeeling and Jalpaiguri districts and Sikkim-I. Journal of Darjeeling Natural History Society 8: 161-177.
- Fraser F.C. 1935a. The dragonfly fauna of Darjeeling and Jalpaiguri districts and Sikkim-II. Journal of Darjeeling Natural History Society 9: 18-31.
- Fraser F.C. 1935b. The dragonfly fauna of Darjeeling and Jalpaiguri districts and Sikkim-III. Journal of Darjeeling Natural History Society 9: 70-84.
- Fraser F.C. 1935c. The dragonfly fauna of Darjeeling and Jalpaiguri districts and Sikkim-IV. Journal of Darjeeling Natural History Society 9: 101-115.
- Fraser F.C. 1935d. The dragonfly fauna of Darjeeling and Jalpaiguri districts and Sikkim-V. Journal of Darjeeling Natural History Society 10: 92-100.
- Fraser F.C. 1936. Fauna of British India Odonata 3. Taylor and Francis Ltd. London.
- Fraser F.C. 1940. The dragonfly fauna of Darjeeling and Jalpaiguri districts and Sikkim-VI. Journal of Darjeeling Natural History Society 14: 112-122, 145-151.
- Papazian, P., Dumont, H.J. & N.J. Mary-Sasal, 2007. The Odonata of the Pacific Ocean Islands of Wallis and Futuna, with special reference to specification in *Ischnura aurora* (Brauer). *Odonatologica* 36(1): 53-62.
- Srivastava V.D. & C. Sinha, 1993. Insecta: Odonata fauna of West Bengal, State Fauna Series, Part 4, Zoological Survey of India, Kolkata, 51- 168.



## Appendix 1

No.	Name of species	Gorumara National Park			Ponds and grasslands at Ramsal village	Murli river near Murli forest camp	Hilly stream in Samsing
		Tiyabon resort and forest	Stream near Mohakal	Chukchuki lake			
	<b>Leptidae</b>						
1	<i>Leptis cf. goroensis</i> Lahiri, 1987	•					
	<b>Calopterygidae</b>						
2	<i>Neurobasis chinensis</i> (Linnaeus, 1758)		•				
3	<i>Vestalis</i> sp.		•				
	<b>Chlorocyphidae</b>						
4	<i>Aristocypha cuneata</i> (Selys, 1853)					•	•
5	<i>Aristocypha quadrimaculata</i> Selys, 1853						•
6	<i>Helicypha biflorata</i> Selys, 1859 [Fig. 21]		•				
	<b>Euphaeidae</b>						
7	<i>Anisopleura</i> sp. [Fig. 22]						•
	<b>Coenagrionidae</b>						
8	<i>Coenagrion</i> sp. ( <i>Coenagrion hisopa</i> sensu Fraser nec (Selys, 1876)). [Fig. 23]	•					
9	<i>Agrionemis clauseni</i> Fraser, 1922	•	•		•		
10	<i>Agrionemis femina</i> (Brauer, 1868) [Fig. 24]					•	
11	<i>Agrionemis lacteola</i> Selys, 1877					•	
12	<i>Agrionemis pygmaea</i> (Rambur, 1842)	•			•		
13	<i>Amphialagma parvum</i> (Selys, 1876) [Fig. 26]	•			•		
14	<i>Ceragrion coromandelianum</i> (Fabricius, 1798)	•			•		•
15	<i>Ceragrion cernuobelium</i> (Brauer, 1865) [Fig. 25]				•		
16	<i>Ceragrion olivaceum</i> Laidlaw, 1914	•					
17	<i>Schnura cf. rubilla</i> (Selys, 1876)						•
18	<i>Marstonagrion aborense</i> (Laidlaw, 1914)						•
19	<i>Paracerion calamorum</i> (Ris, 1916) [Fig. 27]				•		
20	<i>Pseudagrion australisae</i> Selys, 1876				•		
21	<i>Pseudagrion decorum</i> (Rambur, 1842)				•		
22	<i>Pseudagrion microcephalum</i> (Rambur, 1842)				•		
23	<i>Pseudagrion rubriceps</i> Selys, 1876	•			•		
	<b>Platycnemididae</b>						
24	<i>Coelicia renifera</i> (Selys, 1886)						•
25	<i>Coelicia bimaculata</i> Laidlaw, 1914						•
26	<i>Coperia marginipes</i> (Rambur, 1842) [Fig. 28]						
27	<i>Coperia vittata</i> Selys, 1863 [Fig. 29]						
28	<i>Flotanea complani</i> (Fraser, 1922) [Fig. 30]						
29	<i>Onychargia atrocycana</i> (Selys, 1865)						•
	<b>Aeshnidae</b>						
30	<i>Gynacantha drowida</i> Liefelinck, 1960	•					
	<b>Gomphidae</b>						
31	<i>Burmagomphus cf. sivalikensis</i> Laidlaw, 1922						
32	<i>Ictinogomphus rapax</i> (Rambur, 1842)						•
33	<i>Nychogomphus cf. ataricus</i> (Fraser, 1924) [Fig. 31]						•

No.	Name of species	Tiyabon resort and forest	Stream near Mohakal	Chukchuk lake	Ponds and grasslands at Ramsai village	Murti river near Murti forest camp	Hilly stream in Samsing
34	Unidentified Gomphidae 1						•
35	Unidentified Gomphidae 2						•
<b>Libellulidae</b>							
36	<i>Acisoma panorpoides</i> Rambur, 1842				•		
37	<i>Aethriamanta brevipennis</i> [Rambur, 1842]			•			
38	<i>Agrionoptera insignis</i> [Rambur, 1842]		•				
39	<i>Amphithemis vacillans</i> Selys, 1891		•				
40	<i>Brachydiplax chalybea</i> Brauer, 1868				•		
41	<i>Brachythemis contaminata</i> [Fabricius, 1793]	•	•	•	•	•	
42	<i>Brachinopogon geminata</i> [Rambur, 1842]	•					•
43	<i>Crocothemis servilla</i> [Drury, 1770]	•		•			
44	<i>Diplacodes trivialis</i> [Rambur, 1842]	•		•			
45	<i>Hydrobasileus croceus</i> Brauer, 1867				•		•
46	<i>Lathrecista asiatica</i> [Fabricius, 1798]	•					
47	<i>Neurothemis fulvia</i> [Drury, 1773]	•			•		
48	<i>Neurothemis intermedia</i> [Rambur, 1842]	•					
49	<i>Neurothemis tullia</i> [Drury, 1773]				•		
50	<i>Orithetrum chysis</i> [Selys, 1891]		•				
51	<i>Orithetrum glaucum</i> [Brauer, 1865]		•				•
52	<i>Orithetrum luzonicum</i> [Brauer, 1868]						•
53	<i>Orithetrum pruinosum</i> [Burmeister, 1839]	•	•				•
54	<i>Orithetrum sabina</i> [Drury, 1770]	•		•		•	•
55	<i>Orithetrum taeniolatum</i> [Schneider, 1845]						•
56	<i>Pantala flavescens</i> [Fabricius, 1798]	•	•	•	•	•	•
57	<i>Potamarchia congener</i> [Rambur, 1842]		•				
58	<i>Rhodothemis ruta</i> [Rambur, 1842]						
59	<i>Rhyothemis vanegata</i> [Linnaeus, 1763]	•		•	•		
60	<i>Tholymis illariga</i> [Fabricius, 1798]	•					
61	<i>Tramea basilaris</i> [Palisot de Beuvois, 1805]	•					•
62	<i>Trithemis aurora</i> [Burmeister, 1839]						•
63	<i>Trithemis festiva</i> [Rambur, 1842]		•			•	•
64	<i>Trithemis pallidithemis</i> [Kirby, 1889]			•			•
65	<i>Urothemis signata</i> [Rambur, 1842]			•	•		
<b>Macromiidae</b>							
66	<i>Epophthalmia vittata</i> Burmeister, 1839 [Fig. 32]				•		

Table 1: List of Species observed during the Meet (• denotes the presence of the species).



**Figure 19:**  
More than 50 *Hydrobasi-leus croceus* were found to swarm together at a height of about 20-25m. (Photo: Prosenjit Dawn)



**Figure 20:** Two male individuals of *Heliocypha biforata* exhibiting the territorial display in flight. (Photo: Prosenjit Dawn)



Figure 21: *Heliocypha biforata* Selys, 1859 male. (Photo: Prosenjit Dawn)



Figure 22: *Anisopleura* sp. female. (Photo: Prosenjit Dawn)



Figure 23: *Aciagrion* sp. male. (Photo: Prosenjit Dawn)



Figure 24: *Agriocnemis femina* (Brauer, 1868) male. (Photo: Prosenjit Dawn)





Figure 25: *Ceriagrion cerinorubellum* (Brauer, 1865) male. (Photo: Prosenjit Dawn)



Figure 26: *Amphiallagma parvum* (Selys, 1876) male. (Photo: Prosenjit Dawn)



Figure 27: *Paracercion calamorum* (Ris, 1916) male. (Photo: Prosenjit Dawn)



Figure 28: *Coper marginipes* (Rambur, 1842) male. (Photo: Prosenjit Dawn)



Figure 29: *Coper vittata* Selys, 1863 male. (Photo: Prosenjit Dawn)




Figure 30: *Elatoneura campioni* (Fraser, 1922) male. (Photo: Prosenjit Dawn)



Figure 31: *Nychogomphus cf. duaricus* (Fraser, 1924) male. (Photo: Prosenjit Dawn)



Figure 32: *Epophthalmia vittata* Burmeister, 1839 male. (Photo: Prosenjit Dawn)





## INSTRUCTION TO AUTHORS

Faunistic studies of South-East Asian and Pacific islands Odonata is a journal of the International Dragonfly Fund (IDF). It is referred to as the journal in the remainder of these instructions. Transfer of copyright to IDF is considered to have taken place implicitly once a paper has been published in the journal.

The journal publishes original papers only. By original is meant papers that: a) have not been published elsewhere before, and b) the scientific results of the paper have not been published in their entirety under a different title and/or with different wording elsewhere. The republishing of any part of a paper published in the journal must be negotiated with the Editorial Board and can only proceed after mutual agreement.

Papers reporting studies financially supported by the IDF will be reviewed with priority, however, authors working with Odonata from the focal area (as defined on the back page of the front cover) are encouraged to submit their manuscripts even if they have not received any funds from IDF.

Manuscripts submitted to the journal should preferably be in English; alternatively German or French will also be accepted. Every manuscript should be checked by a native speaker of the language in which it is written; if it is not possible for the authors to arrange this, they must inform the Editorial Board on submission of the paper. Authors are encouraged, if possible, to include a version of the abstract in the primary language of the country in which their study was made.

Authors can choose the best way for them to submit their manuscripts between these options: a) via e-mail to the publisher, or b) on a CD, DVD or any other IBM-compatible device. Manuscripts should be prepared in Microsoft Word for Windows.

While preparing the manuscript authors should consider that, although the journal gives some freedom in the style and arrangements of the sections, the editors would like to see the following clearly defined sections: Title (with authors names, physical and e-mail addresses), Abstract, Introduction, Material & Methods, Results, Discussion, Acknowledgments and References. This is a widely used scheme by scientists that everyone should be familiar with. No further instructions are given here, but every author should check the style of the journal.

Authors are advised to avoid any formatting of the text. The manuscripts will be stylised according to the font type and size adopted by the journal. However, check for: a) all species names must be given in italic, b) the authority and year of publication are required on the first appearance of a species name in the text, but not thereafter, and c) citations and reference list must be arranged following the format below.

Reference cited in the text should read as follows: Tillyard (1924), (Tillyard 1924), Swezey & Williams (1942).

The reference list should be prepared according to the following standard:

Swezey, O. & F. Williams, 1942. Dragonflies of Guam. Bernice P. Bishop Museum Bulletin 172: 3-6.

Tillyard, R., 1924. The dragonflies (Order Odonata) of Fiji, with special reference to a collection made by Mr. H.W. Simmonds, F.E.S., on the Island of Viti Levu. Transactions of the Entomological Society London 1923 III-IV: 305-346.

Citations of internet sources should include the date of access.

The manuscript should end with a list of captions to the figures and tables. The latter should be submitted separately from the text preferably as graphics made using one of the Microsoft Office products or as a high resolution picture saved as a .jpg .tif or .ps file. Pictures should be at least 11 cm wide and with a minimum 300 dpi resolution, better 360 dpi. Line drawings and graphics could have 1200 dpi for better details. If you compose many pictures to one figure, please submit the original files as well. Please leave some space in the upper left corner of each picture, to insert a letter (a, b, c...) later. Hand-made drawings should be scanned and submitted electronically. Printed figures sent by the post could be damaged, in which case authors will be asked to resubmit them.

Manuscripts not arranged according to these instructions may also be accepted, but in that case their publication will be delayed until the journal's standards are achieved.

