

Journal of the International Dragonfly Fund

1-18

Rory A. Dow, Chee Yen Choong & Yong Foo Ng

New records of Odonata from Kedah, Malaysia in September 2016, with a checklist of species recorded from the state

published 13.12.2017

No. 23 ISSN 2195-4534


The International Dragonfly Fund (IDF) is a scientific society founded in 1996 for the improvement of odonatological knowledge and the protection of species.

Internet: http://www.dragonflyfund.org/

This series intends to contribute to the knowledge of the regional Odonata fauna of the Southeas-tern Asian and Pacific regions to facilitate cost-efficient and rapid dissemination of faunistic data.

Southeast Asia or Southeastern Asia is a subregion of Asia, consisting of the countries that are geo-graphically south of China, east of India, west of New Guinea and north of Australia. Southeast Asia consists of two geographic regions: Mainland Southeast Asia (Indochina) and Maritime Southeast Asia.

Pacific Islands comprise of Micronesian, Melanesian and Polynesian Islands.



Editorial Work: Martin Schorr, Milen Marinov and Rory Dow

Layout: Martin Schorr

IDF-home page: Holger Hunger

Printing: Colour Connection GmbH, Frankfurt

Impressum: Publisher: International Dragonfly Fund e.V., Schulstr. 7B,

54314 Zerf, Germany. E-mail: oestlap@online.de

Responsible editor: Martin Schorr

Cover picture: Zygonyx iris

Photographer: Chee Yen Choong

# New records of Odonata from Kedah, Malaysia in September 2016, with a checklist of species recorded from the state

Rory A. Dow<sup>1</sup>, Chee Yen Choong<sup>2</sup> & Yong Foo Ng<sup>3</sup>

<sup>1</sup>Naturalis Biodiversity Center, P.O. Box 9517, 2300 RA Leiden, The Netherlands. Email: rory.dow230@yahoo.co.uk

<sup>2</sup>Centre for Insect Systematics, Universiti Kebangsaan Malaysia, 43600 UKM Bangi, Selangor, Malaysia.

Email: cychoong@ukm.edu.my

<sup>3</sup>Centre for Insect Systematics, Universiti Kebangsaan Malaysia, 43600 UKM Bangi, Selangor, Malaysia cychoong@ukm.edu.my

### **Abstract**

The results of a short collecting trip to Kedah in the north-west of Peninsular Malaysia, made in September 2016, are reported. 64 species were collected, 13–14 of these are new records for the state and 28–29 are new records for the mainland of Kedah. A checklist of the Odonata recorded from Kedah (including Langkawi Island) is given in an appendix. At least 126 species of Odonata are now known from the state.

## Bahasa Melayu Abstract

Hasil pengutipan daripada kerja lapangan yang singkat di bahagian barat laut Kedah, Semenanjung Malaysia dalam bulan September 2016 dilaporkan di sini. Sebanyak 64 spesies telah berjaya dikutip. 13–14 spesies daripada ini merupakan rekod baharu untuk negeri Kedah dan 28–29 spesies merupakan rekod baharu untuk tanah besar Kedah. Satu senarai semakan Odonata untuk negeri Kedah (termasuk Pulau Langkawi) diberikan dalam satu lampiran. Sekurang-kurangnya 126 spesies Odonata diketahui wujud di negeri Kedah.

Key words: Odonata checklist, Kedah state, Langkawi Island, Malaysia, new records

### Introduction

Kedah is in the north-west of Peninsular Malaysia, it shares a border with Thailand, and has Perlis to its north and Perak to its south. Fig. 1 shows the location of Kedah in Peninsular Malaysia. Although better studied than some states in Peninsular Malaysia, it is still insufficiently studied. We are aware of records of Odonata from the mainland of Kedah in: Asahina (1966, 1967), Brooks (1981), Che Salmah et al. (2000), Choong et al. (2006), Dow (2010), Dow et al. (2017), Fraser (1942), Laidlaw (1902a, 1902b, 1931, 1932, 1934), Lieftinck (1933, 1954, 1965a, 1965b), Novelo-Gutierrez & Che Salmah (2006), Wahizatul Afzan & Che Salmah (2005). Additionally, records from Langkawi Island can be

found in Butler (2011), Butler & Kohler (2013), Dow et al. (2016), Farizawati et al. (2014), Fraser (1935, 1942), Inoue (1974), Lieftinck (1954), and Tsuda & Kitagawa (1987), and records from the somewhat confusingly named Pulau Perak can be found in Norma-Rashid et al. (2008). These papers contain records of 112–113 species (four taxa omitted from this figure are discussed in notes to the checklist in the appendix). Che Salmah et al. (2004) and Che Salmah & Wahizatul Afzan (2004) also include records from Kedah, but they are mixed with records from other states in such a way that it is not possible to determine which species has been recorded in which state. Moreover, some of the species listed in Che Salmah & Wahizatul Afzan (2004) certainly do not occur in Malaysia, e.g. the Indian species Coeliccia fraseri Laidlaw, 1932 and Protosticta hearseyi Fraser, 1922; presumably these records are the result of using inappropriate literature for identification. It is also possible that there are additional relevant publications in the Malaysian and/or Japanese literature of which we are unaware. The holotypes of Drepanosticta rahmani Dow, Choong & Ng, 2017, Teinobasis kirbyi Laidlaw, 1902 and Leptogomphus risi Laidlaw, 1932 are from Kedah.

In September 2016 the first two authors made a short trip to Kedah, staying in the town of Kulim and making day trips from there to various sites. Collecting was carried out on 12-16 September. Unfortunately the second author became ill shortly after our arrival and was only able to conduct limited field work. Even so, and despite the short duration of the trip (only five days sampling), a number of new records were made for Kedah, and a new species was discovered (see Dow et al. 2017). A checklist of the Odonata now known from Kedah is given in an appendix.



Figure 1. Map of Peninsular Malaysia showing position of Kedah. Image made using GoogleEarth.




Figure 2. Map showing the main sampling sites. Image made using GoogleEarth..



Figure 3–6. Habitat at some of the sampled locations (all photos by C.Y. Choong): (3) main stream at Hutan Lipur Ulu Paip (location 1); (4) main stream at Hutan Lipur Bukit Hijau (location 4); (5) main stream at Hutan Lipur Sungai Sedim; (6) lower part of stream system in hills between Baling and Gulai (Location 9).

## Odonata recorded in September 2016

#### Locations

The main sampling locations are indicated in Fig. 2.

- 1. Main stream at Hutan Lipur Ulu Paip (12.xi; Fig. 3).
- 2. Tributary streams at Hutan Lipur Ulu Paip (12.ix).
- 3. Pond along road outside Hutan Lipur Ulu Paip (12.ix).
- 4. Main stream at Hutan Lipur Bukit Hijau (13.ix, Fig. 4).
- 5. Tributary streams at Hutan Lipur Bukit Hijau (13.ix).
- 6. Rice fields near Hutan Lipur Bukit Hijau (13.ix).
- 7. Small stream near canopy walk at Hutan Lipur Sungai Sedim (14.ix). Fig. 5 shows the main stream at this location, which was difficult to sample.
- 8. Small streams reached by walking several km along an old, overgrown, concrete road at Hutan Lipur Sungai Sedim (14.ix).
- 9. Stream system in hills between Baling and Gulai (15–16.ix; Fig. 6).

### Species list

First records from mainland Kedah are indicated by a \*, new records for Kedah including Langkawi are indicated by a \*\*. The first two authors' names are abbreviated as RAD and CYC. 3+9 indicates a pair in tandem.

### Zygoptera

**LESTIDAE** 

Lestes praemorsus decipiens Kirby, 1893\*

3 - 3. 12.ix, CYC.

#### **PLATYSTICTIDAE**

Drepanosticta fontinalis Lieftinck, 1937

- **2** 17 ನನ, 12.ix, RAD.
- **4** − ♂, 13.ix, CYC; ♂, 13.ix, RAD.
- **5** 4 ♂♂, ♀, 13.ix, RAD.
- 8 11 ♂♂, 14.ix, RAD.
- 9 ♂, 16.ix, CYC.

Drepanosticta rahmani Dow, Choong & Ng, 2017

See Dow et al. (2017) for details. Location 9.

Drepanosticta silenus Laidlaw, 1934\*\*

Until now this species had only been known from records from Bukit Larut (Maxwell Hill) in Perak (Laidlaw 1934, Donnelly 1998).

9 – ♂, 16.ix, RAD.

### **ARGIOLESTIDAE**

Podolestes orientalis Selys, 1862\*\*

8 – 2 ♂♂, 14.ix, RAD.

### CALOPTERYGIDAE

Echo modesta Laidlaw, 1902\*\*

Fig. 7 shows a female.

**2** – ♂, 12.ix, RAD.

9 – 3 &&, 15.ix, RAD; &, 16.ix, CYC; &, 16.ix, RAD.

Neurobasis chinensis (Linnaeus, 1758)

1 - 3, 12.ix, CYC; 3, 12.ix, RAD.

**4** − ♂, 13.ix, CYC.

Vestalis amethystina Lieftinck, 1965

2 - 2 33, 12.ix, CYC; 2 33, 12.ix, RAD.

**5** – ♂, 13.ix, RAD.

8 – ♂, 14.ix, RAD.

9 - 2 ♂♂, 15.ix, RAD.

Vestalis amoena Hagen in Selys, 1853

See the note to this species in the checklist in the appendix; it is not clear whether or not it has been recorded from Kedah before now.

**4** – ♂, 13.ix, CYC; ♂, 13.ix, RAD.



Figure 7: Echo modesta female. Photo by C.Y. Choong.

### **CHLOROCYPHIDAE**

Aristocypha fenestrella (Rambur, 1842)

- **4** ♂, 13.ix, CYC; 2 ♂♂, 13.ix, RAD.
- 8 ♂, 14.ix, RAD.
- 9 ♂, 15.ix, CYC; 2 ♂♂, 15.ix, RAD; ♀, 16.ix, RAD.

Heliocypha biforata (Selys, 1859)

- 1 3, 12.ix, CYC; 3, 12.ix, RAD.
- 2 ♂, 12.ix, RAD.

Heliocypha perforata (Percheron, 1835)

- 1 3, 12.ix, CYC; 3, 12.ix, RAD.
- 4 3, 13.ix, CYC; 3, 13.ix, RAD.

Libellago lineata (Burmeister, 1839)

1 - ♂, 12.ix, CYC; ♂, 12.ix, RAD.

Sundacypha petiolata (Selys, 1859)\*\*

2 – ♂, 12.ix, RAD.

### **DEVADATTIDAE**

Devadatta argyoides (Selys, 1859)

- 2 3 ♂♂, 12.ix, RAD.
- **5** 3, 13.ix, CYC; 3 33, 13.ix, RAD.
- **7** − ♂, 14.ix, RAD.
- **9** ♂, 15.ix, CYC; 3 ♂♂, ♀, 15.ix, RAD; 2 ♂♂, ♀, 16.ix, RAD.

### **EUPHAEIDAE**

Dysphaea dimidiata Selys, 1853 \*\*

- 1 3, 12.ix, CYC; 3, 12.ix, RAD.
- **4** − 2 ♂♂, 13.ix, RAD.

Euphaea ochracea Selys, 1859

- **4** 3, 13.ix, CYC; 3 33, 13.ix, RAD.
- 8 ♂, 14.ix, RAD.
- 9 3, 15.ix, RAD; 3, 16.ix, CYC.
- 6 ♂, 29.vi, CYC.
- 7 4 33, 29.vi, CYC; 3 33, 29.vi, RAD.
- 9 3, 30.vi, CYC; 3, 30.vi, RAD.

### **PHILOSINIDAE**

Rhinagrion viridatuma Fraser, 1938

See the comments on this species in Dow et al. (2016) and Kosterin (2014)

**4** − ♂, 13.ix, RAD.

#### PLATYCNEMIDIDAE

Calicnemia chaseni (Laidlaw, 1928)\*\*

- 8 ♂, 14.ix, RAD.
- 9 ♂, 15.ix, RAD; ♂, 16.ix, RAD.

Coeliccia albicauda (Förster in Laidlaw, 1907)\*\*

- 2 3, 12.ix, RAD.
- **5** ♂, ♂+♀, 13.ix, RAD.
- 9 ♂, 15.ix, RAD.

Coeliccia didyma (Selys, 1863)

9 – ♂, 16.ix, RAD.

Coeliccia erici Laidlaw, 1917\*\*

Material of this species, and its confused history, will be discussed in Dow, Choong & Ng (in prep.). Location **9**.

Coeliccia octogesima (Selys, 1863)\*\*

Not only is this the first record of this species from Kedah, it also appears to be the most northern record yet known.

**5** − 4 ♂♂, 13.ix, RAD.

Copera marginipes (Rambur, 1842)

Fig. 8 shows two males attempting to form tandem with the same female.

2 - 3, 12.ix, CYC; 3, 12.ix, RAD.

Copera vittata (Selys, 1863)\*

8 – ♂, 14.ix, RAD.

Indocnemis orang (Förster in Laidlaw, 1907)\*\*

- **4** − ♂, ♀, 13.ix, CYC; 2 ♂♂, 13.ix, RAD.
- **5** ♂, 13.ix, RAD.
- **9** ♂, 16.ix, RAD.

Prodasineura humeralis (Selys, 1860)\*

- 1 3, 12.ix, CYC; 3, 12.ix, RAD.
- **9** ♂, 15.ix, CYC.

Prodasineura laidlawi (Förster in Laidlaw, 1907)

- **2** 2 ♂♂, 12.ix, CYC; ♂, ♂+♀, 12.ix, RAD.
- **4** − ♂, 13.ix, CYC.
- 9 3, 16.ix, CYC.

Prodasineura notostigma (Selys, 1860)

8 – ♂, 14.ix, RAD.



Figure 8: Copera marginipes triple tandem. Photo by C.Y. Choong.

### COENAGRIONIDAE

Argiocnemis species\*\*

**4** – ♂, 13.ix, RAD.

Ceriagrion cerinorubellum (Brauer, 1865)

**3** − ♂, 12.ix, RAD.

Ischnura senegalensis (Rambur, 1842)

**6** – ♂, 13.ix, RAD.

Pseudagrion australasiae Selys, 1876\*

3 – ♂, 12.ix, RAD.

Pseudagrion microcephalum (Rambur, 1842)

3 – ♂, 12.ix, CYC.

Pseudagrion pruinosum (Burmeister, 1839)

**1** – ♂, 12.ix, CYC; ♂, 12.ix, RAD.

Pseudagrion williamsoni Fraser, 1922\*\*

**3** – 3, 12.ix, CYC; 2 33, 12.ix, RAD.

### **Anisoptera**

**GOMPHIDAE** 

Gomphidia abbotti Williamson, 1907\*

Common on the main stream at Hutan Lipur Ulu Paip between 10 and 11 in the morning on the day the location was visited.

1 - 2 33, 12.ix, CYC; 2 33, 12.ix, RAD.

Gomphidictinus perakensis (Laidlaw, 1902)

This species was common at several of the sites visited during our visit.




Figure 9: Nepogomphus fruhstorferi male. Photo by C.Y. Choong.

- **4** ♂, 13.ix, CYC; ♂, 13.ix, RAD.
- 8 2 33, 14.ix, RAD.
- 9 2 ♂♂, 15.ix, RAD.

Nepogomphus fruhstorferi (Lieftinck, 1934)\*\*

Fig. 9 shows a male.

- **4** ♂, 13.ix, CYC; ♀, 1.ix, RAD.
- 9 ♂, 15.ix, RAD.

Paragomphus capricornis (Förster, 1914)\*

1 – ♀. 12.ix. RAD.

### MACROMIIDAE

Macromia cf westwoodii Selys, 1874\*

The status of the form of M. westwoodii occurring in Peninsular Malaysia is not entirely clear.

**9** – 2 ♂♂, 15.ix, RAD; ♀, 16.ix, RAD.

### LIBELLULIDAE

Acisoma panorpoides Rambur, 1842

- 3 ♂, 12.ix, CYC; ♂, 12.ix, RAD.
- **6** ♂, 13.ix, RAD.

Aethriamanta gracilis (Brauer, 1878)

3 – ♂, 12.ix, RAD.

Brachydiplax chalybea Brauer, 1868

**6** – ♂, 13.ix, RAD.

Brachythemis contaminata (Fabricius, 1793)

**3** – ♂, 12.ix, CYC.

Cratilla metallica (Brauer, 1878)

**4** − ♂, 13.ix, CYC.

Crocothemis servillia (Drury, 1770)

**6** – ♂, 13.ix, RAD.

Diplacodes trivialis (Rambur, 1842)

- **6** ♀, 13.ix, RAD.
- **7** ♀, 14.ix, CYC.

Lyriothemis biappendiculata (Selys, 1878)\*

- 1 ♀, 12.ix, RAD.
- 9 ♂, 16.ix, CYC.

Neurothemis fluctuans (Fabricius, 1793)

- **3** ♂, 12.ix, CYC; ♂, 12.ix, RAD.
- **6** ♂, 13.ix, RAD.

Neurothemis fulvia (Drury, 1773)

**4** – ♀, 13.ix, CYC.

Onychothemis culminicola Förster, 1904\*\*

1 - 2 33, 12.ix, CYC; 3, 12.ix, RAD.

**4** − ♂, 13.ix, CYC.

Onychothemis testacea Laidlaw, 1902

**1** − ♂, 12.ix, RAD.

Orthetrum glaucum (Brauer, 1865)

**4** – ♂, 13.ix, RAD.

Orthetrum luzonicum (Brauer, 1868)

3 - ♂, 12.ix, CYC.

Orthetrum sabina (Drury, 1770)

**6** – ♂, 13.ix, RAD.

Potamarcha congener (Rambur, 1842)\*

**6** – ♂, 13.ix, RAD.

Rhyothemis phyllis (Sulzer, 1776)

**3** – ♂, 12.ix, RAD.

6 – ♂. 13.ix. RAD.

Rhyothemis triangularis Kirby, 1889\*

3 – ♂, 12.ix, RAD.

Tholymis tillarga (Fabricius, 1798)

**6** – ♀, 13.ix, RAD.

Trithemis festiva (Rambur, 1842)

**1** − ♂, 12.ix, RAD.

9 – ♂, 15.ix, CYC.

Tyriobapta torrida Kirby, 1889\*

**4** − ♂, 13.ix, RAD.

8 – ♂, 14.ix, RAD.

Zygonyx iris Selys, 1869\*

Fig. 10 shows a male.

1 – 2 33, 12.ix, CYC; 3, 12.ix, RAD.

**4** − ♂, 13.ix, CYC; ♂, 13.ix, RAD.

**9** – 3, 15.ix, CYC; 3, 15.ix, RAD; 3, 16.ix, CYC.

Zygonyx ida Selys, 1869

9 – ♂, 16.ix, CYC.

### Discussion

We were able to collect 64 species during our brief trip to Kedah; one of these was a new species (*Drepanosticta rahmani*). No less than 13–14 (taking account of the



Figure 10: Zygonyx iris male. Photo by C.Y. Choong.

uncertainty over whether or not Vestalis amoena has been recorded in Kedah before) of these are first records for the state, and 28–29 are new records for mainland Kedah. It is remarkable that almost half of the species collected had not been recorded in mainland Kedah until now (although possibly there are records in the Malaysian or maybe Japanese literature that we are not aware of). With the new records, 126 species have been recorded in Kedah, 95 from the mainland and 85 from Langkawi, one species (Rhodothemis rufa (Rambur, 1842)) has only been recorded from Pulau Perak.

## **Acknowledgements**

We are grateful to Forestry Department Peninsular Malaysia for granting us a permit to collect Odonata in Forest Reserves. The Mohamed bin Zayed Species Conservation Fund provided the grant that supported the field work during reported here. The Economic Planning Unit of Malaysia gave the first author permission to conduct research in Peninsular Malaysia.

#### References

Asahina, S., 1966. Malayan dragonflies taken by Mr. E. Kawase in 1960/1962. Kontyû 34: 18–21.

Asahina, S., 1967. A Revision of the Asiatic Species of the Damselflies of the Genus Ceriagrion (Odonata, Agrionidae). Japanese Journal of Zoology 15(3): 255–334.

- Brooks, S.J., 1981. Dragonfly records from West Malaysia. Notulae odonatologicae 1(7): 113–114.
- Butler, S.G., 2011. Description of the last instar larvae of *Ictinogomphus acutus* (Laidlaw) from Sarawak, with a key to the larvae of the congeneric species (Anisoptera: Gomphidae). Odonatologica 40(2): 123–129.
- Butler, S.G. & H.-U. Kohler, 2013. A preliminary list of Odonata from Langkawi Island, Straits of Malacca. Notulae Odonatologicae 8(1): 4–8.
- Che Salmah, M.R., A. Abu Hassan & Z.S. Ameilia, 2004. Odonate communities (Odonata: Insecta) in a tropical river basin, Malaysia. Wetland Science 2(1): 1–9.
- Che Salmah, M.R., S.T.S. Hassan & A. Abu Hassan, 2000. Local movement and feeding pattern of adult Neurothemis tullia (Drury) (Odonata: Libellulidae) in a rain fed rice field. Tropical Ecology 41(2): 233–241.
- Che Salmah, M.R. & A. Wahizatul Afzan, 2004. Distribution of Odonata (Insecta) in various ecosystems in northern Peninsular Malaysia. Wetland Science 2(3): 184–191.
- Choong, C.Y., A.G. Idris & S. Azman, 2006. Dragonflies (Order: Odonata) of Gunung Jerai. In: Ismail, S.M., W.Y.W. Ahmad, J.M. Som, Y. Muda & A. Latiff (Eds.) Hutan Simpan Gunung Jerai, Kedah. Pengurusan Hutan, Persekitaran Fizikal dan Kepelbagaian Biologi. 378–381.
- Donnelly, N. [T.W.], 1998. Back to Thailand and Malaysia Farangpo 98. Malangpo 15: 137-142.
- Dow, R.A., 2010. A review of the Teinobasis of Sundaland, with the description of Teinobasis cryptica sp. nov. from Malaysia (Odonata: Coenagrionidae). International Journal of Odonatology 13(2): 205–230, plate II.
- Dow, R.A., C.Y. Choong & Y.F. Ng, 2016. Records of Odonata from Perlis, Malaysia in August 2015, with a checklist of species recorded from the state. Faunistic Studies in South–East Asian and Pacific Island Odonata 16: 1–22.
- Dow, R.A., C.Y. Choong & Y.F. Ng, 2017. *Drepanosticta rahmani* sp. nov., from Kedah, Malaysia (Odonata: Zygoptera: Platystictidae). Zootaxa 4338(1): 44–50.
- Farizawati, S., Fauzi, M., Ruslan, M.Y., Norma–Rashid, Y., Ng, Y.F. & Idris, A.G., 2014. The Diversity of Odonates in Five Islands within the West Coast of Peninsular Malaysia. Academic Journal of Entomology 7(3): 102–108.
- Fraser, F.C., 1935. New Oriental Dragonflies (Order Odonata). Records of the Indian Museum 37(3): 321–333.
- Fraser, F.C., 1942. Dr. Raymond Wheeler's Collection of Odonata from the Federated Malay States, with the Descriptions of a New Genus and Two New Species. The Proceedings of the Royal Entomological Society of London, Series B. Taxonomy 11(7): 95–104.
- Inoue, K., 1974. Malaysian dragonflies taken by Mr. Kiyoshi Ohkubo. Gracile 17: 11–14.
- Kosterin, O.E. 2014. Odonata of the south-west and north-east of Cambodia as studied in early rainy season of 2013. International Dragonfly Fund Report 67: 1–94.

- Laidlaw, F.F., 1902a. On a collection of dragonflies made by members of the Skeat expedition in the Malay Peninsula in 1899-1900. Proceedings of the Zoological Society of London 1902: 63-92, plates V-VI.
- Laidlaw, F.F., 1902b. On a collection of dragonflies made by members of the Skeat expedition in the Malay Peninsula in 1899–1900. Part II Coenagrioninae. Proceedings of the Zoological Society of London 1902: 381–389.
- Laidlaw, F.F., 1931. Dragonflies of the Malay Peninsula with descriptions of new species. Journal of the Federated Malay Museums 16(3/4): 175–233.
- Laidlaw, F.F., 1932. Notes on some Malaysian dragonflies, with descriptions of new species. Bulletin of the Raffles Museum, Singapore 7: 95–98.
- Laidlaw, F.F., 1934. A Note of the dragonfly fauna (Odonata) of Mount Kinabalu and of some other mountain areas of Malaysia, with a description of some new or little known species. Journal of the Federated Malay States Museums 17(3): 549–561.
- Lieftinck, M.A., 1933. Notes on the Larvae of Two Interesting Gomphidae (Odon.) from the Malay Peninsula. Bulletin of the Raffles Museum 7: 101–115, plates II–IV.
- Lieftinck, M.A., 1954. Handlist of Malaysian Odonata. A catalogue of the dragonflies of the Malay Peninsula, Sumatra, Java and Borneo, including the adjacent small islands. Treubia (Suppl.) 22: i–xiii + 1–202.
- Lieftnck, M.A., 1965a. Some Odonata of the genus *Drepanosticta* Laidlaw, chiefly from the Malay peninsula (Platystictidae). Zoologische Mededelingen 40(21): 171–186.
- Lieftinck, M.A., 1965b. The species–group of Vestalis amoena Selys, 1853, in Sundaland (Odonata, Calopterygidae). Tijdschrift voor Entomologie 108(11): 325–364.
- Norma-Rashid, Y., Sofian–Azirun, M., Rosli, R. & Rosli, H., 2008. Dragonflies on the islands in the Straits of Malacca. Malaysia Journal of Science 27 (3): 105–111.
- Novelo–Gutiérrez, R. & M.R. Che Salmah, 2006. The larva of *Macromia cincta* Rambur, with a key to the known *Macromia* larvae of the Malaysian Peninsula (Anisoptera: Macromiidae). Odonatologica 35(1): 61–66.
- Orr, A.G., 2005. A pocket guide: Dragonflies of Peninsular Malaysia and Singapore. Natural History Publications (Borneo), Kota Kinabalu, vi + 125 pp.
- Tsuda, S. and Kitagawa, K., 1987. Odonata of southeast Asia collected by the late Mr. M. Iwasaki. Part 1. Odonata of Malay Peninsula (West Malaysia) and Singapore. Gracile 38: 1–7.
- Wahizatul–Afzan, A. & Che Salmah, M.R., 2005. Adult dragonfly communities (Odonata: Insecta) in a tropical rivers of the northern peninsular Malaysia: species composition, biotope and host plant preferences. Wetland Science 3(3): 167–175.

## Appendix: Checklist of Odonata recorded from Kedah

Where the first record from either Kedah as a whole or mainland Kedah is not in this publication, a citation to the first record is made in square parenthesis after the species name. **M**: mainland. **L**: Langkawi Island.

## Zygoptera

### Lestidae

Lestes praemorsus decipiens Kirby, 1894 [M: this publication; L: Butler & Kohler 2013]

Platylestes heterostylus Lieftinck, 1932 [L: Butler & Kohler 2013]

Orolestes wallacei (Kirby, 1889) [M: Laidlaw 1902a as Lestes ridleyi Laidlaw, 1902]

### Platystictidae

Drepanosticta fontinalis Lieftinck, 1937 [M: Lieftinck 1965a; note 1]

Drepanosticta rahmani Dow, Choong & Ng, 2017 [M: Dow et al. 2017]

Drepanosticta silenus Laidlaw, 1934 [M: this publication]

Protosticta foersteri Laidlaw, 1902 [M: Laidlaw 1902b]

### Argiolestidae

Podolestes orientalis Selys, 1862 [M: this publication]

### Calopterygidae

Echo modesta Laidlaw, 1902 [M: this publication; L: Fraser 1935]

Neurobasis chinensis (Linnaeus, 1758) [M: Laidlaw 1931; L: Butler & Kohler 2013]

Vestalis amethystina Lieftinck, 1965 [M: Lieftinck 1965b]

Vestalis amoena Hagen in Selys, 1853 [M: this publication?; note 2]

Vestalis gracilis (Rambur, 1842) [M: Laidlaw 1931; L: Tsuda & Kitagawa 1987]

### Chlorocyphidae (note 3)

Aristocypha fenestrella (Rambur, 1842) [M: Choong et al. 2006]

Heliocypha biforata (Selys, 1859) [M: this publication; L: Inoue 1974]

Heliocypha perforata (Percheron, 1835) [M: Laidlaw 1902a as Rhinocypha inas Laidlaw, 1902]

Libellago lineata (Burmeister, 1839) [M: Fraser 1942]

Sundacypha petiolata (Selys, 1859) [M: this publication]

### Devadattidae

Devadatta argyoides (Selys, 1859) [M: Laidlaw 1931]

### Euphaeidae

Dysphaea dimidiata Selys, 1853 [M: this publication]

Euphaea ochracea Selys, 1859 [M: Choong et al. 2006]

#### Philosinidae

Rhinagrion viridatuma Fraser, 1938 [M: this publication; L: Butler & Kohler 2013 as R. mima (Karsch, 1891)]

### Platycnemididae

Calicnemia chaseni (Laidlaw, 1928) [M: this publication]

Coeliccia albicauda (Förster in Laidlaw, 1907) [M: this publication]

Coeliccia didyma (Selys, 1863) [M: Choong et al. 2006 as Coeliccia nemoricola Laid-

law, 1912; L: Tsuda & Kitagawa 1987]

Coeliccia erici Laidlaw, 1917 [M: this publication]

Coeliccia octogesima (Selys, 1863) [M: this publication]

Copera marginipes (Rambur, 1842) [M: Wahizatul Afzan & Che Salmah 2005; L: Tsuda & Kitagawa 1987]

Copera vittata (Selys, 1863) [M: this publication; L: Butler & Kohler 2013]

Indocnemis orang (Förster in Laidlaw, 1907) [M: this publication]

Onychargia atrocyana Selys, 1865 [M: Laidlaw 1934]

Prodasineura humeralis (Selys, 1860) [M: this publication; L: Butler & Kohler 2013]

Prodasineura laidlawi (Förster in Laidlaw, 1907) [M: Lieftinck 1954; L: Butler & Kohler 2013]

Prodasineura notostigma (Selys, 1860) [M: Laidlaw 1934]

Pseudocopera ciliata (Selys, 1863) [**M:** Wahizatul Afzan & Che Salmah 2005; **L:** Butler & Kohler 2013]

### Coenagrionidae

Aciagrion borneense Ris. 1911 [M: Laidlaw 1934; L: Butler & Kohler 2013]

Aciagrion hisopa (Selys, 1876) [L: Tsuda & Kitagawa 1987]

Agriocnemis femina (Brauer, 1868) [M: Asahina 1966; L: Tsuda & Kitagawa 1987]

Agriocnemis minima Selys, 1877 [M: Asahina 1966 as A. d'abreui Fraser, 1919]

Agriocnemis nana (Laidlaw, 1914) [**M:** Wahizatul Afzan & Che Salmah 2005; **L:** Tsuda & Kitagawa 1987]

Agriocnemis pygmaea (Rambur, 1842) [M: Asahina 1966; L: Fraser 1942]

Argiocnemis rubescens rubeola Selys, 1877 [M: Asahina 1966; L: Tsuda & Kitagawa 1987]

Argiocnemis species [M: this publication]

Ceriagrion auranticum Fraser, 1922 [M: Asahina 1967; L: Butler & Kohler 2013]

Ceriagrion calamineum Lieftinck, 1951 [L: Tsuda & Kitagawa 1987; note 4]

Ceriagrion cerinorubellum (Brauer, 1865) [M: Asahina 1966; L: Asahina 1966]

Ceriagrion olivaceum Laidlaw, 1914 [M: Asahina 1967]

Ceriagrion praetermissum Lieftinck, 1929 [L: Farizawati et al. 2014]

Ischnura senegalensis (Rambur, 1842) [M: Asahina 1966; L: Tsuda & Kitagawa 1987]

Paracercion calamorum (Ris, 1916) [L: Dow et al. 2016]

Pericnemis stictica Hagen in Selys, 1863 [M: Laidlaw 1902b]

Pseudagrion australasiae Selys, 1876 [M: this publication; L: Butler & Kohler 2013]

Pseudagrion microcephalum (Rambur, 1842) [M: Wahizatul Afzan & Che Salmah 2005;

L: Farizawati et al. 2014]

Pseudagrion pruinosum (Burmeister, 1839) [M: Wahizatul Afzan & Che Salmah 2005]

Pseudagrion williamsoni Fraser, 1922 [M: this publication]

Teinobasis kirbyi Laidlaw, 1902 [M: Laidlaw 1902b]

## **Anisoptera**

### Aeshnidae

Anax guttatus (Burmeister, 1839) [L: Tsuda & Kitagawa 1987]

Gynacantha bayadera Selys, 1891 [L: Butler & Kohler 2013]

Heliaeschna idae (Brauer, 1865) [M: Laidlaw 1934]

Indaeschna grubaueri (Förster, 1904) [M: Laidlaw 1902a as Amphiaeschna ampla (Rambur, 1842); L: Butler & Kohler 2013]

### Gomphidae

Acrogomphus malayanus Laidlaw, 1925 [L: Butler & Kohler 2013]

Gomphidia abbotti Williamson, 1907 [M: this publication; L: Butler & Kohler 2013]

Gomphidictinus perakensis (Laidlaw, 1902) [M: Laidlaw 1902a]

Heliogomphus kelantanensis (Laidlaw, 1902) [M: Lieftinck 1933]

Ictinogomphus decoratus melaenops Selys, 1857 [M: Wahizatul Afzan & Che Salmah

2005 as Ictinogomphus rapax (Rambur, 1842); L: Butler 2011]

Leptogomphus risi Laidlaw, 1932 [M: Laidlaw 1932]

Macrogomphus species [L: Butler & Kohler 2013]

Megalogomphus icterops (Martin, 1903) [L: Butler & Kohler 2013]

Merogomphus parvus (Krüger, 1899) [L: Butler & Kohler 2013]

Microgomphus chelifer Selys, 1858 [L: Butler & Kohler 2013]

Nepogomphus fruhstorferi (Lieftinck, 1934) [M: this publication]

Paragomphus capricornis (Förster, 1914) [M: this publication; L: Butler & Kohler 2013]

Sieboldius japponicus Selys, 1854 [M: Laidlaw 1902a as S. grandis Krüger, 1899]

## Macromiidae

Epophthalmia vittigera [L: Butler & Kohler 2013]

Macromia cincta Rambur, 1842 [M: Novelo-Gutierrez & Che Salmah 2006]

Macromia cydippe Laidlaw, 1922 [L: Butler & Kohler 2013]

Macromia ?gerstaeckeri Krüger, 1899 [L: Butler & Kohler 2013]

Macromia of westwoodii Selys, 1874 [M: this publication; L: Butler & Kohler 2013 as 2 westwoodii]

as ?westwoodii]

### Synthemistidae

Idionyx yolanda Selys, 1871 [L: Butler & Kohler 2013]

Macromidia genialis Laidlaw, 1923 [L: Butler & Kohler 2013]

### Libellulidae

Acisoma panorpoides Rambur, 1842 [M: Asahina 1966; L: Butler & Kohler 2013]

Aethriamanta aethra Ris, 1912 [L: Farizawati et al 2014]

Aethriamanta gracilis (Brauer, 1878) [M: Choong et al. 2006; L: Butler & Kohler 2013]

Brachydiplax chalybea Brauer, 1868 [M: Wahizatul Afzan & Che Salmah 2005; L: Butler & Kohler 2013]

Brachydiplax farinosa Krüger, 1902 A [M: Wahizatul Afzan & Che Salmah 2005]

Brachygonia oculata (Brauer, 1878) [M: Fraser 1942]

Brachythemis contaminata (Fabricius, 1793) [M: Asahina 1966; L: Tsuda & Kitagawa 1987]

Camacinia gigantea (Brauer, 1867) [L: Butler & Kohler 2013]

Cratilla lineata (Brauer, 1878) [M: Laidlaw 1934; L: Tsuda & Kitagawa 1987]

Cratilla metallica (Brauer, 1878) [M: Laidlaw 1902a; L: Tsuda & Kitagawa 1987]

Crocothemis servilia (Drury, 1770) [M: Wahizatul Afzan & Che Salmah 2005; L: Tsuda & Kitagawa 1987]

Diplacodes nebulosa (Fabricius, 1793) [L: Tsuda & Kitagawa 1987]

Diplacodes trivialis (Rambur, 1842) [**M:** Wahizatul Afzan & Che Salmah 2005; **L:** Tsuda & Kitagawa 1987]

Hydrobasileus croceus (Brauer, 1867) [L: Butler & Kohler 2013]

Indothemis limbata (Selys, 1891) [L: Butler & Kohler 2013]

Lathrecista asiatica (Fabricius, 1798) [M: Wahizatul Afzan & Che Salmah 2005; L: Tsuda & Kitagawa 1987]

Lyriothemis biappendiculata (Selys, 1878) [M: this publication; L: Butler & Kohler 2013]

Macrodiplax cora (Brauer, 1867) [L: Butler & Kohler 2013]

Neurothemis fluctuans (Fabricius, 1793) [M: Asahina 1966; L: Butler & Kohler 2013]

Neurothemis fulvia (Drury, 1773) [M: Fraser 1942; L: Inoue 1974]

Neurothemis tullia (Drury, 1773) [M: Laidlaw 1934]

Onychothemis culminicola Förster, 1904 [M: this publication]

Onychothemis testacea Laidlaw, 1902 [M: Wahizatul Afzan & Che Salmah 2005]

Orthetrum chrysis (Selys, 1891) [**M:** Wahizatul Afzan & Che Salmah 2005; **L:** Butler & Kohler 2013]

Orthetrum glaucum (Brauer, 1865) [M: Laidlaw 1934; L: Butler & Kohler 2013]

Orthetrum Iuzonicum (Brauer, 1868) [M: Wahizatul Afzan & Che Salmah 2005; L: Butler & Kohler 2013]

Orthetrum sabina (Drury, 1770) [M: Fraser 1942; L: Inoue 1974]

Orthetrum testaceum (Burmeister, 1839) [L: Butler & Kohler 2013]

Pantala flavescens (Fabricius, 1798) [M: Wahizatul Afzan & Che Salmah 2005; L: Butler & Kohler 2013]

Potamarcha congener (Rambur, 1842) [M: this publication; L: Tsuda & Kitagawa 1987]

Pseudothemis jorina Förster, 1904 [L: Butler & Kohler 2013]

Rhodothemis rufa (Rambur, 1842) [Pulau Perak: Norma-Rashid et al. 2008]

Rhyothemis obsolescens Kirby, 1889 [M: Laidlaw 1934]

Rhyothemis phyllis (Sulzer, 1776) [M: Fraser 1942; L: Butler & Kohler 2013]

Rhyothemis triangularis Kirby, 1889 [M: this publication; L: Butler & Kohler 2013]

Tetrathemis irregularis hyalina Kirby, 1889 [L: Butler & Kohler 2013]

Tholymis tillarga (Fabricius, 1798) [M: Asahina 1966; L: Tsuda & Kitagawa 1987]

Tramea transmarina euryale Selys, 1878 [L: Tsuda & Kitagawa 1987]

*Trithemis aurora* (Burmeister, 1839) [**M:** Wahizatul Afzan & Che Salmah 2005; **L:** Butler & Kohler 2013]

Trithemis festiva (Rambur, 1842) [M: Wahizatul Afzan & Che Salmah 2005; L: Tsuda & Kitagawa 1987]

Trithemis pallidinervis (Kirby, 1889) [L: Tsuda & Kitagawa 1987]

Tyriobapta torrida Kirby, 1889 [M: this publication; L: Butler & Kohler 2013]

Urothemis signata insignata (Selys, 1872) [L: Butler & Kohler 2013]

Zygonyx ida Hagen, 1867 [M: Choong et al. 2006 as Z. iris]

Zygonyx iris Selys, 1869 [L: Tsuda & Kitagawa 1987]

Zyxomma obtusum Albarda, 1881 [L: Butler & Kohler 2013]

- **Note 1:** Laidlaw (1934) recorded *Drepanosticta* ? *quadrata* (Selys, 1860) from Kedah. It seems likely that this really refers to *D. fontinalis*; we certainly do not think that it refers to *D. quadrata*.
- **Note 2:** Laidlaw (1902a) recorded Vestalis amoena from Kedah, in reality this could refer to either V. amoena or V. amethystina.
- **Note 3:** Inoue (1974) recorded a *Libellago* sp. from Langkawi, in the absence of further information it is not possible to say to what species this record refers.
- **Note 4:** Ceriagrion coromandelianum (Fabricius, 1798) was listed from Kedah by Laidlaw (1934); this species is not otherwise known to occur in Malaysia and was not listed in either Lieftinck (1954) or Orr (2005). We note that various authors had identified C. calamineum with C. coromandelianum until the former was described, so that this record may refer to C. calamineum.

#### INSTRUCTION TO AUTHORS

Faunistic studies of South-East Asian and Pacific islands Odonata is a journal of the International Dragonfly Fund (IDF). It is referred to as the journal in the remainder of these instructions. Transfer of copyright to IDF is considered to have taken place implicitly once a paper has been published in the journal.

The journal publishes original papers only. By original is meant papers that: a) have not been published elsewhere before, and b) the scientific results of the paper have not been published in their entirety under a different title and/or with different wording elsewhere. The republishing of any part of a paper published in the journal must be negotiated with the Editorial Board and can only proceed after mutual agreement.

Papers reporting studies financially supported by the IDF will be reviewed with priority, however, authors working with Odonata from the focal area (as defined on the back page of the front cover) are encouraged to submit their manuscripts even if they have not received any funds from IDF.

Manuscripts submitted to the journal should preferably be in English; alternatively German or French will also be accepted. Every manuscript should be checked by a native speaker of the language in which it is written; if it is not possible for the authors to arrange this, they must inform the Editorial Board on submission of the paper. Authors are encouraged, if possible, to include a version of the abstract in the primary language of the country in which their study was made.

Authors can choose the best way for them to submit their manuscripts between these options: a) via e-mail to the publisher, or b) on a CD, DVD or any other IBM-compatible device. Manuscripts should be prepared in Microsoft Word for Windows.

While preparing the manuscript authors should consider that, although the journal gives some freedom in the style and arrangements of the sections, the editors would like to see the following clearly defined sections: Title (with authors names, physical and e-mail addresses), Abstract, Introduction, Material & Methods, Results, Discussion, Acknowledgments and References. This is a widely used scheme by scientists that everyone should be familiar with. No further instructions are given here, but every author should check the style of the journal.

Authors are advised to avoid any formatting of the text. The manuscripts will be stylised according to the font type and size adopted by the journal. However, check for: a) all species names must be given in italic, b) the authority and year of publication are required on the first appearance of a species name in the text, but not thereafter, and c) citations and reference list must be arranged following the format below.

Reference cited in the text should read as follows: Tillyard (1924), (Tillyard 1924), Swezey & Williams (1942).

The reference list should be prepared according to the following standard:

Swezey, O. & F. Williams, 1942. Dragonflies of Guam. Bernice P. Bishop Museum Bulletin 172: 3-6.

Tillyard, R., 1924. The dragonflies (Order Odonata) of Fiji, with special reference to a collection made by Mr. H.W. Simmonds, F.E.S., on the Island of Viti Levu. Transactions of the Entomological Society London 1923 III-IV: 305-346.

Citations of internet sources should include the date of access.

The manuscript should end with a list of captions to the figures and tables. The latter should be submitted separately from the text preferably as graphics made using one of the Microsoft Office products or as a high resolution picture saved as a .jpg .tif or .ps file. Pictures should be at least 11 cm wide and with a minimum 300 dpi resolution, better 360 dpi. Line drawings and graphics could have 1200 dpi for better details. If you compose many pictures to one figure, please submit the original files as well. Please leave some space in the upper left corner of each picture, to insert a letter (a, b, c...) later. Hand-made drawings should be scanned and submitted electronically. Printed figures sent by the post could be damaged, in which case authors will be asked to resubmit them.

Manuscripts not arranged according to these instructions may also be accepted, but in that case their publication will be delayed until the journal's standards are achieved.

