

Supplementary Table 3. Previously published genome size (2C-values), derived Cx values (based on the 2C threshold for diploids) and GC content estimates of the subfamily Bromelioideae and Puyoideae. Values marked with * were used for the analyses of this study. Nomenclature follows Gouda et al. (cont. updated) except the genus *Ananas* and *Pseudananas*.

BROMELIOIDEAE					
Taxon	2C [pg]	SD	Derived Cx [pg]	GC content [%]	Reference
<i>Aechmea</i> Ruiz & Pav.					
<i>A. angustifolia</i> Poepp. Endl.	0.69*	±0.005	0.345*		Müller et al. 2019
<i>A. aquilega</i> (Salisb.) Griseb	1.08*	±0.013	0.540*		Gitaí et al., 2014
<i>A. bracteata</i> (Sw.) Griseb.	0.81*	±0.005	0.405*		Müller et al. 2019
<i>A. coelestis</i> (K.Koch) E.Morren	0.96*	±0.004	0.480*		Müller et al. 2019
<i>A. eurycorymbus</i> Harms	2.19*	±0.013	0.5473*		Gitaí et al., 2014
<i>A. fendleri</i> André ex Mez	1.25*	±0.018	0.625*		Gitaí et al., 2014
<i>A. filicaulis</i> (Griseb.) Mez	1.67 1.77	±0.020 ±0.002		44.00	Šmarda et al., 2014 Müller et al. 2019
<i>A. fosteriana</i> L.B.Sm.	0.93*	±0.001	0.465*		Müller et al. 2019
<i>A. mexicana</i> Baker	0.76	±0.009			Müller et al. 2019
<i>A. nudicaulis</i> (L.) Griseb. var. <i>cuspidata</i> Baker	0.78	±0.000		35.12	Favoreto et al., 2012
<i>A. racinæ</i> L.B.Sm	0.88	±0.016			Müller et al. 2019
<i>A. ramosa</i> Mart. ex Schult.f. & Schult.f.	1.37*	±0.018	0.685*	38.63*	Favoreto et al., 2012
<i>A. victoriana</i> L.B.Sm.	0.92*	±0.002	0.460*		Müller et al. 2019
<i>A. weilbachii</i> Didr.	1.10*	±0.001	0.550*		Müller et al. 2019
<i>Ananas</i> Mill.					
<i>A. bracteatus</i> (Lindl.) Schult. & Schult.f.	0.92				Arumuganathan & Earle, 1991
<i>A. comosus</i> (L.) Merr.	1.09 0.93				Arumuganathan & Earle, 1991 Müller et al. 2019
<i>Billbergia</i> Thunb.					
<i>B. amoena</i> (Lodd.) Lindl.	0.74	±0.002			Müller et al. 2019
<i>B. brasiliensis</i> L.B.Sm. [also published as <i>Billbergia velascana</i> Cárdenas]	0.91* 0.90	±0.005 ±0.011	0.455*		Müller et al. 2019 Müller et al. 2019
<i>B. euphemiae</i> E.Morren	0.89*	±0.014	0.445*	37.42*	Favoreto et al., 2012
<i>B. horrida</i> Regel	0.77*	±0.005	0.385*		Favoreto et al., 2012
<i>B. horrida</i> Regel var. <i>tigrina</i> hort. ex Baker	0.85*	±0.014	0.425*		Gitaí et al., 2014
<i>B. magnifica</i> Mez	0.95*	±0.014	0.475*		Müller et al. 2019
<i>B. nutans</i> H.Wendl. ex Regel	0.7517	±0.0567			Ramírez-Morillo & Brown, 2001
<i>B. pallidiflora</i> Liebm.	1.03*	±0.009	0.515*		Gitaí et al., 2014
<i>B. pyramidalis</i> (Sims) Lindl.	0.90*	±0.003	0.450*		Müller et al. 2019
<i>B. tweedieana</i> Baker	0.95	±0.014		33.95	Favoreto et al., 2012
<i>B. viridiflora</i> H.Wendl.	0.66*	±0.002	0.330*		Müller et al. 2019
<i>Bromelia</i> L.					
<i>B. antiacantha</i> Bertol.	0.81*	±0.014	0.405*	38.43*	Favoreto et al., 2012
<i>Canistropis</i> Mez					
<i>C. billbergioides</i> (Schult.f.) Leme [also published as <i>Nidularium billbergioides</i> (Schult. & Schult.f.) L.B.Sm.]	1.00 1.02	±0.005		37.36	Favoreto et al., 2012 Müller et al. 2019
<i>Cryptanthus</i> Otto & A.Dietr.					
<i>C. acaulis</i> (Lindl.) Beer	1.38*	±0.0283			Ramírez-Morillo & Brown, 2001

	1.38			Cruz et al., 2020
<i>C. alagoanus</i> Leme & J.A.Siqueira	0.78			Cruz et al., 2020
<i>C. arelii</i> H.Luther	0.98			Cruz et al., 2020
<i>C. bahianus</i> L.B.Sm.	0.75	±0.0683		Ramírez-Morillo & Brown, 2001
<i>C. beuckeri</i> E.Morren	1.458*	±0.0179		Ramírez-Morillo & Brown, 2001
	1.46			Cruz et al., 2020
<i>C. bromelioides</i> Otto & A.Dietr.	1.52			Cruz et al., 2020
<i>C. burle-marxii</i> Leme	1.32			Cruz et al., 2020
<i>C. capitellatus</i> Leme & L.Kollmann	1.36			Cruz et al., 2020
<i>C. colnagoi</i> Rauh & Leme	1.34			Cruz et al., 2020
<i>C. coriaceus</i> Leme	0.84			Cruz et al., 2020
<i>C. correia-araujoi</i> Leme	1.66			Cruz et al., 2020
<i>C. delicatus</i> Leme	1.46			Cruz et al., 2020
<i>C. dianae</i> Leme	1.36			Cruz et al., 2020
<i>C. giganteus</i> Leme & A.P.Fontana	1.38			Cruz et al., 2020
<i>C. lutherianus</i> I.Ramirez	0.90			Cruz et al., 2020
<i>C. lyman-smithii</i> Leme	1.38			Cruz et al., 2020
<i>C. marginatus</i> L.B.Sm.	1.26*	±0.005		Gitaí et al., 2014
	1.40			Cruz et al., 2020
<i>C. maritimus</i> L.B.Sm.	1.48			Cruz et al., 2020
<i>C. pickelii</i> L.B.Sm.	1.28			Cruz et al., 2020
<i>C. praetextus</i> E.Morren ex Baker	1.35*	±0.014		Gitaí et al., 2014
<i>C. reisii</i> Leme	0.88			Cruz et al., 2020
<i>C. sinuosus</i> L.B.Sm.	1.04			Cruz et al., 2020
<i>C. teretifolius</i> Leme	1.40			Cruz et al., 2020
<i>C. ubairensis</i> I.Ramirez	1.00			Cruz et al., 2020
<i>C. venecianus</i> Leme & L.Kollmann	1.32			Cruz et al., 2020
<i>C. viridipetalus</i> Leme	1.34			Cruz et al., 2020
<i>C. zonatus</i> (Vis.) Beer	1.04			Cruz et al., 2020
<i>C. sp.</i>	1.33*	±0.015	38.78*	Favoreto et al., 2012
<i>Edmundoa</i> Leme				
<i>E. lindenii</i> (Regel) Leme	1.32*	±0.010	0.662*	Gitaí et al., 2014
<i>Fascicularia</i> Mez				
<i>F. bicolor</i> (Ruiz & Pav.) Mez ssp. <i>bicolor</i>	1.05*	±0.010	0.527*	Gitaí et al., 2014
<i>F. bicolor</i> (Ruiz & Pav.) Mez ssp. <i>canaliculata</i> E.C.Nelson & Zizka	1.12*	±0.008	0.560*	Gitaí et al., 2014
<i>Forzzaea</i> Leme, S. Heller & Zizka				
<i>F. leopoldo-horstii</i> Leme, S.Heller & Zizka	1.14			Cruz et al., 2020
<i>Greigia</i> Regel				
<i>G. sphacelata</i> (Ruiz & Pav.) Regel	1.56*	±0.004	0.779*	Gitaí et al., 2014
<i>G. sp.</i>	1.35*	±0.019	0.664*	Gitaí et al., 2014
<i>Hohenbergia</i> Schult. & Schult.f.				
<i>H. aechmeoides</i> Leme [published as <i>Hohenberiga aechmeoides</i> Leme]	1.48			Gonçalves-Oliveira et al., 2020
<i>H. arcuata</i> Leme & M.Machado	1.42			Gonçalves-Oliveira et al., 2020
<i>H. augusta</i> (Vell.) E.Morren [published as <i>Hohenbergia augusta</i> Mez]	1.23			Gonçalves-Oliveira et al., 2020
	1.24			
<i>H. barbarespina</i> Leme & Fraga	1.37			Gonçalves-Oliveira et al., 2020
<i>H. belemii</i> L.B.Sm & Read	1.32			Gonçalves-Oliveira et al., 2020
<i>H. blanchetii</i> (Baker) Mez [published as <i>Hohenbergia blanchetti</i> É.Mooren]	1.40			Gonçalves-Oliveira et al., 2020
<i>H. burle-marxii</i> Leme & W.Till	1.39			Gonçalves-Oliveira et al., 2020
<i>H. capitata</i> Schult. & Schult.f. [published as <i>Hohenbergia capitata</i> Schult.f.]	1.50			Gonçalves-Oliveira et al., 2020

<i>H. castellanosii</i> L.B.Sm. & Read	1.19 1.23 1.24				Gonçalves-Oliveira et al., 2020
<i>H. catingae</i> Ule	1.29 1.41 1.42 1.48				Gonçalves-Oliveira et al., 2020
<i>H. edmundoi</i> L.B.Sm & Read	1.36				Gonçalves-Oliveira et al., 2020
<i>H. flava</i> Leme & C.C.Paula	1.39				Gonçalves-Oliveira et al., 2020
<i>H. horrida</i> Harms	1.17				Gonçalves-Oliveira et al., 2020
<i>H. igatuensis</i> Leme	1.41				Gonçalves-Oliveira et al., 2020
<i>H. lanata</i> E.Pereira & Moutinho	1.46				Gonçalves-Oliveira et al., 2020
<i>H. lativaginata</i> J.R.Macié & Louzada	1.22				Gonçalves-Oliveira et al., 2020
<i>H. lemei</i> H.Luther & K.Norton	1.32				Gonçalves-Oliveira et al., 2020
<i>H. leopoldo-horstii</i> E.Gross, Rauh & Leme	1.333				Gonçalves-Oliveira et al., 2020
<i>H. littoralis</i> L.B.Sm.	0.78				Gonçalves-Oliveira et al., 2020
<i>H. loredanoana</i> Leme & L.Kollmann	1.30				Gonçalves-Oliveira et al., 2020
<i>H. magnispina</i> Leme	1.27				Gonçalves-Oliveira et al., 2020
<i>H. minor</i> L.B.Sm.	1.28				Gonçalves-Oliveira et al., 2020
<i>H. pennae</i> E.Pereira	1.39				Gonçalves-Oliveira et al., 2020
<i>H. ramageana</i> Mez	1.42				Gonçalves-Oliveira et al., 2020
<i>H. reconcavensis</i> Leme & Fraga	1.35				Gonçalves-Oliveira et al., 2020
<i>H. ridleyi</i> (Baker) Mez	1.34				Gonçalves-Oliveira et al., 2020
<i>H. rosea</i> L.B.Sm & Read	1.33				Gonçalves-Oliveira et al., 2020
<i>H. stellata</i> Schult. & Schult.f.	1.33 1.34 1.36				Gonçalves-Oliveira et al., 2020
<i>H. utriculosa</i> Ule	1.52				Gonçalves-Oliveira et al., 2020
<i>H. vestita</i> L.B.Sm.	1.17				Gonçalves-Oliveira et al., 2020
<i>Hoplocryptanthus</i> (Mez) Leme, S. Heller & Zizka					
<i>H. caracensis</i> Leme, S.Heller & Zizka	0.78				Cruz et al., 2020
<i>H. ferrarius</i> Leme, S.Heller & Zizka	0.82				Cruz et al., 2020
<i>H. glaziovii</i> Leme, S.Heller & Zizka	0.80				Cruz et al., 2020
<i>H. regius</i> Leme, S.Heller & Zizka	0.78				Cruz et al., 2020
<i>H. schwackeanus</i> (Mez) Leme, S. Heller & Zizka [published as <i>Cryptanthus schwackeanus</i> Mez]	0.71*	±0.0141			Ramírez-Morillo & Brown, 2001
<i>H. schwackeanus</i> (Mez) Leme, S.Heller & Zizka	0.82				Cruz et al., 2020
<i>H. tiradentesensis</i> Leme, S.Heller & Zizka	0.80				Cruz et al., 2020
<i>Neoregelia</i> L.B.Sm.					
<i>N. carolinae</i> (Beer) L.B.Sm	0.96*	±0.028	0.480*		Müller et al. 2019
<i>N. aff. simulans</i> L.B.Sm.	0.98*	±0.004	0.490*	37.24*	Favoreto et al., 2012
<i>N. zonata</i> L.B.Sm.	0.93*	±0.001	0.465*		Müller et al. 2019
<i>Nidularium</i> Lem.					
<i>N. camposportoi</i> (L.B.Sm.) Wand. & B.A.Moreira [published as <i>Nidularium campos-portoi</i> (L.B.Sm.) Leme]	1.03*	±0.003	0.515*		Müller et al. 2019
<i>N. innocentii</i> Lem.	1.12*	±0.005	0.560*		Müller et al. 2019
<i>Ochagavia</i> Phil.					
<i>O. elegans</i> Phil.	1.12*	±0.015	0.557*		Gitaí et al., 2014
<i>Orthophytum</i> Beer					
<i>O. foliosum</i> L.B.Sm.	1.88*		0.313*		Müller et al. 2019
<i>O. saxicola</i> (Ule) L.B.Sm.	0.64*	±0.0141	0.320*		Ramírez-Morillo & Brown, 2001

Portea K.Koch.					
<i>P. petropolitana</i> (Wawra) Mez	1.31*	± 0.011	0.655*		Gitaí et al., 2014
Pseudananas Hassl. ex Harms					
<i>P. sagenarius</i> (Arruda) Camargo [published as <i>Ananas macrodontes</i> E.Morren]	1.00	± 0.021		39.24	Favoreto et al., 2012
Quesnelia Gaudich.					
<i>Q. arvensis</i> (Vell.) Mez	0.88	± 0.001			Gitaí et al., 2014
Rokautskyia Leme, S. Heller & Zizka					
<i>R. latifolia</i> Leme, S.Heller & Zizka	1.10				Cruz et al., 2020
<i>R. microglazioui</i> Leme, S.Heller & Zizka	0.94				Cruz et al., 2020
<i>R. odoratissima</i> Leme, S.Heller & Zizka	1.10				Cruz et al., 2020
<i>R. pseudoglaacioui</i> Leme, S.Heller & Zizka	1.00				Cruz et al., 2020
<i>R. pseudoscaposa</i> Leme, S.Heller & Zizka	1.04				Cruz et al., 2020
<i>R. sanctaluciae</i> Leme, S.Heller & Zizka	1.00				Cruz et al., 2020
<i>R. scaposa</i> Leme, S.Heller & Zizka	1.04				Cruz et al., 2020
<i>R. whitmanii</i> Leme, S.Heller & Zizka	1.06				Cruz et al., 2020
Ronnbergia					
<i>R. deleonii</i> L.B.Sm.	1.38*	± 0.024			Müller et al. 2019
<i>R. veitchii</i> (Baker) Aguirre-Santoro [published as <i>Aechmea veitchii</i> Baker]	0.77*	± 0.008	0.385*		Müller et al. 2019
<i>R. weberbaueri</i> (Harms) Aguirre-Santoro [published as <i>Aechmea weberbaueri</i> Harms]	1.32	± 0.008			Müller et al. 2019
<i>R. wuelfinghoffii</i> (E.Gross.) Aguirre-Santoro [published as <i>Aechmea wuelfinghoffii</i> E.Gross]	1.03*	± 0.002	0.515*		Müller et al. 2019
Sincoraea Ule					
<i>S. amoena</i> Leme, S.Heller & Zizka	1.04				Cruz et al., 2020
<i>S. heleniceae</i> Leme, S.Heller & Zizka	1.08				Cruz et al., 2020
<i>S. ophiuroides</i> Leme, S.Heller & Zizka	1.02				Cruz et al., 2020
<i>S. ulei</i> Leme, S.Heller & Zizka	1.50				Cruz et al., 2020
xHohenmea B.R.Silva & L.F.Sousa					
<i>xH. itaiipuana</i> B.R.Silva & L.F.Sousa	1.63				Gonçalves-Oliveira et al., 2020
PUYOIDEAE					
Puya Molina					
<i>P. coerulea</i> Miers	1.02	± 0.013			Gitaí et al., 2014
<i>P. mirabilis</i> (Mez) L.B.Sm.	0.88				Ebert & Till, 1997
	0.99	± 0.010			Gitaí et al., 2014
<i>P. raimondii</i> Harms	1.13*				Sgorbati et al., 2007
<i>P. stenothyrsa</i> (Baker) Mez	0.94				Ebert & Till, 1997

References

Arumuganathan, K., and Earle, E.D. (1991). Nuclear DNA content of some important plant species.

Plant Mol. Biol. Rep. 9: 208-218. doi: 10.1007/BF02672069

Cruz, G.A.S, Mendonça Filho, J.R., Vasconcelos, S., Gitaí, J., Salabert De Campos, J.M., Facio

Viccini, L., et al. (2020). Genome size evolution and chromosome numbers of species of the

cryptanthoid complex (Bromelioideae, Bromeliaceae) in a phylogenetic framework. Bot. J. Linn. Soc. 192, 887-899. doi: 10.1093/botlinnean/boz103

Ebert, I., and Till, W. (1997). "Nuclear genome size in Pitcairnioideae (Bromeliaceae) with emphasis on the genus *Pitcairnia*," in Abstracts, Angiosperm genome size discussion meeting, ed. Anonymous (Kew: Royal Botanical Gardens).

Favoreto, F.C., Carvalho, C.R., Lima, A.B.P., Ferreira, A., and Clarindo, W.R. (2012). Genome size and base composition of Bromeliaceae species assessed by flow cytometry. Plant Syst. Evol. 298, 1185-1193. doi: 10.1007/s00606-012-0620-x

Gitaí, J., Paule, J., Zizka, G., Schulte, K., and Benko-Iseppon, A.M.. (2014). Chromosome numbers and DNA content in Bromeliaceae: additional data and critical review. Bot. J. Lin. Soc. 176, 349-368. doi: 10.1111/boj.12211

Gonçalves-Oliveira, R.C., Ximenes, A.F., Oliveira, A.R., Vasconcelos, S., Carvalho-Filho, N., Oliveira, G., et al. (2020). Genome size and chromosome number conservation contrasting with karyotype diversity in *Hohenbergia* (Bromelioideae, Bromeliaceae). Bot. J. Linn. Soc. 192, 900-909. doi: 10.1093/botlinnean/boz094

Gouda, E.J., D. Butcher, C.S. Gouda. cont. updated. Encyclopaedia of Bromeliads, Version 4.
Available online at: <http://bromeliad.nl/encyclopedia>

Müller, L.L.B., Zotz, G., and Albach, D.C. (2019). Bromeliaceae subfamilies show divergent trends of genome size evolution. Sci. Rep. 9, 5136. doi: 10.1038/s41598-019-41474-w

Ramírez-Morillo, I.M., and Brown, G.K. (2001). The origin of the low chromosome number in *Cryptanthus* (Bromeliaceae). Syst. Bot. 26, 722-726. doi: 10.1043/0363-6445-26.4.722

Šmarda, P., Bureš, P., Horová, L., Leitch, I.J., Mucina, L., Pacini, E., et al. (2014). Ecological and evolutionary significance of genomic GC content diversity in monocots. Proc. Natl. Acad. Sci. U.S.A. 111, E4096-E4102. doi: 10.1073/pnas.1321152111