

TNF inhibits catecholamine production from induced sympathetic neuron-like cells in rheumatoid arthritis and osteoarthritis *in vitro*

Markus Herrmann¹, Sven Anders², Rainer H. Straub MD^{1*}, Zsuzsa Jenei-Lanzl PhD^{1,3}

¹Laboratory of Experimental Rheumatology and Neuroendocrine Immunology, Department of Internal Medicine, University Hospital Regensburg, Germany

²Department of Orthopedic Surgery, University Hospital Regensburg, Asklepios Clinic Bad Abbach, Kaiser Karl V Allee 3, 93077, Bad Abbach, Germany

³Dr. Rolf M. Schwiete Research Unit for Osteoarthritis, Orthopedic University Hospital Friedrichsheim gGmbH, Frankfurt/Main, Germany

***Corresponding author**

Rainer H. Straub

Laboratory of Experimental Rheumatology and Neuroendocrine Immunology

Department of Internal Medicine I

University Hospital Regensburg

Biopark I, Am Biopark 9

93053 Regensburg, Germany

Phone: +49 941 944 7120

Email: rainer.straub@ukr.de

Supplementary Figure 1. Rabbit IgG isotype control pictures of OA and RA iTHs as applied in the context of immunofluorescent staining given in Figures 3.

