



Article

Molecular Mechanism of Autosomal Recessive Long QT-Syndrome 1 without Deafness

Annemarie Oertli ^{1,†}, Susanne Rinné ^{1,†}, Robin Moss ^{2,3}, Stefan Kääh ^{4,5}, Gunnar Seemann ^{2,3},
Britt-Maria Beckmann ^{4,6,‡} and Niels Decher ^{1,*,‡} 

- ¹ Institute for Physiology and Pathophysiology, Vegetative Physiology, Philipps-University of Marburg, 35037 Marburg, Germany; annemariekuhn@web.de (A.O.); rinne@staff.uni-marburg.de (S.R.)
² Institute for Experimental Cardiovascular Medicine, University Heart Center Freiburg–Bad Krozingen, Medical Center-University of Freiburg, 79110 Freiburg, Germany;
kai.robin.moss@universitaets-herzzentrum.de (R.M.); gunnar.seemann@universitaets-herzzentrum.de (G.S.)
³ Faculty of Medicine, University of Freiburg, 79110 Freiburg, Germany
⁴ Department of Medicine I, University Hospital, LMU Munich, 80336 Munich, Germany;
Stefan.Kaab@med.uni-muenchen.de (S.K.); beckmann@med.uni-frankfurt.de (B.-M.B.)
⁵ Deutsches Zentrum für Herz-Kreislaufkrankungen (DZHK), Partner Site Munich, 80636 Munich, Germany
⁶ Institut für Rechtsmedizin, Forensische Molekularpathologie, Universitätsklinikum Frankfurt, 60590 Frankfurt, Germany
* Correspondence: decher@staff.uni-marburg.de; Tel.: +49-6421-2862148
† These authors contributed equally to this work.
‡ These authors are co-senior authors.



Citation: Oertli, A.; Rinné, S.; Moss, R.; Kääh, S.; Seemann, G.; Beckmann, B.-M.; Decher, N. Molecular Mechanism of Autosomal Recessive Long QT-Syndrome 1 without Deafness. *Int. J. Mol. Sci.* **2021**, *22*, 1112. <https://doi.org/10.3390/ijms22031112>

Academic Editor: Emanuela Leonarda
Received: 30 December 2020
Accepted: 19 January 2021
Published: 23 January 2021

Publisher's Note: MDPI stays neutral with regard to jurisdictional claims in published maps and institutional affiliations.



Copyright: © 2021 by the authors. Licensee MDPI, Basel, Switzerland. This article is an open access article distributed under the terms and conditions of the Creative Commons Attribution (CC BY) license (<https://creativecommons.org/licenses/by/4.0/>).

Abstract: *KCNQ1* encodes the voltage-gated potassium (Kv) channel *KCNQ1*, also known as *KvLQT1* or *Kv7.1*. Together with its β -subunit *KCNE1*, also denoted as *minK*, this channel generates the slowly activating cardiac delayed rectifier current I_{Ks} , which is a key regulator of the heart rate dependent adaptation of the cardiac action potential duration (APD). Loss-of-function mutations in *KCNQ1* cause congenital long QT1 (LQT1) syndrome, characterized by a delayed cardiac repolarization and a prolonged QT interval in the surface electrocardiogram. Autosomal dominant loss-of-function mutations in *KCNQ1* result in long QT syndrome, called Romano–Ward Syndrome (RWS), while autosomal recessive mutations lead to Jervell and Lange-Nielsen syndrome (JLNS), associated with deafness. Here, we identified a homozygous *KCNQ1* mutation, c.1892_1893insC (p.P631fs*20), in a patient with an isolated LQT syndrome (LQTS) without hearing loss. Nevertheless, the inheritance trait is autosomal recessive, with heterozygous family members being asymptomatic. The results of the electrophysiological characterization of the mutant, using voltage-clamp recordings in *Xenopus laevis* oocytes, are in agreement with an autosomal recessive disorder, since the I_{Ks} reduction was only observed in homomeric mutants, but not in heteromeric I_{Ks} channel complexes containing wild-type channel subunits. We found that *KCNE1* rescues the *KCNQ1* loss-of-function in mutant I_{Ks} channel complexes when they contain wild-type *KCNQ1* subunits, as found in the heterozygous state. Action potential modellings confirmed that the recessive c.1892_1893insC LQT1 mutation only affects the APD of homozygous mutation carriers. Thus, our study provides the molecular mechanism for an atypical autosomal recessive LQT trait that lacks hearing impairment.

Keywords: *KCNQ1*; LQTS; potassium channel; electrophysiology

1. Introduction

There are two important potassium currents responsible for the late phase of repolarization of the cardiac action potential (AP): the rapid delayed rectifier potassium current (I_{Kr}) and the slow rectifier potassium current (I_{Ks}) [1,2]. The stimulation of beta-adrenergic receptors in the heart increases heart rate and contractility and shortens the duration of the AP by activating I_{Ks} [3]. The I_{Ks} current is generated by *KCNQ1* channels, which co-assemble with their cardiac β -subunit *KCNE1*, also known as *minK* [4]. *KCNQ1* channels,

also denoted as Kv7.1, belong to the family of voltage-dependent potassium (Kv) channels which contain six transmembrane segments, with cytoplasmic N- and C-termini [5]. Four α -subunits assemble to a tetrameric structure with a pore-loop located between the S5 and S6 segments [6] that contain the selectivity filter of the channel [7]. In contrast, the *KCNE1* gene encodes for a single transmembrane domain with an extracellular N-terminus and a cytoplasmic C-terminal end. All KCNE family members (KCNE1-5) assemble with KCNQ1 and modify the channel characteristics [8]. In the heart, KCNE1 is the primary accessory subunit of KCNQ1 [9], causing drastic changes to the electrophysiological characteristics of the KCNQ1 channel: current amplitudes are increased, activation is delayed and voltage-dependence of activation is shifted to more positive potentials [4,5]. In addition, KCNE1 is essential for the targeting of KCNQ1 to the surface membrane and its respective stability in the plasma membrane. This becomes evident as in *MinK* (*KCNE1*) knock-out mice, no KCNQ1 immunostaining of the luminal membrane of dark cells of the vestibular system was observed [10]. KCNE1 mutations were identified to be clinically relevant in the human heart and inner ear [10].

Besides the heart, KCNQ1 is expressed in several tissues including kidney, pancreas, lung, placenta, colon, spleen, prostate, peripheral blood leukocytes, small intestine and the stria vascularis of the cochlear duct [11,12]. *KCNQ1* knockout mice presented symptoms such as deafness, balance problems, and morphological anomalies in the internal ear and in the gastrointestinal tract [13]. The KCNQ1/KCNE channel complex is highly relevant for the physiology of the inner ear and a variety of epithelial tissues [14]. For instance, KCNQ1 and KCNE1 were detected in the apical membranes of the marginal cells of the stria vascularis and the dark cells of the vestibular end-organs [15–17].

Congenital long QT syndrome (LQTS) is an inherited channelopathy characterized by a prolonged QT-interval and arrhythmias caused by delayed repolarization [18]. The prevalence of LQTS is 1:2000 in apparently healthy newborns [19]. The clinical manifestations of LQTS are divided into two main categories: arrhythmias and electrocardiogram (ECG) alterations. The cardiac events are due to runs of torsades de pointes ventricular tachycardias, eventually leading to syncope or even to cardiac arrest and sudden cardiac death. *KCNQ1* mutations are the most common with a rate of 39–49% [20–22]. Depending on the type and location of the *KCNQ1* mutation, it can cause rare autosomal recessive Jervell and Lange-Nielsen Syndrome (JLNS) or autosomal dominant Long QT Syndrome 1, also denoted as Romano–Ward Syndrome (RWS) (a less severe form) [23,24]. Homozygous or biallelic compound mutations in *KCNQ1* or *KCNE1* genes were initially reported to lead to JLNS, which is characterized by congenital deafness combined with syncopal attacks and sudden death due to prolonged frequency corrected QT interval (QTc) [25,26]. In the inner ear, I_{Ks} maintains the homeostasis of potassium in the endolymph. The inner ear seems to tolerate I_{Ks} channel deficits much better than the heart. Thus, it was shown that only *KCNQ1* protein levels lower than 10% of the normal level lead to JLNS [23]. Patients with autosomal recessive LQT1 do not suffer from hearing loss. Genetically, they show homozygous or biallelic missense compound mutations [24]. Patients with autosomal recessive LQT1 and JLNS patients present a high rate of cardiac events. The lack of hearing loss in autosomal recessive LQT1 patients is not accompanied by an improved prognosis [24]. *MinK* knock-out mice display inner ear defects with deafness similar to what is observed in patients with JLNS, again highlighting the relevance of the KCNQ1/KCNE1 channel complex in the inner ear [27]. Lastly, autosomal dominant LQT1, known as RWS, is a clinically mild form with genetically heterozygous missense, nonsense, exon skipping and frameshift mutations [23]. However, some *KCNQ1* variants can cause both JLNS (in an autosomal recessive manner) and RWS (in an autosomal dominant manner) [28]. The extent of the loss-of-function determines the symptoms of a *KCNQ1* mutation carrier. A complete loss-of-function results in JLNS and, according to the proportion of the remaining functional *KCNQ1* channels, autosomal recessive LQT1 or autosomal dominant LQT1 is developed [23].

The present study focuses on the identification of the molecular mechanisms of the *KCNQ1* variant, p.P631fs*20 (c.1892_1893insC), leading to rare autosomal recessive LQTS without hearing impairment. The identical mutation was first reported in 1999 by Neyroud et al. [29], and a few years later, Novotny et al. published a case report about a 7-year-old boy suffering from recurrent syncope and carrying this variant in a homozygous manner [30]. However, previous electrophysiological characterization did not reveal why the mutation is only detrimental in the homozygous state. To this end, we performed voltage-clamp experiments in *Xenopus laevis* oocytes expressing the variant alone or together with wild-type *KCNQ1*, in the absence and presence of the cardiac I_{Ks} subunit *KCNE1*. Here, we found that *KCNE1* rescues the trafficking defect of the heterozygous complex of wild-type and mutant *KCNQ1* channel complexes, while it cannot rescue the impaired function of the homomeric p.P631fs*20 (c.1892_1893insC) mutant channel complex.

2. Results

2.1. Identification of a Homozygous *KCNQ1* Gene Mutation in a Family with LQTS without Hearing Impairment

Since the age of three, the index patient (IP) (PID 1889) suffered from syncope, triggered by exercise, sleep or QT prolonging medication. Based on these symptoms, combined with a repeated prolonged QTc interval (Figure 1A) (560 ms under β -blocker-therapy) and a positive family history, we diagnosed LQTS. Notably, the IP did not suffer from hearing loss and she was also diagnosed with multiple sclerosis. To probe for inherited LQTS, we used a candidate gene approach and identified a *KCNQ1* variant. The insertion of cytidine (c.1892_1893insC) leads to a frameshift starting from amino acid P631 and a premature stop codon at position 651 (p.P631fs*20, denoted as *KCNQ1*^{P631fs*20}) (Figure 1B,C); *KCNQ1*^{WT} comprises 676 amino acids. The frameshift in the *KCNQ1* amino acid sequence is located in the cytoplasmic C-terminus of the *KCNQ1* channel protein (Figure 1C). The variant was absent from the Exome Variant Server (EVS) (<https://evs.gs.washington.edu/EVS/>) and not found by the gnomAD database (former ExAC browser) (<https://gnomad.broadinstitute.org/>). Referring to the Protein Variation Effect Analyzer (PROVEAN)-score (<http://provean.jcvi.org/index.php>), *KCNQ1*^{P631fs*20} was predicted to be deleterious.

The IP was homozygous for the *KCNQ1* c.1892_1893insC variant and both parents were heterozygous and asymptomatic mutation carriers (Figure 1D). Additionally, one paternal uncle (QTc 383 ms) and a paternal aunt (QTc 412 ms) were heterozygous for the mutation. The sister died at the age of 21 because of sudden cardiac death (SCD), and she was, in contrast to the IP, deaf. Her brother died when he was 2 years old due to SCD. Two daughters and a son of the IP were heterozygous carriers of the mutation and asymptomatic (QTc 421 ms, 397 ms and 413 ms, respectively) (Figure 1D).




Figure 1. Identification of the homozygous $KCNQ1^{P631fs*20}$ mutation in a patient with LQTS without hearing loss. (A) 12-lead ECG recording of the IP (PID1889) showing a QTc of 560 ms and a heart rate of 60 bpm under beta-blocker therapy. Paper speed was 50 mm/sec. (B) Partial nucleotide and amino acid sequence illustrated from amino acid 630 to 651 for $KCNQ1^{WT}$ (top) and the $KCNQ1^{P631fs*20}$ variant (bottom). The cytidine insertion is highlighted in red as well as the novel amino acid sequence due to the frame shift. Please note that the mutated sequence shows a premature stop codon at amino acid position 651. (C) Cartoon showing the topology of the $KCNQ1$ channel α -subunit. The location of the mutation is indicated by a red circle and the resulting novel amino acid sequence due to the frameshift is provided in red. (D) Pedigree of the family of the IP (marked by a red arrow) with a $KCNQ1^{P631fs*20}$ mutation as a result of a nucleotide insertion mutation (c.1892_1893insC). Filled symbols indicate patients and family members with a previous diagnosis of LQTS with or without symptoms. Squares and circles represent male and female subjects, respectively. In the top right of the symbol, genetic information is given: “+/-” heterozygous mutation carrier, “+/+” homozygous mutation carrier, “N” no genetic information was available. Below the symbols, information about certain symptomatic or further diseases are given. “SCD” sudden cardiac death. In addition, the QTc is shown for the IP. If there are no symptoms, it was marked with “asymp.”. Symbols with a line through mark deceased subjects, and the age and cause of death is indicated below.

2.2. Electrophysiological Characterization of the $KCNQ1^{P631fs*20}$ Variant Revealed a Loss-of-Function in Both Homozygous and Heterozygous States

To characterize the electrophysiological properties of $KCNQ1^{P631fs*20}$, the variant was expressed in *Xenopus laevis* oocytes and compared to oocytes injected with $KCNQ1$ wild-type ($KCNQ1^{WT}$) copy RNA (cRNA) using voltage-clamp recordings. We observed a significant reduction in the current amplitude for the homomeric variant (resembling the homozygous state) (Figure 2A–C). Analyzing the outward currents at +40 mV, the current amplitude was reduced by about 65% compared to $KCNQ1^{WT}$ (resembling healthy controls) (Figure 2C). Injecting similar amounts of $KCNQ1^{WT}$ and $KCNQ1^{P631fs*20}$ (resembling the heterozygous state) led to a significant current reduction (Figure 2A–C); however, it was less pronounced than observed for the variant expressed alone. Current reduction analyzed at +40 mV was about 40% compared to the $KCNQ1^{WT}$ (Figure 2C). We also injected half the amount of $KCNQ1^{WT}$ cRNA to mimic a haploinsufficiency, and found currents comparable to the heterozygous state (Figure 2A–C). Thus, heteromeric channels that contain the mutation suffer from a non-dominant negative but clear loss-of-function.




Figure 2. Functional effects of the $KCNQ1^{P631fs*20}$ mutation. (A) Representative current traces of oocytes injected with $KCNQ1^{WT}$ 100% (14.5 ng/oocyte), $KCNQ1^{P631fs*20}$ (14.5 ng/oocyte), $KCNQ1^{WT}$ 50% (7.25 ng/oocyte) or $KCNQ1^{WT}$ plus $KCNQ1^{P631fs*20}$ (7.25 ng/oocyte each), respectively. Voltage was stepped from -60 to $+60$ mV in 20 mV steps lasting 3000 ms, from a holding potential of -80 mV. (B) Current voltage relationships obtained by blotting the current at the end of each voltage step for each voltage applied normalized to $KCNQ1^{WT}$. In order to obtain the current-voltage relationship (I/V curve), all wild-type recordings were normalized to the value at $+60$ mV. The data of all the other constructs were also divided by the average current amplitude of the wild-type at $+60$ mV of the respective recording day. (C) Current amplitudes analyzed at $+40$ mV and normalized to $KCNQ1^{WT}$ (100%). All the data, also that of the wild-type recordings, were divided by the average current amplitude of the wild-type at $+40$ mV of the respective recording day. Numbers of oocytes recorded are indicated within the bar graphs. Values are expressed as means \pm S.E.M. Error bars represent S.E.M. values. Significance was assessed using two tailed Student's t-test. Asterisks indicate significance: ***, $p < 0.001$.

2.3. In Electrophysiological Recordings, KCNE1 Rescued the $KCNQ1^{P631fs*20}$ Loss-of-Function Present in the Heteromeric State with $KCNQ1^{WT}$

Since in cardiac tissue $KCNQ1$ is expressed in a complex with its β -subunit $KCNE1$ to form the I_{Ks} , we also examined the electrophysiology of the $KCNQ1^{P631fs*20}$ mutant co-expressed with $KCNE1$. As *Xenopus laevis* oocytes express an endogenous $KCNQ1$ channel ($xKCNQ1$), we first recorded the “endogenous xI_{Ks} ” by only injecting $KCNE1$ (Figure 3A) to determine the $xKCNQ1/KCNE1$ background for the subsequent analysis of the exogenously expressed h $KCNQ1/KCNE1$ channel complex. Here, we observed a significant reduction in the current amplitude for $KCNQ1^{P631fs*20}$ injected together with $KCNE1$ (resembling the homozygous state) compared to $KCNQ1^{WT}$ plus $KCNE1$ (Figure 3A–C). The outward current analyzed at $+40$ mV was reduced by about 65% (Figure 3C). The current amplitude was comparable to that of the endogenous xI_{Ks} background. Surprisingly, injecting similar amounts of $KCNQ1^{WT}$ and $KCNQ1^{P631fs*20}$ together with $KCNE1$ (resembling the heterozygous state) led to no current reduction (Figure 3A–C), in contrast to the current reduction observed for the heterozygous state without $KCNE1$ (Figure 2A–C). As a control, we also injected half the amount of $KCNQ1^{WT}$ cRNA together with $KCNE1$ to mimic a haploinsufficiency (Figure 3A–C), and found that the current reduction was less pronounced, with only 35% at $+40$ mV compared to $KCNQ1^{WT}$ expressed with $KCNE1$ (healthy control) (Figure 3C). In summary, the electrophysiological data indicate that there are strongly reduced I_{Ks} currents for homozygous $KCNQ1^{P631fs*20}$ carriers as observed in our IP, and no I_{Ks} current reduction for heterozygous mutation carriers. Thus, $KCNQ1^{P631fs*20}$ channels assembled without $KCNE1$ also display a current reduction in the heterozygous state (Figure 2), which is rescued by $KCNE1$ (Figure 3). Since, in the heart, $KCNQ1$ is predominantly assembled together with $KCNE1$, these results may explain the LQTS and symptoms of the IP, and the absence of a phenotype of heterozygous mutation carriers.




Figure 3. Functional effects of the $KCNQ1^{P631fs*20}$ mutant co-expressed with $KCNE1$. (A) Representative current traces of oocytes injected with $KCNQ1^{WT}$ (8.1 ng/oocyte), $KCNQ1^{P631fs*20}$ (8.1 ng/oocyte), $KCNQ1^{WT}$ 50% (4 ng/oocyte) or $KCNQ1^{WT}$ plus $KCNQ1^{P631fs*20}$ (4 ng/oocyte each), all together with 0.1 ng $KCNE1$ cRNA or only $KCNE1$ cRNA (0.1 ng/oocyte), to obtain *Xenopus* (x) $KCNQ1/KCNE1$ (xI_{Ks}) channels. Voltage was stepped from -40 to $+40$ mV in 20 mV increments lasting 7000 ms, from a holding potential of -80 mV. (B) Current voltage relationships obtained by blotting the current at the end of each voltage step for each voltage applied normalized to $KCNQ1^{WT}$ plus $KCNE1$. In order to obtain the current-voltage relationship (I/V curve), all wild-type recordings were normalized to the value at $+60$ mV. The data of all the other constructs were also divided by the average current amplitude of the wild-type at $+60$ mV of the respective recording day. (C) Current amplitudes analyzed at $+40$ mV and normalized to $KCNQ1^{WT}$ plus $KCNE1$. All the data, also that of the wild-type recordings, were divided by the average current amplitude of the wild-type at $+40$ mV of the respective recording day. (A–C) All data are presented without subtraction of $xKCNQ1$ background currents. Numbers of oocytes recorded are indicated within the bar graphs. Values are expressed as means \pm S.E.M. Error bars represent S.E.M. values. Significance was assessed using two tailed Student's *t*-test. Asterisks indicate significance: ***, $p < 0.001$.

2.4. The $KCNQ1^{P631fs*20}$ Mutant does Not alter the Gating of Homomeric and Heteromeric I_{Ks} Channel Complexes

Next, we analyzed whether the $KCNQ1^{P631fs*20}$ mutation affects the voltage-dependence of activation of I_{Ks} channel complexes containing $KCNE1$ (Figure 4). The voltage-dependence of the homomeric channel complex of $KCNQ1^{P631fs*20}$ and $KCNE1$ was similar to that of the human or *Xenopus* wild-type I_{Ks} channels (Figure 4A), with no significant changes in the voltage of half-maximal activation ($V_{1/2}$) (Figure 4B). Additionally, the “ $KCNE1$ rescued” heteromeric I_{Ks} channel complexes, containing $KCNQ1^{WT}$, $KCNQ1^{P631fs*20}$ and $KCNE1$ subunits, did not have a significantly altered voltage-dependence (Figure 4A,B), indicating that the loss-of-function and the rescue of the $KCNQ1^{P631fs*20}$ currents primarily results from changes in the trafficking of the I_{Ks} channel complex by $KCNE1$.




Figure 4. Voltage-dependence of activation of the $KCNQ1^{P631fs*20}$ mutant co-expressed with KCNE1. (A) Voltage-dependence of activation for $KCNQ1+KCNE1$, $xKCNQ1+KCNE1$, $KCNQ1^{P631fs*20}+KCNE1$ and $KCNQ1^{WT}/KCNQ1^{P631fs*20}+KCNE1$. Recordings were performed with the protocol as described in Figure 3. The tail currents recorded after the 7000 ms pulse were normalized to the respective maximal tail current of each recording to obtain the conductance/voltage (G/V) curves. (B) Normalized tail currents were fitted to a Boltzmann function. The voltage of half-maximal activation ($V_{1/2}$) of $KCNQ1+KCNE1$, $xKCNQ1+KCNE1$, $KCNQ1^{P631fs*20}+KCNE1$ and $KCNQ1^{WT}/KCNQ1^{P631fs*20}+KCNE1$ are illustrated, together with the numbers of oocytes analyzed.

2.5. Action Potential Modelling Predicts a QT Prolongation Exclusively Present In Homozygous Patients

Action potential simulations were performed using the O'Hara-Rudy ventricular cell model [31] after the subtraction of the endogenous xI_{Ks} current, recorded by the injection of KCNE1 alone (Figures 3C and 5A). Thus, simulations were carried out with a completely reduced conductivity of the I_{Ks} channel for the homozygous state and with an 11% reduction for the heterozygous state (Figure 5A). For the homozygous mutation carriers, the simulations predicted a prolongation of the action potential duration (APD). In contrast, the simulations predicted that the APD of the heterozygous patients was not or only very mildly prolonged (Figure 5B), resulting in APD₉₀ values of 232 ms for homozygous patients and 222 ms for heterozygous patients (APD₉₀ for wild type was 221 ms), respectively (Figure 5C). Taken together, the computational simulations confirmed that the APD, and thus the QT prolongation, occurs in homozygous, but not in heterozygous mutation carriers.




Figure 5. Action potential modelling with the homozygous or heterozygous $KCNQ1^{P631fs*20}$ mutation in the I_{Ks} channel complex. (A) Mean current amplitudes of 100% $KCNQ1^{WT}$ injected together with KCNE1, 50% $KCNQ1^{WT}$ injected together with KCNE1, 100% $KCNQ1^{P631fs*20}$ injected together with KCNE1 or 50% $KCNQ1^{WT}$ plus 50% $KCNQ1^{P631fs*20}$ injected together with KCNE1. For all data, the endogenous

xI_{Ks} current recorded by injection of KCNE1 alone was subtracted. Data were normalized to $KCNQ1^{WT}$ 100%. (B) Action potential simulations using the O'Hara-Rudy ventricular cell model. Conductivity of the I_{Ks} channel was set to 100% for wild-type (black), 0% for the homozygous (red, dashed line) and 89% for the heterozygous state (blue, dotted line) as calculated in (A). (C) APD at 90% repolarization (APD₉₀) was evaluated for each configuration after the simulations achieved a steady state.

3. Discussion

In the present study, we describe the inheritance of the *KCNQ1* mutation p.P631fs*20 (c.1892_1893insC) in a family with homozygous and heterozygous mutation carriers over three generations. The homozygous IP presented a pronounced prolongation of the QT interval (QTc 560 ms) with several syncopal events, but without hearing loss. The identical mutation was first reported in 1999 by Neyroud et al. [29]. Here, the heterozygous IP experienced several stress-induced syncopes since the age of three, and her QTc interval was 450 ms [29]. Despite the fact that the father of the IP shared the heterozygous mutation, he was asymptomatic. Nevertheless, the authors proposed RWS, meaning an autosomal dominant inheritance. Our data clearly indicate that the heterozygous state does not cause a loss-of-function or act in a dominant-negative manner, as typically observed for RWS mutations. Most importantly, no further genetic analysis of the family was described in the study by Neyroud et al., thus an additional mutation responsible for the symptoms of the heterozygous patient cannot be excluded.

A few years later, Novotny et al. published a case of a 7-year-old boy suffering from recurrent syncopes and a prolonged QTc interval (up to 700 ms). This boy carried the variant we analyzed in our study in a homozygous manner [30]. The heterozygous mutation carriers of the family of our IP (PID 1889) were asymptomatic, which is in line with the family reported by Novotny et al., where the heterozygous parents of the young boy were also asymptomatic. Both our IP as well as the boy in this case report had no hearing loss. Novotny et al. proposed recessive LQT1 syndrome due to $KCNQ1^{P631fs*20}$. However, in the study of Neyroud et al., as well as in the study by Novotny et al., no electrophysiological characterization of the mutation was performed. Finally, Sato et al. presented a family with two heterozygous mutations, insC1893-1894 (P631fs/19) (named P631fs/20 in our study) and delGTA1783-1785 (delV595), causing autosomal-recessive compound heterozygous LQT1 [32]. The patients carrying the two compound mutations suffered from syncopes and had no hearing loss. The heterozygous family members carrying one of the *KCNQ1* mutations (insC1893-1894 (P631fs/19) or delGTA1783-1785 (delV595)) showed no significant QT prolongation, implying that both mutations lacked a dominant negative effect. This is in line with the lack of phenotype in the heterozygous family members of our IP. In whole cell patch clamp recordings, both mutations (P631fs/19 or delV595) showed a strong current reduction when expressed alone with KCNE1 (homozygous state); however, in the heterozygous state, meaning the expression of each mutant together with $KCNQ1^{WT}$ and KCNE1, the current was not significantly altered [32]. Expressing both mutants together with KCNE1 (as in the compound heterozygous patients), current was strongly reduced [32]. Sato et al. did not perform electrophysiological experiments without KCNE1, and thus the authors could not find the loss-of-function in the heteromeric state, which is present in the absence of KCNE1, and the rescue of heteromeric channel complexes composed of mutant and wild-type channel subunits. As the compound heterozygous patients described by Sato et al. carried the mutation we identified, together with a second mutation, we speculate that KCNE1 can only rescue heteromeric channels containing wild-type *KCNQ1* channel subunits.

The study from Sato et al. demonstrated a decreased expression of $KCNQ1^{P631fs*19}$ channels caused by intracellular aggregation and trafficking defects. The new amino acid sequence resulting from the frame shift contains two potential newly generated endoplasmic reticulum (ER) retention signals (R⁶³³GR and R⁶⁴⁶LR). Thus, the authors proposed a retention of $KCNQ1^{P631fs*19}$ in the ER [32]. Thus, the question arises, why can the mis-

trafficking of the channel and the associated current reduction be exclusively found in the homozygous state? $KCNQ1^{P631fs*19}$ mutants still form heteromeric complexes with $KCNQ1^{WT}$ subunits, but they did not efficiently retain the $KCNQ1^{WT}$ subunits [32]. On the other hand, Sato et al. described that the $KCNQ1^{P631fs*19}$ mutation severely affected the expression and stability of full-length $KCNQ1$ protein in transfected cells, and proposed, based on these observations, that the number of mutant $KCNQ1$ subunits was fewer than the number of $KCNQ1^{WT}$ subunits in the ER, where the subunit assembly occurred. Thus, the amount of functional heteromeric channels at the plasma membrane may not be altered. However, in this study, $KCNE1$ was not co-transfected studying the trafficking and heteromerization of $KCNQ1$ channels. Our data indicate that $KCNE1$ is able to fully restore the mistrafficking of heteromeric complexes of wild-type and mutant channels since we did not observe a current reduction for the heterozygous state (in the presence of $KCNE1$). We did not perform any cell surface expression or trafficking experiments for $KCNQ1^{P631fs*20}$ and $KCNQ1^{WT}$ (with and without $KCNE1$), as the mutation has been previously described as trafficking deficient [32]. Since $KCNE1$ rescues the loss-of-function caused by a mistrafficking of this mutant, while the resulting I_{Ks} channel complex has a normal voltage-dependent gating, the most straightforward explanation is that it rescues the trafficking phenotype, especially as $KCNE1$ is known to modulate the surface expression of $KCNQ1$ [10,33–35]. Yet, as we did not obtain any primary data to confirm this theory, the mechanism of a rescued trafficking defect is currently only a hypothesis.

Bianchi et al. described a $KCNQ1$ mutation (c.1630_1636delCAGTACTinsGTTGAGAT), denoted as $\Delta 544$, which is a deletion/insertion mutation leading to a novel amino acid sequence and a premature stop codon at position 651 [36]. The homozygous mutation carrier of $\Delta 544$ suffered from JLNS and the heterozygous mutation carriers presented a mild phenotype [29]. In voltage-clamp recordings of the mutated I_{Ks} current (co-expression of homomeric $KCNQ1^{\Delta 544}$ and $KCNE1$), Bianchi et al. showed a current reduction down to the level of endogenous xI_{Ks} (injection of $KCNE1$ only) [36]. These data are comparable to our findings with homomeric $KCNQ1^{P631fs*20}$ co-expressed with $KCNE1$, while our patient did not suffer from hearing loss. In contrast to our data with the $KCNQ1^{P631fs*20}$ variant, the co-expression of $KCNQ1^{WT}$ and $KCNQ1^{\Delta 544}$ (resembling the heterozygous state) in the presence of $KCNE1$ led to a current reduction of about 50%, albeit not in a dominant negative manner. We did not observe such a reduction in I_{Ks} for $KCNQ1^{P631fs*20}$ in the heterozygous state when $KCNE1$ is present. These differences may explain the mild phenotype of the heterozygous $KCNQ1^{\Delta 544}$ mutation carriers and the absence of symptoms in heterozygous $KCNQ1^{P631fs*20}$ carriers. Both mutations, $KCNQ1^{P631fs*20}$ as well as $KCNQ1^{\Delta 544}$, share a premature termination of the amino acid chain at the terminal C-terminus and present a severe phenotype in a homozygous manner. Interestingly, $KCNQ1^{\Delta 544}$ presents, as well as $KCNQ1^{P631fs*20}$, retention motifs due to the deletion/insertion. However, Bianchi et al. showed the expression of $KCNQ1^{\Delta 544}$ at the plasma membrane, thus ER retention may not be the mechanism for the current reduction observed for $KCNQ1^{\Delta 544}$, whereas a localization of the mutant in the ER and adjacent organelle, presumably the Golgi apparatus, was described for $KCNQ1^{P631fs*20}$ [32]. Whether these mechanistic differences may explain the different clinical spectrum of LQTS in these families currently remains unclear. Our IP and other homozygous $KCNQ1^{P631fs*20}$ mutation carriers previously described do not suffer from hearing loss. In our voltage-clamp experiments, homomeric $KCNQ1^{P631fs*20}$ was expressed together with $KCNE1$, as it is the case in the inner ear and the heart of our IP. Bhuiyan et al. postulated that a minimum of functional channels prevents hearing loss [23]. The remaining I_{Ks} current may maintain the potassium homeostasis in the endolymph, ensuring proper inner ear function, while it may not be sufficient for normal electric activity of the heart. Why the inner ear is not or only very rarely affected, such as in the case of the sister of our IP, despite the massive current reduction that we have observed in the *Xenopus* oocyte expression system, remains unclear. The trafficking defect of the $KCNQ1^{P631fs*20}$ mutation may manifest to a different extent depending on the tissue.

Our data are in agreement with an autosomal recessive inheritance trait of the $KCNQ1^{P631fs*20}$ mutation. The mechanism behind this unusual recessive genetic LQTS trait only became obvious by co-expression studies in the absence and presence of KCNE1. Expressing the heterozygous $KCNQ1^{P631fs*20}/KCNQ1^{WT}$ channel complex without KCNE1 led to a current reduction of about 39%, comparable to a haploinsufficiency. This loss-of-function in the heterozygous state was completely rescued by KCNE1, while the homomeric mutant channel complexes were not recovered. Recessive inheritance can be explained by a KCNE1 mediated rescue, which is selective for heteromeric channels that contain wild-type and mutant channel subunits as found in heterozygous patients.

4. Materials and Methods

4.1. Clinical Evaluation

In 2003, the *KCNQ1* mutation p.P631fs*20 (c.1892_1893insC) was identified in the Index patients (IP) at the age of 39 years. The diagnosis of LQTS was set by cardiologists specialized on inherited arrhythmia syndromes at the Hospital of the Ludwig Maximilians University of Munich. Detailed medical history focused on LQTS relevant comorbidities and the family history for 3 generations. Additionally, 12-lead resting ECGs were taken. According to Goldenberg et al. [37], we considered a QT interval corrected for the heart rate using Bazett's formula (QTc) ≤ 450 ms in men and ≤ 460 ms in women to be normal. Informed consent prior to genetic investigations was collected for all patients in this study, complying with the ethical standards of the 1964 Declaration of Helsinki and its latest revision at that time.

4.2. Molecular Biology

Blood samples of the IP were taken and genomic DNA was extracted. All exons and intronic splice sites of the genes *KCNQ1*, *KCNH2*, *KCNE1*, *KCNE2* and *SCN5A* were PCR amplified (primer sequences available upon request) and sequenced. For *KCNQ1*, obtained DNA sequence information was compared with the *KCNQ1* wild-type sequence (NM_000218.3). Human *KCNQ1* was cloned into the pSP64T vector. QuikChange Site-Directed Mutagenesis Kit (Agilent Technologies, Santa Clara, CA, USA) was used to introduce the mutation into human *KCNQ1* cDNA. cDNA was linearized with EcoRI (Thermo Fisher Scientific, Waltham, MA, USA) and cRNA was in vitro synthesized using the mMESSEMGEMMACHINE[®]SP6 kit (Thermo Fisher Scientific, Waltham, MA, USA). cRNA concentration was quantified using a spectrophotometer (NanoDrop, Thermo Fisher Scientific, Waltham, MA, USA) and quality was controlled by agarose gel electrophoresis.

4.3. Electrophysiology

cRNA was injected into *Xenopus laevis* oocytes. Oocytes were taken from ovarian lobes of anesthetized *Xenopus laevis* toads. Anesthesia was performed with 2 g/l tricaine-methanesulfonate (SIGMA-Aldrich, Missouri, USA). Subsequently, oocytes were treated with collagenase type II (2 mg/mL, Worthington, USA) in OR2 solution (NaCl 82.5 mM, KCl 2 mM, MgCl₂ 1 mM, HEPES 5 mM, pH 7.4; all from SIGMA-Aldrich, Missouri, USA) for 120 min to remove residual connective tissue. Isolated oocytes were stored at 18 °C in ND96 recording solution (NaCl 96 mM, KCl 2 mM, CaCl₂ 1.8 mM, MgCl₂ 1 mM, HEPES 5 mM, pH 7.5) supplemented with Na-pyruvate (275 mg/L), theophylline (90 mg/L) and gentamicin (50 mg/L) (all from SIGMA-Aldrich, Missouri, USA). Oocytes were injected with 50.6 nl of *KCNQ1* cRNA or *KCNQ1* and *KCNE1* cRNA in equal parts. Standard two electrode voltage-clamp experiments were performed at room temperature (21–22 °C) with an Axoclamp 900A amplifier, a Digidata 1440A and pClamp10 software (Axon instruments, CA, USA) 72 h after cRNA injection. The microelectrodes were made from glass pipettes (Science products, Hofheim, Germany) pulled with a DMZ-Universal puller (Zeitz, Germany) and had a resistance of 0.2–1.0 M Ω when filled with 3 M KCl (SIGMA-Aldrich, Missouri, USA). The following voltage protocols were used: 3000 ms pulses were applied in 20 mV steps from –60 mV to +60 mV, or 7000 ms pulses were

applied in 20 mV steps from -40 mV to $+40$ mV for recording oocytes expressing the KCNQ1 channels, or KCNQ1 and the KCNE1 subunit, respectively.

4.4. Action Potential Modelling

Action potential simulations were obtained using the O'Hara-Rudy ventricular cell model [31] in its endocardial configuration. The set of differential equations was solved with a fixed time step of $10 \mu\text{s}$ using the Rush-Larsen method [38] for all gating variables and the forward Euler method for all other variables at a pacing frequency of 1 Hz. The conductivity of the gks channel was then reduced by 100% for the homozygous state and 11% for the heterozygous state, based on the respective measured mean current amplitudes. To ensure all configurations had reached limit cycle, 500 beats were simulated. APD_{90} was then calculated based on the last beat.

4.5. Data Analysis

All electrophysiology data were acquired with Clampex (Molecular Devices, Sunnyvale, CA, USA) and analyzed with Clampfit 10 (Molecular Devices, Sunnyvale, CA, USA), Origin (OriginLab Corp, Northampton, MS, USA) and Excel (Microsoft Corp, Seattle, WA, USA). All values are expressed as means \pm S.E.M. Error bars in all figures represent S.E.M. values. Significance was assessed using two tailed Student's *t*-test. Asterisks indicate significance: *, $p < 0.05$; **, $p < 0.01$; ***, $p < 0.001$.

Author Contributions: Conceptualization, B.-M.B. and N.D.; methodology, S.K., B.-M.B., G.S. and N.D.; software, G.S.; validation, S.R., B.-M.B. and N.D.; formal analysis, A.O., R.M. and S.R.; investigation, A.O., R.M. and S.R.; resources, S.K. and N.D.; data curation, S.R., B.-M.B. and N.D.; writing—original draft preparation, A.O., S.R. and N.D.; writing—review and editing, S.R., B.-M.B. and N.D.; visualization, A.O., R.M., G.S., B.-M.B. and S.R.; supervision, S.K., B.-M.B., G.S. and N.D.; project administration, B.-M.B. and N.D.; funding acquisition, S.K. and N.D. All authors have read and agreed to the published version of the manuscript.

Funding: This research was funded by the German Federal Ministry of Education and Research (BMBF) in the context of the German National Genom Research Network (NGFN) by grants to Stefan Käab (01GS0499).

Institutional Review Board Statement: This study was conducted according to the guidelines of the Declaration of Helsinki, and approved by the Institutional Review Board (or Ethics Committee) of the University of München and by the local ethics commission of the Regierungspräsidium Giessen (V54-19c 20 15 h 02 MR 20/28 Nr.A 4/2013).

Informed Consent Statement: Informed consent was obtained from all subjects involved in the study.

Data Availability Statement: Data is contained within the article. Raw data are available from the corresponding author on reasonable request.

Acknowledgments: We would like to thank Susanne Bamerny for her support.

Conflicts of Interest: The authors declare no conflict of interest. The funders had no role in the design of the study; in the collection, analyses, or interpretation of data; in the writing of the manuscript, or in the decision to publish the results.

References

1. Sanguinetti, M.C.; Jurkiewicz, N.K. Two components of cardiac delayed rectifier K^+ current. Differential sensitivity to block by class III antiarrhythmic agents. *J. Gen. Physiol.* **1990**, *96*, 195–215. [[CrossRef](#)] [[PubMed](#)]
2. Tsien, R.W.; Noble, D. A transition state theory approach to the kinetics of conductance changes in excitable membranes. *J. Membr. Biol.* **1969**, *1*, 248–273. [[CrossRef](#)]
3. Marx, S.O.; Kurokawa, J.; Reiken, S.; Motoike, H.; D'Armiento, J.; Marks, A.R.; Kass, R.S. Requirement of a macromolecular signaling complex for beta adrenergic receptor modulation of the KCNQ1-KCNE1 potassium channel. *Science* **2002**, *295*, 496–499. [[CrossRef](#)]
4. Sanguinetti, M.C.; Curran, M.E.; Zou, A.; Shen, J.; Spector, P.S.; Atkinson, D.L.; Keating, M.T. Coassembly of K(V)LQT1 and minK (IsK) proteins to form cardiac I(Ks) potassium channel. *Nature* **1996**, *384*, 80–83. [[CrossRef](#)]

5. Barhanin, J.; Lesage, F.; Guillemare, E.; Fink, M.; Lazdunski, M.; Romey, G. K(V)LQT1 and IsK (minK) proteins associate to form the I(Ks) cardiac potassium current. *Nature* **1996**, *384*, 78–80. [[CrossRef](#)]
6. Snyders, D.J. Structure and function of cardiac potassium channels. *Cardiovasc. Res.* **1999**, *42*, 377–390. [[CrossRef](#)]
7. MacKinnon, R. Pore loops: An emerging theme in ion channel structure. *Neuron* **1995**, *14*, 889–892. [[CrossRef](#)]
8. Wrobel, E.; Tapken, D.; Seebohm, G. The KCNE Tango—How KCNE1 Interacts with Kv7.1. *Front. Pharmacol.* **2012**, *3*, 142. [[CrossRef](#)]
9. Bendahhou, S.; Marionneau, C.; Haurogne, K.; Larroque, M.M.; Derand, R.; Szuts, V.; Escande, D.; Demolombe, S.; Barhanin, J. In Vitro molecular interactions and distribution of KCNE family with KCNQ1 in the human heart. *Cardiovasc. Res.* **2005**, *67*, 529–538. [[CrossRef](#)]
10. Warth, R.; Barhanin, J. The multifaceted phenotype of the knockout mouse for the KCNE1 potassium channel gene. *Am. J. Physiol. Regul. Integr. Comp. Physiol.* **2002**, *282*, R639–R648. [[CrossRef](#)]
11. Chouabe, C.; Neyroud, N.; Guicheney, P.; Lazdunski, M.; Romey, G.; Barhanin, J. Properties of KvLQT1 K⁺ channel mutations in Romano-Ward and Jervell and Lange-Nielsen inherited cardiac arrhythmias. *EMBO J.* **1997**, *16*, 5472–5479. [[CrossRef](#)] [[PubMed](#)]
12. Neyroud, N.; Tesson, F.; Denjoy, I.; Leibovici, M.; Donger, C.; Barhanin, J.; Faure, S.; Gary, F.; Coumel, P.; Petit, C.; et al. A novel mutation in the potassium channel gene KVLQT1 causes the Jervell and Lange-Nielsen cardioauditory syndrome. *Nat. Genet.* **1997**, *15*, 186–189. [[CrossRef](#)] [[PubMed](#)]
13. Casimiro, M.C.; Knollmann, B.C.; Ebert, S.N.; Vary, J.C., Jr.; Greene, A.E.; Franz, M.R.; Grinberg, A.; Huang, S.P.; Pfeifer, K. Targeted disruption of the *Kcnq1* gene produces a mouse model of Jervell and Lange-Nielsen Syndrome. *Proc. Natl. Acad. Sci. USA* **2001**, *98*, 2526–2531. [[CrossRef](#)]
14. Bleich, M.; Warth, R. The very small-conductance K⁺ channel KvLQT1 and epithelial function. *Pflüg. Arch. Eur. J. Physiol.* **2000**, *440*, 202–206. [[CrossRef](#)]
15. Forge, A.; Wright, T. The molecular architecture of the inner ear. *Br. Med. Bull.* **2002**, *63*, 5–24. [[CrossRef](#)]
16. Nicolas, M.; Dememes, D.; Martin, A.; Kupersmidt, S.; Barhanin, J. KCNQ1/KCNE1 potassium channels in mammalian vestibular dark cells. *Heart Res.* **2001**, *153*, 132–145. [[CrossRef](#)]
17. Vetter, D.E.; Mann, J.R.; Wangemann, P.; Liu, J.; McLaughlin, K.J.; Lesage, F.; Marcus, D.C.; Lazdunski, M.; Heinemann, S.F.; Barhanin, J. Inner ear defects induced by null mutation of the *isk* gene. *Neuron* **1996**, *17*, 1251–1264. [[CrossRef](#)]
18. Moss, A.J. Long QT Syndrome. *JAMA* **2003**, *289*, 2041–2044. [[CrossRef](#)]
19. Schwartz, P.J.; Stramba-Badiale, M.; Crotti, L.; Pedrazzini, M.; Besana, A.; Bosi, G.; Gabbarini, F.; Goulene, K.; Insolia, R.; Mannarino, S.; et al. Prevalence of the congenital long-QT syndrome. *Circulation* **2009**, *120*, 1761–1767. [[CrossRef](#)]
20. Napolitano, C.; Priori, S.G.; Schwartz, P.J.; Bloise, R.; Ronchetti, E.; Nastoli, J.; Bottelli, G.; Cerrone, M.; Leonardi, S. Genetic testing in the long QT syndrome: Development and validation of an efficient approach to genotyping in clinical practice. *JAMA* **2005**, *294*, 2975–2980. [[CrossRef](#)]
21. Splawski, I.; Shen, J.; Timothy, K.W.; Lehmann, M.H.; Priori, S.; Robinson, J.L.; Moss, A.J.; Schwartz, P.J.; Towbin, J.A.; Vincent, G.M.; et al. Spectrum of mutations in long-QT syndrome genes. KVLQT1, HERG, SCN5A, KCNE1, and KCNE2. *Circulation* **2000**, *102*, 1178–1185. [[CrossRef](#)] [[PubMed](#)]
22. Tester, D.J.; Will, M.L.; Haglund, C.M.; Ackerman, M.J. Compendium of cardiac channel mutations in 541 consecutive unrelated patients referred for long QT syndrome genetic testing. *Heart Rhythm* **2005**, *2*, 507–517. [[CrossRef](#)] [[PubMed](#)]
23. Bhuiyan, Z.A.; Wilde, A.A. IKs in heart and hearing, the ear can do with less than the heart. *Circ. Cardiovasc. Genet.* **2013**, *6*, 141–143. [[CrossRef](#)] [[PubMed](#)]
24. Giudicessi, J.R.; Ackerman, M.J. Prevalence and potential genetic determinants of sensorineural deafness in KCNQ1 homozygosity and compound heterozygosity. *Circ. Cardiovasc. Genet.* **2013**, *6*, 193–200. [[CrossRef](#)] [[PubMed](#)]
25. Jervell, A.; Lange-Nielsen, F. Congenital deaf-mutism, functional heart disease with prolongation of the Q-T interval and sudden death. *Am. Heart J.* **1957**, *54*, 59–68. [[CrossRef](#)]
26. Schulze-Bahr, E.; Wang, Q.; Wedekind, H.; Haverkamp, W.; Chen, Q.; Sun, Y.; Rubie, C.; Hordt, M.; Towbin, J.A.; Borggrefe, M.; et al. KCNE1 mutations cause jervell and Lange-Nielsen syndrome. *Nat. Genet.* **1997**, *17*, 267–268. [[CrossRef](#)]
27. Splawski, I.; Timothy, K.W.; Vincent, G.M.; Atkinson, D.L.; Keating, M.T. George, M. Cober Lecturer: Mark T. Keating. Molecular basis of the long-QT syndrome associated with deafness. *Proc. Assoc. Am. Physicians* **1997**, *109*, 504–511. [[PubMed](#)]
28. Nishimura, M.; Ueda, M.; Ebata, R.; Utsuno, E.; Ishii, T.; Matsushita, K.; Ohara, O.; Shimojo, N.; Kobayashi, Y.; Nomura, F. A novel KCNQ1 nonsense variant in the isoform-specific first exon causes both jervell and Lange-Nielsen syndrome 1 and long QT syndrome 1: A case report. *BMC Med. Genet.* **2017**, *18*, 66. [[CrossRef](#)]
29. Neyroud, N.; Richard, P.; Vignier, N.; Donger, C.; Denjoy, I.; Demay, L.; Shkolnikova, M.; Pesce, R.; Chevalier, P.; Hainque, B.; et al. Genomic organization of the KCNQ1 K⁺ channel gene and identification of C-terminal mutations in the long-QT syndrome. *Circ. Res.* **1999**, *84*, 290–297. [[CrossRef](#)]
30. Novotny, T.; Kadlecova, J.; Janousek, J.; Gaillyova, R.; Bittnerova, A.; Florianova, A.; Sisakova, M.; Toman, O.; Chroust, K.; Papousek, I.; et al. The homozygous KCNQ1 gene mutation associated with recessive Romano-Ward syndrome. *Pacing Clin. Electrophysiol.* **2006**, *29*, 1013–1015. [[CrossRef](#)]
31. O'Hara, T.; Virag, L.; Varro, A.; Rudy, Y. Simulation of the undiseased human cardiac ventricular action potential: Model formulation and experimental validation. *PLoS Comput. Biol.* **2011**, *7*, e1002061. [[CrossRef](#)]

32. Sato, A.; Arimura, T.; Makita, N.; Ishikawa, T.; Aizawa, Y.; Ushinohama, H.; Aizawa, Y.; Kimura, A. Novel mechanisms of trafficking defect caused by KCNQ1 mutations found in long QT syndrome. *J. Biol. Chem.* **2009**, *284*, 35122–35133. [[CrossRef](#)] [[PubMed](#)]
33. Xu, X.; Kanda, V.A.; Choi, E.; Panaghie, G.; Roepke, T.K.; Gaeta, S.A.; Christini, D.J.; Lerner, D.J.; Abbott, G.W. MinK-dependent internalization of the IKs potassium channel. *Cardiovasc. Res.* **2009**, *82*, 430–438. [[CrossRef](#)] [[PubMed](#)]
34. Xu Parks, X.; Qudsi, H.; Braun, C.; Lopes, C.M.B. The auxiliary subunit KCNE1 regulates KCNQ1 channel response to sustained calcium-dependent PKC activation. *PLoS ONE* **2020**, *15*, e0237591. [[CrossRef](#)]
35. Krumerman, A.; Gao, X.; Bian, J.S.; Melman, Y.F.; Kagan, A.; McDonald, T.V. An LQT mutant minK alters KvLQT1 trafficking. *Am. J. Of Physiol.-Cell Physiol.* **2004**, *286*, C1453–C1463. [[CrossRef](#)] [[PubMed](#)]
36. Bianchi, L.; Priori, S.G.; Napolitano, C.; Surewicz, K.A.; Dennis, A.T.; Memmi, M.; Schwartz, P.J.; Brown, A.M. Mechanisms of I(Ks) suppression in LQT1 mutants. *Am. J. Physiol.-Heart Circ. Physiol.* **2000**, *279*, H3003–H3011. [[CrossRef](#)]
37. Goldenberg, I.; Zareba, W.; Moss, A.J. Long QT Syndrome. *Curr. Probl. Cardiol.* **2008**, *33*, 629–694. [[CrossRef](#)]
38. Rush, S.; Larsen, H. A practical algorithm for solving dynamic membrane equations. *IEEE Trans. Bio-Med. Eng.* **1978**, *25*, 389–392. [[CrossRef](#)]