## Impact parameter sensitive study of inner-shell atomic processes in $Xe^{54+}$ , $Xe^{52+} \rightarrow Xe$ collisions

A Gumberidze<sup>1\*</sup>, C Kozhuharov<sup>1</sup>, R T Zhang<sup>1,2</sup>, S Trotsenko<sup>1</sup>, Y S Kozhedub<sup>3</sup>, R D DuBois<sup>4</sup>, H F. Beyer<sup>1</sup>, K-H Blumenhagen<sup>5,6</sup>, C Brandau<sup>1,7</sup>, A Bräuning-Demian<sup>1</sup>, W Chen<sup>1,8</sup>, O Forstner<sup>5,6</sup>, B Gao<sup>1,9</sup>, T Gassner<sup>1</sup>, R E Grisenti<sup>1,10</sup>, S Hagmann<sup>1,10</sup>, P-M Hillenbrand<sup>1</sup>, P Indelicato<sup>11</sup>, A Kumar<sup>12</sup>, M Lestinsky<sup>1</sup>, Yu A Litvinov<sup>1</sup>, N Petridis<sup>1</sup>, D Schury<sup>1,13</sup>, U Spillmann<sup>1</sup>, C Trageser<sup>1,7</sup>, M Trassinelli<sup>13</sup>, X Tu<sup>1,9</sup> and Th Stöhlker<sup>1,5,6</sup>

<sup>1</sup>GSI Helmholtzzentrum für Schwerionenforschung, 64291 Darmstadt, Germany
<sup>2</sup>Columbia Astrophysics Laboratory, Columbia University, New York, NY 10027, USA
<sup>3</sup>Department of Physics, St. Petersburg State University, St. Petersburg 198504, Russia
<sup>4</sup>Department of Physics, Missouri University of Science and Technology, Rolla, Missouri 65409, USA
<sup>5</sup>Helmholtz-Institut Jena, 07743 Jena, Germany
<sup>6</sup>IOQ, Friedrich-Schiller-Universität Jena, 07743 Jena, Germany
<sup>7</sup>Physikalisches Institut, Justus-Liebig-Universitt Gießen, 35392 Gießen, Germany
<sup>8</sup>Chinese Academy of Sciences, Institute of High Energy Physics, Dongguan 523803, China
<sup>9</sup>Institute of Modern Physics, Chinese Academy of Sciences, 730000, Lanzhou, China
<sup>10</sup>Institut für Kernphysik, Universität Frankfurt, 60486 Frankfurt am Main, Germany
<sup>11</sup>LKB, Sorbonne Université, CNRS, ENS-PSL Research University, Collège de France, F-75005 Paris, France
<sup>12</sup>Nuclear Physics Division, Bhabha Atomic Research Centre, Trombay, Mumbai 400085, India
<sup>13</sup>Institut des NanoSciences de Paris, CNRS, Sorbonne Université, F-75005, Paris, France

Synopsis In this work, we present a pilot experiment in the experimental storage ring (ESR) at GSI devoted to impact parameter sensitive studies of inner shell atomic processes for bare and He-like xenon ions ( $Xe^{54+}$ ,  $Xe^{52+}$ ) colliding with neutral xenon gas atoms. The projectile and target x-rays have been measured at different observation angles for all impact parameters as well as for the impact parameter range of  $\sim 35$  - 70 fm.

In this contribution, we present an impact parameter sensitive study of inner shell atomic processes in symmetric ion-atom collisions. The measurement was performed at the Experimental Storage Ring (ESR) at GSI Darmstadt with bare and He-like xenon ions (Xe<sup>54+</sup>, Xe<sup>52+</sup>) colliding with neutral xenon gas atoms. This choice of the projectile charge states was made in order to compare the effect of a filled K-shell with the empty one. The beam energy (for both charge states) was 50 MeV/u. This value of the beam energy was chosen as a compromise between the adiabaticity of the collision and the reasonable beam lifetime/intensity in the ESR after deceleration. In order to obtain information concerning the impact parameter and, in particular to pick out close collisions which are especially important for observing two-center (or quasi-molecular) effects, the scattered projectile ions which had undergone close collisions with the target atoms were detected by a particle detector (plastic scintillator) mounted in a specially

constructed movable pocket at  $\sim 3.5$  m downstream from the target. In addition to the detector for the scattered projectiles, the x-rays emitted from the interaction zone were observed by an array of semiconductor and scintillator detectors mounted at different angles with respect to the ion beam direction. The physical processes leading to the x-ray emission are: excitation of the projectile/target electrons and electron capture from the neutral target into the highly-charged xenon ions. By looking at the coincidences between the x-rays and the scattered projectiles, we were able to clearly demonstrate the possibility of picking out the characteristic x-rays stemming from the close collisions only. This offers us a new observable for the atomic collision experiments at the ESR storage ring. The experimental results will be presented together with the state-of-theart relativistic calculations [1].

## References

[1] Kozhedub Y S et al 2014 Phys. Rev. A 90 042709

Journal of Physics: Conference Series

<sup>\*</sup>E-mail: a.gumberidze@gsi.de

Content from this work may be used under the terms of the Creative Commons Attribution 3.0 licence. Any further distribution of this work must maintain attribution to the author(s) and the title of the work, journal citation and DOI.