

# A guide to the regulation of selective autophagy receptors

 Andrea Gubas<sup>1</sup>  and Ivan Dikic<sup>1,2,3</sup> 

1 Institute of Biochemistry II, Faculty of Medicine, Goethe University Frankfurt, Germany

2 Buchmann Institute for Molecular Life Sciences, Goethe University Frankfurt, Germany

3 Max Planck Institute of Biophysics, Frankfurt, Germany

## Keywords

autophagy; oligomerisation; phosphorylation; receptor; ubiquitination

## Correspondence

I. Dikic, Institute of Biochemistry II, Faculty of Medicine, Goethe University Frankfurt, Theodor-Stern-Kai 7, 60590 Frankfurt, Germany

Tel: +4969 6301 5964

E-mail: dikic@biochem2.uni-frankfurt.de

(Received 18 January 2021, revised 4 March 2021, accepted 16 March 2021)

doi:10.1111/febs.15824

Autophagy is a highly conserved catabolic process cells use to maintain their homeostasis by degrading misfolded, damaged and excessive proteins, nonfunctional organelles, foreign pathogens and other cellular components. Hence, autophagy can be nonselective, where bulky portions of the cytoplasm are degraded upon stress, or a highly selective process, where preselected cellular components are degraded. To distinguish between different cellular components, autophagy employs selective autophagy receptors, which will link the cargo to the autophagy machinery, thereby sequestering it in the autophagosome for its subsequent degradation in the lysosome. Autophagy receptors undergo post-translational and structural modifications to fulfil their role in autophagy, or upon executing their role, for their own degradation. We highlight the four most prominent protein modifications – phosphorylation, ubiquitination, acetylation and oligomerisation – that are essential for autophagy receptor recruitment, function and turnover. Understanding the regulation of selective autophagy receptors will provide deeper insights into the pathway and open up potential therapeutic avenues.

## Introduction

Cells employ various pathways to maintain their homeostasis. Maintaining the balance between synthesis and removal of proteins, organelles or other cellular components is one of the major tasks of the cellular quality control mechanism. Accumulation of damaged or unwanted cellular components can increase the production of reactive oxygen species or DNA damage, and subsequently lead to cell death [1, 2]. To prevent this, cells have evolved two major pathways – the ubiquitin–proteasome system (UPS) and macroautophagy (hereafter autophagy), both of which are involved in the surveillance and quality control of

proteins and organelles [3,4]. While UPS relies on tagging proteins with ubiquitin to target them for degradation within the barrel-shaped proteasome, autophagy allows the degradation of larger cellular components (including organelles) within the lysosome, mediated by a *de novo*-formed vesicle termed an autophagosome (reviewed in Ref. [3–7]). These pathways are finely regulated, and any disruptions to either of them can lead to neurodegeneration, cancer and other pathologies.

The term ‘autophagy’ was first used by Christian de Duve, in 1963 [8], before gaining significant traction

## Abbreviations

AIM, ATG8-interacting motif; ATG, autophagy-related gene; cvt, cytoplasm-to-vacuole; ER, endoplasmic reticulum; GIM, GABARAP-interacting motif; LIR, LC3-interacting region; LLPS, liquid–liquid phase separation; PAS, pre-autophagosomal structure; PB1, Phox and Bem1; PI3K, phosphatidylinositol (3) kinase; PTMs, post-translational modifications; RHD, reticulon homology domain; SARs, selective autophagy receptors; UPS, ubiquitin–proteasome system.

upon the characterisation of autophagy-related proteins (Atg) in yeast [9,10], and has since been shown to have an essential role in higher eukaryotes. The core autophagy machinery is highly conserved from yeast to human, and it engages five multifunctional systems – (a) ULK1/ULK2 kinase complex (Atg1 in yeast); (b) the class III phosphatidylinositol 3-kinase (PI3K) complex; (c) the LC3/GABARAP proteins and their conjugation machinery (Atg8 in yeast); (d) autophagy-related gene 2 (ATG2; same in yeast) and phosphatidylinositol (e) phosphate effector proteins, including WIPI1–4 proteins (Atg18 and Atg21 in yeast); and (f) the membrane-spanning protein ATG9 (same in yeast; reviewed in Ref. [11,12]). The main event in autophagy is a *de novo* formation of a double membrane called phagophore, at endoplasmic reticulum (ER)-associated sites. The orchestrated action of the autophagy machinery components leads to the expansion of the phagophore, recruitment of the cargo and formation of an autophagosome. Autophagosomes fuse with lysosomes, resulting in the degradation of the cargo and recycling of the nutrients back into the cytosol [11]. Depending on the cargo, autophagy can be selective or

nonselective. In nonselective autophagy, bulk portions of the cytoplasm are sequestered by the phagophore for degradation to maintain the nutrient levels in cells. This type of autophagy is frequently observed upon low nutrient levels during starvation. By contrast, selective autophagy is required for maintaining the number and integrity of cellular organelles, as well as protecting the cell from pathogen invasions. Selected cargo can be mitochondria, portions of ER, ribosomes, peroxisomes, lysosomes, lipid droplets, aggregates, ferritin and others [13,14]. The cargo is distinguished and selected by proteins called cargo receptors, or selective autophagy receptors (SARs), which act as a bridge between the phagophore and cargo (Table 1) [6,13–18]. Generally, cargo receptors are defined and characterised by their ability to bind cargo and facilitate the recruitment of autophagic machinery, mainly through the binding of Atg8/LC3/GABARAP proteins, and finally being degraded with the cargo [19]. The specific action of SARs is regulated by protein modifications, predominantly phosphorylation, ubiquitination, acetylation and oligomerisation, which are the focus of this review.

**Table 1.** Selective autophagy types and corresponding receptors.

Selective type of autophagy	Selected cargo	Species	SARs [references]
Aggrephagy	Aggregates	Yeast	Cue5 [25]
		Plants	NBR1 [112]
		Mammals	NBR1, OPTN, p62, TOLLIP, TAX1BP1 [96, 117–120]
BES1 degradation	BES1	Plants	DSK2A/B [69]
Cvt pathway	Ape1, Ape4, Ams1	Yeast	Atg19, Atg34 [22, 23]
ER-phagy	ER	Yeast	Atg39, Atg40 [26]
		Plants	ATI3, C53 [44, 121]
		Mammals	ATL3, C53, CCPG1, FAM134B, RTN3, Sec62, TEX264, CALCOCO1 [36, 44, 66, 103, 122–126]
Ferritinophagy	Ferritin	Mammals	NCOA4 [56, 127]
Glycophagy	Glycogen	Mammals	STBD1 [128]
Lipophagy	Lipid droplets	Mammals	p62 [129, 130]
Lysophagy	Lysosomes	Mammals	p62, TRIM16 [131, 132]
Midbody removal	Midbody rings	Mammals	NBR1, p62, TRIM16 [133–135]
Mitophagy	Mitochondria	Yeast	Atg32 [27]
		Mammals	AMBRA1, Bcl2L13, BNIP3, Cardiolipin, Ceramide, FKBP8, FUNDC1, NDP52, NIX, NLRX1, OPTN, p62, PHB2, TAX1BP1 [46, 79, 136–147]
NPC-phagy	Nuclear pore complex (NPC)	Yeast	Nup159 [148, 149]
Nucleophagy	Nucleus fragments	Yeast	Atg39 [26]
Pexophagy	Peroxisomes	Yeast	Atg30, Atg36 [28, 29]
		Mammals	NBR1, p62 [150, 151]
Plastid-to-vacuole	Plastids	Plants	ATI1 [152]
Proteaphagy	Proteasome	Plants	RPN10 [153]
Ribophagy	Ribosomes	Mammals	NUFIP1 [154]
Xenophagy	Bacteria and viruses	Mammals	NDP52, OPTN, p62, TAX1BP1 [70, 155–157]
		Plants	NBR1 [158]
Zymophagy	Zymogen granules	Mammals	p62 [159]

## Selective autophagy receptors

The main steps in selective autophagy include recognition of the cargo, coupling of the cargo to the phagophore and degradation of the cargo. In particular, the first two steps depend on SARs – they recognise and bind the cargo, and facilitate formation of the growing phagophore [16].

The first type of selective autophagy was identified in *Saccharomyces cerevisiae*. The cytoplasm-to-vacuole pathway, commonly known as Cvt pathway, delivers the enzymes aminopeptidase 1, aspartyl aminopeptidase and  $\alpha$ -mannosidase (Ape1, Ape4 and Ams1, respectively) into the vacuole [20,21]. The recognition of these enzymes is achieved by Atg19 and Atg34 receptors. Both Atg19 and Atg34 can specifically bind the cargo proteins, Atg11 – the scaffolding subunit of Atg1 complex – and Atg8 proteins, which decorate the pre-autophagosomal structure (PAS), thereby recruiting the cargo into the Cvt pathway [22–24]. Selective degradation of mitochondria, peroxisomes, ER, parts of nuclei and aggregates by autophagy has also been studied in yeast, with each of these degradation pathways employing autophagy receptors. Yeast SARs commonly bind Atg8 and Atg11, with the exception of Cue5, which has shown no interaction with Atg11 [25–32]. In yeast, binding to Atg8 proteins is mediated by a W/F/YxxL/I/V motif, commonly known as Atg8-interacting motif (AIM), characterised by hydrophobic residues on positions 0 and +3, and often surrounded by acidic residues to stabilise the interaction [33].

Higher eukaryotes come with additional layers of complexity to selective autophagy pathways, which have been extensively studied. In mammals, over 30 autophagy receptors have been identified and described (reviewed in Ref. [13,14]). Yeast Atg8 has six homologues in mammals, corresponding to LC3 and GABARAP families – LC3A, LC3B, LC3C and GABARAP, GABARAP L1 and GABARAP L2 (also known as GATE-16). Mammalian SARs bind LC3/GABARAP proteins through conserved AIM-like LC3-interaction region and GABARAP-interacting motif (LIRs and GIMs, respectively) [33–35]. The LIR/GIM region, similarly to AIM, is characterised by hydrophobic residues at positions 0 and +3 and frequently flanked by negatively charged amino acids that contribute to reinforcing the binding [34,35].

Interestingly, whereas the majority of yeast SARs interact with the scaffold protein Atg11, which recruits Atg1 complex to cargo, only some mammalian SARs interact with its functional mammalian homologue, FAK-family-interacting protein of 200 kDa (FIP200), which can drive the recruitment and activation of

ULK1 kinase and the downstream machinery to the selected cargo, thereby facilitating phagophore formation [36–43]. In plants, selective autophagy has not been as extensively studied; however, a number of SARs have been reported to interact with ATG8 in an AIM-dependent manner [44,45].

In addition to ATG8/LC3/GABARAP-binding sites, SARs generally possess cargo-binding domains. These are often ubiquitin-binding domains, since autophagic cargo is frequently ubiquitinated. Ubiquitin-independent cargo recognition involves SARs binding directly to the cargo, as well as SARs recognising sugars and lipids exposed on the cargo (Fig. 1; [46–51], reviewed in Ref. [52]).


## Regulation of autophagy receptors

For its full functionality and to minimise costly energy losses, autophagy flux is tightly regulated (reviewed in Ref. [53–55]). SARs undergo structural and post-translational modifications (PTMs) in order to operate in a specific type of autophagy, to facilitate binding to other proteins or simply to be removed via the proteasome. Below, we outlined the most prominent SAR modifications, such as phosphorylation, ubiquitination, acetylation and oligomerisation (Fig. 2; Table 2), and explain how they affect SARs within autophagy pathways.

### Ubiquitination

Ubiquitination is commonly mentioned in the context of protein degradation, as it serves as a signal for protein removal by either the proteasome or autophagy [6]. SARs are also ubiquitinated for their own degradation. An example is NCOA4, a ferritinophagy receptor, involved in selective degradation of ferritin. Ferritin is a protein that serves to store iron in cells and release it in a controlled fashion when iron levels are low [56,57]. NCOA4 is an iron-sensing protein, which engages its C-terminal helical domain in coordinating iron. As such, during high-iron levels, NCOA4 interacts with and is ubiquitinated by HERC2 ubiquitin ligase, thereby preventing ferritin degradation and iron release [57]. Regulation of ferritinophagy is essential for maintaining iron levels in cells, and any perturbations within this process could lead to pathologies, such as anaemia, or neurodegeneration, such as neurodegeneration with brain iron accumulation (NBIA) [58].

Ubiquitination may not only signal for SAR degradation, but also regulate their function in autophagy. For example, p62, a SAR involved in the regulation of




**Fig. 1.** Types of cargo recognition in selective autophagy. During selective autophagy, cargo is carefully selected in several ways for its degradation within the lysosome. (I) In ubiquitin-dependent autophagy, cargo is polyubiquitinated, which serves as a signal for its degradation. Cargo receptors can simultaneously bind ATG8 proteins and polyubiquitin on the cargo, providing a strong link between autophagy machinery and the cargo itself. (II) Cargo receptors can also bind sugars, such as lectins, that are recruited to the cargo, as is the case during lysophagy, where galectin 3 decorates damaged lysosomes and is able to bind the receptors. (III) SARs can also bind directly to the cargo and deliver it to the forming phagophore through the ATG8 binding. (IV) Some lipids (such as cardiolipin) can bind ATG8s and provide a direct link between cargo and autophagy machinery in a receptor-independent manner. Phagophore elongates, surrounding the cargo. When it seals, it fuses with the lysosome and the cargo is degraded.

redox homeostasis, undergoes nondegradative ubiquitination. During normal cellular conditions, Kelch-like ECH-associated protein 1 (KEAP1) interacts with nuclear factor erythroid 2-related factor 2 (NRF2), a transcription factor involved in gene expression of antioxidant proteins. Through its association with Cullin3 E3 ligase, KEAP1 mediates the ubiquitination of NRF2, targeting it for proteasomal degradation. Upon oxidative stress, it is the task of p62 to sequester KEAP1 within aggregates for autophagic degradation in order to release NRF2, which can then translocate to the nucleus and trigger an antioxidative response (reviewed in Ref. [59]). Pan and colleagues have reported that this process can be blocked by ubiquitination of p62 at Lys7 [60], which would normally form a hydrogen bond with Asp69, driving homodimerisation of p62 via its Phox and Bem1 (PB1) domain [61,62]. PB1-mediated homodimerisation is a prerequisite for its sequestering function. Ubiquitinated p62

can no longer mediate autophagic degradation of KEAP1, suppressing the release of NRF2 and stimulating antioxidative actions to restore redox homeostasis [60].

Conversely, SAR ubiquitination can also be a signal for autophagy activation and cargo recognition. KEAP1/Cullin3 also directly ubiquitinates p62 at Lys420, within its UBA domain. The ubiquitination prevents p62 dimer formation through the UBA domain, thereby enhancing its sequestration ability and interactions with phagophores and LC3/GABARAP proteins during ubiquitin stress [63,64]. A similar ubiquitination pattern has been observed for NBR1, with ubiquitin-stress-triggered ubiquitination likely regulating its interaction with polyubiquitin chains [63]. Furthermore, OPTN stably interacts with HACE1 E3 ubiquitin ligase, which leads to HACE1-mediated ubiquitination of OPTN at Lys193, facilitating the complex formation between OPTN, p62 and



**Fig. 2.** Summary of regulatory modifications reported for SARs. SARs are regulated through post-translational, structural and other modifications, most prominent ones summarised in this scheme. SARs and their modified sites are listed under the regulation modification that affects them.

HACE1, as well as to coupling of cargo to the autophagic machinery. HACE1 is downregulated in various cancers, suggesting that the HACE1-OPTN axis may play a role in tumour suppression [65].

There is also preliminary evidence for functional regulation of ER-phagy receptors, but not much is known about ubiquitination of ER-phagy SARs.

Atlastins 1 and 3 (ATL1 and ATL3), membrane-bound dynamin-like GTPases, have been found to function as ER-phagy receptors, binding GABARAPs and targeting parts of the tubular ER for autophagy [66]. Atlastins are involved in the fusion of ER tubules, which is regulated through ubiquitination by SYVN1 E3 ligase [67].



**Table 2.** Autophagy-associated SAR modifications with corresponding enzymes.

Protein	Species	Residues (enzymes, if known)	References
<b>Phosphorylation</b>			
p62	Mammals	S24 (PKA), S403 (CK2, ULK1, TBK1), S407 (ULK1)	[80–82, 160]
OPTN	Mammals	S177, S473 (both TBK1)	[50, 70]
FUNDC1	Mammals	S13 (CK2/phosphatase PGAM5), Y18 (SRC)	[72, 73]
NIX	Mammals	S34, S35, S212	[71, 102]
BNIP3	Mammals	S17, S24	[161]
FAM134B	Mammals	S151 (CAMK2B)	[83]
Atg19	Yeast	S390, S391, S396 (Hrr25)	[75, 77]
Atg30	Yeast	S71, S112	[31]
Atg32	Yeast	T112, S114 (CK2), S119 (CK2)	[31, 74]
Atg36	Yeast	S31 (N/A), S97 (Hrr25)	[31, 77]
DSK2	Plants	S240, S244 (+more) (BIN2)	[69]
<b>Ubiquitination</b>			
p62	Mammals	K7 (TRIM21), K420 (Cullin3)	[60, 63, 64]
OPTN	Mammals	K193 (HACE1)	[65]
ATL1	Mammals	K285, K287 (SYVN1)	[67]
NCOA4	Mammals	N/A (HERC2)	[57]
Atg19	Yeast	K213, K216 (Ubp3p)	[68]
Cue5	Yeast	N/A (Rsp5)	[25]
<b>Acetylation</b>			
p62	Mammals	K420, K435	[86]
FUNDC1	Mammals	K114, K115 (predicted)	[89]
<b>Oligomerisation</b>			
p62	Mammals	K7, R21, D69, E82, E409, G410	[62, 99]
NBR1	Mammals		[96]
OPTN	Mammals	E50	[43, 162]
NIX	Mammals	G204, G208	[102]
FAM134B	Mammals	G216	[83]
RTN3	Mammals		[103]
Atg19	Yeast		[98]
NBR1	Plants	K11, K19, D60, D73	[112]
<b>UFMylation</b>			
C53	Mammals Plants	N/A (UFL1)	[44]

In yeast, not many examples of ubiquitination events regulating cargo receptor function have been reported. However, Baxter and colleagues have shown that Atg19, the Cvt pathway receptor, interacts with and is ubiquitinated by the E3 ligase Ubp3 at lysines 213 and 216. Downregulation of this ubiquitination results in decreased Cvt targeting [68].

Ubiquitination is one of the most abundant PTMs in cells, and it is likely that many more SARs are regulated in this way. Dysregulation of autophagy can

have pathological consequences, and thus, understanding the regulation of SARs by ubiquitination may provide the basis for therapeutic approaches.

## Phosphorylation

Phosphorylation plays an essential part in regulating autophagic flux and is central to autophagy induction. Moreover, it plays an important role in regulating the ability of SARs to bridge the cargo with autophagic machinery. In most cases, phosphorylation regulates SARs binding to Atg8/LC3/GABARAP [31,69–73]. This has been shown already in yeast, and is conserved across species, including humans. In yeast, pexophagy receptors Atg30, Atg32 and Atg36 interact with Atg8 in an AIM-dependent manner. These interactions are facilitated by phosphorylation at residues upstream of the respective AIM regions [31]. Furthermore, Hrr25 and casein kinase 2 (CK2) have been shown to phosphorylate yeast SARs and facilitate their interaction with the scaffold protein Atg11 [74–78].

Plant autophagy SARs are also modified by phosphorylation. During stress conditions, such as drought and starvation, it is required to strengthen the interaction of SARs, such as DSK2, with ATG8 [69].

In mammals, TBK1 phosphorylates OPTN at Ser177 during xenophagy, which strengthens its interaction with LC3s/GABARAPs and subsequently restricts *Salmonella* growth [70], and at Ser473, which facilitates its interaction with ubiquitin chains during mitophagy [79]. Similarly, phosphorylation of Nix at Ser34 and Ser35 enhances its interaction with LC3s and initiates mitophagy [71]. Conversely, phosphorylation of FUNDC1 at Tyr18 prevents binding to LC3s/GABARAPs, thereby inhibiting mitophagy [72]. During hypoxia, the interaction of FUNDC1 with LC3/GABARAP proteins is determined by the phosphorylation of Ser13, which is regulated by the interplay of a phosphoglycerate mutase (PGAM5) phosphatase and casein kinase 2 [73].

p62 function in autophagy is also regulated by various kinases. During proteotoxic stress, the phosphorylation at Ser407 by ULK1 kinase allows the transition of p62 from dimer to a monomer, promoting its binding to ubiquitin, and allowing its recruitment to protein aggregates [80]. Similarly, phosphorylation at Ser403 by various kinases enhances the binding to ubiquitin chains [80–82], subsequently promoting cargo degradation.

Phosphorylation also positively regulates FAM134B-dependent ER-phagy. CAMK2B kinase phosphorylates FAM134B at Ser151 within its reticulation homology domain (RHD) in order to facilitate

FAM134B dimerisation and membrane fragmentation, thereby making it accessible for autophagosome sequestration [83]. Many other SARs undergo phosphorylation modifications as part of their regulation mechanism, with further studies needed to understand them and identify new ones [79,84,85].

### Acetylation

Although equally essential, it took some time for acetylation to attract attention of autophagy researchers, with new roles now rapidly emerging. Acetylation involves the addition of an acetyl group onto lysines or amino-termini of proteins. During selective autophagy, UBA domain-mediated oligomerisation of p62 and its interaction with polyubiquitinated cargo are regulated by acetylation. The activation of acetyltransferase TIP60 drives p62 acetylation at Lys420 and Lys435, facilitating the binding to polyubiquitinated proteins [86]. Ser403 and Ser407 phosphorylation, found in close proximity to acetylated lysines, regulates a similar function of p62 [80,81]. It is likely that there is a link between the two PTMs, although further research is needed to demonstrate this.

LC3 proteins, which shuttle between the nucleus and the cytosol, are also regulated by acetylation. Upon starvation, LC3 is deacetylated by Sirt1 at lysines 49 and 51, driving its translocation from the nucleus into the cytosol, where it is able to participate in the process of autophagy [87].

Acetylation is implicated in regulating ER-phagy SARs through the modification of ATG9A protein. ATG9A interacts with FAM134B and Sec62, two ER-phagy receptors. This interaction is reduced upon acetylation of ATG9A, which prevents binding of ER-phagy receptors to LC3s and inhibits ER-phagy induction. This can be enforced in transgenic mice that overexpress ATase1 (AT-1) in forebrain neurons, causing a progeria-like phenotype. AT-1 is required for translocating cytosolic acetyl-CoA into the ER lumen [88].

FUNDC1 is also possibly regulated by acetylation, as potential acetylation sites on it have been predicted [89], but more work is required to understand its effect and relevance. Acetylation offers an interesting approach in terms of potential therapeutic targets, as exemplified by AT-1 and ER-phagy. However, our knowledge of this modification in autophagy is still at the very beginning.

### Oligomerisation

Protein function, structure or localisation is not only regulated by PTMs. Oligomerisation has been shown

to have an important part in regulating the induction of autophagy, either by promoting kinase activation, such as oligomerisation of ULK1 [90], or by providing the structural architecture for protein scaffold formation on the phagophore, with class III PI3K complex and ATG16L1 as good examples [91,92].

In yeast, it has recently been shown that oligomerisation of Atg17 is prerequisite for liquid–liquid phase separation (LLPS) of the PAS [93,94]. Moreover, LLPS has been shown to regulate the lysosomal degradation of Ede-1-dependent endocytic protein deposits, which are formed as a consequence of defects in early stages of clathrin-mediated endocytosis. In this case, Ede-1 has a role of cargo receptor, and its polymerisation drives LLPS. Through its binding to Atg8 and Atg11, Ede-1 drives the degradation of the formed condensates [95].

Many SARs contain a region in their sequence that allows oligomerisation. Autophagic cargo is generally larger than that of the proteasome, as it mainly consists of organelles or macromolecules, which can attribute to the necessity of autophagy receptors to oligomerise. Furthermore, oligomerisation of SARs allows better exposure of their domains and moieties, as well as increases their binding avidity. To better fulfil their role as bridges between autophagic machinery and cargo, many SARs homo- and hetero-oligomerise, as shown for p62 and NBR1 [96,97]. Additionally, autophagic cargo itself is known to oligomerise, to allow efficient autophagic uptake, as it is the case with ferritin during ferritinophagy [56].

During the yeast Cvt pathway, the cargo proteins Ape1 and Ams1 form dodecamers and tetramers, respectively, allowing efficient cargo packaging. Ape1, as more abundant cargo, carries Ams1 to facilitate its binding to Atg19, which forms a trimer. Atg19 trimers are incorporated into the Ape1/Ams1 aggregates, thereby controlling the size of the assembled cargo. These large aggregates will be efficiently engulfed by the Cvt vesicle and delivered to the vacuole [98].

In mammals, oligomerisation has been extensively studied in the context of SAR regulation. The effect of PTMs on p62 has frequently been attributed to the regulation of its ability to oligomerise. p62 can oligomerise through its PB1 domain and through its UBA domain, which has different consequences for autophagy [62,99]. p62 polymerisation has been found to be one of the requirements of p62 to undergo phase separation, along with ubiquitin binding through its UBA domain, which drives the concentration and sequestration of the autophagic cargo [49].

It has been shown that OPTN forms homo-hexamers, with its role currently unknown [100]. However,

OPTN E50K mutation, associated with normal tension glaucoma (NTG), has been shown to inhibit oligomerisation and solubility of OPTN, and at the same time strongly enhance its interaction with TBK1 [101].

Recently, Nix has been shown to dimerise, thereby stabilising its interaction with LC3A and its function as a cargo receptor. This dimerisation is regulated by phosphorylation at Ser212 [102].

ER-phagy cargo receptors commonly undergo oligomerisation as part of their regulation mechanism. FAM134B oligomerisation is facilitated by the phosphorylation of its RHD by CAMK2B kinase, regulating ER membrane scission for ER-phagy demands [83]. RTN3L, a long isoform of RTN3 reticulon, can homodimerise and heterodimerise with RTN3S isoform, and both have different functions during ER-phagy [103]. RTN3L homodimers can drive fragmentation of ER tubules, which will be subsequently delivered to the lysosome for degradation [103]. Conversely, RTN3L-RTN3S heterodimers contribute to stabilisation of ER tubules instead [103]. In *Drosophila*, Atlastin has also shown a tendency to oligomerise, in GTP-dependent manner. The formation of this homo-oligomer is required for ER membrane tethering and fusion [104,105], with similar role observed with mammalian Atlastins ATL1, ATL2 and ATL3 [106–111]. However, it remains to be discovered how oligomerisation affects ATL1 and ATL2 receptor functions in mammals.

Plant SARs also rely on oligomerisation for their function. Plant NBR1, a functional hybrid of mammalian NBR1 and p62, undergoes oligomerisation via its PB1 domain, similarly to p62, which is required for its role in autophagy [112]. Oligomerisation has also been observed in other SARs, with its function yet to be completely understood [57,113,114], but likely involved in facilitating the linkage of SARs with the cargo and autophagic machinery.

## Concluding remarks

With various cell types and different organelles, macromolecules, aggregates and pathogens all targeted for degradation by autophagy, across species, cells have developed a large number of SARs, and evolved tight regulatory mechanisms, since most of the SARs also have additional functions to being cargo receptors. PTMs such as phosphorylation, ubiquitination and acetylation have been extensively studied (Fig. 2; Table 2); hence, much is known about their role in regulating SARs. In this review, we summarised the most prominent and best characterised modifications of SARs and how they affect the function of SARs in

their respective pathways. Along with PTMs, translational (reviewed in Ref. [115]) and structural changes can be involved in regulating the SAR function. The orchestrated action of multiple PTMs is a rapidly emerging theme in autophagy and beyond. Such cross-talk of PTMs provides an additional layer of complexity and fine-tuning of regulatory mechanisms. Phosphorylation is frequently reported to regulate SAR binding to ubiquitin chains of the targeted cargo [50,80,81], but it can also regulate ubiquitination of the protein itself. PTMs can regulate the same function of one SAR, as shown with p62, with acetylation and phosphorylation, in addition to oligomerisation, all regulating its binding to ubiquitinated cargo [80,81,86]. Conversely, same cargo receptors can be regulated by several different modifications, all of which will regulate different parts of their function, again with p62 being best characterised example [43,63,80–82,85,86,97].

The question of additional PTMs in relation to SARs remains largely unexplored part of the field, but processes such as methylation, glycosylation, SUMOylation and ISGylation could have an important role in SAR regulation. UFMylation has recently emerged as an interesting process involved in regulating ER-phagy [44,116] (Fig. 2; Table 2), with ER-phagy receptor C53 being the only ER-phagy cargo receptor known to be regulated by UFMylation thus far. This regulation has been shown in plants and is conserved in mammals [44]. The rise of UFMylation within the field provides novel insights into the regulatory mechanism of ER-phagy.

And while particular modifications have been identified for many SARs, the identity of enzymes catalysing said modifications is not always clear. Therefore, it is important to understand the identity of kinases, phosphatases, E3 ligases, deubiquitinases, acetylases, deacetylases and others, especially in the context of pathologies. Since any disorder in the process of autophagy, as well as SARs, has been linked to changes in immune response and, therefore pathologies, such as infection, neurodegeneration and cancer, further research of the regulatory mechanisms of SARs will shed light and open new avenues of therapeutic strategies.

## Acknowledgements

We thank Koraljka Husnjak and Daniela Höller for critical reading of the manuscript. Our research is supported by the Deutsche Forschungsgemeinschaft (DFG, German Research Foundation) – Project Number 259130777 – SFB 1177, the grants from Else



Kroener Fresenius Stiftung and Dr. Rolf M. Schwiете Stiftung.

## Conflict of interest

The authors declare no conflict of interest.

## Author contributions

AG and ID wrote and revised the manuscript and designed the figures.

## References

- Guo C & Zhao Y (2020) Autophagy and DNA damage repair. *Genome Instab Dis* **1**, 172–183.
- Filomeni G, De Zio D & Cecconi F (2015) Oxidative stress and autophagy: the clash between damage and metabolic needs. *Cell Death Differ* **22**, 377–388.
- Ji CH & Kwon YT (2017) Crosstalk and interplay between the ubiquitin-proteasome system and autophagy. *Mol Cells* **40**, 441–449.
- Kocaturk NM & Gozuacik D (2018) Crosstalk between mammalian autophagy and the ubiquitin-proteasome system. *Front Cell Dev Biol* **6**, 128.
- Lamark T & Johansen T (2010) Autophagy: links with the proteasome. *Curr Opin Cell Biol* **22**, 192–198.
- Dikic I (2017) Proteasomal and autophagic degradation systems. *Annu Rev Biochem* **86**, 193–224.
- Dikic I & Elazar Z (2018) Mechanism and medical implications of mammalian autophagy. *Nat Rev Mol Cell Biol* **19**, 349–364.
- Klionsky DJ (2008) Autophagy revisited: a conversation with Christian de Duve. *Autophagy* **4**, 740–743.
- Tsukada M & Ohsumi Y (1993) Isolation and characterization of autophagy-defective mutants of *Saccharomyces cerevisiae*. *FEBS Lett* **333**, 169–174.
- Thumm M, Egner R, Koch B, Schlumpberger M, Straub M, Veenhuis M & Wolf DH (1994) Isolation of autophagocytosis mutants of *Saccharomyces cerevisiae*. *FEBS Lett* **349**, 275–280.
- Mercer TJ, Gubas A & Tooze SA (2018) A molecular perspective of mammalian autophagosome biogenesis. *J Biol Chem* **293**, 5386–5395.
- Farré JC & Subramani S (2016) Mechanistic insights into selective autophagy pathways: lessons from yeast. *Nat Rev Mol Cell Biol* **17**, 537–552.
- Kirkin V & Rogov VV (2019) A diversity of selective autophagy receptors determines the specificity of the autophagy pathway. *Mol Cell* **76**, 268–285.
- Johansen T & Lamark T (2020) Selective autophagy: ATG8 family proteins, LIR motifs and cargo receptors. *J Mol Biol* **432**, 80–103.
- Rogov V, Dötsch V, Johansen T & Kirkin V (2014) Interactions between autophagy receptors and ubiquitin-like proteins form the molecular basis for selective autophagy. *Mol Cell* **53**, 167–178.
- Stolz A, Ernst A & Dikic I (2014) Cargo recognition and trafficking in selective autophagy. *Nat Cell Biol* **16**, 495–501.
- Kimura T, Mandell M & Deretic V (2016) Precision autophagy directed by receptor regulators - emerging examples within the TRIM family. *J Cell Sci* **129**, 881–891.
- Fracchiolla D, Sawa-Makarska J, Zens B, de Ruiter A, Zaffagnini G, Brezovich A, Romanov J, Runggatscher K, Kraft C, Zagrovic B *et al.* (2016) Mechanism of cargo-directed Atg8 conjugation during selective autophagy. *Elife* **5**, e18544.
- Galluzzi L, Baehrecke EH, Ballabio A, Boya P, Bravo-San Pedro JM, Cecconi F, Choi AM, Chu CT, Codogno P, Colombo MI *et al.* (2017) Molecular definitions of autophagy and related processes. *EMBO J* **36**, 1811–1836.
- Lynch-Day MA & Klionsky DJ (2010) The Cvt pathway as a model for selective autophagy. *FEBS Lett* **584**, 1359–1366.
- Yuga M, Gomi K, Klionsky DJ & Shintani T (2011) Aspartyl aminopeptidase is imported from the cytoplasm to the vacuole by selective autophagy in *Saccharomyces cerevisiae*. *J Biol Chem* **286**, 13704–13713.
- Scott SV, Guan J, Hutchins MU, Kim J & Klionsky DJ (2001) Cvt19 is a receptor for the cytoplasm-to-vacuole targeting pathway. *Mol Cell* **7**, 1131–1141.
- Suzuki K, Kondo C, Morimoto M & Ohsumi Y (2010) Selective transport of  $\alpha$ -mannosidase by autophagic pathways: identification of a novel receptor, Atg34p. *J Biol Chem* **285**, 30019–30025.
- Leber R, Silles E, Sandoval IV & Mazon MJ (2001) Yol082p, a novel CVT protein involved in the selective targeting of aminopeptidase I to the yeast vacuole. *J Biol Chem* **276**, 29210–29217.
- Lu K, Psakhye I & Jentsch S (2014) Autophagic clearance of PolyQ proteins mediated by ubiquitin-Atg8 adaptors of the conserved CUET protein family. *Cell* **158**, 549–563.
- Mochida K, Oikawa Y, Kimura Y, Kirisako H, Hirano H, Ohsumi Y & Nakatogawa H (2015) Receptor-mediated selective autophagy degrades the endoplasmic reticulum and the nucleus. *Nature* **522**, 359–362.
- Kanki T, Wang K, Cao Y, Baba M & Klionsky DJ (2009) Atg32 is a mitochondrial protein that confers selectivity during mitophagy. *Dev Cell* **17**, 98–109.
- Farré JC, Manjithaya R, Mathewson RD & Subramani S (2008) PpAtg30 tags peroxisomes for turnover by selective autophagy. *Dev Cell* **14**, 365–376.

- 29 Motley AM, Nuttall JM & Hettema EH (2012) Pex3-anchored Atg36 tags peroxisomes for degradation in *Saccharomyces cerevisiae*. *EMBO J* **31**, 2852–2868.
- 30 Okamoto K, Kondo-Okamoto N & Ohsumi Y (2009) Mitochondria-anchored receptor Atg32 mediates degradation of mitochondria via selective autophagy. *Dev Cell* **17**, 87–97.
- 31 Farré JC, Burkenroad A, Burnett SF & Subramani S (2013) Phosphorylation of mitophagy and pexophagy receptors coordinates their interaction with Atg8 and Atg11. *EMBO Rep* **14**, 441–449.
- 32 Nakatogawa H & Mochida K (2015) Reticulophagy and nucleophagy: new findings and unsolved issues. *Autophagy* **11**, 2377–2378.
- 33 Noda NN, Ohsumi Y & Inagaki F (2010) Atg8-family interacting motif crucial for selective autophagy. *FEBS Lett* **584**, 1379–1385.
- 34 Rogov VV, Stolz A, Ravichandran AC, Rios-Szweid DO, Suzuki H, Kniss A, Löhr F, Wakatsuki S, Dötsch V, Dikic I *et al.* (2017) Structural and functional analysis of the GABARAP interaction motif (GIM). *EMBO Rep* **18**, 1382–1396.
- 35 Wirth M, Zhang W, Razi M, Nyoni L, Joshi D, O'Reilly N, Johansen T, Tooze SA & Mouilleron S (2019) Molecular determinants regulating selective binding of autophagy adapters and receptors to ATG8 proteins. *Nat Commun* **10**, 2055.
- 36 Smith MD, Harley ME, Kemp AJ, Wills J, Lee M, Arends M, von Kriegsheim A, Behrends C & Wilkinson S (2018) CCPG1 Is a Non-canonical autophagy cargo receptor essential for ER-Phagy and pancreatic ER proteostasis. *Dev Cell* **44**, 217–232.
- 37 Hara T, Takamura A, Kishi C, Iemura SI, Natsume T, Guan JL & Mizushima N (2008) FIP200, a ULK-interacting protein, is required for autophagosome formation in mammalian cells. *J Cell Biol* **181**, 497–510.
- 38 Fujita N, Morita E, Itoh T, Tanaka A, Nakaoka M, Osada Y, Umemoto T, Saitoh T, Nakatogawa H, Kobayashi S *et al.* (2013) Recruitment of the autophagic machinery to endosomes during infection is mediated by ubiquitin. *J Cell Biol* **203**, 115–128.
- 39 Gammoh N, Florey O, Overholtzer M & Jiang X (2013) Interaction between FIP200 and ATG16L1 distinguishes ULK1 complex-dependent and-independent autophagy. *Nat Struct Mol Biol* **20**, 144–149.
- 40 Nishimura T, Kaizuka T, Cadwell K, Sahani MH, Saitoh T, Akira S, Virgin HW & Mizushima N (2013) FIP200 regulates targeting of Atg16L1 to the isolation membrane. *EMBO Rep* **14**, 284–291.
- 41 Ravenhill BJ, Boyle KB, von Muhlinen N, Ellison CJ, Masson GR, Otten EG, Foeglein A, Williams R & Randow F (2019) The Cargo receptor NDP52 initiates selective autophagy by recruiting the ULK complex to cytosol-invading bacteria. *Mol Cell* **74**, 320–329.e6.
- 42 Turco E, Witt M, Abert C, Bock-Bierbaum T, Su MY, Trapannone R, Sztacho M, Danieli A, Shi X, Zaffagnini G *et al.* (2019) FIP200 claw domain binding to p62 promotes autophagosome formation at ubiquitin condensates. *Mol Cell* **74**, 330–346.
- 43 Vargas JNS, Wang C, Bunker E, Hao L, Maric D, Schiavo G, Randow F & Youle RJ (2019) Spatiotemporal control of ULK1 activation by NDP52 and TBK1 during selective autophagy. *Mol Cell* **74**, 347–362.e6.
- 44 Stephani M, Picchianti L, Gajic A, Beveridge R, Skarwan E, Hernandez VSM, Mohseni A, Clavel M, Zeng Y, Naumann C *et al.* (2020) A cross-kingdom conserved er-phagy receptor maintains endoplasmic reticulum homeostasis during stress. *Elife* **9**, e58396.
- 45 Stephani M & Dagdas Y (2020) Plant selective autophagy—still an uncharted territory with a lot of hidden gems. *J Mol Biol* **432**, 63–79.
- 46 Chu CT, Ji J, Dagda RK, Jiang JF, Tyurina YY, Kapralov AA, Tyurin VA, Yanamala N, Shrivastava IH, Mohammadyani D *et al.* (2013) Cardiolipin externalization to the outer mitochondrial membrane acts as an elimination signal for mitophagy in neuronal cells. *Nat Cell Biol* **15**, 1197–1205.
- 47 Wauer T, Simicek M, Schubert A & Komander D (2015) Mechanism of phospho-ubiquitin-induced PARKIN activation. *Nature* **524**, 370–374.
- 48 Danieli A & Martens S (2018) P62 - Mediated phase separation at the intersection of the ubiquitin-proteasome system and autophagy. *J Cell Sci* **131**, jcs214304.
- 49 Sun D, Wu R, Zheng J, Li P & Yu L (2018) Polyubiquitin chain-induced p62 phase separation drives autophagic cargo segregation. *Cell Res* **28**, 405–415.
- 50 Richter B, Sliter DA, Herhaus L, Stolz A, Wang C, Beli P, Zaffagnini G, Wild P, Martens S, Wagner SA *et al.* (2016) Phosphorylation of OPTN by TBK1 enhances its binding to Ub chains and promotes selective autophagy of damaged mitochondria. *Proc Natl Acad Sci USA* **113**, 4039–4044.
- 51 Maejima I, Takahashi A, Omori H, Kimura T, Takabatake Y, Saitoh T, Yamamoto A, Hamasaki M, Noda T, Isaka Y *et al.* (2013) Autophagy sequesters damaged lysosomes to control lysosomal biogenesis and kidney injury. *EMBO J* **32**, 2336–2347.
- 52 Khaminets A, Behl C & Dikic I (2016) Ubiquitin-dependent and independent signals in selective autophagy. *Trends Cell Biol* **26**, 6–16.
- 53 Wani WY, Boyer-Guittaut M, Dodson M, Chatham J, Darley-Usmar V & Zhang J (2015) Regulation of autophagy by protein post-translational modification. *Lab Invest* **95**, 14–25.

- 54 McEwan DG & Dikic I (2011) The three musketeers of autophagy: phosphorylation, ubiquitylation and acetylation. *Trends Cell Biol* **21**, 195–201.
- 55 Xie Y, Kang R, Sun X, Zhong M, Huang J, Klionsky DJ & Tang D (2015) Posttranslational modification of autophagy-related proteins in macroautophagy. *Autophagy* **11**, 28–45.
- 56 Mancias JD, Wang X, Gygi SP, Harper JW & Kimmelman AC (2014) Quantitative proteomics identifies NCOA4 as the cargo receptor mediating ferritinophagy. *Nature* **509**, 105–109.
- 57 Mancias JD, Vaites LP, Nissim S, Biancur DE, Kim AJ, Wang X, Liu Y, Goessling W, Kimmelman AC & Harper JW (2015) Ferritinophagy via NCOA4 is required for erythropoiesis and is regulated by iron dependent HERC2-mediated proteolysis. *Elife* **4**, e10308.
- 58 Del Rey MQ & Mancias JD (2019) NCOA4-mediated ferritinophagy: a potential link to neurodegeneration. *Front Neurosci* **13**, 238.
- 59 Katsuragi Y, Ichimura Y & Komatsu M (2016) Regulation of the Keap1–Nrf2 pathway by p62/SQSTM1. *Curr Opin Toxicol* **1**, 54–61.
- 60 Pan JA, Sun Y, Jiang YP, Bott AJ, Jaber N, Dou Z, Yang B, Chen JS, Catanzaro JM, Du C *et al.* (2016) TRIM21 ubiquitylates SQSTM1/p62 and suppresses protein sequestration to regulate redox homeostasis. *Mol Cell* **61**, 720–733.
- 61 Wilson MI, Gill DJ, Perisic O, Quinn MT & Williams RL (2003) PB1 domain-mediated heterodimerization in NADPH oxidase and signaling complexes of atypical protein kinase C with Par6 and p62. *Mol Cell* **12**, 39–50.
- 62 Lamark T, Perander M, Outzen H, Kristiansen K, Øvervatn A, Michaelsen E, Bjørkoy G & Johansen T (2003) Interaction codes within the family of mammalian Phox and Bem1p domain-containing proteins. *J Biol Chem* **278**, 34568–34581.
- 63 Peng H, Yang J, Li G, You Q, Han W, Li T, Gao D, Xie X, Lee BH, Du J *et al.* (2017) Ubiquitylation of p62/sequestosome1 activates its autophagy receptor function and controls selective autophagy upon ubiquitin stress. *Cell Res* **27**, 657–674.
- 64 Lee YJ, Chou TF, Pittman SK, Keith AL, Razani B & Wehl CC (2017) Keap1/Cullin3 modulates p62/SQSTM1 activity via UBA domain ubiquitylation. *Cell Rep* **20**, 1994.
- 65 Liu Z, Chen P, Gao H, Gu Y, Yang J, Peng H, Xu X, Wang H, Yang M, Liu X *et al.* (2014) Ubiquitylation of autophagy receptor optineurin by HACE1 activates selective autophagy for tumor suppression. *Cancer Cell* **26**, 106–120.
- 66 Chen Q, Xiao Y, Chai P, Zheng P, Teng J & Chen J (2019) ATL3 Is a tubular ER-Phagy receptor for GABARAP-mediated selective autophagy. *Curr Biol* **29**, 846–855.
- 67 Zhao Y, Feng Z, Zou Y & Liu Y (2020) The E3 ubiquitin ligase SYVN1 ubiquitinates atlastins to remodel the endoplasmic reticulum network. *iScience* **23**, 101494.
- 68 Baxter BK, Abeliovich H, Zhang X, Stirling AG, Burlingame AL & Goldfarb DS (2005) Atg19p ubiquitylation and the cytoplasm to vacuole trafficking pathway in yeast. *J Biol Chem* **280**, 39067–39076.
- 69 Nolan TM, Brennan B, Yang M, Chen J, Zhang M, Li Z, Wang X, Bassham DC, Walley J & Yin Y (2017) Selective autophagy of BES1 mediated by DSK2 balances plant growth and survival. *Dev Cell* **41**, 33–46.
- 70 Wild P, Farhan H, McEwan DG, Wagner S, Rogov VV, Brady NR, Richter B, Korac J, Waidmann O, Choudhary C *et al.* (2011) Phosphorylation of the autophagy receptor optineurin restricts *Salmonella* growth. *Science (80-)* **333**, 228–233.
- 71 Rogov VV, Suzuki H, Marinković M, Lang V, Kato R & Kawasaki M, Buljubašić M, Šprung M, Rogova N, Wakatsuki S *et al.* (2017) Phosphorylation of the mitochondrial autophagy receptor Nix enhances its interaction with LC3 proteins. *Sci Rep* **7**, 1131.
- 72 Kuang Y, Ma K, Zhou C, Ding P, Zhu Y, Chen Q & Xia B (2016) Structural basis for the phosphorylation of FUNDC1 LIR as a molecular switch of mitophagy. *Autophagy* **12**, 2363–2373.
- 73 Chen G, Han Z, Feng D, Chen Y, Chen L, Wu H, Huang L, Zhou C, Cai X, Fu C *et al.* (2014) A regulatory signaling loop comprising the PGAM5 phosphatase and CK2 controls receptor-mediated mitophagy. *Mol Cell* **54**, 362–377.
- 74 Kanki T, Kurihara Y, Jin X, Goda T, Ono Y, Aihara M, Hirota Y, Saigusa T, Aoki Y, Uchiumi T *et al.* (2013) Casein kinase 2 is essential for mitophagy. *EMBO Rep* **14**, 788–794.
- 75 Pfaffenwimmer T, Reiter W, Brach T, Nogellova V, Papinski D, Schuschnig M, Abert C, Ammerer G, Martens S & Kraft C (2014) Hrr25 kinase promotes selective autophagy by phosphorylating the cargo receptor A tg19. *EMBO Rep* **15**, 862–870.
- 76 Mochida K, Ohsumi Y & Nakatogawa H (2014) Hrr25 phosphorylates the autophagic receptor Atg34 to promote vacuolar transport of  $\alpha$ -mannosidase under nitrogen starvation conditions. *FEBS Lett* **588**, 3862–3869.
- 77 Tanaka C, Tan LJ, Mochida K, Kirisako H, Koizumi M, Asai E, Sakoh-Nakatogawa M, Ohsumi Y & Nakatogawa H (2014) Hrr25 triggers selective autophagy-related pathways by phosphorylating receptor proteins. *J Cell Biol* **207**, 91–105.

- 78 Nakatogawa H (2015) Hrr25: an emerging major player in selective autophagy regulation in *Saccharomyces cerevisiae*. *Autophagy* **11**, 432–433.
- 79 Heo JM, Ordureau A, Paulo JA, Rinehart J & Harper JW (2015) The PINK1-PARKIN mitochondrial ubiquitylation pathway drives a program of OPTN/NDP52 recruitment and TBK1 activation to promote mitophagy. *Mol Cell* **60**, 7–20.
- 80 Lim J, Lachenmayer ML, Wu S, Liu W, Kundu M, Wang R, Komatsu M, Oh YJ, Zhao Y & Yue Z (2015) Proteotoxic stress induces phosphorylation of p62/SQSTM1 by ULK1 to regulate selective autophagic clearance of protein aggregates. *PLoS Genet* **11**, e1004987.
- 81 Matsumoto G, Wada K, Okuno M, Kurosawa M & Nukina N (2011) Serine 403 phosphorylation of p62/SQSTM1 regulates selective autophagic clearance of ubiquitinated proteins. *Mol Cell* **44**, 279–289.
- 82 Pilli M, Arko-Mensah J, Ponpuak M, Roberts E, Master S, Mandell MA, Dupont N, Ornatowski W, Jiang S, Bradfute SB *et al.* (2012) TBK-1 promotes autophagy-mediated antimicrobial defense by controlling autophagosome maturation. *Immunity* **37**, 223–234.
- 83 Jiang X, Wang X, Ding X, Du M, Li B, Weng X, Zhang J, Li L, Tian R, Zhu Q *et al.* (2020) FAM134B oligomerization drives endoplasmic reticulum membrane scission for ER-phagy. *EMBO J* **39**, e102608.
- 84 Ducommun S, Deak M, Zeigerer A, Göransson O, Seitz S, Collodet C, Madsen AB, Jensen TE, Violet B, Foretz M *et al.* (2019) Chemical genetic screen identifies Gapex-5/GAPVD1 and STBD1 as novel AMPK substrates. *Cell Signal* **57**, 45–57.
- 85 Ha S, Jeong SH, Yi K, Chung KM, Hong CJ, Kim SW, Kim EK & Yu SW (2017) Phosphorylation of p62 by AMP-activated protein kinase mediates autophagic cell death in adult hippocampal neural stem cells. *J Biol Chem* **292**, 13795–13808.
- 86 You Z, Jiang WX, Qin LY, Gong Z, Wan W, Li J, Wang Y, Zhang H, Peng C, Zhou T *et al.* (2019) Requirement for p62 acetylation in the aggregation of ubiquitylated proteins under nutrient stress. *Nat Commun* **10**, 5792.
- 87 Huang R, Xu Y, Wan W, Shou X, Qian J, You Z, Liu B, Chang C, Zhou T, Lippincott-Schwartz J *et al.* (2015) Deacetylation of nuclear LC3 drives autophagy initiation under starvation. *Mol Cell* **57**, 456–466.
- 88 Peng Y, Shapiro SL, Banduseela VC, Dieterich IA, Hewitt KJ, Bresnick EH, Kong G, Zhang J, Schueler KL, Keller MP *et al.* (2018) Increased transport of acetyl-CoA into the endoplasmic reticulum causes a progeria-like phenotype. *Aging Cell* **17**, e12820.
- 89 Wang W, Wang M, Ruan Y, Tan J, Wang H, Yang T, Li J & Zhou Q (2019) Ginkgolic acids impair mitochondrial function by decreasing mitochondrial biogenesis and promoting FUNDC1-dependent mitophagy. *J Agric Food Chem* **67**, 10097–10106.
- 90 Nazio F, Strappazon F, Antonioli M, Bielli P, Cianfanelli V, Bordi M, Gretzmeier C, Dengjel J, Piacentini M, Fimia GM *et al.* (2013) MTOR inhibits autophagy by controlling ULK1 ubiquitylation, self-association and function through AMBRA1 and TRAF6. *Nat Cell Biol* **15**, 406–416.
- 91 Young LN, Cho K, Lawrence R, Zoncu R & Hurley JH (2016) Dynamics and architecture of the NRBF2-containing phosphatidylinositol 3-kinase complex I of autophagy. *Proc Natl Acad Sci USA* **113**, 8224–8229.
- 92 Parkhouse R, Ebong IO, Robinson CV & Monie TP (2013) The N-Terminal region of the human autophagy protein ATG16L1 contains a domain that folds into a helical structure consistent with formation of a coiled-coil. *PLoS One* **8**, e76237.
- 93 Ragusa MJ, Stanley RE & Hurley JH (2012) Architecture of the atg17 complex as a scaffold for autophagosome biogenesis. *Cell* **151**, 1501–1512.
- 94 Fujioka Y, Alam JM, Noshiro D, Mouri K, Ando T, Okada Y, May AI, Knorr RL, Suzuki K, Ohsumi Y *et al.* (2020) Phase separation organizes the site of autophagosome formation. *Nature* **578**, 301–305.
- 95 Wilfling F, Lee CW, Erdmann PS, Zheng Y, Sherpa D, Jentsch S, Pfander B, Schulman BA & Baumeister W (2020) A selective autophagy pathway for phase-separated endocytic protein deposits. *Mol Cell* **80**, 764–778.
- 96 Kirkin V, Lamark T, Sou YS, Bjørkøy G, Nunn JL, Bruun JA, Shvets E, McEwan DG, Clausen TH, Wild P *et al.* (2009) A role for NBR1 in autophagosomal degradation of ubiquitinated substrates. *Mol Cell* **33**, 505–516.
- 97 Wurzer B, Zaffagnini G, Fracchiolla D, Turco E, Abert C, Romanov J & Martens S (2015) Oligomerization of p62 allows for selection of ubiquitinated cargo and isolation membrane during selective autophagy. *Elife* **4**, e08941.
- 98 Bertipaglia C, Schneider S, Jakobi AJ, Tarafder AK, Bykov YS, Picco A, Kukulski W, Kosinski J, Hagen WJ, Ravichandran AC *et al.* (2016) Higher-order assemblies of oligomeric cargo receptor complexes form the membrane scaffold of the Cvt vesicle. *EMBO Rep* **17**, 1044–1060.
- 99 Long J, Garner TP, Pandya MJ, Craven CJ, Chen P, Shaw B, Williamson MP, Layfield R & Searle MS (2010) Dimerisation of the UBA domain of p62 inhibits ubiquitin binding and regulates NF- $\kappa$ B signalling. *J Mol Biol* **396**, 178–194.
- 100 Ying H, Shen X, Park B & Yue BYJT (2010) Posttranslational modifications, localization, and protein interactions of optineurin, the product of a glaucoma gene. *PLoS One* **5**, e9168.



- 101 Minegishi Y, Iejima D, Kobayashi H, Chi ZL, Kawase K, Yamamoto T, Seki T, Yuasa S, Fukuda K & Iwata T (2013) Enhanced optineurin E50k-TBK1 interaction evokes protein insolubility and initiates familial primary open-angle glaucoma. *Hum Mol Genet* **22**, 3559–3567.
- 102 Marinković M, Šprung M & Novak I (2020) Dimerization of mitophagy receptor BNIP3L/NIX is essential for recruitment of autophagic machinery. *Autophagy*, 1–12. <https://doi.org/10.1080/15548627.2020.1755120>
- 103 Grumati P, Morozzi G, Hölper S, Mari M, Harwardt MLIE, Yan R, Müller S, Reggiori F, Heilemann M & Dikic I (2017) Full length RTN3 regulates turnover of tubular endoplasmic reticulum via selective autophagy. *Elife* **5**, e25555.
- 104 Pendin D, Toso J, Moss TJ, Andrezza C, Moro S, McNew JA & Daga A (2011) GTP-dependent packing of a three-helix bundle is required for atlastin-mediated fusion. *Proc Natl Acad Sci USA* **108**, 16283–16288.
- 105 Moss TJ, Andrezza C, Verma A, Daga A & McNew JA (2011) Membrane fusion by the GTPase atlastin requires a conserved C-terminal cytoplasmic tail and dimerization through the middle domain. *Proc Natl Acad Sci USA* **108**, 11133–11138.
- 106 Bian X, Klemm RW, Liu TY, Zhang M, Sun S, Sui X, Liu X, Rapoport TA & Hu J (2011) Structures of the atlastin GTPase provide insight into homotypic fusion of endoplasmic reticulum membranes. *Proc Natl Acad Sci USA* **108**, 3976–3981.
- 107 Byrnes LJ, Singh A, Szeto K, Benveniste NM, O'Donnell JP, Zipfel WR & Sondermann H (2013) Structural basis for conformational switching and GTP loading of the large G protein atlastin. *EMBO J* **32**, 369–384.
- 108 Byrnes LJ & Sondermann H (2011) Structural basis for the nucleotide-dependent dimerization of the large G protein atlastin-1/SPG3A. *Proc Natl Acad Sci USA* **108**, 2216–2221.
- 109 Sainia SG, Liu C, Zhang P & Lee TH (2014) Membrane tethering by the atlastin GTPase depends on GTP hydrolysis but not on forming the cross-over configuration. *Mol Biol Cell* **25**, 3942–3953.
- 110 Liu TY, Bian X, Romano FB, Shemesh T, Rapoport TA & Hu J (2015) Cis and trans interactions between atlastin molecules during membrane fusion. *Proc Natl Acad Sci USA* **112**, E1851–E1860.
- 111 O'Donnell JP, Cooley RB, Kelly CM, Miller K, Andersen OS, Rusinova R & Sondermann H (2017) Timing and reset mechanism of GTP hydrolysis-driven conformational changes of atlastin. *Structure* **25**, 997–1010.
- 112 Svenning S, Lamark T, Krause K & Johansen T (2011) Plant NBR1 is a selective autophagy substrate and a functional hybrid of the mammalian autophagic adapters NBR1 and p62/SQSTM1. *Autophagy* **7**, 993–1010.
- 113 Monaco C, Visconti R, Barone MV, Pierantoni GM, Berlingieri MT, De Lorenzo C, Mineo A, Vecchio G, Fusco A & Santoro M (2001) The RFG oligomerization domain mediates kinase activation and re-localization of the RET/PTC3 oncoprotein to the plasma membrane. *Oncogene* **20**, 599–608.
- 114 Walinda E, Morimoto D, Sugase K, Konuma T, Tochio H & Shirakawa M (2014) Solution structure of the ubiquitin-associated (UBA) domain of human autophagy receptor NBR1 and its interaction with ubiquitin and polyubiquitin. *J Biol Chem* **289**, 13890–13902.
- 115 Acevo-Rodríguez PS, Maldonado G, Castro-Obregón S & Hernández G (2020) Autophagy regulation by the translation machinery and its implications in cancer. *Front Oncol* **10**, 322.
- 116 Liang JR, Lingeman E, Luong T, Ahmed S, Muhar M, Nguyen T, Olzmann JA & Corn JE (2020) A Genome-wide ER-phagy screen s key roles of mitochondrial metabolism and ER-resident UFMylation. *Cell* **180**, 1160–1177.
- 117 Korac J, Schaeffer V, Kovacevic I, Clement AM, Jungblut B, Behl C, Terzic J & Dikic I (2013) Ubiquitin-independent function of optineurin in autophagic clearance of protein aggregates. *J Cell Sci* **126**, 580–592.
- 118 Bjørkøy G, Lamark T, Brech A, Outzen H, Perander M, Øvervatn A, Stenmark H & Johansen T (2005) p62/SQSTM1 forms protein aggregates degraded by autophagy and has a protective effect on huntingtin-induced cell death. *J Cell Biol* **171**, 603–614.
- 119 Pankiv S, Clausen TH, Lamark T, Brech A, Bruun JA, Outzen H, Øvervatn A, Bjørkøy G & Johansen T (2007) p62/SQSTM1 binds directly to Atg8/LC3 to facilitate degradation of ubiquitinated protein aggregates by autophagy\*[S]. *J Biol Chem* **282**, 24131–24145.
- 120 Sarraf SA, Shah HV, Kanfer G, Pickrell AM, Holtzclaw LA, Ward ME & Youle RJ (2020) Loss of TAX1BP1-directed autophagy results in protein aggregate accumulation in the brain. *Mol Cell* **80**, 779–795.e10.
- 121 Zhou J, Wang Z, Wang X, Li X, Zhang Z, Fan B, Zhu C & Chen Z (2018) Dicot-specific ATG8-interacting ATI3 proteins interact with conserved UBAC2 proteins and play critical roles in plant stress responses. *Autophagy* **14**, 487–504.
- 122 Khaminets A, Heinrich T, Mari M, Grumati P, Huebner AK, Akutsu M, Liebmann L, Stolz A, Nietzsche S, Koch N *et al.* (2015) Regulation of endoplasmic reticulum turnover by selective autophagy. *Nature* **522**, 354–358.
- 123 Fumagalli F, Noack J, Bergmann TJ, Presmanes EC, Pisoni GB, Fasana E, Fregno I, Galli C, Loi M, Soldà

- T *et al.* (2016) Translocon component Sec62 acts in endoplasmic reticulum turnover during stress recovery. *Nat Cell Biol* **18**, 1173–1184.
- 124 An H, Ordureau A, Paulo JA, Shoemaker CJ, Denic V & Harper JW (2019) TEX264 Is an endoplasmic reticulum-resident ATG8-interacting protein critical for ER remodeling during nutrient stress. *Mol Cell* **74**, 891–908.
- 125 Chino H, Hatta T, Natsume T & Mizushima N (2019) Intrinsically disordered protein TEX264 mediates ER-phagy. *Mol Cell* **74**, 909–921.
- 126 Nthiga TM, Kumar Shrestha B, Sjøttem E, Bruun J, Bowitz Larsen K, Bhujabal Z, Lamark T & Johansen T (2020) CALCOCO 1 acts with VAMP -associated proteins to mediate ER -phagy. *EMBO J* **39**, e103649.
- 127 Dowdle WE, Nyfeler B, Nagel J, Elling RA, Liu S, Triantafellow E, Menon S, Wang Z, Honda A, Pardee G *et al.* (2014) Selective VPS34 inhibitor blocks autophagy and uncovers a role for NCOA4 in ferritin degradation and iron homeostasis in vivo. *Nat Cell Biol* **16**, 1069–1079.
- 128 Jiang S, Wells CD & Roach PJ (2011) Starch-binding domain-containing protein 1 (Stbd1) and glycogen metabolism: identification of the Atg8 family interacting motif (AIM) in Stbd1 required for interaction with GABARAPL1. *Biochem Biophys Res Commun* **413**, 420–425.
- 129 Wang L, Zhou J, Yan S, Lei G, Lee CH & Yin XM (2017) Ethanol-triggered lipophagy requires SQSTM1 in AML12 hepatic cells. *Sci Rep* **7**, 12307.
- 130 Yan Y, Wang H, Wei C, Xiang Y, Liang X, Phang CW & Jiao R (2019) HDAC6 regulates lipid droplet turnover in response to nutrient deprivation via p62-mediated selective autophagy. *J Genet Genomics* **46**, 221–229.
- 131 Papadopoulos C, Kirchner P, Bug M, Grum D, Koerver L, Schulze N, Poehler R, Dressler A, Fengler S, Arhzaouy K *et al.* (2017) VCP /p97 cooperates with YOD 1, UBXD 1 and PLAA to drive clearance of ruptured lysosomes by autophagy. *EMBO J* **36**, 135–150.
- 132 Chauhan S, Kumar S, Jain A, Ponpuak M, Mudd MH, Kimura T, Choi SW, Peters R, Mandell M, Bruun JA *et al.* (2016) TRIMs and galectins globally cooperate and TRIM16 and Galectin-3 co-direct autophagy in endomembrane damage homeostasis. *Dev Cell* **39**, 13–27.
- 133 Pohl C & Jentsch S (2009) Midbody ring disposal by autophagy is a post-abscission event of cytokinesis. *Nat Cell Biol* **11**, 65–70.
- 134 Isakson P, Lystad AH, Breen K, Koster G, Stenmark H & Simonsen A (2013) TRAF6 mediates ubiquitination of KIF23/MKLP1 and is required for midbody ring degradation by selective autophagy. *Autophagy* **9**, 1955–1964.
- 135 Mandell MA, Jain A, Kumar S, Castleman MJ, Anwar T, Eskelinen EL, Johansen T, Prekeris R & Deretic V (2016) TRIM17 contributes to autophagy of midbodies while actively sparing other targets from degradation. *J Cell Sci* **129**, 3562–3573.
- 136 van Humbeeck C, Cornelissen T, Hofkens H, Mandemakers W, Gevaert K, de Strooper B & Vandenberghe W (2011) Parkin interacts with ambral to induce mitophagy. *J Neurosci* **31**, 10249–10261.
- 137 Strappazon F, Nazio F, Corrado M, Cianfanelli V, Romagnoli A, Fimia GM, Campello S, Nardacci R, Piacentini M, Campanella M *et al.* (2015) AMBRA1 is able to induce mitophagy via LC3 binding, regardless of PARKIN and p62/SQSTM1. *Cell Death Differ* **22**, 419–432.
- 138 Novak I, Kirkin V, McEwan DG, Zhang J, Wild P, Rozenknop A, Rogov V, Löhr F, Popovic D, Ochipinti A *et al.* (2010) Nix is a selective autophagy receptor for mitochondrial clearance. *EMBO Rep* **11**, 45–51.
- 139 Zhang Y, Yao Y, Qiu X, Wang G, Hu Z, Chen S, Wu Z, Yuan N, Gao H, Wang J *et al.* (2019) Listeria hijacks host mitophagy through a novel mitophagy receptor to evade killing. *Nat Immunol* **20**, 433–446.
- 140 Wong YC & Holzbaur ELF (2014) Optineurin is an autophagy receptor for damaged mitochondria in parkin-mediated mitophagy that is disrupted by an ALS-linked mutation. *Proc Natl Acad Sci USA* **111**, E4439–E4448.
- 141 Wei Y, Chiang WC, Sumpter R, Mishra P & Levine B (2017) Prohibitin 2 is an inner mitochondrial membrane mitophagy receptor. *Cell* **168**, 224–238.
- 142 Murakawa T, Yamaguchi O, Hashimoto A, Hikoso S, Takeda T, Oka T, Yasui H, Ueda H, Akazawa Y, Nakayama H *et al.* (2015) Bcl-2-like protein 13 is a mammalian Atg32 homologue that mediates mitophagy and mitochondrial fragmentation. *Nat Commun* **6**, 419–432.
- 143 Hanna RA, Quinsay MN, Orogo AM, Giang K, Rikka S & Gustafsson ÅB (2012) Microtubule-associated protein 1 light chain 3 (LC3) interacts with Bnip3 protein to selectively remove endoplasmic reticulum and mitochondria via autophagy. *J Biol Chem* **287**, 19094–19104.
- 144 Sentelle RD, Senkal CE, Jiang W, Ponnusamy S, Gencer S, Panneer Selvam S, Ramshesh VK, Peterson YK, Lemasters JJ, Szulc ZM *et al.* (2012) Ceramide targets autophagosomes to mitochondria and induces lethal mitophagy. *Nat Chem Biol* **8**, 831–838.
- 145 Bhujabal Z, Birgisdottir ÅB, Sjøttem E, Brenne HB, Øvervatn A, Habisov S, Kirkin V, Lamark T & Johansen T (2017) FKBP8 recruits LC3A to mediate Parkin-independent mitophagy. *EMBO Rep* **18**, 947–961.

- 146 Liu L, Feng D, Chen G, Chen M, Zheng Q, Song P, Ma Q, Zhu C, Wang R, Qi W *et al.* (2012) Mitochondrial outer-membrane protein FUNDC1 mediates hypoxia-induced mitophagy in mammalian cells. *Nat Cell Biol* **14**, 177–185.
- 147 Lazarou M, Sliter DA, Kane LA, Sarraf SA, Wang C, Burman JL, Sideris DP, Fogel AI & Youle RJ (2015) The ubiquitin kinase PINK1 recruits autophagy receptors to induce mitophagy. *Nature* **524**, 309–314.
- 148 Lee CW, Wilfling F, Ronchi P, Allegretti M, Mosalaganti S, Jentsch S, Beck M & Pfander B (2020) Selective autophagy degrades nuclear pore complexes. *Nat Cell Biol* **22**, 159–166.
- 149 Tomioka Y, Kotani T, Kirisako H, Oikawa Y, Kimura Y, Hirano H, Ohsumi Y & Nakatogawa H (2020) TORC1 inactivation stimulates autophagy of nucleoporin and nuclear pore complexes. *J Cell Biol* **219**, e201910063.
- 150 Deosaran E, Larsen KB, Hua R, Sargent G, Wang Y, Kim S, Lamark T, Jauregui M, Law K, Lippincott-Schwartz J *et al.* (2013) NBR1 acts as an autophagy receptor for peroxisomes. *J Cell Sci* **126**, 939–952.
- 151 Zhang J, Tripathi DN, Jing J, Alexander A, Kim J, Powell RT, Dere R, Tait-Mulder J, Lee JH, Paull TT *et al.* (2015) ATM functions at the peroxisome to induce pexophagy in response to ROS. *Nat Cell Biol* **17**, 1259–1269.
- 152 Michaeli S, Honig A, Levanony H, Peled-Zehavi H & Galili G (2014) Arabidopsis ATG8-INTERACTING PROTEIN1 is involved in autophagy-dependent vesicular trafficking of plastid proteins to the vacuole. *Plant Cell* **26**, 4084–4101.
- 153 Marshall RS, Li F, Gemperline DC, Book AJ & Vierstra RD (2015) Autophagic degradation of the 26S proteasome is mediated by the dual ATG8/Ubiquitin receptor RPN10 in *Arabidopsis*. *Mol Cell* **58**, 1053–1066.
- 154 Wyant GA, Abu-Remaileh M, Frenkel EM, Laqtom NN, Dharamdasani V, Lewis CA, Chan SH, Heinze I, Ori A & Sabatini DM (2018) Nufip1 is a ribosome receptor for starvation-induced ribophagy. *Science* (80-) **360**, 751–758.
- 155 Thurston TLM (2009) The tbk1 adaptor and autophagy receptor ndp52 restricts the proliferation of ubiquitin-coated bacteria. *Nat Immunol* **10**, 1215–1221.
- 156 Zheng YT, Shahnazari S, Brech A, Lamark T, Johansen T & Brumell JH (2009) The adaptor protein p62/SQSTM1 targets invading bacteria to the autophagy pathway. *J Immunol* **183**, 5909–5916.
- 157 Tumbarello DA, Manna PT, Allen M, Bycroft M, Arden SD, Kendrick-Jones J & Buss F (2015) The autophagy receptor TAX1BP1 and the molecular motor myosin VI are required for clearance of *Salmonella typhimurium* by autophagy. *PLoS Pathog* **11**, e1005174.
- 158 Hafrén A, Macia JL, Love AJ, Milner JJ, Drucker M & Hofius D (2017) Selective autophagy limits cauliflower mosaic virus infection by NBR1-mediated targeting of viral capsid protein and particles. *Proc Natl Acad Sci USA* **114**, E2026–E2035.
- 159 Vaccaro MI (2012) Zymophagy: selective autophagy of secretory granules. *Int. J Cell Biol* **2012**, 1–7.
- 160 Christian F, Krause E, Houslay MD & Baillie GS (2014) PKA phosphorylation of p62/SQSTM1 regulates PB1 domain interaction partner binding. *Biochim Biophys Acta Mol Cell Res* **1843**, 2765–2774.
- 161 Zhu Y, Massen S, Terenzio M, Lang V, Chen-Lindner S, Eils R, Novak I, Dikic I, Hamacher-Brady A & Brady NR (2013) Modulation of serines 17 and 24 in the LC3-interacting region of Bnip3 determines pro-survival mitophagy versus apoptosis. *J Biol Chem* **288**, 1099–1113.
- 162 Gao J, Ohtsubo M, Hotta Y & Minoshima S (2014) Oligomerization of optineurin and its oxidative stress- or E50K mutation-driven covalent cross-linking: possible relationship with glaucoma pathology. *PLoS One* **9**, e101206.