

A journal of world insect systematics

INSECTA MUNDI

0975

Updated list of intercepted Coccidae
(Hemiptera: Coccomorpha) at South Korean ports of entry
and potential invasive species to South Korea

Soo-Jung Suh

Plant Pest Control Division/APQA
177, Hyeoksin 8-ro, Gimcheon-si, Gyeongsangbuk-do, South Korea 39660

Date of issue: February 3, 2023

Center for Systematic Entomology, Inc., Gainesville, FL

Suh S-J. 2023. Updated list of intercepted Coccidae (Hemiptera: Coccomorpha) at South Korean ports of entry and potential invasive species to South Korea. *Insecta Mundi* 0975: 1–7.

Published on February 3, 2023 by
Center for Systematic Entomology, Inc.
P.O. Box 141874
Gainesville, FL 32614-1874 USA
<http://centerforsystematicentomology.org/>

INSECTA MUNDI is a journal primarily devoted to insect systematics, but articles can be published on any non-marine arthropod. Topics considered for publication include systematics, taxonomy, nomenclature, checklists, faunal works, and natural history. *Insecta Mundi* will not consider works in the applied sciences (i.e. medical entomology, pest control research, etc.), and no longer publishes book reviews or editorials. *Insecta Mundi* publishes original research or discoveries in an inexpensive and timely manner, distributing them free via open access on the internet on the date of publication.

Insecta Mundi is referenced or abstracted by several sources, including the Zoological Record and CAB Abstracts. *Insecta Mundi* is published irregularly throughout the year, with completed manuscripts assigned an individual number. Manuscripts must be peer reviewed prior to submission, after which they are reviewed by the editorial board to ensure quality. One author of each submitted manuscript must be a current member of the Center for Systematic Entomology.

Guidelines and requirements for the preparation of manuscripts are available on the *Insecta Mundi* website at <http://centerforsystematicentomology.org/insectamundi/>

Chief Editor: David Plotkin, insectamundi@gmail.com
Assistant Editor: Paul E. Skelley, insectamundi@gmail.com
Layout Editor: Robert G. Forsyth
Editorial Board: Davide Dal Pos, Oliver Keller, M. J. Paulsen
Founding Editors: Ross H. Arnett, Jr., J. H. Frank, Virendra Gupta, John B. Heppner, Lionel A. Stange, Michael C. Thomas, Robert E. Woodruff
Review Editors: Listed on the *Insecta Mundi* webpage

Printed copies (ISSN 0749-6737) annually deposited in libraries

Florida Department of Agriculture and Consumer Services, Gainesville, FL, USA
The Natural History Museum, London, UK
National Museum of Natural History, Smithsonian Institution, Washington, DC, USA
Zoological Institute of Russian Academy of Sciences, Saint-Petersburg, Russia

Electronic copies (Online ISSN 1942-1354) in PDF format

Archived digitally by Portico
Florida Virtual Campus: <http://purl.fcla.edu/fcla/insectamundi>
University of Nebraska-Lincoln, Digital Commons: <http://digitalcommons.unl.edu/insectamundi/>
Goethe-Universität, Frankfurt am Main: <http://nbn-resolving.de/urn/resolver.pl?urn:nbn:de:hebis:30:3-135240>

Copyright held by the author(s). This is an open access article distributed under the terms of the Creative Commons, Attribution Non-Commercial License, which permits unrestricted non-commercial use, distribution, and reproduction in any medium, provided the original author(s) and source are credited. <http://creativecommons.org/licenses/by-nc/3.0/>

Updated list of intercepted Coccidae (Hemiptera: Coccomorpha) at South Korean ports of entry and potential invasive species to South Korea

Soo-Jung Suh

Plant Pest Control Division/APQA
177, Hyeoksin 8-ro, Gimcheon-si, Gyeongsangbuk-do, South Korea 39660
suhjs97@korea.kr

Abstract. An updated list is given of 25 species of soft scale insects (Hemiptera: Coccidae) which have been intercepted on plants imported into South Korea during the period of 1996 to 2021. Information on the number of interceptions, host plants, distribution and origin of species intercepted at South Korean ports of entry is provided. In addition, data on intercepted species was analyzed to determine potential invasive species of soft scales that could threaten South Korean plants.

Key words. Exotic species, plant trade, quarantine, soft scales.

ZooBank registration. urn:lsid:zoobank.org:pub:4EB0B411-5611-4F24-B004-922E76F024DD

Introduction

Soft scale insects (Hemiptera: Coccomorpha: Coccidae) occur in all zoogeographical regions of the world. The Coccidae is the third largest family of scale insects (Hemiptera: Coccomorpha) with 1225 species in 178 genera known worldwide (García Morales et al. 2022). Biologically, soft scales which are plant feeders, can occur on just about any part of the plant but are especially common on the stems, leaves and fruits (Miller et al. 2014). Many species are significant economic pests of fruits and also landscaping plants (Hamon and Williams 1984; Gill 1988; Miller et al. 2014; Kondo and Watson 2022). Invasive species of insects which are not native, represent an increasing concern to South Korea since globalization has been facilitating the increase in plant trade, promoting the long-distance movement of pests attached on these plants (Mazzeo et al. 2014). If non-native species are able to establish in a different environment, they often become economic pests, resulting in yield losses, diminished product quality, increased production costs, especially in the application of chemical control measures which often have a deleterious effect on the environment, humans and the ecology of the agroecosystem (Huber et al. 2002; Miller and Miller 2003). Many soft scales are serious pests, particularly those that are invasive species. In South Korea there are 31 species (2.5%) of the soft scales known worldwide. Of these, seven species (22.6%) are likely invaders, namely *Ceroplastes rubens* Maskell, *Coccus ficicola* Choi and Lee, *Coccus hesperidum* Linnaeus, *Eucalymnatus tessellatus* (Signoret), *Parasaissetia nigra* (Nietner), *Saissetia coffeae* (Walker) and *Saissetia miranda* (Cockerell and Parrott). In South Korea, soft scales are usually found on imported plants that are grown inside greenhouses (Paik 1972; Paik 2000; Choi and Lee 2017; 2018). Recently species such as *C. ficicola* and *S. miranda* were discovered on banyan figs (*Ficus benghalensis* L.; Moraceae) in South Korean greenhouses (Choi and Lee 2017; 2018). The brown soft scale, *C. hesperidum*, is a serious pest not only outdoors, but also in greenhouses and internal landscape environments (Paik 1972; Paik 2000; Kwon and Han 2003; Kwon et al. 2005; Lee and Choi 2019).

To prevent the introductions of harmful insects along these pathways, countries apply biosecurity measures to imported plant goods and products, including visual inspections at the points of entry (Saccaggi et al. 2021). Therefore, being able to compile and update information on intercepted soft scales will help to improve the inspection procedures, to detect, identify and mitigate the damage caused by exotic invasive species. This paper provides an updated list of soft scales intercepted on imported plants into South Korea during the period of 1996 to 2021 based on records in the Pest Information System (PIS) database of South Korea. In addition, it deals with potential invasive species of soft scales that could threaten South Korean natural and agricultural environments.

Materials and Methods

Data on soft scales intercepted at ports of entry to South Korea on imported plants between 1996 and 2021 were extracted from the Pest Information System (PIS), a database developed by the Animal Plant Quarantine Agency (APQA). In total, 403 specimens of soft scales were intercepted during phytosanitary inspections on plants imported into South Korea from 1996 to 2021. The checklist contains the identification of soft scales intercepted at the ports of entry to the level of species or genus depending upon the quality of the sample and the life stage as well as a summary of their distribution and hosts based on data provided by Danzig (1980), Kawai (1980), Hamon and Williams (1984), Gill (1988), Kosztarab and Kozár (1988), Williams and Watson (1990), Tang (1991) and Kosztarab (1996). Of the total of specimens that were detected, approximately 73.2% (295 specimens), 14.1% (57) and 12.7% (51) of them were identified to species, genus and family level, respectively. It is difficult to determine the zoogeographical area of origin for some species. In some cases, the criteria such as the first reported location of the species, its current distribution, the distribution of what appears to be its closest relatives and the natural distribution of its primary host plants were considered to determine its probable origin. Slide-mounted specimens examined are deposited in the Collection of Plant Quarantine Technology Center (PQTC), APQA.

Results

A total of 25 species of scale insects in 11 genera were intercepted on plants imported from 24 countries into South Korea during the period of 1996 to 2021 (Table 1). Of soft scales detected at ports of entry during this survey, 41.9% (13 species) were not known to occur in South Korea at the time they were detected and 40% (10 species) are categorized as quarantine pests in South Korea. A list of the six species most frequently intercepted at South Korea ports of entry are *Coccus hesperidum* Linnaeus (62.2%), *Coccus viridis* (Green) (6.3%), *Saissetia oleae* (Olivier) (2.6%), *Ceroplastes ceriferus* (Fabricius) (2.6%), *Parasaissetia nigra* (Nietner) (1.7%) and *Pulvinaria psidii* Maskell (1.4%). Each of these species were intercepted more than five times at South Korean ports of entry and three species *C. hesperidum*, *Cer. ceriferus* and *P. nigra* have been reported to occur in the environment of South Korea. In terms of plants, *Skimmia* (cut flowers) constitutes 32.7% (115 times) of the interceptions and it is the most common intercepted host plant followed by *Musa* (leaves, fruits) at 7.4% (26) and *Citrus* (leaves, seedlings, fruits) at 5.4% (19).

The interception records from the past 26 years also were searched to determine which intercepted species of soft scales pose the greatest threat. While the possibility that other soft scales could invade South Korea exists, the following five species are considered to be the most likely candidates for next invasions into the South Korea. The stellate scale, *Ceroplastes stellifer* (Westwood) was first described on *Paphiopedilum niveum* (Orchidaceae) from Thailand. This species was intercepted once on *Dypsis* sp. (Arecaceae) from Malaysia and on *Garcinia* sp. (Clusiaceae) from Thailand at the South Korean ports of entry. Its host range (22 plant families) and distribution (59 countries) are relatively wide and it is considered a potential pest on crops and several ornamentals in Florida (Hamon and Williams 1984; García Morales et al. 2022; Peronti and Kondo 2022). The green scale, *Coccus viridis* (Green), was first described on coffee from Sri Lanka (Green 1889; García Morales et al. 2022). This species was found 22 times on plants such as *Camellia* (Theaceae) from Malaysia, *Dypsis* from China, *Heptapleurum* (Araliaceae) from China, Indonesia, Malaysia, *Ixora* (Rubiaceae) from Thailand, *Jatropha* (Euphorbiaceae) from Indonesia, *Gardenia* (Rubiaceae) from Indonesia, *Polyscias* (Araliaceae) from China, Indonesia, Malaysia and *Zamioculcas* (Araceae) from Taiwan in quarantine inspections. It is a serious pest of coffee, citrus and other crops in many tropical areas (Hamon and Williams 1984; Kondo et al. 2022). Almost all of its known hosts are from subtropical and tropical habitats. Therefore, it is likely to cause concern in greenhouses if this pest is introduced to South Korea. Plants used for propagation should be carefully examined for pests since the soft scale can often survive for longer periods of time on leaves. *Pulvinaria psidii* Maskell was first reported on a plant belonging to the genus *Psidium* (Myrtaceae) from Hawaii. This species was intercepted five times on plants such as *Coffea* sp. (Rubiaceae) from Vietnam, *Heptapleurum* (Araliaceae) from Costa Rica, *Litchi* (Sapindaceae) from China and *Nephelium* (Sapindaceae) from China. It is reported to be a pest of mango, *Mangifera* sp. (Anacardiaceae), in Egypt (Nada et al. 1990). The iceplant scale, *Pulvinariella mesembryanthemi* (Vallot) is a Palearctic species

and was intercepted once on seedlings of *Mesembryanthemum* (Aizoaceae) from Australia. It is reported as a potential pest to Aizoaceae in California (García Morales et al. 2022). The black scale, *Saissetia oleae* (Olivier), was intercepted nine times on *Citrus* (Rutaceae) from Chile, New Zealand, *Durio* (Malvaceae) from Thailand, *Ficus* (Moraceae) from Japan, Malaysia and *Olea* (Oleaceae) from Italy, Netherlands at quarantine inspections. It is considered to be a major pest of citrus in many countries, and of olive in the Mediterranean region (Bodenhimer 1951; Bartlett 1978; Hamon and Williams 1984; Gill 1988; García Morales et al. 2022; Gavrillov-Zimin et al. 2022). Its hosts are primarily from subtropical and tropical habitats; however, citrus and olives, two of its known hosts can grow out of doors in the southern region of South Korea. Therefore, it is considered a potential threat to plants if introduced into Korea.

Discussion

The introduction of even a single species is of concern to South Korean environment, either in greenhouses or outdoor setting. If a pest can enter South Korea, over time there is a strong likelihood for establishment (Huber et al. 2002). Furthermore, scale insects reported as invasive pests in South Korea are sometimes only recognized after their populations have exploded to the point where they cause economic damage to plants. As a result, more appropriate and cost-effective quarantine procedures must be developed. One step in this procedure is to regularly update a list of pests intercepted on imported plants, identify potential invasive species which pose the greatest threat by analyzing information on intercepted pests and keep them under constant surveillance. Unfortunately, approximate 26.8% of the intercepted soft scales in South Korea are immatures which have not yet developed the diagnostic morphological characters required for confident identification or are older, poor quality adult female specimens which have these characteristics obscured and difficult to distinguish and evaluate (PIS 2022). Hence, the use of molecular tools like DNA barcodes may be useful in the identification of invasive scale insects and could be used to develop more effective pest management options for regulating pest species.

Acknowledgments

I am grateful to Drs. Gregory A. Evans (USDA/APHIS/NIS, Washington, DC, USA) and Takumasa Kondo (Corporación Colombiana de Investigación Agropecuaria – Agrosavia, Centro de Investigación Palmira, Palmira, Valle, Colombia) for reviewing the manuscript.

Table 1. List of soft scale insects intercepted at South Korean ports of entry between 1996 to 2021. Codes for the zoographical regions recognized are as follows: NA, Nearctic; NT, Neotropical; PA, Palearctic; OR, Oriental; AU, Australasian. [Total Int: number of intercepted records of soft scale species, In Kor: distributed in South Korea, Reg Orig: abbreviation of the zoographical region from where the species was first described, *: potential invasive species to South Korea.]

Species	Total Int	In Kor	Host Genus	Consignment origin	Reg Orig
<i>Ceroplastes ceriferus</i> (Fabricius)	9	yes	<i>Berzelia</i>	South Africa	OR
			<i>Camellia</i>	Japan	
			<i>Ficus</i>	China, Vietnam	
			<i>Magnolia</i>	Japan	
			<i>Yucca</i>	China	
			undetermined plant	Taiwan	
<i>Ceroplastes cirripediformis</i> Comstock	1	no	undetermined plant	Costa Rica	NA

Species	Total Int	In Kor	Host Genus	Consignment origin	Reg Orig
<i>Ceroplastes japonicus</i> Green	3	yes	<i>Chaenomeles</i>	Japan	PA
			<i>Euonymus</i>	Japan	
			undetermined plant	Italy	
<i>Ceroplastes rubens</i> Maskell	4	yes	<i>Mangifera</i>	Thailand	AU
			undetermined plant	Japan, Sri Lanka	
<i>Ceroplastes rusci</i> (Linnaeus)	2	no	<i>Pandanus</i>	Vietnam	PA
			undetermined plant	Israel	
* <i>Ceroplastes stellifer</i> (Westwood)	2	no	<i>Dyopsis</i>	Malaysia	OR
			<i>Garcinia</i>	Thailand	
<i>Ceroplastes</i> sp.	8	—	<i>Camellia</i>	China	—
			<i>Eucalyptus</i>	Indonesia	
			<i>Podocarpus</i>	Malaysia	
			undetermined plant	Israel, Italy, Malaysia, Sri Lanka	
<i>Coccus hesperidum</i> Linnaeus	219	yes	<i>Agave</i>	Poland	PA
			<i>Asplenium</i>	Japan	
			<i>Banksia</i>	South Africa	
			<i>Berzelia</i>	South Africa	
			<i>Citrus</i>	Chile, Thailand	
			<i>Cordyline</i>	Indonesia	
			<i>Cymbidium</i>	China, Japan	
			<i>Dracaena</i>	Indonesia	
			<i>Durio</i>	Thailand	
			<i>Dyopsis</i>	China	
			<i>Eryngium</i>	Thailand, Vietnam	
			<i>Eucalyptus</i>	Indonesia	
			<i>Ficus</i>	China	
			<i>Garcinia</i>	Singapore	
			<i>Gaultheria</i>	USA	
			<i>Heliconia</i>	Thailand	
			<i>Heptapleurum</i>	China, Malaysia, Sri Lanka, Taiwan	
			<i>Howea</i>	China	
			<i>Ixora</i>	Thailand	
			<i>Jasminum</i>	Thailand	
<i>Mangifera</i>	Indonesia				
<i>Musa</i>	Colombia, Ecuador, Philippines, Thailand, Vietnam				
<i>Philodendron</i>	Indonesia				
<i>Pittosporum</i>	Indonesia, New Zealand				

Species	Total Int	In Kor	Host Genus	Consignment origin	Reg Orig
			<i>Polyscias</i>	Indonesia	
			<i>Skimmia</i>	Netherlands	
			<i>Tradescantia</i>	Sri Lanka	
			<i>Vaccinium</i>	Chile	
			<i>Wrightia</i>	Thailand	
			<i>Zamioculcas</i>	Sri Lanka, Taiwan	
			undetermined plant	Australia, China, Japan, Netherlands, Singapore, Thailand, Vietnam	
<i>Coccus longulus</i> (Douglas)	1	no	undetermined plant	Vietnam	PA
* <i>Coccus viridis</i> (Green)	22	no	<i>Camellia</i>	Malaysia	OR
			<i>Cymbidium</i>	Japan	
			<i>Dracaena</i>	Philippines	
			<i>Dypsis</i>	China	
			<i>Heptapleurum</i>	China, Indonesia, Malaysia	
			<i>Ixora</i>	Thailand	
			<i>Jatropha</i>	Indonesia	
			<i>Gardenia</i>	Indonesia	
			<i>Litchi</i>	China	
			<i>Musa</i>	Vietnam	
			<i>Polyscias</i>	China, Indonesia, Malaysia	
			<i>Synsepalum</i>	Japan	
			<i>Zamioculcas</i>	Taiwan	
<i>Coccus</i> sp.	38	—	<i>Actinidia</i>	Chile	—
			<i>Aloe</i>	Italy	
			<i>Garcinia</i>	Thailand	
			<i>Gaultheria</i>	China, Netherlands, USA	
			<i>Helleborus</i>	Indonesia, Italy, Netherlands, Philippines	
			<i>Jasminum</i>	Vietnam	
			<i>Leucadendron</i>	South Africa	
			<i>Musa</i>	Philippines, Thailand	
			<i>Prunus</i>	Thailand, Uzbekistan	
			<i>Skimmia</i>	Netherlands	
			<i>Vaccinium</i>	USA	
			<i>Veronica</i>	Netherlands	
			undetermined plant	Madagascar, Malaysia	
<i>Discochiton cocophyllae</i> (Banks)	1	no	<i>Cocos</i>	Philippines	OR
<i>Discochiton expansum</i> (Green)	1	no	<i>Dypsis</i>	Malaysia	OR

Species	Total Int	In Kor	Host Genus	Consignment origin	Reg Orig
<i>Drepanococcus chiton</i> (Green)	1	no	<i>Dimocarpus</i>	Vietnam	OR
<i>Milviscutulus mangiferae</i> (Green)	3	no	<i>Cordyline</i> <i>Heptapleurum</i>	Indonesia Costa Rica	OR
<i>Milviscutulus</i> sp.	1	—	<i>Cordyline</i>	Sri Lanka	—
<i>Parasaissetia nigra</i> (Nietner)	6	yes	<i>Agave</i> <i>Citrus</i> <i>Durio</i> <i>Musa</i> <i>Philodendron</i> undetermined tree	Indonesia Thailand Thailand Philippines Japan Thailand	OR
<i>Parthenolecanium</i> sp.	1	—	<i>Pittosporum</i>	Italy	—
<i>Protopulvinaria pyriformis</i> (Cockerell)	2	no	undetermined tree	Indonesia, Italy	NT
* <i>Pulvinaria psidii</i> Maskell	5	no	<i>Coffea</i> <i>Heptapleurum</i> <i>Litchi</i> <i>Nephelium</i> undetermined plant	Vietnam Costa Rica China China Vietnam	AU
<i>Pulvinaria</i> sp.	2	—	<i>Dimocarpus</i> undetermined plant	Vietnam New Zealand	—
* <i>Pulvinariella mesembryanthemi</i> (Vallot)	1	no	<i>Mesembryanthemum</i>	Australia	PA
<i>Saissetia coffeae</i> (Walker)	3	yes	<i>Asplenium</i>	Japan	OR
* <i>Saissetia oleae</i> (Olivier)	9	no	<i>Citrus</i> <i>Durio</i> <i>Ficus</i> <i>Olea</i>	Chile, New Zealand Thailand Japan, Malaysia Italy, Netherlands	PA
<i>Saissetia</i> sp.	7	—	<i>Codiaeum</i> <i>Cymbidium</i> <i>Durio</i> <i>Eryngium</i> undetermined plant	Sri Lanka Japan China, Thailand Vietnam Indonesia	—

Literature Cited

- Bartlett BR. 1978.** Coccidae. p. 57–73. In: Clausen CP (ed.). Introduced parasites and predators of arthropod pests and weeds: a world review, agriculture handbook no. 480. Agricultural Research Service, United States Department of Agriculture; Washington, DC. 545 p.
- Bodenheimer FS. 1951.** Citrus entomology in the Middle East with special reference to Egypt, Iran, Iraq, Palestine, Syria, Turkey. W. Junk Publishers; The Hague, Netherlands. 663 p.

- Choi J, Lee S. 2017.** Taxonomic review of the tribe Saissetiini (Hemiptera: Coccidae) in Korea Journal of Asia-Pacific Entomology 20: 101–111.
- Choi J, Lee S. 2018.** Review of the genus *Coccus* Linnaeus from Korea, with description of a new species (Hemiptera, Cocomorpha, Coccidae). ZooKeys 734: 121–135.
- Danzig EM. 1980.** Coccoids of the Far East USSR (Homoptera, Coccinea): phylogenetic analysis of Coccids in the world fauna. Nauka Publishers; Leningrad, Russia. 367 p.
- García Morales M, Denno BD, Miller DR, Miller GL, Ben-Dov Y, Hardy NB. 2022.** ScaleNet: A literature-based model of scale insect biology and systematics. Available at <http://scalenet.info>. (Last accessed November 2022.)
- Gavrilov-Zimin IA, Kondo T, Watson GW. 2022.** Chapter 4.4.25: *Saissetia* spp. p. 331–337. In: Kondo T, Watson GW (eds.). Encyclopedia of scale insect pests. CABI; Wallingford, UK. 608 p.
- Gill RJ. 1988.** The scale insects of California: part 1. the soft scales (Homoptera: Coccoidea: Coccidae). California Department of Food and Agriculture; Sacramento, California. 132 p.
- Green EE. 1889.** Descriptions of two new species of *Lecanium* from Ceylon. Entomologist's Monthly Magazine 25: 248–250.
- Hamon AB, Williams ML. 1984.** The soft scale insects of Florida (Homoptera: Coccoidea: Coccidae). Arthropods of Florida and neighboring land areas Vol. 11. Florida Department of Agriculture and Consumer Services, Division of Plant Industry; Gainesville, Florida. 194 p.
- Huber DM, Hugh-Jones ME, Rust MK, Sheffield SR, Simberloff D, Taylor CR. 2002.** Invasive pest species: impacts on agricultural production, natural resources, and the environment. Issue paper no. 20. Council for Agricultural Science and Technology; Ames, Iowa. 18 p.
- Kawai S. 1980.** Scale insects of Japan in colors. National Agricultural Education Association; Tokyo, Japan. 455 p.
- Kondo T, Watson GW. 2022.** Chapter 5: Collection, preservation, slide-mounting, labelling and vouchering of scale insects. p. 548–558. In: Kondo T, Watson GW (eds.). Encyclopedia of scale insect pests. CABI; Wallingford, UK. 608 p.
- Kondo T, Watson GW, Gavrilov-Zimin IA. 2022.** Chapter 4.4.4: *Coccus* spp. p. 248–256. In: Kondo T, Watson GW (eds.). Encyclopedia of scale insect pests. CABI; Wallingford, UK. 608 p.
- Kosztarab MP, Kozár F. 1988.** Scale insects of central Europe. Akadémiai Kiadó; Budapest, Hungary. 456 p.
- Kosztarab MP. 1996.** Scale insects of Northeastern North America; identification, biology and distribution. Virginia Museum of Natural History; Martinsburg, Virginia. 650 p.
- Kwon GM, Han MJ. 2003.** Scale insects (Stenorrhyncha) occurred on fruit trees in Korea. Korean Journal of Applied Entomology 42(4): 279–288.
- Kwon GM, Han MJ, Choi DR. 2005.** Scale insects (Stenorrhyncha) occurring on flowering plants in Korea. Korean Journal of Applied Entomology 44(1): 51–59.
- Lee S, Choi J. 2019.** Insect fauna of Korea 9.6 Coccidae. National Institute of Biological Resources; Seoul, South Korea. 111 p.
- Mazzeo G, Longo S, Pellizzari G, Porcelli F, Suma P, Russo A. 2014.** Exotic scale insects (Coccoidea) on ornamental plants in Italy: a never-ending story. Acta Zoologica Bulgarica Suppl. 6: 55–61.
- Miller D, Rung A, Parikh G, Venable G, Redford AJ, Evans GA, Gill RJ. 2014.** Scale Insects, Edition 2. USDA APHIS Identification Technology Program (ITP). Fort Collins, CO. Available at <http://idtools.org/id/scales/>. (Last accessed November 2022.)
- Miller GL, Miller DR. 2003.** Invasive soft scales (Hemiptera: Coccidae) and their threat to U.S. agriculture. Proceedings of the Entomological Society of Washington 105(4): 832–846.
- Nada S, Abd Rabo S, Hussein GED. 1990.** Scale insects infesting mango trees in Egypt (Homoptera: Coccoidea). Proceedings of the Sixth International Symposium of Scale Insects Studies, Cracow, Poland: 133–134.
- Paik JC. 2000.** Economic insects of Korea 6, Homoptera (Coccinea). Insecta Koreana Suppl. 13. National Institute of Agriculture, Science and Technology; Seoul, South Korea. 193 p.
- Paik WH. 1972.** Scale insects found in the greenhouse in Korea. Korean Journal Plant Protection 11: 1–4.
- Peronti ALBG, Kondo T. 2022.** Chapter 4.4.3: *Ceroplastes* spp. p. 230–246. In: Kondo T, Watson GW (eds.). Encyclopedia of scale insect pests. CABI; Wallingford, UK. 608 p.
- PIS. 2022.** Pest Information System (internal database Plant Quarantine Technology Center/ APQA, South Korea). Available at <https://doi.org/10.110.128.00>. (Last accessed March 2022.)
- Saccaggi DL, Arendse M, Wilson JRU, Terblanche JS. 2021.** Contaminant organisms recorded on plant product imports to South Africa 1994–2019. Scientific Data 8(83): 1–11.
- Tang FT. 1991.** The Coccidae of China. Shanxi United Universities Press; Taiyuan, China. 377 p.
- Williams DJ, Watson GW. 1990.** The scale insects of the tropical South Pacific region. Pt. 3: The soft scales (Coccidae) and other families. CAB International; Wallingford, UK. 267 p.

Received November 10, 2022; accepted December 17, 2022.

Review editor Joe Eger.

