



Υ suppression at forward rapidity in Pb–Pb collisions at $\sqrt{s_{NN}} = 5.02$ TeV

ALICE Collaboration*



ARTICLE INFO

Article history:

Received 23 May 2018

Received in revised form 28 October 2018

Accepted 12 November 2018

Available online 29 December 2018

Editor: M. Doser

ABSTRACT

Inclusive $\Upsilon(1S)$ and $\Upsilon(2S)$ production have been measured in Pb–Pb collisions at the centre-of-mass energy per nucleon–nucleon pair $\sqrt{s_{NN}} = 5.02$ TeV, using the ALICE detector at the CERN LHC. The Υ mesons are reconstructed in the centre-of-mass rapidity interval $2.5 < y < 4$ and in the transverse-momentum range $p_T < 15$ GeV/c, via their decays to muon pairs. In this Letter, we present results on the inclusive $\Upsilon(1S)$ nuclear modification factor R_{AA} as a function of collision centrality, transverse momentum and rapidity. The $\Upsilon(1S)$ and $\Upsilon(2S)$ R_{AA} , integrated over the centrality range 0–90%, are $0.37 \pm 0.02(\text{stat}) \pm 0.03(\text{syst})$ and $0.10 \pm 0.04(\text{stat}) \pm 0.02(\text{syst})$, respectively, leading to a ratio $R_{AA}^{\Upsilon(2S)} / R_{AA}^{\Upsilon(1S)}$ of $0.28 \pm 0.12(\text{stat}) \pm 0.06(\text{syst})$. The observed $\Upsilon(1S)$ suppression increases with the centrality of the collision and no significant variation is observed as a function of transverse momentum and rapidity.

© 2018 Published by Elsevier B.V. This is an open access article under the CC BY license (<http://creativecommons.org/licenses/by/4.0/>). Funded by SCOAP³.

1. Introduction

A detailed study of the properties of the Quark–Gluon Plasma (QGP) [1] is the main goal of heavy-ion experiments at ultra-relativistic energies [2–6]. Quarkonia, *i.e.* bound states of charm or bottom quark–antiquark pairs, are sensitive probes of color deconfinement, due to the Quantum-Chromo Dynamics Debye screening mechanism [7–9] leading to quarkonium suppression. Moreover, the various quarkonium states have different binding energies and therefore different dissociation temperatures in a QGP, leading to sequential suppression [7,10]. Theory estimates [11] indicate that bottomonium formation may occur before QGP thermalization [12] because of the large bottom quark mass. In this situation, a quantitative description of the influence of the medium on the bound states becomes challenging. While the dissociation temperatures vary significantly between different models [8,9], it is commonly accepted that the widths of the spectral functions of the bottomonium states increase compared to the widths in vacuum, due to the high temperature of the surrounding medium [13]. Finally, taking into account that feed-down processes from higher-mass resonances (around 40% for the $\Upsilon(1S)$ and 30% for the $\Upsilon(2S)$ [9]) are not negligible, the evaluation of the medium temperature via bottomonium measurements remains a complex endeavour.

The first studies of quarkonium production in heavy-ion collisions were devoted to charmonium states, and a suppression of their yields was observed at the SPS [14–16], at RHIC [17,18] and

at the LHC [19–22]. The weaker J/ψ suppression observed at LHC energies, where the centre-of-mass energy per nucleon–nucleon pair ($\sqrt{s_{NN}}$) is one order of magnitude larger than at RHIC, can be explained by means of a competitive (re)generation mechanism, which occurs during the deconfined phase and/or at the hadronization stage [23–26]. This production mechanism strongly depends on the (re)combination probability of deconfined quarks present in the medium and thus on the initial number of produced $c\bar{c}$ pairs. The effect has been found to be more important at low p_T and in the most central collisions [22,20,27].

The high-energy collisions delivered by the LHC allow for a detailed study of bottomonium states. For bottomonium production, perturbative calculations of production rates in elementary nucleon–nucleon collisions are more reliable than for charmonium yields due to the higher mass of the bottom quark with respect to charm. Since the number of produced $b\bar{b}$ pairs in central heavy-ion collisions amount to a few pairs per event at the LHC, the probability for (re)generation of bottomonia through (re)combination is much smaller than in the case of charmonia.

The $\Upsilon(1S)$ nuclear modification factor R_{AA} is quantified as the ratio of the $\Upsilon(1S)$ yield in nucleus–nucleus collisions to the production cross section measured in pp collisions scaled by the nuclear overlap function (T_{AA}). The latter is obtained via the Glauber model [28,29]. A strong suppression of the $\Upsilon(1S)$ state in Pb–Pb collisions has been observed at $\sqrt{s_{NN}} = 2.76$ TeV by ALICE [30] and CMS [31,32] in the rapidity ranges $2.5 < y < 4$ and $|y| < 2.4$, respectively. The suppression increases with the centrality of the collision, reaching about 60% and 80% for the most central collisions at mid [32] and forward rapidity [30], respectively. Moreover,

* E-mail address: alice-publications@cern.ch.

the $\Upsilon(2S)$ suppression reaches about 90% and for $\Upsilon(3S)$ data are compatible with a complete suppression [32]. As a function of p_T the $\Upsilon(1S) R_{AA}$, measured for $p_T < 20 \text{ GeV}/c$ by CMS [32], is compatible with a constant value. When considering the y -dependence resulting from the comparison of ALICE and CMS results, there is an indication for a stronger suppression at forward y . Transport models [26,33] as well as an anisotropic hydro-dynamical model [34] fairly reproduce the experimental observations of CMS, while they tend to overestimate the R_{AA} values measured by ALICE.

The bottomonium suppression due to the QGP should be disentangled from the suppression due to Cold Nuclear Matter (CNM) effects, such as the nuclear modification of the parton distribution functions due to shadowing [35,36], as well as parton energy loss [37]. These effects on the bottomonium production were studied in p-Pb collisions by ALICE [38] and LHCb [39], who reported for the $\Upsilon(1S)$ a nuclear modification factor slightly lower than unity at forward rapidity and compatible with unity at backward rapidity, although with significant uncertainties. Recently, ATLAS results indicate a significant suppression of the $\Upsilon(1S)$ for $p_T < 40 \text{ GeV}/c$ around mid-rapidity [40]. Additional measurements at forward/backward rapidity with higher statistics, are needed to fully constrain the models and perform a meaningful extrapolation of CNM effects to Pb-Pb collisions.

In this Letter we present the first results on the $\Upsilon(1S)$ and $\Upsilon(2S) R_{AA}$ measured by the ALICE Collaboration in Pb-Pb collisions at $\sqrt{s_{NN}} = 5.02 \text{ TeV}$. The pp reference cross sections used in the R_{AA} calculations have been determined by an interpolation procedure based on various ALICE [41,42] and LHCb [43,44] results at different energies. The nuclear modification factor for the $\Upsilon(1S)$ is presented as a function of the centrality of the collision and also differentially in p_T and rapidity. For the $\Upsilon(2S)$, an R_{AA} value integrated over the centrality of the collision is quoted. Finally, the results are compared to theoretical calculations.

2. Experimental apparatus and data sample

An extensive description of the ALICE apparatus can be found in [45,46]. The analysis presented in this Letter is based on muons detected at forward rapidity ($2.5 < y < 4$)¹ with the muon spectrometer [47]. The detectors relevant for Υ measurements in Pb-Pb collisions are described below.

The Silicon Pixel Detector, corresponding to the two innermost layers of the Inner Tracking System [48], is used for the primary vertex determination. The inner and outer layer cover the pseudo-rapidity ranges $|\eta| < 2$ and $|\eta| < 1.4$, respectively.

The V0 scintillator hodoscopes [49] provide the centrality estimate. They are made of two arrays of scintillators placed in the pseudo-rapidity ranges $2.8 < \eta < 5.1$ and $-3.7 < \eta < -1.7$. The logical AND of the signals from the two hodoscopes constitutes the Minimum Bias (MB) trigger. The MB trigger is fully efficient for the studied 0–90% most central collisions.

The Zero Degree Calorimeters (ZDC) are installed at $\pm 112.5 \text{ m}$ from the nominal interaction point along the beam line. Each of the two ZDCs is composed of two sampling calorimeters designed for detecting spectator protons, neutrons and nuclear fragments. The evaluation of the signal amplitude of the ZDCs allows for the rejection of events corresponding to an electromagnetic interaction of the colliding Pb nuclei [50].

The muon spectrometer covers the pseudorapidity range $-4 < \eta < -2.5$. It is composed of a front absorber, which filters muons

upstream of the muon tracker, consisting of five tracking stations with two planes of cathode-pad chambers each, and of a dipole magnet providing a $3 \text{ T}\cdot\text{m}$ integrated magnetic field. Downstream of the tracking system, a 1.2 m thick iron wall stops efficiently the punch-through hadrons. The muon trigger system is located downstream of the iron wall and consists of two stations, each one equipped with two planes of Resistive Plate Chambers (RPC), with an efficiency higher than 95% [51]. The muon-trigger system is able to deliver single and dimuon triggers selecting muons with p_T larger than a programmable threshold, via an algorithm based on the RPC spatial information [52]. Throughout its entire length, a conical absorber shields the muon spectrometer against secondary particles produced by the interaction of primary particles in the beam pipe.

The trigger condition used for data taking is a dimuon-Minimum Bias ($\mu\mu\text{-MB}$) trigger formed by the logical AND of the MB trigger and an unlike-sign dimuon trigger with a p_T threshold of $1 \text{ GeV}/c$ for each of the two muons.

The centrality estimation is performed using a Glauber fit to the sum of the signal amplitudes of the V0 scintillators [53–55]. Centrality ranges are given as percentages of the total hadronic Pb-Pb cross section. In addition to the centrality, the Glauber model allows an estimate of the average number of participant nucleons (N_{part}), of the average number of binary collisions (N_{coll}) and of the nuclear overlap function (T_{AA}), for each centrality interval [56]. In the present analysis, the data sample corresponds to an integrated luminosity $L_{\text{int}} \approx 225 \text{ }\mu\text{b}^{-1}$ in the centrality interval 0–90% that has been divided into four centrality classes: 0–10%, 10–30%, 30–50% and 50–90%.

3. Data analysis

The evaluation of R_{AA} is performed through the following expression:

$$R_{AA} = \frac{N^\Upsilon}{\text{BR}_{\Upsilon \rightarrow \mu^+ \mu^-} \cdot (A \times \varepsilon)_{\Upsilon \rightarrow \mu^+ \mu^-} \cdot N_{\mu\mu\text{-MB}} \cdot F_{\text{norm}} \cdot \sigma_{\text{pp}}^\Upsilon \cdot \langle T_{AA} \rangle}, \quad (1)$$

where N^Υ is the number of detected resonance decays to muon pairs, while $\text{BR}_{\Upsilon \rightarrow \mu^+ \mu^-} = (2.48 \pm 0.05)\%$ and $(1.93 \pm 0.17)\%$ are the branching ratios for the dimuon decay of $\Upsilon(1S)$ and $\Upsilon(2S)$, respectively [57]. The $(A \times \varepsilon)_{\Upsilon \rightarrow \mu^+ \mu^-}$ factor is the product of acceptance and detection efficiency for the Υ state under study. The normalization factor $N_{\mu\mu\text{-MB}} \cdot F_{\text{norm}}$ is the product of the number of analyzed $\mu\mu\text{-MB}$ events and the inverse of the probability to obtain an unlike-sign dimuon trigger in a MB-triggered event [22]. A dataset of $1.5 \cdot 10^9$ minimum bias equivalent events, $N_{\mu\mu\text{-MB}} \cdot F_{\text{norm}}$, has been used for bottomonium measurements. Finally, $\sigma_{\text{pp}}^\Upsilon$ is the reference pp cross section and $\langle T_{AA} \rangle$ represents the nuclear overlap function [55].

The signal yields are evaluated by performing fits to the $\mu^+ \mu^-$ invariant mass distributions. In order to improve the purity of the dimuon sample a set of selection criteria [30] has been applied on the muon tracks, including the request of the matching between the tracks reconstructed in the trigger and tracking detectors of the muon spectrometer and a cut on the track transverse momentum ($p_T > 2 \text{ GeV}/c$). The latter cut has a small effect ($\sim 2\%$) on the number of detected resonances. The raw Υ yields are extracted using the sum of three extended Crystal Ball (CB) functions [58], one for each of $\Upsilon(1S)$, $\Upsilon(2S)$ and $\Upsilon(3S)$. The extended CB function consists of a Gaussian core with non-Gaussian tails on both sides to take into account the radiative contributions of the Υ production and the absorber effects of muon energy loss in the low mass tail, whereas the high mass tail is attributed to the multiple

¹ In the ALICE reference frame, the muon spectrometer covers a negative η range and consequently a negative y range. We have chosen to present our results with a positive y notation.




Fig. 1. Red and magenta solid lines correspond to $\Upsilon(1S)$ and $\Upsilon(2S)$ signal functions, respectively. The contribution from $\Upsilon(3S)$ yield is compatible with zero. Dotted blue lines represent the background (left) and residual background (right), respectively. The sum of the various functions is also shown as a solid blue line.

Coulomb scattering in the front absorber and the momentum resolution of the tracking chambers. The background is fitted with the sum of two exponential functions (see left panel of Fig. 1). Since the signal-to-background (S/B) ratio is low in the tail regions of the extended CB functions, the tail parameters are fixed to values obtained from the Monte Carlo (MC) simulation. The mass position and the width parameters of the $\Upsilon(1S)$ are left free for the integrated spectrum (*i.e.* centrality class 0–90%, $p_T < 15 \text{ GeV}/c$ and $2.5 < y < 4$). Whereas for the signal extraction as a function of centrality, the mass position and width ($160 \pm 15 \text{ MeV}/c^2$) of the $\Upsilon(1S)$ are fixed to the values obtained in the fit to the centrality-integrated (0–90%) mass spectrum. Finally, for studies as a function of p_T and y , the mass position and the width obtained for the centrality-integrated mass spectrum are scaled according to their evolution observed in the MC. Due to the poor S/B ratio for the higher mass states, the values of the mass of the $\Upsilon(2S)$ and $\Upsilon(3S)$ are fixed to the PDG mass differences with respect to the $\Upsilon(1S)$, and the ratio of $\Upsilon(2S)$ ($\Upsilon(3S)$) to $\Upsilon(1S)$ widths is fixed to values from the MC simulation, *i.e.* 1.03 (1.06). In the fit shown in Fig. 1 only signals corresponding to the $\Upsilon(1S)$ and $\Upsilon(2S)$ are visible, since the $\Upsilon(3S)$ contribution is compatible with zero events. Alternatively, the combinatorial background is modeled with the event-mixing method. In this approach, an invariant mass dimuon spectrum is constructed by pairing muons from different events with similar multiplicities as described in [22]. The combinatorial background is then subtracted from the raw dimuon spectrum (right panel of Fig. 1) and the resulting distribution is fitted with the sum of three extended CB and an exponential function to account for the residual background. Finally, the number of detected Υ resonances, N^Υ , is obtained as the average [58] of the fitting methods described above (and also below in the discussion on signal systematics), leading to $N^{\Upsilon(1S)} = 1126 \pm 53(\text{stat}) \pm 47(\text{syst})$ and $N^{\Upsilon(2S)} = 77 \pm 33(\text{stat}) \pm 17(\text{syst})$.

The measured Υ yields, N^Υ , are corrected for the detector acceptance and efficiency using MC simulations. Since the occupancy of the detector varies with the centrality of the collisions, the generated Υ decays are embedded into real MB events to simulate the various particle multiplicity scenarios as in data. The p_T and y distributions of the generated Υ are obtained from existing pp measurements [59–61] using the interpolation procedure described in [62]. The EKS98 nuclear shadowing parameterization [35] is used to include an estimate of CNM effects. Since available data favor a small or null polarization for $\Upsilon(1S)$ [63–66], an unpolarized production is assumed. The variations of the performance of

the tracking and triggering systems throughout the data-taking period as well as the residual misalignment of the tracking chambers are taken into account in the simulation. The $A \times \varepsilon$ values, for the range $p_T < 15 \text{ GeV}/c$, $2.5 < y < 4$ and the 0–90% centrality class are 0.263 and 0.264 for the $\Upsilon(1S)$ and $\Upsilon(2S)$, respectively, with a negligible statistical uncertainty. A decrease of 2% is observed in $A \times \varepsilon$ for the 0–10% central collisions with respect to the 50–90% sample due to the higher occupancy in the most central events. The $A \times \varepsilon$ is higher by 20% in $3 < y < 3.5$ compared to the values at $2.5 < y < 3$ and $3.5 < y < 4$ mainly due to the geometric acceptance of the detector, whereas it has no variation as a function of p_T . The systematic uncertainty on $A \times \varepsilon$ is discussed below.

The systematic uncertainty on the signal extraction is evaluated using various functions for modelling the background shape, as well as adopting two fitting ranges, *i.e.* $(7\text{--}14) \text{ GeV}/c^2$ and $(7.5\text{--}14.5) \text{ GeV}/c^2$. The tail parameters of the signal functions have been varied using estimates provided by two MC particle transport models: GEANT4 [67] and GEANT3 [68]. In the centrality, p_T or y differential studies, the mass position and width are also varied by amounts, which correspond to the uncertainties on the mass position and the width returned by the fit to the centrality-integrated invariant mass spectrum. The ratio of $\Upsilon(2S)$ ($\Upsilon(3S)$) to $\Upsilon(1S)$ widths is varied from 1 (1) to 1.06 (1.12). The values of N^Υ and their statistical uncertainties are obtained by taking the average of N^Υ and of the corresponding statistical uncertainties from the various fits. This procedure is applied to both fits of the raw and combinatorial-background-subtracted spectra. The systematic uncertainties are estimated as the root mean square of the distribution of N^Υ obtained from the various fits. The effect induced by the $p_T > 2 \text{ GeV}/c$ cut on single muons on the $A \times \varepsilon$ -corrected Υ yields was estimated by varying that cut by $\pm 0.2 \text{ GeV}/c$ in the MC. A $\pm 2\%$ maximum variation on $N^\Upsilon/(A \times \varepsilon)$ was observed and included in the systematic uncertainties.

Various sources contribute to the systematic uncertainties of $A \times \varepsilon$, such as the p_T and y shapes of the input distributions for the MC simulations, the trigger efficiency, the track reconstruction efficiency and finally the matching efficiency between tracks in the muon tracking and triggering chambers. Various sets of simulations are produced with different Υ input p_T and y distributions, obtained from empirical parameterizations and/or extrapolations of available data sets at different energies. The maximum relative difference of $A \times \varepsilon$ for the various shapes is taken as the systematic uncertainty due to the input MC. In order to calculate the systematic uncertainty on trigger efficiency, the trigger response function

Table 1

Summary of the systematic uncertainties for the R_{AA} calculation. Type I (II) refers to correlated (uncorrelated) systematic uncertainties.

Sources	$\Upsilon(1S)$				$\Upsilon(2S)$
	Centrality	y	p_T	Integrated	Integrated
Signal extraction	4.3–6.1%(II)	4.2–6.8%(II)	5.2–8.7%(II)	4.1%	21.7%
Muon p_T cut	0.3–2.4%(II)	0.1–1.2%(II)	0.1–2.4%(II)	0.7%	0.7%
Input MC	0.9%(I)	0.6–2.6%(II)	1–1.4%(II)	0.9%	0.9%
Tracker efficiency	3%(I) and 0–1%(II)	1%(I) and 3%(II)	1%(I) and 3%(II)	3%	3%
Trigger efficiency	3%(I)	1.4–3.7%(II)	0.4–2.6%(II)	3%	3%
Matching efficiency	1%(I)	1%(II)	1%(II)	1%	1%
Centrality	0.2–2.4%(II)	–	–	–	–
F_{norm}	0.5%(I)	0.5%(I)	0.5%(I)	0.5%	0.5%
$\langle T_{AA} \rangle$	3.1–5.3%(II)	3.2%(I)	3.2%(I)	3.2%	3.2%
$\text{BR}_{\Upsilon \rightarrow \mu^+ \mu^-} \cdot \sigma_{\text{pp}}^{\Upsilon}$	6.3%(I)	6.6–11.3%(II)	5.5–11.5%(II)	6.3%	7.5%

for single muons is evaluated using either MC or data. The two response functions are then separately applied to simulations of an Υ sample and the difference obtained for the Υ reconstruction efficiency is taken as systematic uncertainty. The systematic uncertainty on the tracking efficiency is obtained starting from an evaluation of the single muon tracking efficiency in MC and data. This evaluation is performed via a procedure, detailed in [22], based on the redundancy of the tracking chamber information. The dimuon tracking efficiency is then obtained by combining the single muon efficiencies and the systematic uncertainty is taken as the difference of the values obtained with the procedure based on MC and data. The muon tracks for data analysis are chosen based on a selection on the χ^2 of the matching between a track segment in the trigger system with a track in the tracking chambers. The matching systematics are obtained by varying the χ^2 selection cut in data and MC and comparing the effects on the muon reconstruction efficiency [22].

The systematic uncertainty on the centrality measurement is evaluated by varying the V0 signal amplitude by $\pm 0.5\%$ corresponding to 90% of the hadronic cross section in Pb–Pb collisions, used as anchor point to define the centrality classes. The systematic uncertainty on the evaluation of $\sigma_{\text{pp}}^{\Upsilon}$ is detailed in the next section. Finally, the systematic uncertainty evaluation of F_{norm} and $\langle T_{AA} \rangle$ are described in [22] and [53], respectively. The different systematic uncertainty sources on the R_{AA} calculation are summarized in Table 1. If the above mentioned systematic uncertainty is correlated as a function of centrality, p_T or y , it is quoted as correlated (type I) systematic uncertainty, otherwise it is treated as uncorrelated (type II).

4. Proton–proton reference cross sections

The pp reference cross section for $\Upsilon(1S)$ and $\Upsilon(2S)$ production are computed by means of an interpolation procedure as described for $\Upsilon(1S)$ in [69]. The energy interpolation for the Υ cross section, as a function of rapidity and for the p_T and y integrated result, uses the measurements of Υ production cross sections in pp collisions at $\sqrt{s} = 7$ and 8 TeV by ALICE [41,42] and at $\sqrt{s} = 2.76, 7$ and 8 TeV by LHCb [43,44]. The interpolation is performed by using various empirical functions and, in addition, the shape of the energy dependence of the bottomonium cross sections calculated using two theoretical models, i.e. the Leading Order Colour Evaporation Model (LO-CEM) [70] and the Fixed Order Next-to-Leading Logs (FONLL) model [71]. The latter gives cross sections for open beauty, which is here used as a proxy to study the evolution of the bottomonium cross section [69]. The energy interpolation for the $\Upsilon(1S)$ cross section as a function of p_T is based on LHCb measurements only, since the p_T coverage of the results of this analysis ($p_T < 15$ GeV/c) is more extended than that of the corresponding ALICE pp data ($p_T < 12$ GeV/c). The result of the interpolation procedure gives $\text{BR}_{\Upsilon(1S) \rightarrow \mu^+ \mu^-} \cdot \sigma_{\text{pp}}^{\Upsilon(1S)} = 1221 \pm 77(\text{syst})$ pb

Table 2

The interpolated branching ratio times cross section of $\Upsilon(1S)$ for the p_T and y bins under study. The quoted uncertainties are systematic.

p_T (GeV/c)	y	$\text{BR}_{\Upsilon(1S) \rightarrow \mu^+ \mu^-} \cdot \sigma_{\text{pp}}^{\Upsilon(1S)}$ (pb)
[0–2]		226 ± 26
[2–4]		361 ± 20
[4–6]	[2.5–4]	288 ± 24
[6–15]		311 ± 23
	[2.5–3]	506 ± 57
[0–15]	[3–3.5]	415 ± 28
	[3.5–4]	288 ± 24

and $\text{BR}_{\Upsilon(2S) \rightarrow \mu^+ \mu^-} \cdot \sigma_{\text{pp}}^{\Upsilon(2S)} = 302 \pm 23(\text{syst})$ pb assuming unpolarized quarkonia and integrating over the ranges $2.5 < y < 4$ and $p_T < 15$ GeV/c. The uncertainties correspond to the quadratic sum of two terms. The first term dominates the total uncertainty on the interpolated value and reflects the statistical and systematic uncertainties on the data points used in the interpolation procedure. The second term is related to the spread among the interpolated cross sections obtained by using either the empirical functions or the energy dependence estimated from the theoretical models mentioned above. The numerical values obtained from the interpolation procedure are summarized in Table 2 for the various kinematic ranges used in the analysis.

5. Results

The nuclear modification factors for inclusive $\Upsilon(1S)$ and $\Upsilon(2S)$ production in Pb–Pb collisions at $\sqrt{s_{\text{NN}}} = 5.02$ TeV for the ranges $p_T < 15$ GeV/c, $2.5 < y < 4$ and the 0–90% centrality class are $R_{AA}^{\Upsilon(1S)} = 0.37 \pm 0.02(\text{stat}) \pm 0.03(\text{syst})$ and $R_{AA}^{\Upsilon(2S)} = 0.10 \pm 0.04(\text{stat}) \pm 0.02(\text{syst})$, respectively. The ratio $R_{AA}^{\Upsilon(2S)} / R_{AA}^{\Upsilon(1S)}$ is $0.28 \pm 0.12(\text{stat}) \pm 0.06(\text{syst})$. Since the decay kinematics of the two Υ states is very similar, most of the systematic uncertainty sources entering the ratio cancel out except those on the signal extraction and on the pp cross section, which are the dominant contributions to the total systematic uncertainty. The measurements show a strong suppression for both bottomonium states with the more weakly bound state being significantly more suppressed. The ratio between the $\Upsilon(1S)$ R_{AA} at $\sqrt{s_{\text{NN}}} = 5.02$ TeV and 2.76 TeV [30] is $1.23 \pm 0.21(\text{stat}) \pm 0.19(\text{syst})$. The sources of systematic uncertainties entering the calculation of the ratio are considered uncorrelated, except for the $\langle T_{AA} \rangle$ component, whose uncertainty cancels out. The ratio is compatible with unity within uncertainties.

The centrality, p_T and y dependences of the $\Upsilon(1S)$ R_{AA} at forward rapidity at $\sqrt{s_{\text{NN}}} = 5.02$ TeV are shown in Fig. 2. A decrease of R_{AA} with increasing centrality is observed down to $R_{AA}^{\Upsilon(1S)} = 0.34 \pm 0.03(\text{stat}) \pm 0.02(\text{syst})$ for the 0–10% most central collisions. No significant p_T -dependence is observed up to $p_T = 15$ GeV/c




Fig. 2. Inclusive $\Upsilon(1S)$ R_{AA} as a function of centrality (top), p_T (left) and y (right) at forward rapidity at $\sqrt{s_{NN}} = 5.02$ TeV. ALICE results at $\sqrt{s_{NN}} = 2.76$ TeV as a function of centrality and y are shown for comparison [30]. The vertical error bars and the boxes represent the statistical and uncorrelated systematic uncertainties, respectively. The relative correlated uncertainty is shown as boxes at unity. ALICE $\Upsilon(1S)$ R_{AA} measurements at $\sqrt{s_{NN}} = 5.02$ TeV are compared to predictions from two transport models [33,72] and one hydro-dynamical model [34] as a function of centrality (top), p_T (left) and y (right). See text for details on the models.

within uncertainties. The nuclear modification factor shows no significant dependence on rapidity. The $\Upsilon(1S)$ R_{AA} as a function of centrality and rapidity measured by ALICE at $\sqrt{s_{NN}} = 2.76$ TeV [30] are also shown in Fig. 2. Similar trends can be observed at both collision energies.

The inclusive $\Upsilon(1S)$ R_{AA} measurements are compared in Fig. 2 to several calculations: two transport models (TM) [33,72] and one hydro-dynamical model [34]. To describe the quarkonium motion in the medium, both transport codes use a rate-equation approach which accounts for both suppression and (re)generation mechanisms in the QGP. In the TM1 model [33] the evolution of the thermal medium is based on a thermal-fireball expansion while the TM2 model [72] uses a 2+1 dimensional version of the ideal hydrodynamic equations. The two models use different rate equations and both models include a feed-down contribution from higher-mass bottomonia to the $\Upsilon(1S)$. In TM2, two sets of feed-down fractions are assumed. Finally, the $\Upsilon(1S)$ production cross section in pp collisions at $\sqrt{s} = 5.02$ TeV in the rapidity range $2.5 < y < 4$ is taken as $d\sigma_{pp}^{\Upsilon(1S)}/dy = 28.8$ nb in TM1 and $d\sigma_{pp}^{\Upsilon(1S)}/dy = 30$ nb in TM2. Those values deviate by about 2σ (TM1) and 1.4σ (TM2) from the result obtained using the pp interpolation method reported in the previous section. TM1 predictions are shown as bands accounting for shadowing effects as calculated in [36]. The upper limit shown in Fig. 2 corresponds to the extreme case of the absence of shadowing while the lower limit

reflects a reduction of 30% due to shadowing. The TM1 model implements the feed-down fractions reported in [9]. In the TM2 model, the shadowing parameterization is based on EKS98 [35] and the band edges correspond to two different sets of feed-down fractions (27% from χ_b ; 11% from $\Upsilon(2S + 3S)$ and 37% from χ_b ; 12% from $\Upsilon(2S + 3S)$) adopted by the authors. In the third model [34], a thermal suppression of the bottomonium states is calculated using a complex-valued heavy-quark potential parametrized by means of lattice QCD and embedded in a medium evolving according to 3+1d anisotropic hydrodynamics. In this recent study, the R_{AA} shows no sensitivity to the plasma shear viscosity-to-entropy density ratio ($4\pi\eta/s$) parameter of the hydro evolution, which is therefore set to $4\pi\eta/s = 2$ consistent with particle spectra fits. The band of the model quantifies the heavy-quark potential uncertainty, which has been estimated by including a $\pm 15\%$ variation of the Debye mass of the QCD medium that is tuned by a fit to the real-part of the lattice in-medium heavy-quark potential. Furthermore, the predictions shown are referring to the initial momentum-space anisotropy parameter $\xi_0 = 0$, which corresponds to a perfectly isotropic QGP at the starting point of the hydro-dynamical evolution at $\tau_0 = 0.3$ fm/c. Finally, this model accounts for feed-down contributions but it includes neither a (re)generation mechanism nor CNM effects. The centrality dependence of the $\Upsilon(1S)$ R_{AA} is fairly reproduced by the model calculations in the top panel of Fig. 2. The data are best described by TM1 when (re)generation is included and by TM2 when (re)generation is not taken

into account. The hydro-dynamical model describes the trend of the data, the fact that the data lie on the upper edge of the uncertainty band for $N_{\text{part}} > 70$ could indicate a smaller Debye mass and thus a stronger heavy-quark potential. The data as a function of p_T (bottom left panel of Fig. 2) can be described with or without the (re)generation scenario of the TM1 model while showing agreement with the hydro-dynamical model for the upper edge of the uncertainty band. Finally, the y -dependence of the $\Upsilon(1S)$ R_{AA} is described, within uncertainties, by the hydro-dynamical model in the bottom right panel of Fig. 2 despite the possibly different trend between data and calculations.

The low $\Upsilon(1S)$ R_{AA} reported in this Letter raises the important question whether direct $\Upsilon(1S)$ are suppressed at LHC energies or only the feed-down contribution from higher mass states. However, the large uncertainties of the current measurements of CNM effects [38–40] prevent a firm conclusion.

6. Summary

The nuclear modification factors of inclusive $\Upsilon(1S)$ and $\Upsilon(2S)$ production at forward rapidity ($2.5 < y < 4$) and for $p_T < 15 \text{ GeV}/c$ in Pb–Pb collisions at $\sqrt{s_{\text{NN}}} = 5.02 \text{ TeV}$ have been measured using the ALICE detector. The observed $\Upsilon(1S)$ suppression increases with the centrality of the collision and no significant variation is observed as a function of transverse momentum or rapidity. A larger suppression of the $\Upsilon(2S)$ bound state compared to the ground state is also reported. Transport and dynamical model calculations reproduce qualitatively the centrality and kinematic dependence of the $\Upsilon(1S)$ nuclear modification factor.

Acknowledgements

The ALICE Collaboration would like to thank all its engineers and technicians for their invaluable contributions to the construction of the experiment and the CERN accelerator teams for the outstanding performance of the LHC complex. The ALICE Collaboration gratefully acknowledges the resources and support provided by all Grid centres and the Worldwide LHC Computing Grid (WLCG) collaboration. The ALICE Collaboration acknowledges the following funding agencies for their support in building and running the ALICE detector: A. I. Alikhanyan National Science Laboratory (Yerevan Physics Institute) Foundation (ANSL), State Committee of Science and World Federation of Scientists (WFS), Armenia; Austrian Academy of Sciences and Österreichische Nationalstiftung für Forschung, Technologie und Entwicklung, Austria; Ministry of Communications and High Technologies, National Nuclear Research Center, Azerbaijan; Conselho Nacional de Desenvolvimento Científico e Tecnológico (CNPq), Universidade Federal do Rio Grande do Sul (UFRGS), Financiadora de Estudos e Projetos (Finep) and Fundação de Amparo à Pesquisa do Estado de São Paulo (FAPESP), Brazil; Ministry of Science & Technology of China (MSTC), National Natural Science Foundation of China (NSFC) and Ministry of Education of China (MOEC), China; Ministry of Science and Education, Croatia; Ministry of Education, Youth and Sports of the Czech Republic, Czech Republic; The Danish Council for Independent Research – Natural Sciences, the Carlsberg Foundation and Danish National Research Foundation (DNRF), Denmark; Helsinki Institute of Physics (HIP), Finland; Commissariat à l’Energie Atomique (CEA) and Institut National de Physique Nucléaire et de Physique des Particules (IN2P3) and Centre National de la Recherche Scientifique (CNRS), France; Bundesministerium für Bildung, Wissenschaft, Forschung und Technologie (BMBF) and GSI Helmholtzzentrum für Schwerionenforschung GmbH, Germany; General Secretariat for Research and Technology, Ministry of Education, Research and

Religions, Greece; National Research Development and Innovation Office, Hungary; Department of Atomic Energy, Government of India (DAE), Department of Science and Technology, Government of India (DST), University Grants Commission, Government of India (UGC) and Council of Scientific and Industrial Research (CSIR), India; Indonesian Institute of Sciences, Indonesia; Centro Fermi - Museo Storico della Fisica e Centro Studi e Ricerche Enrico Fermi and Istituto Nazionale di Fisica Nucleare (INFN), Italy; Institute for Innovative Science and Technology, Nagasaki Institute of Applied Science (IIST), Japan Society for the Promotion of Science (JSPS) KAKENHI and Japanese Ministry of Education, Culture, Sports, Science and Technology (MEXT), Japan; Consejo Nacional de Ciencia (CONACYT) y Tecnología, through Fondo de Cooperación Internacional en Ciencia y Tecnología (FONCICYT) and Dirección General de Asuntos del Personal Académico (DGAPA), Mexico; Nederlandse Organisatie voor Wetenschappelijk Onderzoek (NWO), Netherlands; The Research Council of Norway, Norway; Commission on Science and Technology for Sustainable Development in the South (COMSATS), Pakistan; Pontificia Universidad Católica del Perú, Peru; Ministry of Science and Higher Education and National Science Centre, Poland; Korea Institute of Science and Technology Information and National Research Foundation of Korea (NRF), Republic of Korea; Ministry of Education and Scientific Research, Institute of Atomic Physics and Romanian National Agency for Science, Technology and Innovation, Romania; Joint Institute for Nuclear Research (JINR), Ministry of Education and Science of the Russian Federation and National Research Centre Kurchatov Institute, Russia; Ministry of Education, Science, Research and Sport of the Slovak Republic, Slovakia; National Research Foundation of South Africa, South Africa; Centro de Aplicaciones Tecnológicas y Desarrollo Nuclear (CEADEN), Cubaenergía, Cuba and Centro de Investigaciones Energéticas, Medioambientales y Tecnológicas (CIEMAT), Spain; Swedish Research Council (VR) and Knut & Alice Wallenberg Foundation (KAW), Sweden; European Organization for Nuclear Research, Switzerland; National Science and Technology Development Agency (NSDTA), Suranaree University of Technology (SUT) and Office of the Higher Education Commission under NRU project of Thailand, Thailand; Turkish Atomic Energy Agency (TAEK), Turkey; National Academy of Sciences of Ukraine, Ukraine; Science and Technology Facilities Council (STFC), United Kingdom; National Science Foundation of the United States of America (NSF) and United States Department of Energy, Office of Nuclear Physics (DOE NP), United States of America.

References

- [1] E.V. Shuryak, Quark-gluon plasma and hadronic production of leptons, photons and pions, *Phys. Lett. B* 78 (1978) 150, *Yad. Fiz.* 28 (1978) 796.
- [2] BRAHMS Collaboration, I. Arsene, et al., Quark gluon plasma and color glass condensate at RHIC? The Perspective from the BRAHMS experiment, *Nucl. Phys. A* 757 (2005) 1–27, arXiv:nucl-ex/0410020 [nucl-ex].
- [3] PHOBOS Collaboration, B.B. Back, et al., The PHOBOS perspective on discoveries at RHIC, *Nucl. Phys. A* 757 (2005) 28–101, arXiv:nucl-ex/0410022 [nucl-ex].
- [4] STAR Collaboration, J. Adams, et al., Experimental and theoretical challenges in the search for the quark gluon plasma: the STAR Collaboration’s critical assessment of the evidence from RHIC collisions, *Nucl. Phys. A* 757 (2005) 102–183, arXiv:nucl-ex/0501009 [nucl-ex].
- [5] PHENIX Collaboration, K. Adcox, et al., Formation of dense partonic matter in relativistic nucleus–nucleus collisions at RHIC: experimental evaluation by the PHENIX collaboration, *Nucl. Phys. A* 757 (2005) 184–283, arXiv:nucl-ex/0410003 [nucl-ex].
- [6] B. Müller, J. Schukraft, B. Wyslouch, First results from Pb+Pb collisions at the LHC, *Annu. Rev. Nucl. Part. Sci.* 62 (2012) 361–386, arXiv:1202.3233 [hep-ex].
- [7] T. Matsui, H. Satz, J/ψ suppression by quark-gluon plasma formation, *Phys. Lett. B* 178 (1986) 416–422.
- [8] N. Brambilla, et al., Heavy quarkonium: progress, puzzles, and opportunities, *Eur. Phys. J. C* 71 (2011) 1534, arXiv:1010.5827 [hep-ph].
- [9] A. Andronic, et al., Heavy-flavour and quarkonium production in the LHC era: from proton-proton to heavy-ion collisions, *Eur. Phys. J. C* 76 (3) (2016), arXiv:1506.03981 [nucl-ex].

- [10] S. Digal, P. Petreczky, H. Satz, Quarkonium feed down and sequential suppression, Phys. Rev. D 64 (2001) 094015, arXiv:hep-ph/0106017 [hep-ph].
- [11] B. Kroupa, R. Ryblewski, M. Strickland, Bottomonia suppression in 2.76 TeV Pb–Pb collisions, Phys. Rev. C 92 (6) (2015) 061901, arXiv:1507.03951 [hep-ph].
- [12] M. Martinez, M. Strickland, Measuring QGP thermalization time with dileptons, Phys. Rev. Lett. 100 (2008) 102301, arXiv:0709.3576 [hep-ph].
- [13] Y. Burnier, O. Kaczmarek, A. Rothkopf, Static quark–antiquark potential in the quark–gluon plasma from lattice QCD, Phys. Rev. Lett. 114 (8) (2015) 082001, arXiv:1410.2546 [hep-lat].
- [14] NA38 Collaboration, M.C. Abreu, et al., J/ψ , ψ' and Drell–Yan production in S–U interactions at 200 GeV per nucleon, Phys. Lett. B 449 (1999) 128–136, <http://hal.in2p3.fr/in2p3-00002434>.
- [15] NA50 Collaboration, B. Alessandro, et al., A new measurement of J/ψ suppression in Pb–Pb collisions at 158–GeV per nucleon, Eur. Phys. J. C 39 (2005) 335–345, arXiv:hep-ex/0412036 [hep-ex].
- [16] NA60 Collaboration, R. Arnaldi, et al., J/ψ production in indium–indium collisions at 158–GeV/nucleon, Phys. Rev. Lett. 99 (2007) 132302.
- [17] PHENIX Collaboration, A. Adare, et al., J/ψ suppression at forward rapidity in Au+Au collisions at $\sqrt{s_{NN}} = 200$ GeV, Phys. Rev. C 84 (2011) 054912, arXiv: 1103.6269 [nucl-ex].
- [18] STAR Collaboration, B.I. Abelev, et al., J/ψ production at high transverse momentum in p+p and Cu+Cu collisions at $\sqrt{s_{NN}} = 200$ GeV, Phys. Rev. C 80 (2009) 041902, arXiv:0904.0439 [nucl-ex].
- [19] ALICE Collaboration, B. Abelev, et al., J/ψ suppression at forward rapidity in Pb–Pb collisions at $\sqrt{s_{NN}} = 2.76$ TeV, Phys. Rev. Lett. 109 (2012) 072301, arXiv: 1202.1383 [hep-ex].
- [20] ALICE Collaboration, B.B. Abelev, et al., Centrality, rapidity and transverse momentum dependence of J/ψ suppression in Pb–Pb collisions at $\sqrt{s_{NN}} = 2.76$ TeV, Phys. Lett. B 734 (2014) 314–327, arXiv:1311.0214 [nucl-ex].
- [21] CMS Collaboration, V. Khachatryan, et al., Suppression and azimuthal anisotropy of prompt and nonprompt J/ψ production in PbPb collisions at $\sqrt{s_{NN}} = 2.76$ TeV, Eur. Phys. J. C 77 (4) (2017) 252, arXiv:1610.00613 [nucl-ex].
- [22] ALICE Collaboration, J. Adam, et al., J/ψ suppression at forward rapidity in Pb–Pb collisions at $\sqrt{s_{NN}} = 5.02$ TeV, Phys. Lett. B 766 (2017) 212–224, arXiv: 1606.08197 [nucl-ex].
- [23] P. Braun-Munzinger, J. Stachel, (Non)thermal aspects of charmonium production and a new look at J/ψ suppression, Phys. Lett. B 490 (2000) 196–202, arXiv:nucl-th/0007059 [nucl-th].
- [24] R.L. Thews, M. Schrödter, J. Rafelski, Enhanced J/ψ production in deconfined quark matter, Phys. Rev. C 63 (2001) 054905, arXiv:hep-ph/0007323 [hep-ph].
- [25] X. Zhao, R. Rapp, Medium modifications and production of charmonia at LHC, Nucl. Phys. A 859 (2011) 114–125, arXiv:1102.2194 [hep-ph].
- [26] K. Zhou, N. Xu, Z. Xu, P. Zhuang, Medium effects on charmonium production at ultrarelativistic energies available at the CERN Large Hadron Collider, Phys. Rev. C 89 (5) (2014) 054911, arXiv:1401.5845 [nucl-th].
- [27] ALICE Collaboration, J. Adam, et al., Differential studies of inclusive J/ψ and $\psi(2S)$ production at forward rapidity in Pb–Pb collisions at $\sqrt{s_{NN}} = 2.76$ TeV, J. High Energy Phys. 05 (2016) 179, arXiv:1506.08804 [nucl-ex].
- [28] M.L. Miller, K. Reiglers, S.J. Sanders, P. Steinberg, Glauber modeling in high energy nuclear collisions, Annu. Rev. Nucl. Part. Sci. 57 (2007) 205–243, arXiv: nucl-ex/0701025 [nucl-ex].
- [29] C. Loizides, J. Kämä, D. d’Enterria, Precision Monte Carlo Glauber predictions at present and future nuclear colliders, arXiv:1710.07098 [nucl-ex].
- [30] ALICE Collaboration, B.B. Abelev, et al., Suppression of $\Upsilon(1S)$ at forward rapidity in Pb–Pb collisions at $\sqrt{s_{NN}} = 2.76$ TeV, Phys. Lett. B 738 (2014) 361–372, arXiv:1405.4493 [nucl-ex].
- [31] CMS Collaboration, S. Chatrchyan, et al., Observation of sequential Υ suppression in Pb–Pb collisions, Phys. Rev. Lett. 109 (2012) 222301, arXiv:1208.2826 [nucl-ex].
- [32] CMS Collaboration, V. Khachatryan, et al., Suppression of $\Upsilon(1S)$, $\Upsilon(2S)$ and $\Upsilon(3S)$ production in PbPb collisions at $\sqrt{s_{NN}} = 2.76$ TeV, Phys. Lett. B 770 (2017) 357–379, arXiv:1611.01510 [nucl-ex].
- [33] X. Du, R. Rapp, M. He, Color screening and regeneration of bottomonia in high-energy heavy-ion collisions, Phys. Rev. C 96 (5) (2017) 054901, arXiv: 1706.08670 [hep-ph].
- [34] B. Kroupa, A. Rothkopf, M. Strickland, Bottomonium suppression using a lattice QCD vetted potential, Phys. Rev. D 97 (1) (2018) 016017, arXiv:1710.02319 [hep-ph].
- [35] K.J. Eskola, V.J. Kolhinen, C.A. Salgado, The scale dependent nuclear effects in parton distributions for practical applications, Eur. Phys. J. C 9 (1999) 61–68, arXiv:hep-ph/9807297 [hep-ph].
- [36] K.J. Eskola, H. Paukkunen, C.A. Salgado, EPS09: a new generation of NLO and LO nuclear parton distribution functions, J. High Energy Phys. 04 (2009) 065, arXiv:0902.4154 [hep-ph].
- [37] F. Arleo, S. Peigne, Heavy-quarkonium suppression in p+A collisions from parton energy loss in cold QCD matter, J. High Energy Phys. 03 (2013) 122, arXiv: 1212.0434 [hep-ph].
- [38] ALICE Collaboration, B.B. Abelev, et al., Production of inclusive $\Upsilon(1S)$ and $\Upsilon(2S)$ in p–Pb collisions at $\sqrt{s_{NN}} = 5.02$ TeV, Phys. Lett. B 740 (2015) 105–117, arXiv: 1410.2234 [nucl-ex].
- [39] LHCb Collaboration, R. Aaij, et al., Study of Υ production and cold nuclear matter effects in pPb collisions at $\sqrt{s_{NN}} = 5$ TeV, J. High Energy Phys. 07 (2014) 094, arXiv:1405.5152 [nucl-ex].
- [40] ATLAS Collaboration, M. Aaboud, et al., Measurement of quarkonium production in proton–lead and proton–proton collisions at 5.02 TeV with the ATLAS detector, Eur. Phys. J. C 78 (3) (2018), arXiv:1709.03089 [nucl-ex].
- [41] ALICE Collaboration, B.B. Abelev, et al., Measurement of quarkonium production at forward rapidity in pp collisions at $\sqrt{s} = 7$ TeV, Eur. Phys. J. C 74 (8) (2014) 2974, arXiv:1403.3648 [nucl-ex].
- [42] ALICE Collaboration, J. Adam, et al., Inclusive quarkonium production at forward rapidity in pp collisions at $\sqrt{s} = 8$ TeV, Eur. Phys. J. C 76 (4) (2016) 184, arXiv:1509.08258 [hep-ex].
- [43] LHCb Collaboration, R. Aaij, et al., Measurement of Υ production in pp collisions at $\sqrt{s} = 2.76$ TeV, Eur. Phys. J. C 74 (4) (2014) 2835, arXiv:1402.2539 [hep-ex].
- [44] LHCb Collaboration, R. Aaij, et al., Forward production of Υ mesons in pp collisions at $\sqrt{s} = 7$ and 8 TeV, J. High Energy Phys. 11 (2015) 103, arXiv: 1509.02372 [hep-ex].
- [45] ALICE Collaboration, K. Aamodt, et al., The ALICE experiment at the CERN LHC, J. Instrum. 3 (2008) S08002.
- [46] ALICE Collaboration, B.B. Abelev, et al., Performance of the ALICE experiment at the CERN LHC, Int. J. Mod. Phys. A 29 (2014) 1430044, arXiv:1402.4476 [nucl-ex].
- [47] ALICE Collaboration, K. Aamodt, et al., Rapidity and transverse momentum dependence of inclusive J/ψ production in pp collisions at $\sqrt{s} = 7$ TeV, Phys. Lett. B 704 (2011) 442, arXiv:1105.0380 [hep-ex].
- [48] ALICE Collaboration, K. Aamodt, et al., Alignment of the ALICE inner tracking system with cosmic-ray tracks, J. Instrum. 5 (2010) P03003, arXiv:1001.0502 [physics.ins-det].
- [49] ALICE Collaboration, E. Abbas, et al., Performance of the ALICE VZERO system, J. Instrum. 8 (2013) P10016, arXiv:1306.3130 [nucl-ex].
- [50] ALICE Collaboration, B. Abelev, et al., Measurement of the cross section for electromagnetic dissociation with neutron emission in Pb–Pb collisions at $\sqrt{s_{NN}} = 2.76$ TeV, Phys. Rev. Lett. 109 (2012) 252302, arXiv:1203.2436 [nucl-ex].
- [51] ALICE Collaboration, F. Bossù, M. Gagliardi, M. Marchisone, Performance of the RPC-based ALICE muon trigger system at the LHC, J. Instrum. 7 (12) (2012) T12002, <http://stacks.iop.org/1748-0221/7/i=12/a=T12002>.
- [52] R. Arnaldi, et al., Design and performance of the ALICE muon trigger system, Nucl. Phys., Proc. Suppl. 158 (2006) 21–24.
- [53] ALICE Collaboration, B. Abelev, et al., Centrality determination of Pb–Pb collisions at $\sqrt{s_{NN}} = 2.76$ TeV with ALICE, Phys. Rev. C 88 (4) (2013) 044909, arXiv:1301.4361 [nucl-ex].
- [54] ALICE Collaboration, J. Adam, et al., Centrality dependence of the charged-particle multiplicity density at midrapidity in Pb–Pb collisions at $\sqrt{s_{NN}} = 5.02$ TeV, Phys. Rev. Lett. 116 (22) (2016) 222302, arXiv:1512.06104 [nucl-ex].
- [55] ALICE Collaboration, Centrality determination in heavy ion collisions, ALICE-PUBLIC-2018-011, 2018.
- [56] D.G. d’Enterria, Hard scattering cross-sections at LHC in the Glauber approach: from pp to pA and AA collisions, arXiv:nucl-ex/0302016 [nucl-ex].
- [57] C. Patrignani, et al., Particle Data Group, Chin. Phys. C 40 (10) (2016) 100001 and 2017 update.
- [58] ALICE Collaboration, Quarkonium signal extraction in ALICE, ALICE-PUBLIC-2015-006, <https://cds.cern.ch/record/2060096>.
- [59] CDF Collaboration, D. Acosta, et al., Υ production and polarization in $p\bar{p}$ collisions at $\sqrt{s} = 1.8$ –TeV, Phys. Rev. Lett. 88 (2002) 161802.
- [60] LHCb Collaboration, R. Aaij, et al., Measurement of Υ production in pp collisions at $\sqrt{s} = 7$ TeV, Eur. Phys. J. C 72 (2012), arXiv:1202.6579 [hep-ex].
- [61] CMS Collaboration, V. Khachatryan, et al., Upsilon production cross-section in pp collisions at $\sqrt{s} = 7$ TeV, Phys. Rev. D 83 (2011) 112004, arXiv:1012.5545 [hep-ex].
- [62] F. Bossù, Z.C. del Valle, A. de Falco, M. Gagliardi, S. Grigoryan, G. Martínez García, Phenomenological interpolation of the inclusive J/ψ cross section to proton–proton collisions at 2.76 TeV and 5.5 TeV, arXiv:1103.2394 [nucl-ex].
- [63] D0 Collaboration, V.M. Abazov, et al., Measurement of the polarization of the $\Upsilon(1S)$ and $\Upsilon(2S)$ states in $p\bar{p}$ collisions at $\sqrt{s} = 1.96$ –TeV, Phys. Rev. Lett. 101 (2008) 182004, arXiv:0804.2799 [hep-ex].
- [64] CDF Collaboration, T. Altonen, et al., Measurements of angular distributions of muons from Υ meson decays in $p\bar{p}$ collisions at $\sqrt{s} = 1.96$ TeV, Phys. Rev. Lett. 108 (2012) 151802, arXiv:1112.1591 [hep-ex].
- [65] CMS Collaboration, S. Chatrchyan, et al., Measurement of the $\Upsilon(1S)$, $\Upsilon(2S)$ and $\Upsilon(3S)$ polarizations in pp collisions at $\sqrt{s} = 7$ TeV, Phys. Rev. Lett. 110 (8) (2013) 081802, arXiv:1209.2922 [hep-ex].
- [66] LHCb Collaboration, R. Aaij, et al., Measurement of the Υ polarizations in pp collisions at $\sqrt{s} = 7$ and 8 TeV, J. High Energy Phys. 12 (2017), arXiv:1709.01301 [hep-ex].
- [67] GEANT4 Collaboration, S. Agostinelli, et al., GEANT4: a simulation toolkit, Nucl. Instrum. Methods, Sect. A 506 (2003) 250–303.

- [68] R. Brun, F. Bruyant, F. Carminati, S. Giani, M. Maire, A. McPherson, G. Patrick, L. Urban, GEANT: Detector Description and Simulation Tool; Oct 1994, CERN Program Library, CERN, Geneva, 1993, <http://cds.cern.ch/record/1082634>, Long Writeup W5013.
- [69] ALICE and LHC Collaboration, Reference pp cross-sections for $\Upsilon(1S)$ studies in proton-lead collisions at $\sqrt{s_{NN}} = 5.02$ TeV and comparisons between ALICE and LHCb results, <http://cds.cern.ch/record/1748460>, ALICE-PUBLIC-2014-002, LHCb-CONF-2014-003.
- [70] V. Cheung, R. Vogt, Polarization of prompt J/ψ and $\Upsilon(1S)$ production in the color evaporation model, Phys. Rev. D 96 (5) (2017) 054014, arXiv:1706.07686 [hep-ph].
- [71] M. Cacciari, M. Greco, P. Nason, The p_T spectrum in heavy flavor hadroproduction, J. High Energy Phys. 05 (1998) 007, arXiv:hep-ph/9803400 [hep-ph].
- [72] K. Zhou, N. Xu, P. Zhuang, Υ production in heavy ion collisions at LHC, Nucl. Phys. A 931 (2014) 654–658, arXiv:1408.3900 [hep-ph].

ALICE Collaboration

- S. Acharya ¹³⁹, F.T. Acosta ²⁰, D. Adamová ⁹³, J. Adolfsson ⁸⁰, M.M. Aggarwal ⁹⁸, G. Aglieri Rinella ³⁴, M. Agnello ³¹, N. Agrawal ⁴⁸, Z. Ahammed ¹³⁹, S.U. Ahn ⁷⁶, S. Aiola ¹⁴⁴, A. Akindinov ⁶⁴, M. Al-Turany ¹⁰⁴, S.N. Alam ¹³⁹, D.S.D. Albuquerque ¹²¹, D. Aleksandrov ⁸⁷, B. Alessandro ⁵⁸, R. Alfaro Molina ⁷², Y. Ali ¹⁵, A. Alice ^{10,53,27}, A. Alkin ², J. Alme ²², T. Alt ⁶⁹, L. Altenkamper ²², I. Altsybeev ¹¹¹, M.N. Anaam ⁶, C. Andrei ⁴⁷, D. Andreou ³⁴, H.A. Andrews ¹⁰⁸, A. Andronic ^{142,104}, M. Angeletti ³⁴, V. Anguelov ¹⁰², C. Anson ¹⁶, T. Antičić ¹⁰⁵, F. Antinori ⁵⁶, P. Antonioli ⁵³, R. Anwar ¹²⁵, N. Apadula ⁷⁹, L. Aphecetche ¹¹³, H. Appelshäuser ⁶⁹, S. Arcelli ²⁷, R. Arnaldi ⁵⁸, O.W. Arnold ^{103,116}, I.C. Arsene ²¹, M. Arslanbekov ¹⁰², B. Audurier ¹¹³, A. Augustinus ³⁴, R. Averbeck ¹⁰⁴, M.D. Azmi ¹⁷, A. Badalà ⁵⁵, Y.W. Baek ^{60,40}, S. Bagnasco ⁵⁸, R. Bailhache ⁶⁹, R. Bala ⁹⁹, A. Baldissari ¹³⁵, M. Ball ⁴², R.C. Baral ⁸⁵, A.M. Barbano ²⁶, R. Barbera ²⁸, F. Barile ⁵², L. Barioglio ²⁶, G.G. Barnaföldi ¹⁴³, L.S. Barnby ⁹², V. Barret ¹³², P. Bartalini ⁶, K. Barth ³⁴, E. Bartsch ⁶⁹, N. Bastid ¹³², S. Basu ¹⁴¹, G. Batigne ¹¹³, B. Batyunya ⁷⁵, P.C. Batzing ²¹, J.L. Bazo Alba ¹⁰⁹, I.G. Bearden ⁸⁸, H. Beck ¹⁰², C. Bedda ⁶³, N.K. Behera ⁶⁰, I. Belikov ¹³⁴, F. Bellini ³⁴, H. Bello Martinez ⁴⁴, R. Bellwied ¹²⁵, L.G.E. Beltran ¹¹⁹, V. Belyaev ⁹¹, G. Bencedi ¹⁴³, S. Beole ²⁶, A. Bercuci ⁴⁷, Y. Berdnikov ⁹⁶, D. Berenyi ¹⁴³, R.A. Bertens ¹²⁸, D. Berzana ^{34,58}, L. Betev ³⁴, P.P. Bhaduri ¹³⁹, A. Bhasin ⁹⁹, I.R. Bhat ⁹⁹, H. Bhatt ⁴⁸, B. Bhattacharjee ⁴¹, J. Bhom ¹¹⁷, A. Bianchi ²⁶, L. Bianchi ¹²⁵, N. Bianchi ⁵¹, J. Bielčík ³⁷, J. Bielčíková ⁹³, A. Bilandžić ^{116,103}, G. Biro ¹⁴³, R. Biswas ³, S. Biswas ³, J.T. Blair ¹¹⁸, D. Blau ⁸⁷, C. Blume ⁶⁹, G. Boca ¹³⁷, F. Bock ³⁴, A. Bogdanov ⁹¹, L. Boldizsár ¹⁴³, M. Bombara ³⁸, G. Bonomi ¹³⁸, M. Bonora ³⁴, H. Borel ¹³⁵, A. Borissov ^{18,142}, M. Borri ¹²⁷, E. Botta ²⁶, C. Bourjau ⁸⁸, L. Bratrud ⁶⁹, P. Braun-Munzinger ¹⁰⁴, M. Bregant ¹²⁰, T.A. Broker ⁶⁹, M. Broz ³⁷, E.J. Bruckner ⁴³, E. Bruna ⁵⁸, G.E. Bruno ^{34,33}, D. Budnikov ¹⁰⁶, H. Buesching ⁶⁹, S. Bufalino ³¹, P. Buhler ¹¹², P. Buncic ³⁴, O. Busch ^{131,i}, Z. Buthelezi ⁷³, J.B. Butt ¹⁵, J.T. Buxton ⁹⁵, J. Cabala ¹¹⁵, D. Caffarri ⁸⁹, H. Caines ¹⁴⁴, A. Caliva ¹⁰⁴, E. Calvo Villar ¹⁰⁹, R.S. Camacho ⁴⁴, P. Camerini ²⁵, A.A. Capon ¹¹², F. Carena ³⁴, W. Carena ³⁴, F. Carnesecchi ^{27,10}, J. Castillo Castellanos ¹³⁵, A.J. Castro ¹²⁸, E.A.R. Casula ⁵⁴, C. Ceballos Sanchez ⁸, S. Chandra ¹³⁹, B. Chang ¹²⁶, W. Chang ⁶, S. Chapelard ³⁴, M. Chartier ¹²⁷, S. Chattopadhyay ¹³⁹, S. Chattopadhyay ¹⁰⁷, A. Chauvin ^{103,116}, C. Cheshkov ¹³³, B. Cheynis ¹³³, V. Chibante Barroso ³⁴, D.D. Chinellato ¹²¹, S. Cho ⁶⁰, P. Chochula ³⁴, T. Chowdhury ¹³², P. Christakoglou ⁸⁹, C.H. Christensen ⁸⁸, P. Christiansen ⁸⁰, T. Chujo ¹³¹, S.U. Chung ¹⁸, C. Cicalo ⁵⁴, L. Cifarelli ^{10,27}, F. Cindolo ⁵³, J. Cleymans ¹²⁴, F. Colamaria ⁵², D. Colella ^{65,34,52}, A. Collu ⁷⁹, M. Colocci ²⁷, M. Concas ^{58,ii}, G. Conesa Balbastre ⁷⁸, Z. Conesa del Valle ⁶¹, J.G. Contreras ³⁷, T.M. Cormier ⁹⁴, Y. Corrales Morales ⁵⁸, P. Cortese ³², M.R. Cosentino ¹²², F. Costa ³⁴, S. Costanza ¹³⁷, J. Crkovská ⁶¹, P. Crochet ¹³², E. Cuautle ⁷⁰, L. Cunqueiro ^{142,94}, T. Dahms ^{103,116}, A. Dainese ⁵⁶, S. Dani ⁶⁶, M.C. Danisch ¹⁰², A. Danu ⁶⁸, D. Das ¹⁰⁷, I. Das ¹⁰⁷, S. Das ³, A. Dash ⁸⁵, S. Dash ⁴⁸, S. De ⁴⁹, A. De Caro ³⁰, G. de Cataldo ⁵², C. de Conti ¹²⁰, J. de Cuveland ³⁹, A. De Falco ²⁴, D. De Gruttola ^{10,30}, N. De Marco ⁵⁸, S. De Pasquale ³⁰, R.D. De Souza ¹²¹, H.F. Degenhardt ¹²⁰, A. Deistung ^{104,102}, A. Deloff ⁸⁴, S. Delsanto ²⁶, C. Deplano ⁸⁹, P. Dhankher ⁴⁸, D. Di Bari ³³, A. Di Mauro ³⁴, B. Di Ruzza ⁵⁶, R.A. Diaz ⁸, T. Dietel ¹²⁴, P. Dillenseger ⁶⁹, Y. Ding ⁶, R. Divià ³⁴, Ø. Djupsland ²², A. Dobrin ³⁴, D. Domenicis Gimenez ¹²⁰, B. Dönigus ⁶⁹, O. Dordic ²¹, L.V.R. Doremaleen ⁶³, A.K. Dubey ¹³⁹, A. Dubla ¹⁰⁴, L. Ducroux ¹³³, S. Dudi ⁹⁸, A.K. Duggal ⁹⁸, M. Dukhishyam ⁸⁵, P. Dupieux ¹³², R.J. Ehlers ¹⁴⁴, D. Elia ⁵², E. Endress ¹⁰⁹, H. Engel ⁷⁴, E. Epple ¹⁴⁴, B. Erazmus ¹¹³, F. Erhardt ⁹⁷, M.R. Ersdal ²², B. Espagnon ⁶¹, G. Eulisse ³⁴, J. Eum ¹⁸, D. Evans ¹⁰⁸, S. Evdokimov ⁹⁰, L. Fabbietti ^{103,116}, M. Faggin ²⁹, J. Faivre ⁷⁸, A. Fantoni ⁵¹, M. Fasel ⁹⁴, L. Feldkamp ¹⁴², A. Feliciello ⁵⁸, G. Feofilov ¹¹¹, A. Fernández Téllez ⁴⁴, A. Ferretti ²⁶, A. Festanti ^{29,34}, V.J.G. Feuillard ¹⁰², J. Figiel ¹¹⁷, M.A.S. Figueredo ¹²⁰, S. Filchagin ¹⁰⁶, D. Finogeev ⁶², F.M. Fionda ²², G. Fiorenza ⁵², F. Flor ¹²⁵, M. Floris ³⁴, S. Foertsch ⁷³, P. Foka ¹⁰⁴, S. Fokin ⁸⁷, E. Fragiocomo ⁵⁹, A. Francescon ³⁴, A. Francisco ¹¹³, U. Frankenfeld ¹⁰⁴, G.G. Fronze ²⁶, U. Fuchs ³⁴, C. Furet ⁷⁸, A. Furs ⁶², M. Fusco Girard ³⁰, J.J. Gaardhøje ⁸⁸,

- M. Gagliardi ²⁶, A.M. Gago ¹⁰⁹, K. Gajdosova ⁸⁸, M. Gallio ²⁶, C.D. Galvan ¹¹⁹, P. Ganoti ⁸³, C. Garabatos ¹⁰⁴, E. Garcia-Solis ¹¹, K. Garg ²⁸, C. Gargiulo ³⁴, P. Gasik ^{116,103}, E.F. Gauger ¹¹⁸, M.B. Gay Ducati ⁷¹, M. Germain ¹¹³, J. Ghosh ¹⁰⁷, P. Ghosh ¹³⁹, S.K. Ghosh ³, P. Gianotti ⁵¹, P. Giubellino ^{104,58}, P. Giubilato ²⁹, P. Glässel ¹⁰², D.M. Goméz Coral ⁷², A. Gomez Ramirez ⁷⁴, V. Gonzalez ¹⁰⁴, P. González-Zamora ⁴⁴, S. Gorbunov ³⁹, L. Görlich ¹¹⁷, S. Gotovac ³⁵, V. Grabski ⁷², L.K. Graczykowski ¹⁴⁰, K.L. Graham ¹⁰⁸, L. Greiner ⁷⁹, A. Grelli ⁶³, C. Grigoras ³⁴, V. Grigoriev ⁹¹, A. Grigoryan ¹, S. Grigoryan ⁷⁵, J.M. Gronefeld ¹⁰⁴, F. Grossa ³¹, J.F. Grosse-Oetringhaus ³⁴, R. Grossos ¹⁰⁴, R. Guernane ⁷⁸, B. Guerzoni ²⁷, M. Guittiere ¹¹³, K. Gulbrandsen ⁸⁸, T. Gunji ¹³⁰, A. Gupta ⁹⁹, R. Gupta ⁹⁹, I.B. Guzman ⁴⁴, R. Haake ³⁴, M.K. Habib ¹⁰⁴, C. Hadjidakis ⁶¹, H. Hamagaki ⁸¹, G. Hamar ¹⁴³, M. Hamid ⁶, J.C. Hamon ¹³⁴, R. Hannigan ¹¹⁸, M.R. Haque ⁶³, J.W. Harris ¹⁴⁴, A. Harton ¹¹, H. Hassan ⁷⁸, D. Hatzifotiadou ^{53,10}, S. Hayashi ¹³⁰, S.T. Heckel ⁶⁹, E. Hellbär ⁶⁹, H. Helstrup ³⁶, A. Herghhelegiu ⁴⁷, E.G. Hernandez ⁴⁴, G. Herrera Corral ⁹, F. Herrmann ¹⁴², K.F. Hetland ³⁶, T.E. Hilden ⁴³, H. Hillemanns ³⁴, C. Hills ¹²⁷, B. Hippolyte ¹³⁴, B. Hohlweier ¹⁰³, D. Horak ³⁷, S. Hornung ¹⁰⁴, R. Hosokawa ^{131,78}, J. Hota ⁶⁶, P. Hristov ³⁴, C. Huang ⁶¹, C. Hughes ¹²⁸, P. Huhn ⁶⁹, T.J. Humanic ⁹⁵, H. Hushnud ¹⁰⁷, N. Hussain ⁴¹, T. Hussain ¹⁷, D. Hutter ³⁹, D.S. Hwang ¹⁹, J.P. Iddon ¹²⁷, S.A. Iga Buitron ⁷⁰, R. Ilkaev ¹⁰⁶, M. Inaba ¹³¹, M. Ippolitov ⁸⁷, M.S. Islam ¹⁰⁷, M. Ivanov ¹⁰⁴, V. Ivanov ⁹⁶, V. Izucheev ⁹⁰, B. Jacak ⁷⁹, N. Jacazio ²⁷, P.M. Jacobs ⁷⁹, M.B. Jadhav ⁴⁸, S. Jadlovska ¹¹⁵, J. Jadlovsky ¹¹⁵, S. Jaelani ⁶³, C. Jahnke ^{120,116}, M.J. Jakubowska ¹⁴⁰, M.A. Janik ¹⁴⁰, C. Jena ⁸⁵, M. Jercic ⁹⁷, O. Jevons ¹⁰⁸, R.T. Jimenez Bustamante ¹⁰⁴, M. Jin ¹²⁵, P.G. Jones ¹⁰⁸, A. Jusko ¹⁰⁸, P. Kalinak ⁶⁵, A. Kalweit ³⁴, J.H. Kang ¹⁴⁵, V. Kaplin ⁹¹, S. Kar ⁶, A. Karasu Uysal ⁷⁷, O. Karavichev ⁶², T. Karavicheva ⁶², P. Karczmarczyk ³⁴, E. Karpechev ⁶², U. Kebschull ⁷⁴, R. Keidel ⁴⁶, D.L.D. Keijdener ⁶³, M. Keil ³⁴, B. Ketzer ⁴², Z. Khabanova ⁸⁹, A.M. Khan ⁶, S. Khan ¹⁷, S.A. Khan ¹³⁹, A. Khanzadeev ⁹⁶, Y. Kharlov ⁹⁰, A. Khatun ¹⁷, A. Khuntia ⁴⁹, M.M. Kielbowicz ¹¹⁷, B. Kileng ³⁶, B. Kim ¹³¹, D. Kim ¹⁴⁵, D.J. Kim ¹²⁶, E.J. Kim ¹³, H. Kim ¹⁴⁵, J.S. Kim ⁴⁰, J. Kim ¹⁰², M. Kim ^{102,60}, S. Kim ¹⁹, T. Kim ¹⁴⁵, T. Kim ¹⁴⁵, S. Kirsch ³⁹, I. Kisel ³⁹, S. Kiselev ⁶⁴, A. Kisiel ¹⁴⁰, J.L. Klay ⁵, C. Klein ⁶⁹, J. Klein ^{34,58}, C. Klein-Bösing ¹⁴², S. Klewin ¹⁰², A. Kluge ³⁴, M.L. Knichel ³⁴, A.G. Knospe ¹²⁵, C. Kobdaj ¹¹⁴, M. Kofarago ¹⁴³, M.K. Köhler ¹⁰², T. Kollegger ¹⁰⁴, N. Kondratyeva ⁹¹, E. Kondratyuk ⁹⁰, A. Konevskikh ⁶², M. Konyushikhin ¹⁴¹, O. Kovalenko ⁸⁴, V. Kovalenko ¹¹¹, M. Kowalski ¹¹⁷, I. Králík ⁶⁵, A. Kravčáková ³⁸, L. Kreis ¹⁰⁴, M. Krivda ^{65,108}, F. Krizek ⁹³, M. Krüger ⁶⁹, E. Kryshen ⁹⁶, M. Krzewicki ³⁹, A.M. Kubera ⁹⁵, V. Kučera ^{60,93}, C. Kuhn ¹³⁴, P.G. Kuijer ⁸⁹, J. Kumar ⁴⁸, L. Kumar ⁹⁸, S. Kumar ⁴⁸, S. Kundu ⁸⁵, P. Kurashvili ⁸⁴, A. Kurepin ⁶², A.B. Kurepin ⁶², A. Kuryakin ¹⁰⁶, S. Kushpil ⁹³, J. Kvapil ¹⁰⁸, M.J. Kweon ⁶⁰, Y. Kwon ¹⁴⁵, S.L. La Pointe ³⁹, P. La Rocca ²⁸, Y.S. Lai ⁷⁹, I. Lakomov ³⁴, R. Langoy ¹²³, K. Lapidus ¹⁴⁴, A. Lardeux ²¹, P. Larionov ⁵¹, E. Laudi ³⁴, R. Lavicka ³⁷, R. Lea ²⁵, L. Leardini ¹⁰², S. Lee ¹⁴⁵, F. Lehas ⁸⁹, S. Lehner ¹¹², J. Lehrbach ³⁹, R.C. Lemmon ⁹², I. León Monzón ¹¹⁹, P. Lévai ¹⁴³, X. Li ¹², X.L. Li ⁶, J. Lien ¹²³, R. Lietava ¹⁰⁸, B. Lim ¹⁸, S. Lindal ²¹, V. Lindenstruth ³⁹, S.W. Lindsay ¹²⁷, C. Lippmann ¹⁰⁴, M.A. Lisa ⁹⁵, V. Litichevskyi ⁴³, A. Liu ⁷⁹, H.M. Ljunggren ⁸⁰, W.J. Llope ¹⁴¹, D.F. Lodato ⁶³, V. Loginov ⁹¹, C. Loizides ^{94,79}, P. Loncar ³⁵, X. Lopez ¹³², E. López Torres ⁸, A. Lowe ¹⁴³, P. Luettig ⁶⁹, J.R. Luhder ¹⁴², M. Lunardon ²⁹, G. Luparello ⁵⁹, M. Lupi ³⁴, A. Maevskaya ⁶², M. Mager ³⁴, S.M. Mahmood ²¹, A. Maire ¹³⁴, R.D. Majka ¹⁴⁴, M. Malaev ⁹⁶, Q.W. Malik ²¹, L. Malinina ^{75,iii}, D. Mal'Kevich ⁶⁴, P. Malzacher ¹⁰⁴, A. Mamontov ¹⁰⁶, V. Manko ⁸⁷, F. Manso ¹³², V. Manzari ⁵², Y. Mao ⁶, M. Marchisone ^{129,73,133}, J. Mareš ⁶⁷, G.V. Margagliotti ²⁵, A. Margotti ⁵³, J. Margutti ⁶³, A. Marín ¹⁰⁴, C. Markert ¹¹⁸, M. Marquard ⁶⁹, N.A. Martin ¹⁰⁴, P. Martinengo ³⁴, J.L. Martinez ¹²⁵, M.I. Martínez ⁴⁴, G. Martínez García ¹¹³, M. Martinez Pedreira ³⁴, S. Masciocchi ¹⁰⁴, M. Masera ²⁶, A. Masoni ⁵⁴, L. Massacrier ⁶¹, E. Masson ¹¹³, A. Mastroserio ^{52,136}, A.M. Mathis ^{116,103}, P.F.T. Matuoka ¹²⁰, A. Matyja ^{117,128}, C. Mayer ¹¹⁷, M. Mazzilli ³³, M.A. Mazzoni ⁵⁷, F. Meddi ²³, Y. Melikyan ⁹¹, A. Menchaca-Rocha ⁷², E. Meninno ³⁰, J. Mercado Pérez ¹⁰², M. Meres ¹⁴, C.S. Meza ¹⁰⁹, S. Mhlanga ¹²⁴, Y. Miake ¹³¹, L. Micheletti ²⁶, M.M. Mieskolainen ⁴³, D.L. Mihaylov ¹⁰³, K. Mikhaylov ^{64,75}, A. Mischke ⁶³, A.N. Mishra ⁷⁰, D. Miśkowiec ¹⁰⁴, J. Mitra ¹³⁹, C.M. Mitu ⁶⁸, N. Mohammadi ³⁴, A.P. Mohanty ⁶³, B. Mohanty ⁸⁵, M. Mohisin Khan ^{17,iv}, D.A. Moreira De Godoy ¹⁴², L.A.P. Moreno ⁴⁴, S. Moretto ²⁹, A. Morreale ¹¹³, A. Morsch ³⁴, V. Muccifora ⁵¹, E. Mudnic ³⁵, D. Mühlheim ¹⁴², S. Muhuri ¹³⁹, M. Mukherjee ³, J.D. Mulligan ¹⁴⁴, M.G. Munhoz ¹²⁰, K. Münnig ⁴², M.I.A. Munoz ⁷⁹, R.H. Munzer ⁶⁹, H. Murakami ¹³⁰, S. Murray ⁷³, L. Musa ³⁴, J. Musinsky ⁶⁵, C.J. Myers ¹²⁵, J.W. Myrcha ¹⁴⁰, B. Naik ⁴⁸, R. Nair ⁸⁴, B.K. Nandi ⁴⁸, R. Nania ^{53,10}, E. Nappi ⁵², A. Narayan ⁴⁸, M.U. Naru ¹⁵, A.F. Nassirpour ⁸⁰, H. Natal da Luz ¹²⁰, C. Nattrass ¹²⁸, S.R. Navarro ⁴⁴, K. Nayak ⁸⁵, R. Nayak ⁴⁸, T.K. Nayak ¹³⁹,

- S. Nazarenko ¹⁰⁶, R.A. Negrao De Oliveira ^{69,34}, L. Nellen ⁷⁰, S.V. Nesbo ³⁶, G. Neskovic ³⁹, F. Ng ¹²⁵, M. Nicassio ¹⁰⁴, J. Niedziela ^{140,34}, B.S. Nielsen ⁸⁸, S. Nikolaev ⁸⁷, S. Nikulin ⁸⁷, V. Nikulin ⁹⁶, F. Noferini ^{10,53}, P. Nomokonov ⁷⁵, G. Nooren ⁶³, J.C.C. Noris ⁴⁴, J. Norman ⁷⁸, A. Nyanin ⁸⁷, J. Nystrand ²², H. Oh ¹⁴⁵, A. Ohlson ¹⁰², J. Oleniacz ¹⁴⁰, A.C. Oliveira Da Silva ¹²⁰, M.H. Oliver ¹⁴⁴, J. Onderwaater ¹⁰⁴, C. Oppedisano ⁵⁸, R. Orava ⁴³, M. Oravec ¹¹⁵, A. Ortiz Velasquez ⁷⁰, A. Oskarsson ⁸⁰, J. Otwinowski ¹¹⁷, K. Oyama ⁸¹, Y. Pachmayer ¹⁰², V. Pacik ⁸⁸, D. Pagano ¹³⁸, G. Paić ⁷⁰, P. Palni ⁶, J. Pan ¹⁴¹, A.K. Pandey ⁴⁸, S. Panebianco ¹³⁵, V. Papikyan ¹, P. Pareek ⁴⁹, J. Park ⁶⁰, J.E. Parkkila ¹²⁶, S. Parmar ⁹⁸, A. Passfeld ¹⁴², S.P. Pathak ¹²⁵, R.N. Patra ¹³⁹, B. Paul ⁵⁸, H. Pei ⁶, T. Peitzmann ⁶³, X. Peng ⁶, L.G. Pereira ⁷¹, H. Pereira Da Costa ¹³⁵, D. Peresunko ⁸⁷, E. Perez Lezama ⁶⁹, V. Peskov ⁶⁹, Y. Pestov ⁴, V. Petráček ³⁷, M. Petrovici ⁴⁷, C. Petta ²⁸, R.P. Pezzi ⁷¹, S. Piano ⁵⁹, M. Pikna ¹⁴, P. Pillot ¹¹³, L.O.D.L. Pimentel ⁸⁸, O. Pinazza ^{53,34}, L. Pinsky ¹²⁵, S. Pisano ⁵¹, D.B. Piyarathna ¹²⁵, M. Płoskoń ⁷⁹, M. Planinic ⁹⁷, F. Pliquet ⁶⁹, J. Pluta ¹⁴⁰, S. Pochybova ¹⁴³, P.L.M. Podesta-Lerma ¹¹⁹, M.G. Poghosyan ⁹⁴, B. Polichtchouk ⁹⁰, N. Poljak ⁹⁷, W. Poonsawat ¹¹⁴, A. Pop ⁴⁷, H. Poppenborg ¹⁴², S. Porteboeuf-Houssais ¹³², V. Pozdniakov ⁷⁵, S.K. Prasad ³, R. Preghenella ⁵³, F. Prino ⁵⁸, C.A. Pruneau ¹⁴¹, I. Pshenichnov ⁶², M. Puccio ²⁶, V. Punin ¹⁰⁶, J. Putschke ¹⁴¹, S. Raha ³, S. Rajput ⁹⁹, J. Rak ¹²⁶, A. Rakotozafindrabe ¹³⁵, L. Ramello ³², F. Rami ¹³⁴, R. Raniwala ¹⁰⁰, S. Raniwala ¹⁰⁰, S.S. Räsänen ⁴³, B.T. Rascanu ⁶⁹, V. Ratza ⁴², I. Ravasenga ³¹, K.F. Read ^{128,94}, K. Redlich ^{84,v}, A. Rehman ²², P. Reichelt ⁶⁹, F. Reidt ³⁴, X. Ren ⁶, R. Renfordt ⁶⁹, A. Reshetin ⁶², J.-P. Revol ¹⁰, K. Reygers ¹⁰², V. Riabov ⁹⁶, T. Richert ^{63,80}, M. Richter ²¹, P. Riedler ³⁴, W. Riegler ³⁴, F. Riggi ²⁸, C. Ristea ⁶⁸, S.P. Rode ⁴⁹, M. Rodríguez Cahuantzi ⁴⁴, K. Røed ²¹, R. Rogalev ⁹⁰, E. Rogochaya ⁷⁵, D. Rohr ³⁴, D. Röhrich ²², P.S. Rokita ¹⁴⁰, F. Ronchetti ⁵¹, E.D. Rosas ⁷⁰, K. Roslon ¹⁴⁰, P. Rosnet ¹³², A. Rossi ²⁹, A. Rotondi ¹³⁷, F. Roukoutakis ⁸³, C. Roy ¹³⁴, P. Roy ¹⁰⁷, O.V. Rueda ⁷⁰, R. Rui ²⁵, B. Rumyantsev ⁷⁵, A. Rustamov ⁸⁶, E. Ryabinkin ⁸⁷, Y. Ryabov ⁹⁶, A. Rybicki ¹¹⁷, S. Saarinen ⁴³, S. Sadhu ¹³⁹, S. Sadovsky ⁹⁰, K. Šafařík ³⁴, S.K. Saha ¹³⁹, B. Sahoo ⁴⁸, P. Sahoo ⁴⁹, R. Sahoo ⁴⁹, S. Sahoo ⁶⁶, P.K. Sahu ⁶⁶, J. Saini ¹³⁹, S. Sakai ¹³¹, M.A. Saleh ¹⁴¹, S. Sambyal ⁹⁹, V. Samsonov ^{96,91}, A. Sandoval ⁷², A. Sarkar ⁷³, D. Sarkar ¹³⁹, N. Sarkar ¹³⁹, P. Sarma ⁴¹, M.H.P. Sas ⁶³, E. Scapparone ⁵³, F. Scarlassara ²⁹, B. Schaefer ⁹⁴, H.S. Scheid ⁶⁹, C. Schiaua ⁴⁷, R. Schicker ¹⁰², C. Schmidt ¹⁰⁴, H.R. Schmidt ¹⁰¹, M.O. Schmidt ¹⁰², M. Schmidt ¹⁰¹, N.V. Schmidt ^{94,69}, J. Schukraft ³⁴, Y. Schutz ^{34,134}, K. Schwarz ¹⁰⁴, K. Schweda ¹⁰⁴, G. Scioli ²⁷, E. Scomparin ⁵⁸, M. Šefčík ³⁸, J.E. Seger ¹⁶, Y. Sekiguchi ¹³⁰, D. Sekihata ⁴⁵, I. Selyuzhenkov ^{104,91}, K. Senosi ⁷³, S. Senyukov ¹³⁴, E. Serradilla ⁷², P. Sett ⁴⁸, A. Sevcenco ⁶⁸, A. Shabanov ⁶², A. Shabetai ¹¹³, R. Shahoyan ³⁴, W. Shaikh ¹⁰⁷, A. Shangaraev ⁹⁰, A. Sharma ⁹⁸, A. Sharma ⁹⁹, M. Sharma ⁹⁹, N. Sharma ⁹⁸, A.I. Sheikh ¹³⁹, K. Shigaki ⁴⁵, M. Shimomura ⁸², S. Shirinkin ⁶⁴, Q. Shou ^{6,110}, K. Shtejer ²⁶, Y. Sibiriak ⁸⁷, S. Siddhanta ⁵⁴, K.M. Sielewicz ³⁴, T. Siemarczuk ⁸⁴, D. Silvermyr ⁸⁰, G. Simatovic ⁸⁹, G. Simonetti ^{34,103}, R. Singaraju ¹³⁹, R. Singh ⁸⁵, R. Singh ⁹⁹, V. Singhal ¹³⁹, T. Sinha ¹⁰⁷, B. Sitar ¹⁴, M. Sitta ³², T.B. Skaali ²¹, M. Slupecki ¹²⁶, N. Smirnov ¹⁴⁴, R.J.M. Snellings ⁶³, T.W. Snellman ¹²⁶, J. Song ¹⁸, F. Soramel ²⁹, S. Sorensen ¹²⁸, F. Sozzi ¹⁰⁴, I. Sputowska ¹¹⁷, J. Stachel ¹⁰², I. Stan ⁶⁸, P. Stankus ⁹⁴, E. Stenlund ⁸⁰, D. Stocco ¹¹³, M.M. Storetvedt ³⁶, P. Strmen ¹⁴, A.A.P. Suade ¹²⁰, T. Sugitate ⁴⁵, C. Suire ⁶¹, M. Suleymanov ¹⁵, M. Suljic ^{34,25}, R. Sultanov ⁶⁴, M. Šumbera ⁹³, S. Sumowidagdo ⁵⁰, K. Suzuki ¹¹², S. Swain ⁶⁶, A. Szabo ¹⁴, I. Szarka ¹⁴, U. Tabassam ¹⁵, J. Takahashi ¹²¹, G.J. Tambave ²², N. Tanaka ¹³¹, M. Tarhini ¹¹³, M. Tariq ¹⁷, M.G. Tarzila ⁴⁷, A. Tauro ³⁴, G. Tejeda Muñoz ⁴⁴, A. Telesca ³⁴, C. Terrevoli ²⁹, B. Teyssier ¹³³, D. Thakur ⁴⁹, S. Thakur ¹³⁹, D. Thomas ¹¹⁸, F. Thoresen ⁸⁸, R. Tieulent ¹³³, A. Tikhonov ⁶², A.R. Timmins ¹²⁵, A. Toia ⁶⁹, N. Topilskaya ⁶², M. Toppi ⁵¹, S.R. Torres ¹¹⁹, S. Tripathy ⁴⁹, S. Trogolo ²⁶, G. Trombetta ³³, L. Tropp ³⁸, V. Trubnikov ², W.H. Trzaska ¹²⁶, T.P. Trzcinski ¹⁴⁰, B.A. Trzeciak ⁶³, T. Tsuji ¹³⁰, A. Tumkin ¹⁰⁶, R. Turrisi ⁵⁶, T.S. Tveter ²¹, K. Ullaland ²², E.N. Umaka ¹²⁵, A. Uras ¹³³, G.L. Usai ²⁴, A. Utrobicic ⁹⁷, M. Vala ¹¹⁵, J.W. Van Hoorn ³⁴, M. van Leeuwen ⁶³, P. Vande Vyvre ³⁴, D. Varga ¹⁴³, A. Vargas ⁴⁴, M. Vargyas ¹²⁶, R. Varma ⁴⁸, M. Vasileiou ⁸³, A. Vasiliev ⁸⁷, A. Vauthier ⁷⁸, O. Vázquez Doce ^{103,116}, V. Vechernin ¹¹¹, A.M. Veen ⁶³, E. Vercellin ²⁶, S. Vergara Limón ⁴⁴, L. Vermunt ⁶³, R. Vernet ⁷, R. Vértesi ¹⁴³, L. Vickovic ³⁵, J. Viinikainen ¹²⁶, Z. Vilakazi ¹²⁹, O. Villalobos Baillie ¹⁰⁸, A. Villatoro Tello ⁴⁴, A. Vinogradov ⁸⁷, T. Virgili ³⁰, V. Vislavicius ^{88,80}, A. Vodopyanov ⁷⁵, M.A. Völkl ¹⁰¹, K. Voloshin ⁶⁴, S.A. Voloshin ¹⁴¹, G. Volpe ³³, B. von Haller ³⁴, I. Vorobyev ^{116,103}, D. Voscek ¹¹⁵, D. Vranic ^{104,34}, J. Vrláková ³⁸, B. Wagner ²², H. Wang ⁶³, M. Wang ⁶, Y. Watanabe ¹³¹, M. Weber ¹¹², S.G. Weber ¹⁰⁴, A. Wegryznek ³⁴, D.F. Weiser ¹⁰², S.C. Wenzel ³⁴, J.P. Wessels ¹⁴², U. Westerhoff ¹⁴², A.M. Whitehead ¹²⁴, J. Wiechula ⁶⁹, J. Wikne ²¹,

G. Wilk⁸⁴, J. Wilkinson⁵³, G.A. Willems^{142,34}, M.C.S. Williams⁵³, E. Willsher¹⁰⁸, B. Windelband¹⁰², W.E. Witt¹²⁸, R. Xu⁶, S. Yalcin⁷⁷, K. Yamakawa⁴⁵, S. Yano⁴⁵, Z. Yin⁶, H. Yokoyama^{78,131}, I.-K. Yoo¹⁸, J.H. Yoon⁶⁰, V. Yurchenko², V. Zaccolo⁵⁸, A. Zaman¹⁵, C. Zampolli³⁴, H.J.C. Zanolli¹²⁰, N. Zardoshti¹⁰⁸, A. Zarochentsev¹¹¹, P. Závada⁶⁷, N. Zaviyalov¹⁰⁶, H. Zbroszczyk¹⁴⁰, M. Zhalov⁹⁶, X. Zhang⁶, Y. Zhang⁶, Z. Zhang^{6,132}, C. Zhao²¹, V. Zherebchevskii¹¹¹, N. Zhigareva⁶⁴, D. Zhou⁶, Y. Zhou⁸⁸, Z. Zhou²², H. Zhu⁶, J. Zhu⁶, Y. Zhu⁶, A. Zichichi^{27,10}, M.B. Zimmermann³⁴, G. Zinovjev², J. Zmeskal¹¹², S. Zou⁶

¹ A.I. Alikhanyan National Science Laboratory (Yerevan Physics Institute) Foundation, Yerevan, Armenia² Bogolyubov Institute for Theoretical Physics, National Academy of Sciences of Ukraine, Kiev, Ukraine³ Bose Institute, Department of Physics and Centre for Astroparticle Physics and Space Science (CAPSS), Kolkata, India⁴ Budker Institute for Nuclear Physics, Novosibirsk, Russia⁵ California Polytechnic State University, San Luis Obispo, CA, United States⁶ Central China Normal University, Wuhan, China⁷ Centre de Calcul de l'IN2P3, Villeurbanne, Lyon, France⁸ Centro de Aplicaciones Tecnológicas y Desarrollo Nuclear (CEADEN), Havana, Cuba⁹ Centro de Investigación y de Estudios Avanzados (CINVESTAV), Mexico City and Mérida, Mexico¹⁰ Centro Fermi – Museo Storico della Fisica e Centro Studi e Ricerche 'Enrico Fermi', Rome, Italy¹¹ Chicago State University, Chicago, IL, United States¹² China Institute of Atomic Energy, Beijing, China¹³ Chonbuk National University, Jeonju, Republic of Korea¹⁴ Comenius University Bratislava, Faculty of Mathematics, Physics and Informatics, Bratislava, Slovakia¹⁵ COMSATS Institute of Information Technology (CIIT), Islamabad, Pakistan¹⁶ Creighton University, Omaha, NE, United States¹⁷ Department of Physics, Aligarh Muslim University, Aligarh, India¹⁸ Department of Physics, Pusan National University, Pusan, Republic of Korea¹⁹ Department of Physics, Sejong University, Seoul, Republic of Korea²⁰ Department of Physics, University of California, Berkeley, CA, United States²¹ Department of Physics, University of Oslo, Oslo, Norway²² Department of Physics and Technology, University of Bergen, Bergen, Norway²³ Dipartimento di Fisica dell'Università 'La Sapienza' and Sezione INFN, Rome, Italy²⁴ Dipartimento di Fisica dell'Università and Sezione INFN, Cagliari, Italy²⁵ Dipartimento di Fisica dell'Università and Sezione INFN, Trieste, Italy²⁶ Dipartimento di Fisica dell'Università and Sezione INFN, Turin, Italy²⁷ Dipartimento di Fisica e Astronomia dell'Università and Sezione INFN, Bologna, Italy²⁸ Dipartimento di Fisica e Astronomia dell'Università and Sezione INFN, Catania, Italy²⁹ Dipartimento di Fisica e Astronomia dell'Università and Sezione INFN, Padova, Italy³⁰ Dipartimento di Fisica 'E.R. Caianiello' dell'Università and Gruppo Collegato INFN, Salerno, Italy³¹ Dipartimento DISAT del Politecnico and Sezione INFN, Turin, Italy³² Dipartimento di Scienze e Innovazione Tecnologica dell'Università del Piemonte Orientale and INFN Sezione di Torino, Alessandria, Italy³³ Dipartimento Interateneo di Fisica 'M. Merlin' and Sezione INFN, Bari, Italy³⁴ European Organization for Nuclear Research (CERN), Geneva, Switzerland³⁵ Faculty of Electrical Engineering, Mechanical Engineering and Naval Architecture, University of Split, Split, Croatia³⁶ Faculty of Engineering and Science, Western Norway University of Applied Sciences, Bergen, Norway³⁷ Faculty of Nuclear Sciences and Physical Engineering, Czech Technical University in Prague, Prague, Czech Republic³⁸ Faculty of Science, P.J. Šafárik University, Košice, Slovakia³⁹ Frankfurt Institute for Advanced Studies, Johann Wolfgang Goethe-Universität Frankfurt, Frankfurt, Germany⁴⁰ Gangneung-Wonju National University, Gangneung, Republic of Korea⁴¹ Gauhati University, Department of Physics, Guwahati, India⁴² Helmholtz-Institut für Strahlen- und Kernphysik, Rheinische Friedrich-Wilhelms-Universität Bonn, Bonn, Germany⁴³ Helsinki Institute of Physics (HIP), Helsinki, Finland⁴⁴ High Energy Physics Group, Universidad Autónoma de Puebla, Puebla, Mexico⁴⁵ Hiroshima University, Hiroshima, Japan⁴⁶ Hochschule Worms, Zentrum für Technologietransfer und Telekommunikation (ZTT), Worms, Germany⁴⁷ Horia Hulubei National Institute of Physics and Nuclear Engineering, Bucharest, Romania⁴⁸ Indian Institute of Technology Bombay (IIT), Mumbai, India⁴⁹ Indian Institute of Technology Indore, Indore, India⁵⁰ Indonesian Institute of Sciences, Jakarta, Indonesia⁵¹ INFN, Laboratori Nazionali di Frascati, Frascati, Italy⁵² INFN, Sezione di Bari, Bari, Italy⁵³ INFN, Sezione di Bologna, Bologna, Italy⁵⁴ INFN, Sezione di Cagliari, Cagliari, Italy⁵⁵ INFN, Sezione di Catania, Catania, Italy⁵⁶ INFN, Sezione di Padova, Padova, Italy⁵⁷ INFN, Sezione di Roma, Rome, Italy⁵⁸ INFN, Sezione di Torino, Turin, Italy⁵⁹ INFN, Sezione di Trieste, Trieste, Italy⁶⁰ Inha University, Incheon, Republic of Korea⁶¹ Institut de Physique Nucléaire d'Orsay (IPNO), Institut National de Physique Nucléaire et de Physique des Particules (IN2P3/CNRS), Université de Paris-Sud, Université Paris-Saclay, Orsay, France⁶² Institute for Nuclear Research, Academy of Sciences, Moscow, Russia⁶³ Institute for Subatomic Physics, Utrecht University/Nikhef, Utrecht, Netherlands⁶⁴ Institute for Theoretical and Experimental Physics, Moscow, Russia⁶⁵ Institute of Experimental Physics, Slovak Academy of Sciences, Košice, Slovakia⁶⁶ Institute of Physics, Homi Bhabha National Institute, Bhubaneswar, India⁶⁷ Institute of Physics of the Czech Academy of Sciences, Prague, Czech Republic⁶⁸ Institute of Space Science (ISS), Bucharest, Romania

- ⁶⁹ Institut für Kernphysik, Johann Wolfgang Goethe-Universität Frankfurt, Frankfurt, Germany
⁷⁰ Instituto de Ciencias Nucleares, Universidad Nacional Autónoma de México, Mexico City, Mexico
⁷¹ Instituto de Física, Universidade Federal do Rio Grande do Sul (UFRGS), Porto Alegre, Brazil
⁷² Instituto de Física, Universidad Nacional Autónoma de México, Mexico City, Mexico
⁷³ iThemba LABS, National Research Foundation, Somerset West, South Africa
⁷⁴ Johann-Wolfgang-Goethe Universität Frankfurt Institut für Informatik, Fachbereich Informatik und Mathematik, Frankfurt, Germany
⁷⁵ Joint Institute for Nuclear Research (JINR), Dubna, Russia
⁷⁶ Korea Institute of Science and Technology Information, Daejeon, Republic of Korea
⁷⁷ KTO Karatay University, Konya, Turkey
⁷⁸ Laboratoire de Physique Subatomique et de Cosmologie, Université Grenoble-Alpes, CNRS-IN2P3, Grenoble, France
⁷⁹ Lawrence Berkeley National Laboratory, Berkeley, CA, United States
⁸⁰ Lund University Department of Physics, Division of Particle Physics, Lund, Sweden
⁸¹ Nagasaki Institute of Applied Science, Nagasaki, Japan
⁸² Nara Women's University (NWU), Nara, Japan
⁸³ National and Kapodistrian University of Athens, School of Science, Department of Physics, Athens, Greece
⁸⁴ National Centre for Nuclear Research, Warsaw, Poland
⁸⁵ National Institute of Science Education and Research, Homi Bhabha National Institute, Jatni, India
⁸⁶ National Nuclear Research Center, Baku, Azerbaijan
⁸⁷ National Research Centre Kurchatov Institute, Moscow, Russia
⁸⁸ Niels Bohr Institute, University of Copenhagen, Copenhagen, Denmark
⁸⁹ Nikhef, National institute for subatomic physics, Amsterdam, Netherlands
⁹⁰ NRC Kurchatov Institute IHEP, Protvino, Russia
⁹¹ NRU Moscow Engineering Physics Institute, Moscow, Russia
⁹² Nuclear Physics Group, STFC Daresbury Laboratory, Daresbury, United Kingdom
⁹³ Nuclear Physics Institute of the Czech Academy of Sciences, Řež u Prahy, Czech Republic
⁹⁴ Oak Ridge National Laboratory, Oak Ridge, TN, United States
⁹⁵ Ohio State University, Columbus, OH, United States
⁹⁶ Petersburg Nuclear Physics Institute, Gatchina, Russia
⁹⁷ Physics department, Faculty of science, University of Zagreb, Zagreb, Croatia
⁹⁸ Physics Department, Panjab University, Chandigarh, India
⁹⁹ Physics Department, University of Jammu, Jammu, India
¹⁰⁰ Physics Department, University of Rajasthan, Jaipur, India
¹⁰¹ Physikalisches Institut, Eberhard-Karls-Universität Tübingen, Tübingen, Germany
¹⁰² Physikalisches Institut, Ruprecht-Karls-Universität Heidelberg, Heidelberg, Germany
¹⁰³ Physik Department, Technische Universität München, Munich, Germany
¹⁰⁴ Research Division and ExtreMe Matter Institute EMMI, GSI Helmholtzzentrum für Schwerionenforschung GmbH, Darmstadt, Germany
¹⁰⁵ Rudjer Bošković Institute, Zagreb, Croatia
¹⁰⁶ Russian Federal Nuclear Center (VNIIEF), Sarov, Russia
¹⁰⁷ Saha Institute of Nuclear Physics, Homi Bhabha National Institute, Kolkata, India
¹⁰⁸ School of Physics and Astronomy, University of Birmingham, Birmingham, United Kingdom
¹⁰⁹ Sección Física, Departamento de Ciencias, Pontificia Universidad Católica del Perú, Lima, Peru
¹¹⁰ Shanghai Institute of Applied Physics, Shanghai, China
¹¹¹ St. Petersburg State University, St. Petersburg, Russia
¹¹² Stefan Meyer Institut für Subatomare Physik (SMI), Vienna, Austria
¹¹³ SUBATECH, IMT Atlantique, Université de Nantes, CNRS-IN2P3, Nantes, France
¹¹⁴ Suranaree University of Technology, Nakhon Ratchasima, Thailand
¹¹⁵ Technical University of Košice, Košice, Slovakia
¹¹⁶ Technische Universität München, Excellence Cluster 'Universe', Munich, Germany
¹¹⁷ The Henryk Niewodniczanski Institute of Nuclear Physics, Polish Academy of Sciences, Cracow, Poland
¹¹⁸ The University of Texas at Austin, Austin, TX, United States
¹¹⁹ Universidad Autónoma de Sinaloa, Culiacán, Mexico
¹²⁰ Universidade de São Paulo (USP), São Paulo, Brazil
¹²¹ Universidade Estadual de Campinas (UNICAMP), Campinas, Brazil
¹²² Universidade Federal do ABC, Santo André, Brazil
¹²³ University College of Southeast Norway, Tønsberg, Norway
¹²⁴ University of Cape Town, Cape Town, South Africa
¹²⁵ University of Houston, Houston, TX, United States
¹²⁶ University of Jyväskylä, Jyväskylä, Finland
¹²⁷ University of Liverpool, Liverpool, United Kingdom
¹²⁸ University of Tennessee, Knoxville, TN, United States
¹²⁹ University of the Witwatersrand, Johannesburg, South Africa
¹³⁰ University of Tokyo, Tokyo, Japan
¹³¹ University of Tsukuba, Tsukuba, Japan
¹³² Université Clermont Auvergne, CNRS/IN2P3, LPC, Clermont-Ferrand, France
¹³³ Université de Lyon, Université Lyon 1, CNRS/IN2P3, IPN-Lyon, Villeurbanne, Lyon, France
¹³⁴ Université de Strasbourg, CNRS, IPHC UMR 7178, F-67000 Strasbourg, France, Strasbourg, France
¹³⁵ Université Paris-Saclay, Centre d'Études de Saclay (CEA), IRFU, Department de Physique Nucléaire (DPhN), Saclay, France
¹³⁶ Università degli Studi di Foggia, Foggia, Italy
¹³⁷ Università degli Studi di Pavia, Pavia, Italy
¹³⁸ Università di Brescia, Brescia, Italy
¹³⁹ Variable Energy Cyclotron Centre, Homi Bhabha National Institute, Kolkata, India
¹⁴⁰ Warsaw University of Technology, Warsaw, Poland
¹⁴¹ Wayne State University, Detroit, MI, United States
¹⁴² Westfälische Wilhelms-Universität Münster, Institut für Kernphysik, Münster, Germany
¹⁴³ Wigner Research Centre for Physics, Hungarian Academy of Sciences, Budapest, Hungary
¹⁴⁴ Yale University, New Haven, CT, United States
¹⁴⁵ Yonsei University, Seoul, Republic of Korea

ⁱ Deceased.

ⁱⁱ Dipartimento DET del Politecnico di Torino, Turin, Italy.

ⁱⁱⁱ M.V. Lomonosov Moscow State University, D.V. Skobeltsyn Institute of Nuclear, Physics, Moscow, Russia.

^{iv} Department of Applied Physics, Aligarh Muslim University, Aligarh, India.

^v Institute of Theoretical Physics, University of Wroclaw, Poland.