

Benchmark Continuum Limit Results for Spectroscopy with Stabilized Wilson Fermions

F. Cuteri,^a A. Francis,^b P. Fritsch,^c G. Pederiva,^{d,*} A. Rago,^e A. Shindler,^f A. Walker-Loud^g and S. Zafeiropoulos^h

^a*Institut für Theoretische Physik, Goethe Universität, Max-von-Laue-Str. 1, 60438 Frankfurt, Germany*

^b*Institute of Physics, National Yang Ming Chiao Tung University, 30010 Hsinchu, Taiwan*

^c*School of Mathematics, Trinity College Dublin, Dublin 2, Ireland*

^d*Jülich Supercomputing Centre, Forschungszentrum Jülich GmbH, 52428 Jülich, Germany*

^e*CP³-Origins, University of Southern Denmark, 5230 Odense, Denmark*

^f*FRIB & Physics Department, Michigan State University, East Lansing, MI 48824, USA*

^g*Nuclear Science Division, Lawrence Berkeley National Laboratory, Berkeley, CA 94720, USA*

^h*Aix Marseille Univ, Université de Toulon, CNRS, CPT, Marseille, France.*

E-mail: g.pederiva@fz-juelich.de

The OpenLat initiative presents its results of lattice QCD simulations using Stabilized Wilson Fermions (SWF) using 2+1 quark flavors. Focusing on the SU(3) flavor symmetric point $m_\pi = m_K = 412$ MeV, four different lattice spacings ($a = 0.064, 0.077, 0.094, 0.12$ fm) are used to perform the continuum limit to study cutoff effects.

We present results on light hadron masses; for the determination we use a Bayesian analysis framework with constraints and model averaging to minimize the bias in the analysis.

*The 39th International Symposium on Lattice Field Theory (Lattice2022),
8-13 August, 2022
Bonn, Germany*

*Speaker

1. Introduction

In recent years there has been a growing interest in Stabilized Wilson Fermions (SWF), with the OpenLat initiative playing a crucial role in their development. The SWF is a set of both algorithmic and analytical improvements over standard Wilson-Clover fermions designed to overcome some of their issues, for example SWF allow for simulations at coarse lattice spacing even at small pion masses [1]. The SWF package includes the use of the Stochastic Molecular Dynamics (SMD) algorithm instead of the HMC and the exponential clover action among other things.

The study of SWF has been initiated in the context of master-field simulations [1] and since then more simulations where new encouraging results have been produced [2, 3]. The open lattice initiative (OpenLat), on the other hand, was founded [4] with the goal of generating state-of-the-art QCD ensembles using SWF and share them according to the principles of the open science philosophy with all the LQCD community.

One of the first results that is planned for the newly generated ensembles is that of the light hadron spectrum as benchmark and in order to assess the discretization effects. The spectrum is computed using a Bayesian analysis framework and model averaging, the aim is to reduce the bias introduced from the analysis and establish solid benchmarks.

2. OpenLat Ensembles

The ensemble generation follows a three stages approach, at the end of which a reference publication will be released [5]. With this publication all configurations and metadata are made openly available. The metadata consists of all the observables used for validation during the generation. We also want to include a study of the light hadron spectrum to be used as reference and benchmark for the discretization effects of the SWF ensembles. In figure 1 the parameter space for the ensemble that are being produced and planned is reported, together with their current status.




Figure 1: Overview of the OpenLat ensembles presented here as parametrized by the lattice spacing a and $m_\pi L$. The height of the bars in the histograms represents the number of independent configurations that have been generated. Shaded boxes are ensembles in the tuning state [5, 6]

For this study of hadron spectroscopy, the ensembles collected in Table 1 were used¹.

Ensemble	$L^3 \times T$	t_0/a^2	a [fm]	$N_{cf\ g}$	N_{srcs}
a12m412	$24^3 \times 96$	1.48548(64)	0.12	600	100
a094m412S	$24^3 \times 96$	2.4482(50)	0.094	500	100
a094m412	$32^3 \times 96$	2.43979(89)	0.094	1200	100
a077m412	$48^3 \times 96$	3.6198(30)	0.077	900	100
a064m412	$48^3 \times 96$	5.2588(46)	0.064	1000	100

Table 1: SWF Ensembles used for the analysis. We note that the ensemble a094m412S is labelled differently as it represents a smaller volume compared to the others. $N_{cf\ g}$ represents the number of gauge field configurations analyzed, but not all of them are to be considered independent. N_{srcs} is the number of stochastic sources used to compute the averaged hadron correlators.

3. A Bayesian Analysis Framework for Hadron Masses

One issue with multi-state fitting for hadron correlators is the numerical stability of the algorithm coming from the non-linearity of the fit function. This makes a direct fitting of the hadron correlator data with a sum of exponential not feasible in practice without some redefined method. Furthermore, a naive approach necessarily introduces an arbitrary cut in the number of states to be fitted, which biases the results. In our determination of the hadron spectrum we use an unbiased method based on constrained Bayesian fitting and model averaging to minimize the bias in the results.

3.1 Bayesian Fitting with Constraints

A solution for improving the stability of multi-state fits can be the introduction of educated initial guesses for the fit parameters, as first introduced by Lepage et al. in [7] for the case of correlators from LQCD. These initial values, or *priors*, introduce a bias in the end results of the fit, which needs to be accounted for. In our analysis we computed the correlators in two different ways, using Gaussian smearing only at the source (*PS*) and both at the source and the sink (*SS*). The functional form of the function to be fitted is then:

$$C(t, Z_{P,n}, Z_{S,n}, E_n) = \sum_{n=0}^{n=N_s} Z_{S,n}^2 e^{-E_n t} + \sum_{n=0}^{n=N_s} Z_{S,n} Z_{P,n} e^{-E_n t}, \quad (1)$$

Where N_s is the number of states to be fitted, $Z_{P/S}$ are the amplitudes for the point and smeared sources and E_n are the energy levels. The fit is constrained by the introduction of priors for the fit

¹More details about the generation process can be found in the proceeding of this conference for the talk by A. Francis [5]

parameters and their uncertainties chosen as follows:

$$\begin{aligned}
\tilde{Z}_{P,0} &\rightarrow \text{fixed from data} & \tilde{Z}_{P,i} &= \tilde{Z}_{P,0} \text{ for } i = 1 \dots N_s & (2) \\
\tilde{Z}_{S,0} &\rightarrow \text{fixed from data} & \tilde{Z}_{S,1} &= \tilde{Z}_{S,0}; \tilde{Z}_{S,i} = \frac{\tilde{Z}_{S,0}}{2} \text{ for } i = 2 \dots N_s \\
\sigma_{\tilde{Z}_{P,0}} &\rightarrow \text{fixed from data} & \sigma_{\tilde{Z}_{P,i}} &= 2\sigma_{\tilde{Z}_{P,0}} \text{ for } i = 1 \dots N_s \\
\sigma_{\tilde{Z}_{S,0}} &\rightarrow \text{fixed from data} & \sigma_{\tilde{Z}_{S,1}} &= 2\sigma_{\tilde{Z}_{S,0}}; \sigma_{\tilde{Z}_{S,i}} = \sigma_{\tilde{Z}_{S,0}} \text{ for } i = 2 \dots N_s.
\end{aligned}$$

This choice of priors for the amplitudes $\tilde{Z}_{P,n}$ and $\tilde{Z}_{S,n}$ are motivated by the expectation that smearing should reduce the overlap with higher excited states, thus $\tilde{Z}_{S,n}$ is considered to be suppressed for high n . The ground state values are fixed from data, see figure 2. For the energies, on the other hand, we fix the prior \tilde{E}_0 and its associated $\sigma_{\tilde{E}_0}$ by looking at the effective mass plot. All the other energy priors are set to be ordered in steps of $2m_\pi$ above the ground state, with equal $\sigma_{\tilde{E}_n}$, but fixing the log of the prior, to enforce the ordering of the energies.

$$\begin{aligned}
\tilde{E}_0 &\rightarrow \text{fixed from data} & \ln(\tilde{E}_i - \tilde{E}_{i-1}) &= \ln(2m_\pi) \text{ for } i = 1 \dots N_s & (3) \\
\sigma_{\tilde{E}_0} &\rightarrow \text{fixed from data} & \sigma_{\tilde{E}_i} &= \sigma_{\tilde{E}_0} \text{ for } i = 1 \dots N_s.
\end{aligned}$$

The values for $\sigma_{\tilde{E}_0}$, $\sigma_{\tilde{Z}_{P,0}}$ and $\sigma_{\tilde{Z}_{S,0}}$ are chosen to be 10 times the expected value from the data (the shaded regions in figure 2).




Figure 2: Amplitudes for the point-smear (left) and smear-smear (right) correlators and the respective prior choices. The data is from the $a094m412$ ensemble, with 100 stochastic sources per configuration.

The χ^2 function to minimize is modified to include the contributions from the priors by defining the augmented χ^2 :

$$\chi^2 \rightarrow \chi_{\text{aug}}^2 = \chi^2 + \chi_{\text{prior}}^2 \quad \text{with} \quad \chi_{\text{prior}}^2 = \sum_{N_s} \frac{Z_{P,n} - \tilde{Z}_{P,n}}{\sigma_{\tilde{Z}_{P,n}}^2} + \sum_{N_s} \frac{Z_{S,n} - \tilde{Z}_{S,n}}{\sigma_{\tilde{Z}_{S,n}}^2} + \sum_{N_s} \frac{E_n - \tilde{E}_n}{\sigma_{\tilde{E}_n}^2} \quad (4)$$

3.2 Bayesian Model Averaging

To further determine an unbiased value for the mass of the hadrons, we use a Bayesian model averaging procedure based on the Akaike Information Criterion (AIC) that has been recently proposed in [8] for the case of lattice correlator data with data windowing selection. This procedure further eliminates the need to “manually” check for convergence in the constrained fitting procedure as the number of states increases. In turn, it defines a weight for every fit result, parametrized by the pair t_{min}, N_s that is used to determine the weighted average of the fits.

Given a fit result with parameters $M = A_1, \dots, A_n, E_1, \dots, E_n$ the AIC is:

$$\text{AIC}_M = -2 \log(\text{pr}(M)) + \chi_{\text{aug}}^2 + 2k + 2n \quad (5)$$

where $\text{pr}(M)$ is the model likelihood function of the model, k is the number of fit parameters, including priors, and n is the number of excluded points from the fit, i.e. it depends on changes of t_{min} . The minimum value of the AIC among all models is used to determine the relative likelihood of a given model M_i and some data D as:

$$\text{pr}(M_i|D) = \frac{\exp(\text{AIC}_{\text{min}} - \text{AIC}_i)}{\sum_j \exp(\text{AIC}_{\text{min}} - \text{AIC}_j)} \quad (6)$$

where the denominator is just to fix the normalization to allow a probabilistic interpretation of the quantity. The relative likelihood is then used as weight for computing the weighted sum of a fit parameter $\rho \in M$. The unbiased estimator for the parameter is then its weighted average and its uncertainty is given by:

$$\sigma_\rho = \sum_i \sigma_{\rho,i} \text{pr}(M_i|D) + \sum_i \langle \rho \rangle_i^2 \text{pr}(M_i|D) - \left(\sum_i \langle \rho \rangle_i \text{pr}(M_i|D) \right)^2 \quad (7)$$

There are a few considerations to make regarding this procedure. The first and most important is that it requires very little input, in particular it requires making an educated guess for the priors and attaching to their values a reasonably large uncertainty such that they do not become dominant in the fit; this greatly reduces the bias of the fit. Another interesting feature that can be immediately inferred from Equation (5) is that this method favors models that are simpler, i.e. fit functions with fewer number of states. This sort of built-in Occam's razor is very useful to gain intuition on how many states can be determined from the given set of data a posteriori. At the same time, Equation (5) tells us that this procedure favors models with larger fitting ranges, hence more data, which is a feature that we implicitly desire.

4. Numerical Results

The calculations of the hadron correlators were performed using the `lalibe` [9] software, which builds on top of `Chroma` [10]. The chosen smearing parameters are $N_{\text{smear}} = 32$ and $\sigma = 3.86$, where σ , following the notation used in [11], is $\sqrt{4\alpha N_{\text{smear}}}$ and represents a dimensionless smearing radius.

As outlined in the previous section, a scan over a range of t_{min} is made, while t_{max} is fixed to a value where it does not affect the fit results due to the low signal-to-noise. All the results for $N_S = 1, \dots, 5$ are then combined using model averaging. In Figure 3 we show our results for the `a094m412` ensemble for the proton as an example.

As discussed in [7], all fits should converge for large enough N_S given that there is enough information in the t range to fix the ground state. This is in fact the case: for low N_S we observe that the ground state estimate converges to the results coming from fits with more states for large enough t_{min} , where there is no information on the higher states. The height of the top panel in Figure 3 is set to be the width of the prior of the ground state energy. The model averaging result




Figure 3: Fit results for the proton mass for the a094m412 ensemble.

and error, shown as the gray band in the plot, is indeed compatible with all the converged fits and has, as expected, an uncertainty of the same order of the individual fits.

The lower panel of Figure 3 shows the weights of the different fits as they enter the average. One observes that our procedure based on the AIC selects the fits with low N_S and low t_{\min} . In the case of the proton shown above most of the contribution comes from the $N_S = 2$ fits.

The fitting procedure and model averaging is used on all ensembles for the pion, proton and Ω baryon, which are the only hadrons at the SU(3) flavor symmetric point. First, in Figure 4 we show $\phi_4 = 8t_0(m_K^2 + \frac{1}{2}m_\pi^2)$, to check the mass tunings of the ensembles, according to the imposed quality criteria the value of ϕ_4 should be within the gray band at the 1σ level, which is indeed the case².




Figure 4: Values of ϕ_4 for SWF at the SU(3) flavor-symmetric point.

In Figure 5, we show aggregated results for the proton and Ω mass as a function of a^2 . One notices that the given SWF ensembles show very small cutoff effects for hadronic observables. The

²The authors would like to point out that this figure is different from the one shown at during the conference presentation due to the finding of a mistake in one of the parameters.

continuum extrapolations for the masses of the proton and Ω are $m_p = 1175.0 \pm 6.4$ MeV and $m_\Omega = 1454.7 \pm 16.8$.




Figure 5: Continuum limit extrapolation for the proton (left) and Ω (right) masses for SWF at the SU(3) flavor-symmetric point.

5. Outlook for Studies with SWF

The results for the scaling properties of the hadron spectrum are very encouraging and show the validity and quality of the new SWF ensembles that have been generated by the OpenLat initiative. The analysis presented here is an ongoing work that will be repeated for every ensemble produced by the collaboration. The results will be included in the publication accompanying the public release of the ensembles in the future.

Acknowledgements

The authors acknowledge support from the HPC computing centres hpc-qcd (CERN), HPE Apollo Hawk (HLRS) under grant stabwf/44185, Cori (NERSC), Frontera (TACC), Piz Daint (CSCS), Occigen (CINES), Jean-Zay (IDRIS) and Irène-Joliot-Curie (TGCC) under projects (2020,2021,2022)-A0080511504, (2020,2021,2022)-A0080502271 by GENCI and PRACE project 2021250098. This work also used the DiRAC Extreme Scaling service at the University of Edinburgh, operated by the Edinburgh Parallel Computing Centre on behalf of the STFC DiRAC HPC Facility. DiRAC is part of the UK National e-Infrastructure. AS acknowledges funding support under the National Science Foundation grant PHY-2209185. AF acknowledges support under the Ministry of Science and Technology Taiwan grant 111-2112-M-A49-018-MY2.

References

- [1] A. Francis, P. Fritsch, M. Lüscher and A. Rago, *Master-field simulations of $O(a)$ -improved lattice QCD: Algorithms, stability and exactness*, *Comput. Phys. Commun.* **255** (2020) 107355 [1911.04533].

- [2] P. Fritzscht, J. Bulava, M. Cè, A. Francis, M. Lüscher and A. Rago, *Master-field simulations of QCD*, *PoS LATTICE2021* (2022) 465 [2111.11544].
- [3] M. Cè, M. Bruno, J. Bulava, A. Francis, P. Fritzscht, J.R. Green et al., *Approaching the master-field: Hadronic observables in large volumes*, *PoS LATTICE2021* (2022) 383 [2110.15375].
- [4] F. Cuteri, A. Francis, P. Fritzscht, G. Pederiva, A. Rago, A. Shindler et al., “OPEN LATtice initiative.” <https://openlat1.gitlab.io/>.
- [5] F. Cuteri, A. Francis, P. Fritzscht, G. Pederiva, A. Rago, A. Shindler et al., *Gauge generation and dissemination in openlat*, 2212.07314.
- [6] A.S. Francis, F. Cuteri, P. Fritzscht, G. Pederiva, A. Rago, A. Schindler et al., *Properties, ensembles and hadron spectra with Stabilised Wilson Fermions*, *PoS LATTICE2021* (2022) 118 [2201.03874].
- [7] G.P. Lepage, B. Clark, C.T.H. Davies, K. Hornbostel, P.B. Mackenzie, C. Morningstar et al., *Constrained curve fitting*, *Nucl. Phys. B Proc. Suppl.* **106** (2002) 12 [hep-lat/0110175].
- [8] W.I. Jay and E.T. Neil, *Bayesian model averaging for analysis of lattice field theory results*, *Phys. Rev. D* **103** (2021) 114502 [2008.01069].
- [9] A. Gambhir, D. Brantley, J. Chang, B. Hörz, H. Monge-Camacho, P. Vranas et al., “Lalibe (Lattice Livermore Berkeley).” <https://github.com/callat-qcd/lalibe>.
- [10] SciDAC, LHPC, UKQCD collaboration, *The Chroma software system for lattice QCD*, *Nucl. Phys. B Proc. Suppl.* **140** (2005) 832 [hep-lat/0409003].
- [11] S. Gusken, *A Study of smearing techniques for hadron correlation functions*, *Nucl. Phys. B Proc. Suppl.* **17** (1990) 361.