

Uniqueness and Lipschitz stability in electrical impedance tomography with finitely many electrodes

Bastian Harrach 

Institute for Mathematics, Goethe-University Frankfurt, Frankfurt am Main, Germany

E-mail: harrach@math.uni-frankfurt.de

Received 14 August 2018, revised 3 December 2018

Accepted for publication 7 December 2018

Published 3 January 2019



Abstract

For the linearized reconstruction problem in electrical impedance tomography with the complete electrode model, Lechleiter and Rieder (2008 *Inverse Problems* **24** 065009) have shown that a piecewise polynomial conductivity on a fixed partition is uniquely determined if enough electrodes are being used. We extend their result to the full non-linear case and show that measurements on a sufficiently high number of electrodes uniquely determine a conductivity in any finite-dimensional subset of piecewise-analytic functions. We also prove Lipschitz stability, and derive analogue results for the continuum model, where finitely many measurements determine a finite-dimensional Galerkin projection of the Neumann-to-Dirichlet operator on a boundary part.

Keywords: electrical impedance tomography (EIT), complete electrode model (CEM), uniqueness, Lipschitz stability, localized potentials, monotonicity

1. Introduction

We consider the inverse conductivity problem of determining the coefficient function σ in the elliptic partial differential equation

$$\nabla \cdot (\sigma \nabla u) = 0 \quad \text{in } \Omega \quad (1)$$

from knowledge of boundary measurements of u . The problem arises in electrical impedance tomography (EIT), or electrical resistivity tomography, which is a novel technique to image the conductivity distribution σ inside a subject Ω from electric voltage and current measurements on the subject's boundary $\partial\Omega$, see [1, 11, 13, 20, 21, 26, 49, 50, 71, 73, 75, 77, 80, 87, 89], and the references therein for a broad overview on the developments in EIT.



To model the boundary measurements we consider the continuum model, where we measure the local Neumann-to-Dirichlet (NtD) operator (on a boundary part $\Sigma \subseteq \partial\Omega$)

$$\Lambda(\sigma) : g \mapsto u|_{\Sigma}, \quad \text{where } u \text{ solves (1) with } \sigma \partial_{\nu} u|_{\partial\Omega} = \begin{cases} g & \text{on } \Sigma, \\ 0 & \text{else,} \end{cases}$$

and the more realistic complete electrode model (CEM) with electrodes $E_1, \dots, E_M \subseteq \partial\Omega$ all having the same contact impedance $z > 0$. In the CEM, we measure

$$R_M(\sigma) : (J_1, \dots, J_M) \mapsto (U_1, \dots, U_M),$$

where u solves (1) with

$$\begin{aligned} \sigma \partial_{\nu} u &= 0 && \text{on } \partial\Omega \setminus \bigcup_{m=1}^M E_m, \\ u + z\sigma \partial_{\nu} u &= \text{const.} =: U_m && \text{on } E_m, \quad m = 1, \dots, M, \\ \int_{E_m} \sigma \partial_{\nu} u|_{E_m} ds &= J_m && \text{on } E_m, \quad m = 1, \dots, M. \end{aligned}$$

The question whether full or local Neumann–Dirichlet-measurements uniquely determine the coefficient function σ has become famous under the name Calderón problem [23, 24], and has been intensively studied in the mathematical literature due to its practical relevance for EIT and many other related inverse coefficient problems, see [7, 9, 25, 28, 34, 35, 54, 55, 58, 61–63, 66–68, 76, 84].

In this work we will study the question whether σ can be uniquely and stably reconstructed from a finite number of electrode measurements. A natural discretization is to assume that σ is piecewise constant (or piecewise polynomial) on a given resolution or partition of Ω , so that σ will lie in an *a priori* known finite-dimensional subset \mathcal{F} of piecewise-analytic functions. Moreover, it seems natural to assume that upper and lower bounds on the conductivity are *a priori* known, i.e.

$$\sigma \in \mathcal{F}_{[a,b]} := \{\sigma \in \mathcal{F} : a \leq \sigma(x) \leq b \text{ for all } x \in \Omega\}.$$

Our main result for the continuum model is that a (sufficiently high dimensional) finite-dimensional Galerkin projection $P_{G_N} \Lambda(\sigma) P_{G_N}$ already uniquely determines σ and that Lipschitz stability holds

$$\exists c > 0 : c \|\sigma_1 - \sigma_2\| \leq \|P_{G_N} (\Lambda(\sigma_1) - \Lambda(\sigma_2)) P_{G_N}\|$$

see theorem 2.4.

Under the additional assumption that σ is an *a priori* known smooth function close to the boundary, we then turn to the CEM. We show that a (sufficiently large) finite number of electrodes M suffices to uniquely determine σ with Lipschitz stability

$$\exists c > 0 : c \|\sigma_1 - \sigma_2\| \leq \|R_M(\sigma_1) - R_M(\sigma_2)\|,$$

see theorem 3.1. This shows that the discretized EIT problem is uniquely and stably solvable if enough electrodes are being used, which may be relevant for practical implementations of EIT reconstruction algorithms.

Note that our results are non-constructive, we do not have a practically useful estimate of the Lipschitz constant or the required number of electrodes yet. Also note, that the necessary number of electrodes and the stability constant $c > 0$ depend on the ansatz set $\mathcal{F}_{[a,b]}$. Due to the intrinsic ill-posedness of the non-discretized EIT problem, we can naturally expect that

a larger set $\mathcal{F}_{[a,b]}$ will lead to worse stability constants and a higher required number of electrodes, with $c \rightarrow 0$ and $M \rightarrow \infty$ when $\dim(\text{span } \mathcal{F}) \rightarrow \infty$.

Let us give some more references on related results and the origins of our approach. A recent preprint of Alberti and Santacesaria [2] uses complex geometrical optics solutions to show that (in the continuum model) there exists a finite number of boundary voltages, so that the knowledge of the corresponding boundary currents uniquely determines the conductivity σ and that Lipschitz stability holds. Their result holds in dimension $d \geq 3$ with measurements on the full boundary $\partial\Omega$, σ is assumed to be identically one close to $\partial\Omega$, bounded by *a priori* known constants, and $\frac{\Delta\sqrt{\sigma}}{\sqrt{\sigma}}$ has to belong to an *a priori* known finite-dimensional subspace of L^∞ . Our result in this work requires less restrictive assumptions as we can treat any dimension $d \geq 2$, partial boundary data, and the CEM. But, on the other hand, we require the assumption of piecewise-analyticity which is more restrictive than the assumptions in [2].

For the linearized EIT problem (both, in the continuum model, and with the CEM), Lechleiter and Rieder [70] have shown that a piecewise polynomial conductivity on a fixed partition is uniquely determined if enough electrodes are being used. The main tool in [70] is the theory of localized potentials developed by the author [32] and the convergence of CEM-solutions to solutions of the continuum model shown by Hyvönen, Lechleiter and Hakula [51, 69]. Our result uses similar tools and first treats the non-linear EIT problem with the continuum model using localized potentials [32, 46] and monotonicity estimates between the non-linearized and the linearized problem from Ikehata, Kang, Seo and Sheen [53, 59]. Then we extend the results to the CEM using recent results on the approximation of the continuum model by the CEM from Hyvönen, Garde and Staboulis [30, 52].

The idea of using monotonicity estimates and localized potentials techniques has led to a number of results for inverse coefficient problems [8, 12, 22, 33, 36, 37, 39, 40, 44–46, 48], and several recent works build practical reconstruction methods on monotonicity properties [29–31, 38, 42, 43, 47, 72, 83, 85, 86, 88, 92]. Together with the recent preprints [41, 79], the present work shows that this idea can also be used to obtain Lipschitz stability estimates, which are usually derived from technically more challenging approaches involving Carleman estimates or quantitative unique continuation, see [3–6, 10, 14–19, 27, 56, 57, 60, 64, 65, 74, 78, 81, 90, 91].

The work is organized as follows. In section 2 we treat the continuum model, and show that the NtD operator or a (sufficiently high dimensional) finite-dimensional Galerkin projection uniquely determine the conductivity with Lipschitz stability. We formulate our main results for the continuum model in theorems 2.3 and 2.4 in section 2.1, summarize some known results from the literature in section 2.2, and then prove the theorems in section 2.3. In section 3 we then treat the CEM. Again we first formulate a uniqueness and Lipschitz stability result in theorem 3.1 in section 3.1, then summarize known results from the literature in section 3.2, and finally prove the theorem in section 3.3.

2. Uniqueness and Lipschitz stability from continuous data

2.1. Setting and main results

Let $\Omega \subset \mathbb{R}^d$, $d \geq 2$ be a bounded domain with smooth boundary $\partial\Omega$ and outer normal vector ν . $L_+^\infty(\Omega)$ denotes the subspace of $L^\infty(\Omega)$ -functions with positive essential infima. $H_\diamond^1(\Omega)$ and $L_\diamond^2(\partial\Omega)$ denote the spaces of H^1 - and L^2 -functions with vanishing integral mean on $\partial\Omega$.

For $\sigma \in L_+^\infty(\Omega)$, and a relatively open boundary part $\Sigma \subseteq \partial\Omega$, the local NtD operator $\Lambda(\sigma)$ is defined by

$$\Lambda(\sigma) : L^2_\diamond(\Sigma) \rightarrow L^2_\diamond(\Sigma), \quad g \mapsto u_\sigma^g|_\Sigma,$$

where $u_\sigma^g \in H^1_\diamond(\Omega)$ is the unique solution of

$$\nabla \cdot (\sigma \nabla u_\sigma^g) = 0 \text{ in } \Omega, \quad \sigma \partial_\nu u_\sigma^g|_{\partial\Omega} = \begin{cases} g & \text{on } \Sigma, \\ 0 & \text{else.} \end{cases} \quad (2)$$

This is equivalent to the variational formulation that $u_\sigma^g \in H^1_\diamond(\Omega)$ solves

$$\int_\Omega \sigma \nabla u_\sigma^g \cdot \nabla w \, dx = \int_\Sigma g w|_\Sigma \, ds \quad \text{for all } w \in H^1_\diamond(\Omega). \quad (3)$$

It is well known and easily shown that $\Lambda(\sigma)$ is compact and self-adjoint.

We will consider conductivities that are *a priori* known to belong to a finite dimensional set of piecewise-analytic functions and that are bounded from above and below by *a priori* known constants. To that end, we first define piecewise-analyticity as in [46, definition 2.1]:

Definition 2.1.

- (a) A subset $\Gamma \subseteq \partial O$ of the boundary of an open set $O \subseteq \mathbb{R}^n$ is called a *smooth boundary piece* if it is a C^∞ -surface and O lies on one side of it, i.e. if for each $z \in \Gamma$ there exists a ball $B_\epsilon(z)$ and a function $\gamma \in C^\infty(\mathbb{R}^{n-1}, \mathbb{R})$ such that upon relabeling and reorienting

$$\begin{aligned} \Gamma &= \partial O \cap B_\epsilon(z) = \{x \in B_\epsilon(z) \mid x_n = \gamma(x_1, \dots, x_{n-1})\}, \\ O \cap B_\epsilon(z) &= \{x \in B_\epsilon(z) \mid x_n > \gamma(x_1, \dots, x_{n-1})\}. \end{aligned}$$

- (b) O is said to have *smooth boundary* if ∂O is a union of smooth boundary pieces. O is said to have *piecewise smooth boundary* if ∂O is a countable union of the closures of smooth boundary pieces.
- (c) A function $\kappa \in L^\infty(\Omega)$ is called *piecewise analytic* if there exist finitely many pairwise disjoint subdomains $O_1, \dots, O_M \subset \Omega$ with piecewise smooth boundaries, such that $\overline{\Omega} = \overline{O_1} \cup \dots \cup \overline{O_M}$, and $\kappa|_{O_m}$ has an extension which is (real-)analytic in a neighborhood of $\overline{O_m}$, $m = 1, \dots, M$.

Note that (to the knowledge of the author), it is not clear whether the sum of two piecewise-analytic functions is always piecewise-analytic, i.e. whether the set of piecewise-analytic functions is a vector space. However, this can be guaranteed with a slightly stronger definition of piecewise analyticity (see [67, lemma 1]). Moreover, finite-dimensional vector spaces of piecewise-analytic functions (or subsets thereof) naturally arise as parameter spaces for the inverse conductivity problem, e.g. when we fix a partition of the imaging domain Ω into a finite number of subdomains (e.g. triangles, pixels, or voxels) and the conductivity is assumed to be a polynomial of fixed maximal order on each of these subdomains. Therefore, we make the following definition:

Definition 2.2. A set $\mathcal{F} \subseteq L^\infty(\Omega)$ is called a *finite-dimensional subset of piecewise-analytic functions* if its linear span

$$\text{span } \mathcal{F} = \left\{ \sum_{j=1}^k \lambda_j f_j : k \in \mathbb{N}, \lambda_j \in \mathbb{R}, f_j \in \mathcal{F} \right\} \subseteq L^\infty(\Omega)$$

contains only piecewise-analytic functions and $\dim(\text{span } \mathcal{F}) < \infty$.

Given a finite-dimensional subset \mathcal{F} of piecewise analytic functions and two numbers $b > a > 0$, we denote the set

$$\mathcal{F}_{[a,b]} := \{\sigma \in \mathcal{F} : a \leq \sigma(x) \leq b \text{ for all } x \in \Omega\}.$$

Throughout this paper, the domain Ω , the finite-dimensional subset \mathcal{F} and the bounds $b > a > 0$ are fixed, and the constants in the Lipschitz stability results will depend on them.

Our first result shows Lipschitz stability for the inverse conductivity problem in $\mathcal{F}_{[a,b]}$ when the complete infinite-dimensional NtD-operator is measured.

Theorem 2.3. *There exists $c > 0$ such that*

$$\|\Lambda(\sigma_1) - \Lambda(\sigma_2)\|_{\mathcal{L}(L^2_\diamond(\Sigma))} \geq c \|\sigma_1 - \sigma_2\|_{L^\infty(\Omega)} \quad \text{for all } \sigma_1, \sigma_2 \in \mathcal{F}_{[a,b]}.$$

Proof. Theorem 2.3 will be proven in section 2.3. □

We then turn to the question whether $\sigma \in \mathcal{F}_{[a,b]}$ is already uniquely determined by finitely many boundary measurements in the continuum model. For a (finite- or infinite-dimensional) subspace $G \subseteq L^2_\diamond(\Sigma)$ we denote by

$$P_G : L^2_\diamond(\Sigma) \rightarrow G, \quad P_G g = \begin{cases} g & \text{if } g \in G, \\ 0 & \text{if } g \in G^\perp, \end{cases}$$

the orthogonal projection operator on G with respect to the L^2 -scalar product

$$\langle g, h \rangle := \int_\Sigma gh \, ds \quad \text{for all } g, h \in L^2(\Sigma). \quad (4)$$

Clearly, $P_G = P_G^*$, and if G is finite dimensional with a basis $G = \text{span}(g_1, \dots, g_n)$ then measurements of

$$\langle g_j, \Lambda(\sigma)g_k \rangle \quad j, k = 1, \dots, n$$

determine the Galerkin projection of the NtD operator $P_G \Lambda(\sigma) P_G$, so that this can be regarded as a model for finitely many voltage/current measurements in the continuum model.

Our next result shows that this uniquely determines $\sigma \in \mathcal{F}_{[a,b]}$ (with Lipschitz stability) if the space G is large enough.

Theorem 2.4. *For each sequence of subspaces*

$$G_1 \subseteq G_2 \subseteq \dots \subseteq L^2_\diamond(\Sigma) \quad \text{with} \quad \overline{\bigcup_{n \in \mathbb{N}} G_n} = L^2_\diamond(\Sigma)$$

there exists $N \in \mathbb{N}$, and $c > 0$ such that

$$\|P_{G_n}(\Lambda(\sigma_1) - \Lambda(\sigma_2))P_{G_n}\|_{\mathcal{L}(L^2_\diamond(\Sigma))} \geq c \|\sigma_1 - \sigma_2\|_{L^\infty(\Omega)}$$

for all $\sigma_1, \sigma_2 \in \mathcal{F}_{[a,b]}$, and all $n \geq N$.

In particular, this implies that for all $\sigma_1, \sigma_2 \in \mathcal{F}_{[a,b]}$ and all $n \geq N$

$$P_{G_n} \Lambda(\sigma_1) P_{G_n} = P_{G_n} \Lambda(\sigma_2) P_{G_n} \quad \text{if and only if} \quad \sigma_1 = \sigma_2.$$

Proof. Theorem 2.4 will be proven in section 2.3. □

2.2. Differentiability, monotonicity and localized potentials

In this subsection, we summarize some known results from the literature, that we will use to prove theorem 2.3 and 2.4. As defined in (4), $\langle \cdot, \cdot \rangle$ always denotes the $L^2(\Sigma)$ -scalar product, and $u_\sigma^g \in H_\diamond^1(\Omega)$ denotes the solution of (2) with conductivity $\sigma \in L_+^\infty(\Omega)$ and Neumann data $g \in L_\diamond^2(\Sigma)$ in the following.

Our first tool is that the NtD operator is continuously Fréchet differentiable with respect to the conductivity.

Lemma 2.5.

(a) *The mapping*

$$\Lambda : L_+^\infty(\Omega) \rightarrow \mathcal{L}(L_\diamond^2(\Sigma)), \quad \sigma \mapsto \Lambda(\sigma)$$

is Fréchet differentiable. Its derivative is given by

$$\Lambda'(\sigma) \in \mathcal{L}(L^\infty(\Omega), \mathcal{L}(L_\diamond^2(\Sigma))), \quad (\Lambda'(\sigma)\kappa)g = v|_\Sigma, \quad (5)$$

where $v \in H_\diamond^1(\Omega)$ solves

$$\int_\Omega \sigma \nabla v \cdot \nabla w \, dx = - \int_\Omega \kappa \nabla u_\sigma^g \cdot \nabla w \, dx \quad \text{for all } w \in H_\diamond^1(\Omega).$$

(b) *For all $\sigma \in L_+^\infty(\Omega)$ and $\kappa \in L^\infty(\Omega)$ the operator $\Lambda'(\sigma)\kappa \in \mathcal{L}(L_\diamond^2(\Sigma))$ is self-adjoint and compact, and it fulfills*

$$\langle (\Lambda'(\sigma)\kappa)g, h \rangle = \int_\Omega \sigma \nabla u_\sigma^h \cdot \nabla v \, dx = - \int_\Omega \kappa \nabla u_\sigma^g \cdot \nabla u_\sigma^h \, dx,$$

for all $g, h \in L_\diamond^2(\Sigma)$.

(c) *The mapping*

$$\Lambda' : L_+^\infty(\Omega) \rightarrow \mathcal{L}(L^\infty(\Omega), \mathcal{L}(L_\diamond^2(\Sigma))), \quad \sigma \mapsto \Lambda'(\sigma)$$

is continuous.

Proof. This follows from the variational formulation of the conductivity equation (3), see e.g. [70, section 2] or [30, appendix B]. \square

Our next tool is a monotonicity relation between the NtD-operator and its derivative that goes back to Ikehata, Kang, Seo, and Sheen [53, 59], and has been used in several other works, see the list of works on monotonicity-based methods cited in the introduction.

Lemma 2.6. *For all $\sigma_1, \sigma_2 \in L_+^\infty(\Omega)$ and $g \in L_\diamond^2(\Sigma)$, it holds that*

$$\begin{aligned} \langle (\Lambda'(\sigma_2)(\sigma_1 - \sigma_2))g, g \rangle &= \int_\Omega (\sigma_2 - \sigma_1) |\nabla u_{\sigma_2}^g|^2 \, dx \\ &\leq \langle g, (\Lambda(\sigma_1) - \Lambda(\sigma_2))g \rangle. \end{aligned} \quad (6)$$

Proof. See, e.g. [45, lemma 2.1]. \square

The energy terms $|\nabla u_\sigma^g|^2$ in the monotonicity estimate can be controlled using the technique of localized potentials [32]. Roughly speaking, the energy $|\nabla u_\sigma^g|^2$ can be made arbitrarily

large in a subset $D_1 \subseteq \Omega$ without making it large in another subset $D_2 \subseteq \Omega$ whenever D_1 can be reached from the boundary without passing D_2 .

To formulate this rigorously, we adopt the notation from [46, definitions 2.2 and 2.3] and denote by $\text{int } D$ the topological interior of a subset $D \subseteq \bar{\Omega}$, and by $\text{out}_\Sigma D$ its outer hull, i.e.

$$\text{out}_\Sigma D := \bar{\Omega} \setminus \bigcup \{U \subseteq \bar{\Omega} : U \text{ rel. open, } U \cap \Omega \text{ connected, } U \cap \Sigma \neq \emptyset\}.$$

With this notation, we have the following localized potentials result:

Lemma 2.7. *Let $\sigma \in L_+^\infty(\Omega)$ be piecewise analytic and let $D_1, D_2 \subseteq \bar{\Omega}$ be two measurable sets with*

$$\text{int } D_1 \not\subseteq \text{out}_\Sigma D_2.$$

Then there exists a sequence of currents $(g_m)_{m \in \mathbb{N}} \subset L_\diamond^2(\Sigma)$ such that the corresponding solutions $(u_\sigma^{g_m})_{m \in \mathbb{N}} \subset H_\diamond^1(\Omega)$ fulfill

$$\lim_{m \rightarrow \infty} \int_{D_1} |\nabla u_\sigma^{g_m}|^2 \, dx = \infty \quad \text{and} \quad \lim_{m \rightarrow \infty} \int_{D_2} |\nabla u_\sigma^{g_m}|^2 \, dx = 0.$$

Proof. ([46], theorem 3.6 and section 4.3). □

We will also need the following definiteness property of piecewise-analytic functions from [46]:

Lemma 2.8. *Let $0 \neq \kappa \in L^\infty(\Omega)$ be piecewise-analytic. Then there exist two sets*

$$D_1 = \text{int } D_1 \quad \text{and} \quad D_2 = \text{out}_\Sigma D_2$$

with

$$D_1 = \text{int } D_1 \not\subseteq \text{out}_\Sigma D_2 = D_2$$

and either

- (i) $\kappa|_{\Omega \setminus D_2} \geq 0$ and $\kappa|_{D_1} \in L_+^\infty(D_1)$, or
- (ii) $\kappa|_{\Omega \setminus D_2} \leq 0$ and $-\kappa|_{D_1} \in L_+^\infty(D_1)$.

Proof. ([46], theorem A.1, corollary A.2, and section 4.3). □

2.3. Proof of theorems 2.3 and 2.4

We can now prove theorems 2.3 and 2.4. For the sake of brevity, we write $\|\cdot\|$ for $\|\cdot\|_{\mathcal{L}(L_\diamond^2(\Sigma))}$, $\|\cdot\|_{L^\infty(\Omega)}$ and $\|\cdot\|_{L_\diamond^2(\Sigma)}$ throughout this subsection.

We follow the approach in [41] and first use the monotonicity relation in lemma 2.6 to bound the difference of the non-linear NtD operators by an expression containing their linearized counterparts.

Lemma 2.9. *For all $\sigma_1, \sigma_2 \in \mathcal{F}_{[a,b]}$ with $\sigma_1 \neq \sigma_2$,*

$$\frac{\|\Lambda(\sigma_1) - \Lambda(\sigma_2)\|}{\|\sigma_1 - \sigma_2\|} \geq \inf_{\substack{(\tau_1, \tau_2, \kappa) \\ \in \mathcal{F}_{[a,b]} \times \mathcal{F}_{[a,b]} \times \mathcal{K}}} \sup_{g \in L_\diamond^2(\Sigma), \|g\|=1} f(\tau_1, \tau_2, \kappa, g),$$

where $f : L_+^\infty(\Omega) \times L_+^\infty(\Omega) \times L^\infty(\Omega) \times L_\diamond^2(\Sigma) \rightarrow \mathbb{R}$ is defined by

$$f(\tau_1, \tau_2, \kappa, g) := \max \{ \langle (\Lambda'(\tau_1)\kappa)g, g \rangle, -\langle (\Lambda'(\tau_2)\kappa)g, g \rangle \},$$

and $\mathcal{K} := \{ \kappa \in \text{span } \mathcal{F} : \|\kappa\| = 1 \}$.

Proof. The NtD-operators are self-adjoint so that for all $\sigma_1, \sigma_2 \in L^\infty(\Omega)$

$$\|\Lambda(\sigma_1) - \Lambda(\sigma_2)\| = \sup_{g \in L_\diamond^2(\Sigma), \|g\|=1} |\langle g, (\Lambda(\sigma_1) - \Lambda(\sigma_2))g \rangle|.$$

Using the monotonicity inequality in lemma 2.6 also with interchanged roles of σ_1 and σ_2 , we obtain that for all $\sigma_1, \sigma_2 \in L_+^\infty(\Omega)$, $\sigma_1 \neq \sigma_2$, and all $g \in L_\diamond^2(\partial\Omega)$

$$\begin{aligned} & |\langle g, (\Lambda(\sigma_1) - \Lambda(\sigma_2))g \rangle| \\ &= \max \{ \langle g, (\Lambda(\sigma_2) - \Lambda(\sigma_1))g \rangle, \langle g, (\Lambda(\sigma_1) - \Lambda(\sigma_2))g \rangle \} \\ &\geq \max \{ \langle \Lambda'(\sigma_1)(\sigma_2 - \sigma_1)g, g \rangle, \langle \Lambda'(\sigma_2)(\sigma_1 - \sigma_2)g, g \rangle \} \\ &= \|\sigma_1 - \sigma_2\| \max \left\{ \left\langle \Lambda'(\sigma_1) \frac{\sigma_2 - \sigma_1}{\|\sigma_1 - \sigma_2\|} g, g \right\rangle, \left\langle \Lambda'(\sigma_2) \frac{\sigma_1 - \sigma_2}{\|\sigma_1 - \sigma_2\|} g, g \right\rangle \right\} \\ &= \|\sigma_1 - \sigma_2\| f(\sigma_1, \sigma_2, \frac{\sigma_2 - \sigma_1}{\|\sigma_1 - \sigma_2\|}, g). \end{aligned}$$

Hence,

$$\begin{aligned} \frac{\|\Lambda(\sigma_1) - \Lambda(\sigma_2)\|}{\|\sigma_1 - \sigma_2\|} &= \sup_{g \in L_\diamond^2(\Sigma), \|g\|=1} \frac{|\langle g, (\Lambda(\sigma_1) - \Lambda(\sigma_2))g \rangle|}{\|\sigma_1 - \sigma_2\|} \\ &\geq \sup_{g \in L_\diamond^2(\Sigma), \|g\|=1} f(\sigma_1, \sigma_2, \frac{\sigma_2 - \sigma_1}{\|\sigma_1 - \sigma_2\|}, g) \\ &\geq \inf_{\substack{(\tau_1, \tau_2, \kappa) \\ \in \mathcal{F}_{[a,b]} \times \mathcal{F}_{[a,b]} \times \mathcal{K}}} \sup_{g \in L_\diamond^2(\Sigma), \|g\|=1} f(\tau_1, \tau_2, \kappa, g). \end{aligned} \quad \square$$

Now we use a compactness argument to show that the expression in the lower bound in lemma 2.9 attains its minimum.

Lemma 2.10. *There exists $(\hat{\tau}_1, \hat{\tau}_2, \hat{\kappa}) \in \mathcal{F}_{[a,b]} \times \mathcal{F}_{[a,b]} \times \mathcal{K}$ so that*

$$\inf_{\substack{(\tau_1, \tau_2, \kappa) \\ \in \mathcal{F}_{[a,b]} \times \mathcal{F}_{[a,b]} \times \mathcal{K}}} \sup_{g \in L_\diamond^2(\Sigma), \|g\|=1} f(\tau_1, \tau_2, \kappa, g) = \sup_{g \in L_\diamond^2(\Sigma), \|g\|=1} f(\hat{\tau}_1, \hat{\tau}_2, \hat{\kappa}, g).$$

Proof. Since f is continuous by lemma 2.5, the function

$$(\tau_1, \tau_2, \kappa) \mapsto \sup_{g \in L_\diamond^2(\Sigma), \|g\|=1} f(\tau_1, \tau_2, \kappa, g)$$

is lower semicontinuous and thus attains its minimum over the compact set $\mathcal{F}_{[a,b]} \times \mathcal{F}_{[a,b]} \times \mathcal{K}$. \square

It remains to show that the minimum attained in lemma 2.10 must be positive. To show that we use the localized potentials from lemma 2.7.

Lemma 2.11. *Let $0 \neq \kappa \in L^\infty(\Omega)$ be piecewise-analytic. Then at least one of the following two properties holds true:*

(i) *For all piecewise analytic $\sigma \in L_+^\infty(\Omega)$ there exists $g \in L_\diamond^2(\partial\Omega)$ with*

$$-\langle (\Lambda'(\sigma)\kappa) g, g \rangle = \int_\Omega \kappa |\nabla u_\sigma^g|^2 dx > 0.$$

(ii) *For all piecewise analytic $\sigma \in L_+^\infty(\Omega)$ there exists $g \in L_\diamond^2(\partial\Omega)$ with*

$$-\langle (\Lambda'(\sigma)\kappa) g, g \rangle = \int_\Omega \kappa |\nabla u_\sigma^g|^2 dx < 0.$$

Hence, a fortiori,

$$\sup_{g \in L_\diamond^2(\Sigma), \|g\|=1} f(\tau_1, \tau_2, \kappa, g) > 0 \quad \text{for all } (\tau_1, \tau_2, \kappa) \in \mathcal{F}_{[a,b]} \times \mathcal{F}_{[a,b]} \times \mathcal{K}.$$

Proof. Using the definiteness property of piecewise analytic functions from lemma 2.8 we obtain two sets $D_1 = \text{int } D_1$ and $D_2 = \text{out}_\Sigma D_2$ with

$$D_1 = \text{int } D_1 \not\subseteq \text{out}_\Sigma D_2 = D_2$$

and either

- (i) $\kappa|_{\Omega \setminus D_2} \geq 0$ and $\kappa|_{D_1} \in L_+^\infty(D_1)$, or
- (ii) $\kappa|_{\Omega \setminus D_2} \leq 0$ and $-\kappa|_{D_1} \in L_+^\infty(D_1)$.

Let $(g_m)_{m \in \mathbb{N}} \subset L_\diamond^2(\Sigma)$ be the localized potentials sequence from lemma 2.7. Then, in case (i), we obtain

$$\begin{aligned} -\langle (\Lambda'(\sigma)\kappa) g_m, g_m \rangle &= \int_\Omega \kappa |\nabla u_\sigma^{g_m}|^2 dx \\ &= \int_{D_1} \kappa |\nabla u_\sigma^{g_m}|^2 dx + \int_{D_2} \kappa |\nabla u_\sigma^{g_m}|^2 dx + \int_{\Omega \setminus (D_1 \cup D_2)} \kappa |\nabla u_\sigma^{g_m}|^2 dx \\ &\geq \text{ess inf } \kappa|_{D_1} \int_{D_1} |\nabla u_\sigma^{g_m}|^2 dx - \|\kappa\|_{L^\infty(D_2)} \int_{D_2} |\nabla u_\sigma^{g_m}|^2 dx \rightarrow \infty, \end{aligned}$$

so that $\int_\Omega \kappa |\nabla u_\sigma^{g_m}|^2 dx > 0$ for sufficiently large $m \in \mathbb{N}$.

In case (ii) we obtain

$$\begin{aligned} -\langle (\Lambda'(\sigma)\kappa) g_m, g_m \rangle &= \int_\Omega \kappa |\nabla u_\sigma^{g_m}|^2 dx \\ &= \int_{D_1} \kappa |\nabla u_\sigma^{g_m}|^2 dx + \int_{D_2} \kappa |\nabla u_\sigma^{g_m}|^2 dx + \int_{\Omega \setminus (D_1 \cup D_2)} \kappa |\nabla u_\sigma^{g_m}|^2 dx \\ &\leq -\text{ess inf } (-\kappa)|_{D_1} \int_{D_1} |\nabla u_\sigma^{g_m}|^2 dx + \|\kappa\|_{L^\infty(D_2)} \int_{D_2} |\nabla u_\sigma^{g_m}|^2 dx \rightarrow -\infty, \end{aligned}$$

so that $\int_\Omega \kappa |\nabla u_\sigma^{g_m}|^2 dx < 0$ for sufficiently large $m \in \mathbb{N}$. □

Remark 2.12. It is known (see e.g. [45, corollary 3.5(b)]) that for all piecewise analytic σ , the Fréchet derivative $\Lambda'(\sigma)$ is injective on the set of piecewise analytic functions, i.e. $\Lambda'(\sigma)\kappa \neq 0$ for all piecewise analytic $0 \neq \kappa \in L^\infty(\Omega)$.

Since $\Lambda'(\sigma)\kappa$ is a compact self-adjoint operator, this means that $\Lambda'(\sigma)\kappa$ must possess either a positive or a negative eigenvalue. Lemma 2.11 can be interpreted in the sense, that for each $\kappa \neq 0$ this property is sign-uniform in σ , i.e. for each $\kappa \neq 0$, the operator $\Lambda'(\sigma)\kappa$ either possesses a positive eigenvalue for all σ , or it possesses a negative eigenvalue for all σ (or both properties are fulfilled).

With these preparations we can now show the theorems 2.3 and 2.4.

Proof of theorem 2.3. The assertion follows from lemmas 2.9–2.11 with

$$c := \sup_{g \in L^2_\Sigma(\Sigma), \|g\|=1} f(\hat{\tau}_1, \hat{\tau}_2, \hat{\kappa}, g) > 0.$$

□

Proof of theorem 2.4. Using that

$$\|P_{G_n}(\Lambda(\sigma_1) - \Lambda(\sigma_2))P_{G_n}\| = \sup_{g \in G_n, \|g\|=1} |\langle g, (\Lambda(\sigma_1) - \Lambda(\sigma_2))g \rangle|,$$

we obtain as in lemmas 2.9 and 2.10 that for all $n \in \mathbb{N}$, there exists $(\hat{\tau}_1^{(n)}, \hat{\tau}_2^{(n)}, \hat{\kappa}^{(n)}) \in \mathcal{F}_{[a,b]} \times \mathcal{F}_{[a,b]} \times \mathcal{K}$ so that

$$\frac{\|P_{G_n}(\Lambda(\sigma_1) - \Lambda(\sigma_2))P_{G_n}\|}{\|\sigma_1 - \sigma_2\|} \geq \sup_{g \in G_n, \|g\|=1} f(\hat{\tau}_1^{(n)}, \hat{\tau}_2^{(n)}, \hat{\kappa}^{(n)}, g). \quad (7)$$

The right hand side of (7) is monotonically increasing in $n \in \mathbb{N}$ since the spaces G_n are nested. Hence, the assertion of theorem 2.4 follows, if we can prove that there exists $n \in \mathbb{N}$ with

$$\sup_{g \in G_n, \|g\|=1} f(\tau_1, \tau_2, \kappa, g) > 0 \quad \text{for all } (\tau_1, \tau_2, \kappa) \in \mathcal{F}_{[a,b]} \times \mathcal{F}_{[a,b]} \times \mathcal{K}. \quad (8)$$

We argue by contradiction and assume that this is not the case. Then there exists a sequence $(\tau_1^{(n)}, \tau_2^{(n)}, \kappa^{(n)}) \in \mathcal{F}_{[a,b]} \times \mathcal{F}_{[a,b]} \times \mathcal{K}$ with

$$\sup_{g \in G_n, \|g\|=1} f(\tau_1^{(n)}, \tau_2^{(n)}, \kappa^{(n)}, g) \leq 0 \quad \text{for all } n \in \mathbb{N},$$

which also implies

$$\sup_{g \in G_m, \|g\|=1} f(\tau_1^{(n)}, \tau_2^{(n)}, \kappa^{(n)}, g) \leq 0 \quad \text{for all } n \in \mathbb{N}, n \geq m.$$

After passing to a subsequence if necessary, we can assume by compactness that the sequence $(\tau_1^{(n)}, \tau_2^{(n)}, \kappa^{(n)})$ converges to some element

$$(\hat{\tau}_1, \hat{\tau}_2, \hat{\kappa}) \in \mathcal{F}_{[a,b]} \times \mathcal{F}_{[a,b]} \times \mathcal{K}.$$

Since, for all $m \in \mathbb{N}$, the function

$$(\tau_1, \tau_2, \kappa) \mapsto \sup_{g \in G_m, \|g\|=1} f(\tau_1, \tau_2, \kappa, g)$$

is lower semicontinuous, it follows that

$$\sup_{g \in G_m, \|g\|=1} f(\hat{\tau}_1, \hat{\tau}_2, \hat{\kappa}, g) \leq 0 \quad \text{for all } m \in \mathbb{N}.$$

But, by continuity, this would imply

$$f(\hat{\tau}_1, \hat{\tau}_2, \hat{\kappa}, g) \leq 0 \quad \text{for all } g \in \overline{\bigcup_{m \in \mathbb{N}} G_m} = L^2_\diamond(\Sigma),$$

which contradicts lemma 2.11. This shows that (8) must be true for sufficiently large $n \in \mathbb{N}$ and thus theorem 2.4 is proven. \square

3. Uniqueness and Lipschitz stability from electrode measurements

3.1. Setting and main results

Now we consider the CEM. As before let $\sigma \in L^\infty_+(\Omega)$ denote the conductivity distribution in a smoothly bounded domain $\Omega \subseteq \mathbb{R}^d$, $d \geq 2$. We assume that $M \in \mathbb{N}$ open, connected, mutually disjoint electrodes $E_m \subseteq \partial\Omega$, $m = 1, \dots, M$, are attached to the boundary of Ω with all electrodes having the same contact impedance $z > 0$. When a current with strength $J_m \in \mathbb{R}$ is driven through the m -th electrode (with $\sum_{m=1}^M J_m = 0$), the resulting electrical potential $(u, U) \in H^1(\Omega) \times \mathbb{R}^M$ solves the following equations:

$$\nabla \cdot \sigma \nabla u = 0 \quad \text{in } \Omega, \quad (9)$$

$$\sigma \partial_\nu u = 0 \quad \text{on } \partial\Omega \setminus \bigcup_{m=1}^M E_m, \quad (10)$$

$$u + z\sigma \partial_\nu u = \text{const.} =: U_m \quad \text{on } E_m, \quad m = 1, \dots, M, \quad (11)$$

$$\int_{E_m} \sigma \partial_\nu u|_{E_m} ds = J_m \quad \text{on } E_m, \quad m = 1, \dots, M, \quad (12)$$

where $U = (U_1, \dots, U_M) \in \mathbb{R}^M$ is a vector containing the electric potentials on the electrodes E_1, \dots, E_M .

It can be shown that (9)–(12) possess a solution $(u, U) \in H^1(\Omega) \times \mathbb{R}^M$ and that the solution is unique under the additional gauge (or ground level) condition $U \in \mathbb{R}^M_\diamond$, where \mathbb{R}^M_\diamond is the subspace of vectors in \mathbb{R}^M with zero mean, see e.g. [82]. We can thus define the M -electrode current-to-potential operator

$$R_M(\sigma) : \mathbb{R}^M_\diamond \rightarrow \mathbb{R}^M_\diamond : I = (I_1, \dots, I_M) \mapsto U = (U_1, \dots, U_M),$$

where $(u, U) \in H^1(\Omega) \times \mathbb{R}^M_\diamond$ solves (9)–(12). Note also that (9)–(12) are equivalent to the variational formulation that $(u, U) \in H^1(\Omega) \times \mathbb{R}^M_\diamond$ solves

$$\int_\Omega \sigma \nabla u \cdot \nabla w dx + \sum_{m=1}^M \int_{E_m} \frac{1}{z} (u - U_m)(w - W_m) ds = \sum_{m=1}^M J_m W_m \quad (13)$$

for all $(w, W) \in H^1(\Omega) \times \mathbb{R}^M_\diamond$, see again, [82].

As in the previous section, we will consider conductivities that belong to a finite dimensional subset of piecewise-analytic functions. Additionally, in order to use results from [30] on the approximation properties of the CEM, we assume that the background conductivity is an *a priori* known smooth function in a fixed neighborhood O of the boundary $\partial\Omega$, i.e. we assume that \mathcal{F} is a finite dimensional subset of piecewise-analytic functions, so that there exists $\sigma_0 \in C^\infty(\bar{O})$ with $\sigma|_O = \sigma_0|_O$ for all $\sigma \in \mathcal{F}$. Together with the assumption of *a priori* known bounds, we assume (for $b > a > 0$)

$$\sigma \in \mathcal{F}_{[a,b]} := \{\sigma \in \mathcal{F} : a \leq \sigma(x) \leq b \text{ for all } x \in \Omega\}.$$

We will show that $R_M(\sigma)$ uniquely determines $\sigma \in \mathcal{F}_{[a,b]}$ (with Lipschitz stability) if, roughly speaking, enough electrodes are being used. To make this statement precise, assume that the number of electrodes is increased so that the electrode configurations fulfill the Hyvönen criteria [30, 52]:

(H1) For all electrode configurations

$$E_1^{(M)}, \dots, E_M^{(M)} \subseteq \partial\Omega$$

there exist open, connected, and mutually disjoint sets (called virtual extended electrodes) $\tilde{E}_m^{(M)}$, $m = 1, \dots, M$ with

$$E_m^{(M)} \subseteq \tilde{E}_m^{(M)}, \quad \bigcup_{m=1}^M \overline{\tilde{E}_m^{(M)}} = \partial\Omega,$$

so that

$$h_M := \max_{m=1, \dots, M} \left\{ \sup_{x, y \in \tilde{E}_m^{(M)}} \text{dist}(x, y) \right\} \rightarrow 0 \quad \text{for } M \rightarrow \infty,$$

$$\exists c_E > 0 : \min_{m=1, \dots, M} \frac{|E_m^{(M)}|}{|\tilde{E}_m^{(M)}|} \geq c_E \quad \text{for all } M \in \mathbb{N}.$$

(H2) The operators

$$Q_M : \mathbb{R}^M \rightarrow L^2(\partial\Omega), \quad (J_m)_{m=1}^M \mapsto \sum_{m=1}^M J_m \chi_{\tilde{E}_m^{(M)}}$$

$$P_M : L^2(\partial\Omega) \rightarrow \mathbb{R}^M, \quad g \mapsto \left(\frac{1}{|E_m^{(M)}|} \int_{E_m^{(M)}} g \, ds \right)_{m=1}^M$$

fulfill that

$$\exists C_E > 0 : \|(I - Q_M P_M)f\|_{L^2(\partial\Omega)} \leq C_E h_M \inf_{c \in \mathbb{R}} \|f + c\|_{H^1(\partial\Omega)}$$

for all $M \in \mathbb{N}$ and all $f \in H^1(\partial\Omega)$.

The first criterion implies the natural assumption that the electrode sizes shrink to zero, but always cover a certain fraction of the boundary. The somewhat technical second criterion can be interpreted as a Poincaré-type inequality that is fulfilled for regular enough electrode

shapes, see [30, 52, 69]. Together these criteria guarantee that the electrode measurement approximate all possible continuous measurements in a suitable sense.

Now we can state our main result:

Theorem 3.1. *There exists $N \in \mathbb{N}$ and $c > 0$ such that for all $M \geq N$*

$$\|R_M(\sigma_1) - R_M(\sigma_2)\|_{\mathcal{L}(\mathbb{R}_\diamond^M)} \geq c \|\sigma_1 - \sigma_2\|_{L^\infty(\Omega)} \quad \text{for all } \sigma_1, \sigma_2 \in \mathcal{F}_{[a,b]}.$$

In particular, this implies that for all $\sigma_1, \sigma_2 \in \mathcal{F}_{[a,b]}$ and $M \geq N$

$$R_M(\sigma_1) = R_M(\sigma_2) \quad \text{if and only if} \quad \sigma_1 = \sigma_2.$$

Theorem 3.1 will be proven in section 3.3.

3.2. Differentiability, monotonicity, and approximation of linearized measurements

The electrode measurements $R_M(\sigma)$ fulfill analogue differentiability and monotonicity properties as the NtD-Operators. In the following $\langle \cdot, \cdot \rangle_M$ denotes the Euclidian scalar product in \mathbb{R}^M . For a vector $J = (J_1, \dots, J_M) \in \mathbb{R}_\diamond^M$, we denote by $u_\sigma^{(J)} \in H_\diamond^1(\Omega)$ the solution of the CEM equations (9)–(12) with conductivity $\sigma \in L_+^\infty(\Omega)$ and electrode currents $J_1, \dots, J_m \in \mathbb{R}$.

Lemma 3.2.

(a) *The mapping*

$$R_M : L_+^\infty(\Omega) \rightarrow \mathcal{L}(\mathbb{R}_\diamond^M), \quad \sigma \mapsto R_M(\sigma)$$

is Fréchet differentiable. Its derivative is given by

$$R'_M(\sigma) \in \mathcal{L}(L^\infty(\Omega), \mathcal{L}(\mathbb{R}_\diamond^M)), \quad (R'_M(\sigma)\kappa)J = V,$$

where $(v, V) \in H^1(\Omega) \times \mathbb{R}_\diamond^m$ solves

$$\begin{aligned} & \int_\Omega \sigma \nabla v \cdot \nabla w \, dx + \sum_{m=1}^M \int_{E_m} \frac{1}{z} (v - V_m)(w - W_m) \, ds \\ & = - \int_\Omega \kappa \nabla u_\sigma^{(J)} \cdot \nabla w \, dx \end{aligned} \tag{14}$$

for all $(w, W) \in H^1(\Omega) \times \mathbb{R}_\diamond^m$.

(b) *For all $\sigma \in L_+^\infty(\Omega)$ and $\kappa \in L^\infty(\Omega)$ the operator $R'_M(\sigma)\kappa \in \mathcal{L}(\mathbb{R}_\diamond^M)$ is self-adjoint, and it fulfills*

$$\langle (R'_M(\sigma)\kappa)I, J \rangle_M = - \int_\Omega \kappa \nabla u_\sigma^{(I)} \cdot \nabla u_\sigma^{(J)} \, dx,$$

for all $I, J \in \mathbb{R}_\diamond^M$.

(c) *The mapping*

$$R'_M : L_+^\infty(\Omega) \rightarrow \mathcal{L}(L^\infty(\Omega), \mathcal{L}(\mathbb{R}_\diamond^M)), \quad \sigma \mapsto R'_M(\sigma)$$

is continuous.

Proof. This follows from the variational formulation of the CEM (13), see e.g. [70, section 2] or [30, appendix B]. \square

Lemma 3.3. For all $\sigma_1, \sigma_2 \in L_+^\infty(\Omega)$ and $J \in \mathbb{R}_\diamond^M$, it holds that

$$\begin{aligned} \langle (R'(\sigma_2)(\sigma_1 - \sigma_2))J, J \rangle_M &= \int_\Omega (\sigma_2 - \sigma_1) |\nabla u_{\sigma_2}^{(J)}|^2 dx \\ &\leq \langle (R_M(\sigma_1) - R_M(\sigma_2))J, J \rangle_M. \end{aligned}$$

Proof. ([47], theorem 2). \square

We will also require the following result from Garde and Staboulis [30] that the linearized CEM measurements approximate the linearized NtD operator.

Lemma 3.4. Under the Hyvönen assumptions (H1) and (H2), there exists $C > 0$ such that for all $\sigma \in \mathcal{F}_{[a,b]}$ and $\kappa \in \text{span } \mathcal{F}$

$$\|\Lambda'(\sigma)\kappa - LQ_M(R'(\sigma)\kappa)Q_M^*\|_{\mathcal{L}(L_\diamond^2(\partial\Omega))} \leq Ch_M \|\sigma\|_{L^\infty(\Omega)} \|\kappa\|_{L^\infty(\Omega)},$$

where

$$\begin{aligned} L : L^2(\partial\Omega) &\rightarrow L_\diamond^2(\partial\Omega), \quad g \mapsto g - \frac{1}{|\partial\Omega|} \int_{\partial\Omega} g ds, \\ Q_M^* : L^2(\partial\Omega) &\rightarrow \mathbb{R}^M, \quad g \mapsto \left(\int_{\bar{E}_1^{(M)}} g ds, \dots, \int_{\bar{E}_M^{(M)}} g ds \right). \end{aligned}$$

Moreover, for all $g \in L_\diamond^2(\partial\Omega)$

$$\langle LQ_M(R'(\sigma)\kappa)Q_M^*g, g \rangle = \langle (R'(\sigma)\kappa)Q_M^*g, Q_M^*g \rangle_M.$$

Proof. ([30], theorem 3, proposition 4). \square

3.3. Proof of theorem 3.1

Again, for the sake of brevity, we omit norm subscripts when the choice of the norm is clear from the context. As in section 2.3 we obtain from the monotonicity result lemma 3.3 that

$$\|R_M(\sigma_1) - R_M(\sigma_2)\| \geq \|\sigma_1 - \sigma_2\| \inf_{\substack{(\tau_1, \tau_2, \kappa) \\ \in \mathcal{F}_{[a,b]} \times \mathcal{F}_{[a,b]} \times \mathcal{K}}} \sup_{\substack{J \in \mathbb{R}_\diamond^M \\ \|J\|=1}} f_M(\tau_1, \tau_2, \kappa, J),$$

where $f_M : L_+^\infty(\Omega) \times L_+^\infty(\Omega) \times L^\infty(\Omega) \times \mathbb{R}_\diamond^M \rightarrow \mathbb{R}$ is defined by

$$f_M(\tau_1, \tau_2, \kappa, J) := \max \{ \langle (R'_M(\tau_1)\kappa)J, J \rangle, - \langle (R'_M(\tau_2)\kappa)J, J \rangle \},$$

and $\mathcal{K} := \{ \kappa \in \text{span } \mathcal{F} : \|\kappa\| = 1 \}$.

We compare this with $f : L_+^\infty(\Omega) \times L_+^\infty(\Omega) \times L^\infty(\Omega) \times L_\diamond^2(\partial\Omega) \rightarrow \mathbb{R}$,

$$f(\tau_1, \tau_2, \kappa, g) := \max \{ \langle (\Lambda'(\tau_1)\kappa)g, g \rangle, - \langle (\Lambda'(\tau_2)\kappa)g, g \rangle \},$$

from the continuum model (see lemma 2.9) with $\Sigma = \partial\Omega$.

For all real numbers $r_1, r_2, s_1, s_2 \in \mathbb{R}$ one has that

$$\begin{aligned} \max\{r_1, s_1\} - \max\{r_2, s_2\} &= \frac{1}{2} (r_1 + s_1 + |r_1 - s_1| - r_2 - s_2 - |r_2 - s_2|) \\ &\leq \frac{1}{2} (r_1 + s_1 + |r_1 - s_1 - (r_2 - s_2)| - r_2 - s_2) \\ &= \max\{(r_1 - r_2), (s_1 - s_2)\}. \end{aligned}$$

Hence, we obtain with lemma 3.4 that for all $(\tau_1, \tau_2, \kappa) \in \mathcal{F}_{[a,b]} \times \mathcal{F}_{[a,b]} \times \mathcal{K}$ and $g \in L^2_\diamond(\partial\Omega)$ with $\|g\| = 1$

$$\begin{aligned} &|f(\tau_1, \tau_2, \kappa, g) - f_M(\tau_1, \tau_2, \kappa, Q_M^* g)| \\ &\leq \max\{|\langle (\Lambda'(\tau_1)\kappa - Q_M(R'_M(\tau_1)\kappa) Q_M^*) g, g \rangle|, \\ &\quad |\langle (\Lambda'(\tau_2)\kappa - Q_M(R'_M(\tau_2)\kappa) Q_M^*) g, g \rangle|\} \\ &\leq Ch_M \|\kappa\| \|g\| \max\{\|\tau_1\|, \|\tau_2\|\} \leq Ch_M b, \end{aligned}$$

where $\|\kappa\| = 1$ and $\max\{\|\tau_1\|, \|\tau_2\|\} \leq b$ follows from the definition of \mathcal{K} and $\mathcal{F}_{[a,b]}$.

For all $g \in L^2(\partial\Omega)$ we have that

$$\|Q_M^* g\|^2 = \sum_{m=1}^M \left(\int_{\tilde{E}_m^{(M)}} g \, ds \right)^2 \leq \sum_{m=1}^M |\tilde{E}_m^{(M)}| \int_{\tilde{E}_m^{(M)}} g^2 \, ds \leq |\partial\Omega| \|g\|^2,$$

so that we obtain for all $(\tau_1, \tau_2, \kappa) \in \mathcal{F}_{[a,b]} \times \mathcal{F}_{[a,b]} \times \mathcal{K}$

$$\begin{aligned} \sup_{\substack{J \in \mathbb{R}_\diamond^M \\ \|J\|=1}} f_M(\tau_1, \tau_2, \kappa, J) &\geq |\partial\Omega|^{-1} \sup_{\substack{g \in L^2_\diamond(\partial\Omega) \\ \|g\|=1}} f_M(\tau_1, \tau_2, \kappa, Q_M^* g) \\ &\geq |\partial\Omega|^{-1} \sup_{\substack{g \in L^2_\diamond(\partial\Omega) \\ \|g\|=1}} f(\tau_1, \tau_2, \kappa, g) - |\partial\Omega|^{-1} Ch_M b. \end{aligned}$$

Since the first summand is positive by lemma 2.11, it follows that for sufficiently large numbers of electrodes M

$$\sup_{\substack{J \in \mathbb{R}_\diamond^M \\ \|J\|=1}} f_M(\tau_1, \tau_2, \kappa, J) > 0 \quad \text{for all } (\tau_1, \tau_2, \kappa) \in \mathcal{F}_{[a,b]} \times \mathcal{F}_{[a,b]} \times \mathcal{K}.$$

With the same lower semicontinuity and compactness argument as in the continuum model, this yields

$$\inf_{(\tau_1, \tau_2, \kappa) \in \mathcal{F}_{[a,b]} \times \mathcal{F}_{[a,b]} \times \mathcal{K}} \sup_{\substack{J \in \mathbb{R}_\diamond^M \\ \|J\|=1}} f_M(\tau_1, \tau_2, \kappa, J) > 0,$$

so that the assertion is proven. \square

Acknowledgments

This work is devoted to the memory of Professor Armin Lechleiter who will be deeply missed as a scientist, and as a friend.

ORCID iDs

Bastian Harrach  <https://orcid.org/0000-0002-8666-7791>

References

- [1] Adler A, Gaburro R and Lionheart W 2015 Electrical impedance tomography *Handbook of Mathematical Methods in Imaging* ed O Scherzer (New York: Springer) pp 701–62
- [2] Alberti G S and Santacesaria M 2018 Calderón’s inverse problem with a finite number of measurements preprint (arXiv:1803.04224)
- [3] Alessandrini G, Beretta E and Vessella S 1996 Determining linear cracks by boundary measurements: Lipschitz stability *SIAM J. Math. Anal.* **27** 361–75
- [4] Alessandrini G, de Hoop M V, Gaburro R and Sincich E 2018 Lipschitz stability for a piecewise linear Schrödinger potential from local Cauchy data *Asymptotic Anal.* **108** 115–49
- [5] Alessandrini G, Maarten V, Gaburro R and Sincich E 2017 Lipschitz stability for the electrostatic inverse boundary value problem with piecewise linear conductivities *J. Math. Pures Appl.* **107** 638–64
- [6] Alessandrini G and Vessella S 2005 Lipschitz stability for the inverse conductivity problem *Adv. Appl. Math.* **35** 207–41
- [7] Ammari H and Uhlmann G 2004 Reconstruction of the potential from partial Cauchy data for the Schrödinger equation *Indiana Univ. Math. J.* **53** 169–83
- [8] Arnold L and Harrach B 2013 Unique shape detection in transient eddy current problems *Inverse Problems* **29** 095004
- [9] Astala K and Päivärinta L 2006 Calderón’s inverse conductivity problem in the plane *Ann. Math.* **265**–99
- [10] Bacchelli V and Vessella S 2006 Lipschitz stability for a stationary 2D inverse problem with unknown polygonal boundary *Inverse problems* **22** 1627
- [11] Barber D and Brown B 1984 Applied potential tomography *J. Phys. E: Sci. Instrum.* **17** 723–33
- [12] Barth A, Harrach B, Hyvönen N and Mustonen L 2017 Detecting stochastic inclusions in electrical impedance tomography *Inverse Problems* **33** 115012
- [13] Bayford R 2006 Bioimpedance tomography (electrical impedance tomography) *Annu. Rev. Biomed. Eng.* **8** 63–91
- [14] Beilina L, Cristofol M, Li S and Yamamoto M 2017 Lipschitz stability for an inverse hyperbolic problem of determining two coefficients by a finite number of observations *Inverse Problems* **34** 015001
- [15] Bellassoued M, Jellali D and Yamamoto M 2006 Lipschitz stability for a hyperbolic inverse problem by finite local boundary data *Appl. Anal.* **85** 1219–43
- [16] Bellassoued M and Yamamoto M 2007 Lipschitz stability in determining density and two Lamé coefficients *J. Math. Anal. Appl.* **329** 1240–59
- [17] Beretta E, de Hoop M V, Francini E, Vessella S and Zhai J 2017 Uniqueness and Lipschitz stability of an inverse boundary value problem for time-harmonic elastic waves *Inverse Problems* **33** 035013
- [18] Beretta E, De Hoop M V and Qiu L 2013 Lipschitz stability of an inverse boundary value problem for a Schrödinger-type equation *SIAM J. Math. Anal.* **45** 679–99
- [19] Beretta E and Francini E 2011 Lipschitz stability for the electrical impedance tomography problem: the complex case *Commun. PDE* **36** 1723–49
- [20] Borcea L 2002 Electrical impedance tomography *Inverse problems* **18** R99
- [21] Borcea L 2003 Addendum to ‘Electrical impedance tomography’ *Inverse Problems* **19** 997–8
- [22] Brander T, Harrach B, Kar M and Salo M 2018 Monotonicity and enclosure methods for the p -Laplace equation *SIAM J. Appl. Math.* **78** 742–58
- [23] Calderón A P 1980 On an inverse boundary value problem *Seminar on Numerical Analysis and its Application to Continuum Physics* ed W H Meyer and M A Raupp (Rio de Janeiro: Brazilian Mathematical Society) pp 65–73
- [24] Calderón A P 2006 On an inverse boundary value problem *Comput. Appl. Math.* **25** 133–8
- [25] Caro P and Rogers K M 2016 Global uniqueness for the Calderón problem with Lipschitz conductivities *Forum of Mathematics, Pi* vol 4 (Cambridge: Cambridge University Press)
- [26] Cheney M, Isaacson D and Newell J 1999 Electrical impedance tomography *SIAM Rev.* **41** 85–101
- [27] Cheng J *et al* 2003 Lipschitz stability in the lateral Cauchy problem for elasticity system *J. Math. Kyoto Univ.* **43** 475–501
- [28] Druskin V 1998 On the uniqueness of inverse problems from incomplete boundary data *SIAM J. Appl. Math.* **58** 1591–603

- [29] Garde H 2017 Comparison of linear and non-linear monotonicity-based shape reconstruction using exact matrix characterizations *Inverse Problems Sci. Eng.*
- [30] Garde H and Staboulis S 2017 Convergence and regularization for monotonicity-based shape reconstruction in electrical impedance tomography *Numer. Math.* **135** 1221–51
- [31] Garde H and Staboulis S 2017 The regularized monotonicity method: detecting irregular indefinite inclusions preprint (arXiv:1705.07372)
- [32] Gebauer B 2008 Localized potentials in electrical impedance tomography *Inverse Problems Imaging* **2** 251–69
- [33] Griesmaier R and Harrach B 2018 Monotonicity in inverse medium scattering on unbounded domains *SIAM J. Appl. Math.* **78** 2533–57
- [34] Guillarmou C and Tzou L 2013 The Calderón inverse problem in two dimensions *Inverse Problems and Applications: Inside Out. II (Mathematical Sciences Research Institute Publications vol 60)* (Cambridge: Cambridge University Press) pp 119–66
- [35] Haberman B *et al* 2013 Uniqueness in Calderóns problem with Lipschitz conductivities *Duke Math. J.* **162** 497–516
- [36] Harrach B 2009 On uniqueness in diffuse optical tomography *Inverse Problems* **25** 055010
- [37] Harrach B 2012 Simultaneous determination of the diffusion and absorption coefficient from boundary data *Inverse Problems Imaging* **6** 663–79
- [38] Harrach B, Lee E and Ullrich M 2015 Combining frequency-difference and ultrasound modulated electrical impedance tomography *Inverse Problems* **31** 095003
- [39] Harrach B and Lin Y-H 2018 Monotonicity-based inversion of the fractional Schrödinger equation preprint (arXiv:1711.05641)
- [40] Harrach B, Lin Y-H and Liu H 2018 On localizing and concentrating electromagnetic fields *SIAM J. Appl. Math.* **78** 2558–74
- [41] Harrach B and Meftahi H 2018 Global uniqueness and Lipschitz-stability for the inverse Robin transmission problem preprint (arXiv:1808.01806)
- [42] Harrach B and Minh M N 2016 Enhancing residual-based techniques with shape reconstruction features in electrical impedance tomography *Inverse Problems* **32** 125002
- [43] Harrach B and Minh M N 2018 Monotonicity-based regularization for phantom experiment data in electrical impedance tomography *New Trends in Parameter Identification for Mathematical Models* (Berlin: Springer) pp 107–20
- [44] Harrach B, Pohjola V and Salo M Monotonicity and local uniqueness for the helmholtz equation *Anal. PDE* to appear
- [45] Harrach B and Seo J K 2010 Exact shape-reconstruction by one-step linearization in electrical impedance tomography *SIAM J. Math. Anal.* **42** 1505–18
- [46] Harrach B and Ullrich M 2013 Monotonicity-based shape reconstruction in electrical impedance tomography *SIAM J. Math. Anal.* **45** 3382–403
- [47] Harrach B and Ullrich M 2015 Resolution guarantees in electrical impedance tomography *IEEE Trans. Med. Imaging* **34** 1513–21
- [48] Harrach B and Ullrich M 2017 Local uniqueness for an inverse boundary value problem with partial data *Proc. Am. Math. Soc.* 145 1087–95
- [49] Henderson R and Webster J 1978 An impedance camera for spatially specific measurements of the thorax *IEEE Trans. Biomed. Eng.* **25** 250–4
- [50] Holder D 2005 *Electrical Impedance Tomography: Methods, History and Applications* (Bristol: IOP Publishing)
- [51] Hyvönen N 2004 Complete electrode model of electrical impedance tomography: approximation properties and characterization of inclusions *SIAM J. Appl. Math.* **64** 902–31
- [52] Hyvönen N 2009 Approximating idealized boundary data of electric impedance tomography by electrode measurements *Math. Models Methods Appl. Sci.* **19** 1185–202
- [53] Ikehata M 1998 Size estimation of inclusion *J. Inverse Ill-Posed Problems* **6** 127–40
- [54] Imanuvilov O Y, Uhlmann G and Yamamoto M 2010 The Calderón problem with partial data in two dimensions *J. Am. Math. Soc.* **23** 655–91
- [55] Imanuvilov O Y, Uhlmann G and Yamamoto M 2015 The Neumann-to-Dirichlet map in two dimensions *Adv. Math.* **281** 578–93
- [56] Imanuvilov O Y and Yamamoto M 1998 Lipschitz stability in inverse parabolic problems by the Carleman estimate *Inverse problems* **14** 1229
- [57] Imanuvilov O Y and Yamamoto M 2001 Global Lipschitz stability in an inverse hyperbolic problem by interior observations *Inverse problems* **17** 717

- [58] Isakov V 2007 On uniqueness in the inverse conductivity problem with local data *Inverse Problems Imaging* **1** 95–105
- [59] Kang H, Seo J K and Sheen D 1997 The inverse conductivity problem with one measurement: stability and estimation of size *SIAM J. Math. Anal.* **28** 1389–405
- [60] Kazemi M A and Klibanov M V 1993 Stability estimates for ill-posed Cauchy problems involving hyperbolic equations and inequalities *Appl. Anal.* **50** 93–102
- [61] Kenig C and Salo M 2013 The Calderón problem with partial data on manifolds and applications *Anal. PDE* **6** 2003–48
- [62] Kenig C and Salo M 2014 Recent progress in the calderón problem with partial data *Contemp. Math.* **615** 193–222
- [63] Kenig C E, Sjöstrand J and Uhlmann G 2007 The Calderón problem with partial data *Ann. Math.* **165** 567–91
- [64] Klibanov M V and Pamyatnykh S E 2006 Lipschitz stability of a non-standard problem for the non-stationary transport equation via a Carleman estimate *Inverse Problems* **22** 881
- [65] Klibanov M V and Yamamoto M 2006 Lipschitz stability of an inverse problem for an acoustic equation *Appl. Anal.* **85** 515–38
- [66] Kohn R V and Vogelius M 1984 Determining conductivity by boundary measurements *Commun. Pure Appl. Math.* **37** 289–98
- [67] Kohn R V and Vogelius M 1985 Determining conductivity by boundary measurements II. Interior results *Commun. Pure Appl. Math.* **38** 643–67
- [68] Krupchyk K and Uhlmann G 2016 The Calderón problem with partial data for conductivities with $3/2$ derivatives *Commun. Math. Phys.* **348** 185–219
- [69] Lechleiter A, Hyvönen N and Hakula H 2008 The factorization method applied to the complete electrode model of impedance tomography *SIAM J. Appl. Math.* **68** 1097–121
- [70] Lechleiter A and Rieder A 2008 Newton regularizations for impedance tomography: convergence by local injectivity *Inverse Problems* **24** 065009
- [71] Lionheart W R B 2004 EIT reconstruction algorithms: pitfalls, challenges and recent developments *Physiol. Meas.* **25** 125–42
- [72] Maffucci A, Vento A, Ventre S and Tamburrino A 2016 A novel technique for evaluating the effective permittivity of inhomogeneous interconnects based on the monotonicity property *IEEE Trans. Compon. Packag. Manuf. Technol.* **6** 1417–27
- [73] Martinsen O G and Grimnes S 2011 *Bioimpedance and Bioelectricity Basics* (New York: Academic)
- [74] Meléndez P G 2013 Lipschitz stability in an inverse problem for the main coefficient of a Kuramoto–Sivashinsky type equation *J. Math. Anal. Appl.* **408** 275–90
- [75] Metherall P, Barber D, Smallwood R and Brown B 1996 Three dimensional electrical impedance tomography *Nature* **380** 509–12
- [76] Nachman A I 1996 Global uniqueness for a two-dimensional inverse boundary value problem *Ann. Math.* **143** 71–96
- [77] Newell J, Gisser D G and Isaacson D 1988 An electric current tomograph *IEEE Trans. Biomed. Eng.* **35** 828–33
- [78] Rüländ A and Sincich E 2018 Lipschitz stability for the finite dimensional fractional Calderón problem with finite cauchy data preprint (arXiv:1805.00866)
- [79] Seo J K, Kim K C, Jargal A, Lee K and Harrach B 2018 A learning-based method for solving ill-posed nonlinear inverse problems: a simulation study of lung EIT preprint (arXiv:1810.10112)
- [80] Seo J K and Woo E J 2013 Electrical impedance tomography *Nonlinear Inverse Problems in Imaging* (Hoboken, NJ: Wiley) pp 195–249
- [81] Sincich E 2007 Lipschitz stability for the inverse robin problem *Inverse Problems* **23** 1311
- [82] Somersalo E, Cheney M and Isaacson D 1992 Existence and uniqueness for electrode models for electric current computed tomography *SIAM J. Appl. Math.* **52** 1023–40
- [83] Su Z, Udpa L, Giovinco G, Ventre S and Tamburrino A 2017 Monotonicity principle in pulsed eddy current testing and its application to defect sizing *Int. Applied Computational Electromagnetics Society Symp. (Italy)* (IEEE) pp 1–2
- [84] Sylvester J and Uhlmann G 1987 A global uniqueness theorem for an inverse boundary value problem *Ann. Math.* **153**–69
- [85] Tamburrino A and Rubinacci G 2002 A new non-iterative inversion method for electrical resistance tomography *Inverse Problems* **18** 1809
- [86] Tamburrino A, Sua Z, Ventre S, Udpa L and Udpa S S 2016 Monotonicity based imaging method in time domain eddy current testing *Electromagn. Nondestruct. Eval.* **41** 1

- [87] Uhlmann G 2009 Electrical impedance tomography and Calderón's problem *Inverse problems* **25** 123011
- [88] Ventre S, Maffucci A, Caire F, Le Lostec N, Perrotta A, Rubinacci G, Sartre B, Vento A and Tamburrino A 2017 Design of a real-time eddy current tomography system *IEEE Trans. Magn.* **53** 1–8
- [89] Wexler A, Fry B and Neuman M 1985 Impedance-computed tomography algorithm and system *Appl. Opt.* **24** 3985–92
- [90] Yuan G and Yamamoto M 2007 Lipschitz stability in inverse problems for a Kirchhoff plate equation *Asymptotic Anal.* **53** 29–60
- [91] Yuan G and Yamamoto M 2009 Lipschitz stability in the determination of the principal part of a parabolic equation *ESAIM: Control Optim. Calculus Variations* **15** 525–54
- [92] Zhou L, Harrach B and Seo J K 2018 Monotonicity-based electrical impedance tomography for lung imaging *Inverse Problems* **34** 045005