

Suppl. Table 6. AraG and cytarabine concentrations that reduce B-ALL and T-ALL cell line sensitivity by 50% (IC50).

B-ALL	IC50	
Cell line	AraG ($\mu\text{g/mL}$)	Cytarabine (ng/mL)
697	0.27 ± 0.01	1.23 ± 0.05
BALL-1	4.76 ± 0.50	2.74 ± 0.05
GRANTA-452	6.69 ± 0.70	3.42 ± 0.25
HAL-01	2.15 ± 0.49	0.98 ± 0.04
KARPAS231	26.73 ± 2.62	11.57 ± 0.77
MHH-CALL-4	31.22 ± 2.50	27.07 ± 3.05
MN-60	99.10 ± 1.62	14.95 ± 0.99
NALM-6	3.25 ± 0.28	2.19 ± 0.06
NALM-16	65.19 ± 2.72	8.79 ± 0.42
REH	1.48 ± 0.07	1.31 ± 0.17
ROS-50	90.62 ± 9.05	18.28 ± 3.57
RS4;11	16.42 ± 1.32	5.74 ± 1.89
SEM	35.64 ± 3.71	27.35 ± 3.18
TANOUE	49.14 ± 2.95	27.67 ± 0.60
TOM-1	0.10 ± 0.01	1.38 ± 0.03
T-ALL	IC50	
Cell line	AraG ($\mu\text{g/mL}$)	Cytarabine (ng/mL)
ALL-SIL	1.28 ± 0.16	5.17 ± 0.64
CCRF-CEM	0.43 ± 0.02	3.88 ± 0.70
CTV-1	0.38 ± 0.07	1.72 ± 0.01
HSB-2	0.52 ± 0.05	3.51 ± 0.07
JJHan	0.97 ± 0.19	5.61 ± 0.53
Jurkat	0.96 ± 0.06	6.65 ± 0.52
KE-37	0.32 ± 0.11	1.83 ± 0.25
MOLT-4	0.46 ± 0.01	3.02 ± 0.10
MOLT-16	15.55 ± 1.30	6.48 ± 0.53
P12-ICHIKAWA	0.26 ± 0.01	2.12 ± 0.19
RPMI-8402	0.44 ± 0.01	3.43 ± 0.21