

## REVIEW

# Fibroblast-mediated intercellular crosstalk in the healthy and diseased heart

 Luka Nicin<sup>1,2,3</sup>, Julian U. G. Wagner<sup>1,2,3</sup>, Guillermo Luxán<sup>1,2,3</sup>  and Stefanie Dimmeler<sup>1,2,3</sup> 

1 Institute for Cardiovascular Regeneration, Goethe University, Frankfurt am Main, Germany

2 German Center for Cardiovascular Research (DZHK), Frankfurt am Main, Germany

3 Cardio-Pulmonary Institute (CPI), Frankfurt am Main, Germany

## Correspondence

S. Dimmeler, Institute for Cardiovascular Regeneration, Centre of Molecular Medicine, Goethe University Frankfurt, Theodor Stern Kai 7, 60590 Frankfurt, Germany  
 Tel: +49-69-6301-5158  
 E-mail: dimmeler@em.uni-frankfurt.de

(Received 22 September 2021, revised 28 October 2021, accepted 4 November 2021, available online 1 December 2021)

doi:10.1002/1873-3468.14234

Edited by Eirini Kefalogianni

**Cardiac fibroblasts constitute a major cell population in the heart. They secrete extracellular matrix components and various other factors shaping the microenvironment of the heart. In silico analysis of intercellular communication based on single-cell RNA sequencing revealed that fibroblasts are the source of the majority of outgoing signals to other cell types. This observation suggests that fibroblasts play key roles in orchestrating cellular interactions that maintain organ homeostasis but that can also contribute to disease states. Here, we will review the current knowledge of fibroblast interactions in the healthy, diseased, and aging heart. We focus on the interactions that fibroblasts establish with other cells of the heart, specifically cardiomyocytes, endothelial cells and immune cells, and particularly those relying on paracrine, electrical, and exosomal communication modes.**

**Keywords:** aging; cardiomyocytes; cardiovascular disease; electrical coupling; endothelial cells; exosomes; fibroblasts; heart; immune cells; paracrine signaling


Cardiac fibroblasts constitute a major cell population in the heart defined by their capacity to generate extracellular matrix (ECM). This cell type is crucial for the maintenance of the cardiac structure and the mechanical properties of the heart. Although, historically, the identification of a fibroblast population in heart has proven challenging, because lineage-specific markers for fibroblasts remains controversial [1], the epicardial transcription factor 21 (Tcf21) and the platelet-derived growth factor receptor alpha (PDGFR $\alpha$ ) are commonly used to identify quiescent fibroblasts in the heart [2,3] (Fig. 1). Despite the lack of definitive markers, fibroblasts can be in general identified by the lack of basement membrane, presence of multiple elongated cellular processes, extensive rough endoplasmic

reticulum, and abundant granular cytoplasmic material [4]. It is important to note that other cells in the heart, such as endothelial cells [5] or pericytes [6], can produce ECM and alterations observed in these cells may lead to cell states that can phenotypically and functionally overlap with fibroblasts; thus, exclusion criteria need to be carefully considered to properly define cardiac fibroblasts.

Initial studies estimated cardiac fibroblast number between 27% and 50% of total cells in mouse and rat ventricles, respectively [7,8]. Refinements like the optimization of tissue digestion protocols or the use of different cardiac fibroblast-specific mouse lines have allowed to revisit the cardiac tissue composition and establish that fibroblasts, resident adventitial and

## Abbreviations

ECM, extracellular matrix; FGF, fibroblast-derived growth factor; GM-CSF, granulocyte-macrophage colony-stimulating factor; HIMF, hypoxia-induced mitogenic factor; LIF, leukemia inhibitory factor; MHC, major histocompatibility complex; MIF, macrophage migration inhibitory factor; MMPs, metalloproteases; PRRs, pattern recognition receptors.



**Fig. 1.** Origin and diversity of cardiac fibroblasts. During embryonic development, cardiac fibroblasts (identified by the expression of PDGFR $\alpha$ , TCF21 [2,3]) arise from cardiac endothelial cells, the epicardium, and the neural crest [2,11,12,19]. Upon stimulation by inflammatory or profibrotic cytokines (e.g., TGF $\beta$ ), cardiac fibroblasts transit to an activated state, which can be identified using marker genes and regulators such as TGF $\beta$ , POSTN, SMAD 2/3/4 [24,25,27–29], and can further differentiate to myofibroblasts that show a strong contractile and matrix-secreting phenotype [30].

interstitial, contribute to approximately 10% of all cells [9]. Nevertheless, the study of fibroblast biology remains crucial for the correct understanding of cardiac health, disease, and aging.

Cardiac fibroblasts are of heterogeneous origin, arising from different sources in the heart including the epicardium, the neural crest, and the endothelium (Fig. 1). Nonetheless, it has been reported that this developmental heterogeneity does not predict different pathological responses of cardiac fibroblasts [10]. Several reports have shown that cardiac fibroblasts originate from the epicardium, the mesothelium that covers the heart [11,12]. In the mouse, cellular progenitors of fibroblasts have been described to invade the myocardium from the epicardium around embryonic day 13.5 [13,14] and are characterized by the expression of Tcf21 [15], Wt1 [16], and Tbx18 [17]. It is important to note that the epicardium can give rise to different cell types in the heart [11,17,18] and that the heterogeneous gene expression levels of the above markers has been linked to the specification into different lineages.

In particular, the expression of Tcf21 in a subset of epicardial cells is crucial for their differentiation into fibroblasts [14].

The neural crest is comprised of a heterogeneous population of migratory cells that originate from the dorsal part of the neural tube and give rise to a variety of cell types in different organs [19], including cardiac fibroblasts [20]. The neural crest plays a crucial role in the development of the outflow region of the heart [21], giving rise to valve mesenchyme. Neural crest-derived fibroblasts, identified using the *Pax3-Cre* transgenic model [22], have been detected within the myocardium, predominately in the right atrium [10].

Finally, the embryonic endothelium constitutes a third source of cardiac fibroblasts [2] and gives rise to 10–20% of the fibroblasts resident in the heart, in particular those located in the ventricular septum and right ventricle [10,23].

Cardiac fibroblasts are a plastic cell population capable of activation and differentiation after injury to the heart, being this ischemic, mechanical, or

inflammatory (Fig. 1). Activation results in increased ECM deposition and leads to fibrosis. Lineage tracing, and later single-cell RNA sequencing have revealed that activated fibroblasts express high levels of Periostin (Postn) [24,25]. Furthermore, targeted ablation of Periostin in activated fibroblasts resulted in reduced fibrosis, although it induced localized cardiomyocyte hypertrophy [25]. Activated fibroblasts express several matrix metalloproteases (MMPs) that degrade the ECM and mediate the activation of TGF $\beta$ , thus paving the way for pathological fibrosis. The cytokine TGF $\beta$ , the best characterized fibrogenic growth factor, plays a crucial role in fibroblast activation [26]. It induces the phosphorylation of SMAD2 and SMAD3, which subsequently translocate to the nucleus together with SMAD4 to promote the expression of fibroblasts activation-specific target genes [27–29].

After activation, cardiac fibroblasts additionally transdifferentiate into myofibroblasts in a step-wise way with intermediate phenotypes [30]. The appearance of myofibroblasts is a hallmark of the cardiac fibrotic response [31]. Myofibroblasts are characterized by the expression of alpha smooth muscle actin ( $\alpha$ SMA) in stress fibers and are able to exert contractile force [25,32]. Furthermore, myofibroblasts change their ECM profile expressing molecules like Fibronectin (FN) [33] or Thrombospondin-1 (TSP-1) [27]. This transdifferentiation can work in both directions, as myofibroblasts can differentiate back to quiescent fibroblasts once the fibrotic scar has been stabilized [34]. After injury, myofibroblasts migrate to the site of injury in response to the secretion of different growth factors and hormones, such as TGF $\beta$  [35], PDGF [36], Angiotensin II [37], Aldosterone [38,39] or Endothelin-1 [40], and others, where they orchestrate pathological remodeling [32,41].

Several nonfibroblast cells have been reported to give rise to activated fibroblasts or myofibroblasts upon cardiac injury. Bone marrow-derived cells, perivascular cells or endothelial cells have all been reported to be the primary source of newly activated fibroblasts and myofibroblasts in the heart after injury. However, these findings remain controversial as recent reports using highly refined genetic markers have not confirmed these results (reviewed in [42]). Nonetheless, single-cell RNA sequencing has recently shown that endothelial cells transiently acquire mesenchymal fibroblast-like features after myocardial infarct [43].

Fibroblasts have a high capacity to modulate and affect the environment that surrounds them. Below, we will explore the interaction of fibroblasts with the other cells in the heart.


## Communication between fibroblasts and cardiomyocytes

Cardiomyocytes are the cardiac-specific contractile units, accounting for 65% to 80% of the volume of the adult mammalian heart [44,45]. In the healthy heart, cardiomyocytes are organized in layers of aligned cells in a complex anisotropic structure, with distinct mechanical and biochemical interactions guiding cardiomyocyte organization and homeostasis [46–48]. The communication between cardiomyocytes and other cardiac cell types, especially cardiac fibroblasts, plays a crucial role in regulating cardiomyocyte function in particularly upon acute or chronic stress conditions. Interestingly, transcriptomic profiling has revealed more similarities between cardiac fibroblasts and cardiomyocytes than between other cell types [49]. Furthermore, specific age- and disease-dependent signatures occur with a similar kinetic in cardiac fibroblasts and cardiomyocytes, suggesting an interrelated stress response and possibly close communication between these two cell types [50]. The interactions between cardiac fibroblasts and cardiomyocytes involve paracrine signaling [51] (Fig. 2A), mechanical stimuli [52] (Fig. 2B), electrotonic coupling via gap junctions [48,53] (Fig. 2C), and exosome-mediated crosstalk [54] (Fig. 2D).

## Fibroblast-cardiomyocyte cell communication in the diseased heart

### Cytokines, hormones, and growth factors

Paracrine signaling was investigated as a common route of communication between cardiac fibroblasts and cardiomyocytes in the heart, allowing for indirect crosstalk without direct cell to cell contacts [55,56]. Several paracrine signaling pathways impact both, cardiac fibroblasts activation to myofibroblast triggering fibrosis, and cardiomyocyte function. Angiotensin II (Ang II) and the cytokine TGF $\beta$  were among the first paracrine effector identified in the context of cardiac fibrosis. Cardiac fibroblasts are a major source of the hormone Ang II [57], which has an important role in ventricular remodeling after cardiac injury [58,59]. Ang II also has more direct arrhythmogenic actions by inducing cardiomyocyte swelling [60], which is known to impair the gap junction permeability in adult rat ventricular cardiomyocytes [61]. Inhibition of Ang II generation or downstream interference with the Ang II receptors prevents cardiac hypertrophy and interstitial fibrosis and normalizes intercellular communication under disease states [62,63]. TGF $\beta$  is induced under



**Fig. 2.** Fibroblast–cardiomyocyte crosstalk in diseased and aged heart. (A) Crosstalk by paracrine signals and cytokines. Upon TGF $\beta$ -stimulation cardiac fibroblasts differentiate to myofibroblasts, which release paracrine factors such as TGF $\beta$ , AngII, and IL6 that induce cardiomyocyte hypertrophy and cardiac fibrosis [51,57,67,68]. Fibroblast-derived factors might also be cardiac protective, as IL33 is attenuating hypertrophy in cardiomyocytes [71]. Cardiomyocytes can also signal back to fibroblasts by releasing LIF, CT-1, and calcitonin, which stimulates fibroblast proliferation and matrix protein production [75–77]. (B) Crosstalk via extracellular matrix components. Upon injury, cardiac fibroblasts increase the secretion of matrix components, such as collagens, proteoglycans, and other glycoproteins. The increase in matrix components grants cardiac stability but also makes the extracellular environment more rigid, which alters cardiomyocyte integrin-signaling and mechanosensing [85]. (C) Crosstalk via electric coupling. Cardiomyocytes are electrically coupled via gap junctions (Connexin 43, Cx43) that reside within the intercalated disks [91,92]. Under ischemic conditions, Cx43 might translocate from the intercalated disks at the polar ends of the cell to the lateral surfaces, therefore increasing the likelihood of cardiomyocyte to fibroblast/myofibroblast coupling associated with the risk of arrhythmia [93]. (D) Exosome-mediated communication. Fibroblasts can release miR-21-3p-laden exosomes. Cardiomyocytes take up such exosomes and through the action of miR-21-3p AngII is released thereby leading to cardiomyocyte hypertrophy and fibrosis [99]. Cardiomyocytes can signal back to fibroblasts and exert both fibrotic and protective effects. By releasing miR-208-carrying exosomes, cardiomyocytes can stimulate the fibroblast-to-myofibroblast transition [101]. In turn, miR-29b and miR-455-laden exosomes may attenuate fibrosis [103].

diseased conditions in fibroblasts and other cardiac cell types and acts as the main driver for myofibroblast activation [64,65]. TGF $\beta$ -dependent activation of cardiac fibroblasts to myofibroblasts induces cardiac fibrosis, cardiomyocyte hypertrophy while reducing action potentials and conductivity in neighboring cardiomyocytes [51].

Moreover, pro-inflammatory cytokines, especially interleukin-6 (IL-6), are important mediators in the communication process. IL-6 is a pleiotropic cytokine that has distinct biological functions in the heart [66], with crucial contributions to aging- and disease-dependent processes including cardiomyocyte hypertrophy and fibrosis [67,68]. Enhanced secretion of IL-6 by cardiac fibroblasts effects multiple crosstalk pathways by inducing myofibroblast proliferation and differentiation, cardiomyocyte apoptosis, and angiogenesis of endothelial cells [69]. Recently, Kumar *et al.* reported that the hypoxia-induced mitogenic factor (HIMF) acts as an upstream regulator of IL-6 in cardiomyocyte to cardiac fibroblast paracrine crosstalk [70].

Other fibroblast-derived cytokines may also elicit beneficial effects. Thus, interleukin 33 (IL-33), which is produced by fibroblasts inhibits cardiomyocyte hypertrophy and fibrosis after pressure overload, suggesting that IL-33 may function as a paracrine signal produced by fibroblasts to modulate cardiomyocyte responses to hypertrophic stimuli [71]. Nevertheless, multiple other cardiac cell types can contribute to the secretion of the paracrine factors as discussed above in a complex network of heterocellular crosstalk.

Wnt signaling plays a major role in cardiac fibroblast–cardiomyocyte crosstalk. Inhibition of Wnt signaling globally after myocardial infarction *in vivo* reduced fibrosis, ameliorated cardiomyocyte recovery, and improved cardiac function [72,73]. However, cardiomyocyte-specific blockade of Wnt signaling induced a contradictory response, by increasing fibrosis and impairing cardiomyocyte recovery after myocardial infarction [74]. Taken together, these observations demonstrate that Wnt signaling is a major regulator of cardiac fibroblast–cardiomyocyte communication, with distinct responses in global versus cardiomyocyte-specific inhibition.

In addition, cytokines and hormones can also be produced by cardiomyocytes and signal to fibroblasts. For example, leukemia inhibitory factor (LIF) and cardiotrophin-1 (CT-1) can regulate fibroblast proliferation and cardiomyocyte hypertrophy likely amplifying the injury response. LIF induces hypertrophy in cardiomyocytes while inhibiting myofibroblast activation and collagen deposition *in vitro* [75]. CT-1 enhances cardiomyocyte hypertrophy but promotes fibroblast

proliferation [76]. In addition, atrial cardiomyocytes produce the hormone calcitonin that induces neighboring collagen-producing cardiac fibroblasts to stimulate proliferation and further secretion of ECM proteins [77]. Interestingly, this study demonstrated that human patients with atrial fibrillation showed increased levels of myocardial calcitonin compared to healthy individuals, suggesting that this paracrine crosstalk mechanism could have therapeutic implications [77].

### Extracellular matrix

The mechanical properties of the cardiac tissue environment are mainly regulated by the ECM, which surrounds individual cardiomyocytes and provides structural support for cardiomyocyte organization within the myocardium [78]. With cardiac fibroblasts regulating stiffness of the cardiac microenvironment, they also regulate the mechanical stimuli on the neighboring cardiomyocytes. Two distinct models can be used to describe the cardiomyocyte mechanosensing machinery: The localized model suggests stretch signals near the plasma membrane, whereas the decentralized model considers force generation at the cell surface, which is then transmitted to other parts of the cell [79]. In addition to structural support, physical ECM–cardiomyocyte interactions have been shown to provide a strong mechanical link from the intracellular contractile apparatus to the surrounding ECM, an interaction established by the dystrophin glycoprotein complex [80]. The transmembrane protein dystroglycan connects the cytoskeleton of the cardiomyocytes to the ECM, contributing to cardiac homeostasis [81]. Apart from electrical conductivity, various studies demonstrated that culturing cardiac fibroblasts or cardiomyocytes on stiffer substrates results in increased expression of cell-cell contact proteins. Pulsatile stretch induces an upregulation of adhesion proteins, such as N-cadherin, in both cardiac fibroblasts and cardiomyocytes [82,83].

Fibroblast-derived extracellular matrix proteins may also control cardiomyocyte responses to injury resulting in reparative or regenerative processes [84]. Interestingly, embryonic fibroblasts express a pattern of extracellular matrix proteins that induce cardiomyocyte proliferation by activating  $\beta$ 1-integrin signaling [85]. This activity was lost in adult heart-derived fibroblasts, suggesting that changes in the fibroblast–cardiomyocyte communication may contribute to the loss of regenerative capacities in the adult heart. Furthermore, in a model of pressure overload, activated cardiac fibroblast exhibit a protective role by preserving the ECM network, thereby blocking inflammation and decreasing cardiomyocyte injury,

by Smad-mediated pathway that suppresses matrix-degrading proteins [86].

### Electrical coupling

In addition to their role as main drivers of fibrotic processes by exacerbated collagen deposition and myocardial stiffening, cardiac fibroblast can couple through gap junctions with each other and with cardiomyocyte, maintaining the scar regions electrically conductive [87]. Although activated myofibroblasts can have a positive impact on conduction across disarranged cardiomyocyte networks [88], computer-modeling and experimental studies demonstrated their detrimental effects on cardiac function by directly depolarizing cardiomyocyte, decreasing action potential conduction and promoting ectopic electrical activity [89,90]. Gap junctions are the main mechanism of electrotonic coupling in the heart, with connexin 43 (Cx43) being the most prominent gap junction protein expressed in both cardiomyocytes and cardiac fibroblasts [91,92]. During ischemic conditions, cardiomyocytes undergo structural remodeling, with Cx43 translocating from the intercalated disks at the polar ends of the cell to the lateral surfaces, therefore increasing the likelihood of cardiomyocyte to myofibroblast coupling associated with the risk of arrhythmia [93]. Increased expression of Cx43 in cardiac fibroblasts after cardiac injury was associated with augmented coupling with neighboring cardiomyocytes [94]. In contrast, cardiomyocytes from injured hearts had reduced Cx43 levels, which has been shown to result in slowed conduction velocity and thereby contributing to the risk for arrhythmias [95]. This study was supported by *ex vivo* experiments where isolated myofibroblasts from infarcted hearts cocultured with cardiomyocytes decreased the conduction velocities and duration of the action potential [55]. The differential regulation of Cx43 expression in cardiac fibroblasts and cardiomyocytes was proposed to be mediated via the activation of  $\beta$ -adrenergic receptors upon cardiac injury, which results in reduced Cx43 expression in cardiomyocytes, but increased expression in fibroblasts [96]. Such interaction can be studied in heterocellular cardiac tissue models allowing to quantify electric interactions between fibroblasts and cardiomyocytes *in vitro* by cell type-specific optogenetic manipulation of membrane potential [97]. Funken *et al.* demonstrated that under basal conditions, fibroblast membrane potential alterations had minor effects on cardiomyocytes. However, after TGF $\beta$ 1 stimulation and differentiation of fibroblasts toward myofibroblasts, fibroblast-specific depolarization or hyperpolarization directly modulated cardiomyocyte membrane potential and accelerated or blocked spontaneous beating, respectively. Together, multiple studies

have demonstrated the role of electrical coupling of activated myofibroblasts and cardiomyocyte in aging and disease [98], but the extent of cardiac fibroblast–cardiomyocyte interactions in the healthy heart and the role they play in the maintenance of electrophysiological conduction remains to be elucidated.

### Exosomes

Finally, exosomes are well recognized to play essential roles in mediating intercellular crosstalk. They transport molecules that regulate the physiological and pathophysiological processes of the recipient cells [54]. Analysis of the exosome content of cardiac fibroblast by RNA sequencing showed a high abundance of the 3'-p passenger 'star' strand of the profibrotic microRNA miR-21 in exosomes of Ang II-treated cardiac fibroblast. miR-21-3p containing exosomes were shown to be taken up by cardiomyocytes via a endocytosis-dependent manner and augments cardiomyocyte miR-21-3p expression, which reduces the expression of cardiomyocyte-specific structural proteins, potentially contributing to cardiomyocyte hypertrophy [99]. Additionally, other reports confirmed exosome-mediated interaction and show that cardiac fibroblast-derived exosomes augment Ang II production in cardiomyocytes resulting in the induction of cardiac pathological hypertrophy [100]. On the other hand, cardiomyocyte-derived exosomes also can control the function of cardiac fibroblast, as demonstrated by the induction of fibrosis and myofibroblast activation via cardiomyocyte-derived, miR-208 loaded exosomes [101]. However, cardiomyocyte-derived exosomes may also exhibit anti-fibrogenic roles by attenuating fibrosis and activating angiogenesis via Hsp20 [102], miR-29b, and miR-455 [103]. Further studies are needed to identify the specific mechanisms of crosstalk between cardiac fibroblasts and cardiomyocytes via miRNA-loaded exosomes, especially in homeostasis versus disease conditions [54].

### Fibroblast–cardiomyocyte communication in the aged heart

Fibroblasts contribute to age-associated alterations in the heart, and cardiac fibrosis is a hallmark of the aging heart. It is not entirely clear whether fibroblasts are the primary triggers of aging or if they are involved secondary to senescence-associated changes in the cardiac microenvironment. Fibroblasts are activated during aging and showed the most profound change in gene expression as compared to other cell types of the aging mouse heart [104]. Aged fibroblasts are characterized by changed expression patterns of inflammatory, extracellular matrix organization angiogenesis, and osteogenic genes [104].

This affects the microenvironment and possibly may contribute to increased substrate stiffness. Since increased substrate stiffness triggers contractile dysfunction associated with telomere shortening one may speculate that changes in the extracellular matrix environment by aged heart fibroblasts may support a phenotype of accelerated aging [105]. Moreover, studies demonstrated that aging of the heart is accompanied by disturbed expression patterns and distribution of Cx43, with age-dependent decrease of Cx43 [106], which may be linked to increased risk for arrhythmic events [107].

### Communication between fibroblasts and endothelial cells


Fibroblasts are closely interacting with endothelial cells in the heart [108]. Endothelial cells form the inner surface of the cardiac chambers and line the entire macro- and microcirculatory system that supplies the heart with blood. The vasculature thereby ensures tissue oxygenation and supply of nutrients. In addition, endothelial cell-derived paracrine signals, so-called ‘angiocrine mediators’, contribute to the vascular niche, which control organ homeostasis, repair, and regeneration in various organs such as liver, lung, and bones [109]. However, less is known whether and how such angiocrine mediators control cardiac disease and aging and to what extent cardiac fibroblasts are involved in maintaining or disturbing the vascular niche.

By controlling the extracellular matrix and by providing growth factors, fibroblasts play a critical role in the growth and stabilization of capillaries. Most prominently, fibroblast-derived growth factors (FGF) are well-known pro-angiogenic factors. While the effect of other mesenchymal cells, such as mesenchymal stromal cells, have been extensively studied and various studies demonstrated that they can be used as cell therapeutic strategy to provide pro-angiogenic factors, matrix proteins, and exosomes [110–112], the specific interactions of intrinsic cardiac fibroblasts with endothelial cells in the tissue has not been deeply explored. Interestingly, *in silico* prediction of cellular communication based on single-nuclei-RNA-sequencing data sets of healthy hearts revealed that endothelial cells show a high number of incoming signals arising from fibroblasts both in the healthy as well as in the diseased and aging heart [104].

### Fibroblast-endothelial cell communication in the diseased heart

Particularly after myocardial injury or stress, myofibroblasts are the most ligand providing cell type to

signal to endothelial cells [113]. Especially, a distinct type of activated fibroblasts that expressed Wnt inhibitory factors (WntX-fibroblast) was detected in the border zone at day 3 postinfarction and was proposed to interfere with cardiac endothelial cells. This ‘WntX-fibroblast’ population expressed paracrine factors, such as *Wif1*, *Timp3*, *Ptn*, *Mdk*, *Apoe*, *Fbln1*, *Igf1*, and *Rspo3*, which were corresponding to endothelial-expressed receptors [113]. All these factors are known to regulate angiogenesis [114–117], which suggests a potential role of the subpopulation of WntX-fibroblasts in regulating the revascularization of the border zone upon infarction. While most of the expressed genes are pro-angiogenic, some inhibit vessel growth and may also have a negative impact on the heart. For example, *Wif1* inhibits tumor angiogenesis via interfering with Wnt and Vegf pathways [118,119] (Fig. 3A). Especially, after myocardial infarction *Wif1* expression was found to be induced in cardiomyocytes, but not in fibroblasts or endothelial cells and its deletion was reported to further induce abnormal chamber remodeling upon myocardial infarction in mice [120]. By contrast, cardiomyocytes-specific *Wif1* induction rather caused dilated cardiomyopathy *in vivo* [121], indicating the need of a well-balanced *Wif1* expression to positively contribute to cardiac repair. However, to what extent the newly identified WntX-fibroblast population plays a role in cardiac remodeling, is still debatable. In advanced postinfarct cardiac remodeling, myofibroblasts further contribute to the fibroblast-endothelial crosstalk by expressing matrix proteins like *Postn*, *Fn1*, and *Col8a1* (Fig. 3A). These factors can interact on endothelial cell expressed receptors and might thereby regulate endothelial cell adhesion and angiogenesis [113]. Of note, such bioinformatical approaches ignore anatomical and spatial information, hence immunofluorescence detections were used to confirm the close proximity between *Pdgfra+* fibroblasts and *CD31+* endothelial cells [113]. Despite the histological proximity, robust experimental approaches proofing that the specific fibroblast subpopulations and fibroblast-derived factors control endothelial cell phenotypes and functions are still lacking. However, first *in vitro* approaches revealed that fibroblast not only secrete paracrine factors, but also release microRNA-loaded exosomes to communicate with endothelial cells. In a profibrotic setting, it was shown that mouse cardiac endothelial cell function is impaired when cultured with exosomes that were isolated from TGF $\beta$ -pretreated fibroblasts [122]. These findings are of potential interest since the up-take of fibroblast-derived miR in endothelial cells might change their phenotype toward mesenchymal cells.



**Fig. 3.** Fibroblast-endothelial cell crosstalk in diseased and aged heart. (A) In the diseased heart, a Wnt-expressing subpopulation of activated fibroblasts (WntX-Fibroblast) releases paracrine factors, such as Wif1, Timp3, Ptn, Mdk, Apoe, Fbn1, Igf1, and Rspo3, that control angiogenesis [113]. In addition, myofibroblasts release matrix proteins that further promote angiogenesis and endothelial cell adhesion [113]. (B) In the aging heart, endothelial cells secrete pro-inflammatory and profibrotic factors (ET-1, angiotensin II, TGF $\beta$ , TNF $\alpha$ , and IL-6) that provide a stimulatory environment for fibroblast activation [126–128]. Cardiac fibroblasts also signal back to endothelial cells. By secreting Serpin E1, they control angiogenesis in the aging heart [104].

### Fibroblast–endothelial cell communication in the aged heart

Similar to pathophysiological states, cardiac aging is also associated with fibroblast activation and fibrosis [123], as well as vascular remodeling [124]. Aged heart-derived endothelial cells were found to switch their basement membrane from a laminin  $\beta$ 2 to a laminin  $\beta$ 1 rich matrix in both cardiac aging and disease. *In vitro* studies show that this change in matrix proteins has an autocrine influence on endothelial cells and control cell adhesion, autophagy, and inflammation [125]. Since laminins are also critical regulators of fibroblasts (at least in the lung), one may consider also an interaction with laminin  $\beta$ 1 and  $\beta$ 2 on fibroblasts. Advanced age and cardiac disease are additionally associated with an increased expression of endothelial-derived pro-inflammatory factors, such as ET-1 [126], Angiotensin II [127], TGF $\beta$ , TNF $\alpha$  and IL-6 [128] (Fig. 3B). These factors can create a profibrotic environment, stimulating fibroblast activation and hence vascular fibrosis, although fibroblasts show an age-related decline in their response to growth factors such as TGF $\beta$  and EGF [129,130] (Fig. 3B).

Recent studies confirm bioinformatic predictions, suggesting that fibroblasts signal back to endothelial cells in the aging heart: aged heart fibroblast express angiogenesis-controlling genes including Serpin E1 (also known as plasminogen activator inhibitor-1

(PAI-1)) (Fig. 3B). Supernatant of fibroblasts isolated from aged hearts, which were enriched in Serpin E1, impaired endothelial cells *in vitro* [104]. Since Serpin E1 was identified to be a senescence-associated secretory factor [131,132], a potential role for senescent fibroblasts in mediating endothelial impairment in the aging heart should be additionally considered in this context. However, one should note that fibroblast senescence may also beneficially affect injury response in the young mouse heart since it limits the expansion of profibrotic fibroblasts [133]. In addition, in neonatal mice, fibroblast senescence was reported to rather support heart regeneration upon myocardial infarction, as fibroblast proliferation and thereby fibrosis were reduced [134]. Therefore, targeting fibroblast senescence as therapeutic strategy may be considered with caution.

### Communication between fibroblasts and immune cells

Under homeostatic conditions, nonmyocytes in the heart mostly exhibit quiescent phenotypes, but during exposure to stress by aging or disease, a variety of cell types change their phenotype and contribute to remodeling processes. Here, various distinct immune cell types, such as T lymphocytes, monocytes, and macrophages, infiltrate into the myocardium and have diverse spatially and temporally regulated functions



[135,136]. Recently published single-cell RNA-sequencing data of the human heart demonstrated that immune cells constitute between 5% and 10% of all cardiac cell types, revealing various immune cell states with distinct clusters [44]. In the same study, bioinformatically predicted cell-cell interactions identified specific crosstalk patterns between cardiac fibroblasts and immune cells. Here, monocyte-derived and antigen-producing macrophages were predicted to crosstalk with cardiac fibroblasts via CD74-MIF, an interaction contributing to fibrosis if inhibited [137] (Fig. 4A). Additionally, a recently published bioinformatic cell-cell communication study of the heart showed that cardiac M1 macrophages and myofibroblasts exhibited a large number of outgoing communication signals [113]. Other cardiac fibroblasts-immune cell interactions demonstrated that upon cardiac injury myofibroblasts secrete the factor GM-CSF instructing resident macrophages to recruit other inflammatory cells and triggering myocardial inflammation [138,139] (Fig. 4B). Moreover, cardiac fibroblasts modulate inflammatory processes at various levels by interfering with chemotaxis, infiltration, and migration of inflammatory cells. In recently conducted studies, cardiac fibroblasts have been demonstrated to be directly involved in responding to damage by pattern recognition receptors (PRRs), which in turn causes a feed-forward inflammatory response via NF- $\kappa$ B, MAPK8, and p38 stress signaling pathways triggering the release of pro-inflammatory cytokines [140,141] (Fig. 4B).


Additionally, the crosstalk between cardiac fibroblasts and immune cells was shown to occur in bidirectional fashion, with cardiac infiltrating Ly6C<sup>hi</sup> macrophages mediating the overexpression of profibrotic genes, stimulating cardiac fibroblast proliferation and collagen production [142] (Fig. 4C). Macrophages also communicate via the release of exosomes containing miRNAs, specifically miR-155, which regulates inflammation and cardiac injury upon myocardial infarction by transfer of miR-155 to cardiac fibroblasts via macrophage derived exosomes [143]. Another miRNA, miR-21, has recently been shown to modulate macrophage-cardiac fibroblast crosstalk by regulating cardiac macrophages in their paracrine, profibrotic secretome toward cardiac fibroblasts, controlling cardiac remodeling, and function [144]. Recent studies highlight the different roles of tissue-resident macrophages versus circulating bone marrow-derived monocytes. Dick *et al.* [145] reported that within six months, 80% of the resident CCR2<sup>+</sup> macrophages and about 25% of the CCR2-MHC-II<sup>hi</sup> macrophages were replaced by cells derived from circulating monocytes, whereas CCR2-MHC-II<sup>lo</sup> showed

little replacement. It would be interesting to explore how these different populations interact with fibroblast after cardiac injury.

Moreover, neutrophils are among the first immune cells infiltrating into the damaged myocardium. Activated neutrophils release oxygen species (ROS) generated via the NADPH oxidase, exerting profibrotic processes in the myocardial environment [146,147] (Fig. 4C). Although it is challenging to exclude confounding effects of ROS released by other cardiac cell types, it has been shown that NADPH oxidase 2 (Nox2)-deficient mice show an attenuated fibrotic response under pressure overload, also confirmed by reduction of collagen and MMP-2 production [148]. Neutrophils also communicate via the release of granules containing myeloperoxidase, MMP, elastase, cathepsins, and others. These granules can degrade connective tissue released by activated cardiac fibroblasts during fibrosis [149]. Several other immune cells, for example, eosinophils, modulate cardiac function upon injury. After myocardial infarction, eosinophil counts within the heart are increased in the infarct region, and they exhibit an important cardioprotective function in protecting cardiomyocytes from apoptosis and blocking cardiac fibrosis by inhibiting TGF- $\beta$ -induced cardiac fibroblast activation [150] (Fig. 4C). The trafficking of eosinophils to the heart was also investigated in myocarditis, showing that high levels of eotaxin (CCL11) in fibroblasts attracts eosinophils, which expressing the eotaxin receptor CCR3 [151]. Here, fibroblasts are the main source of CCL11 in the heart, underlining the central role of cardiac fibroblasts in regulating cardiac inflammation via paracrine signaling [151]. Together, these studies demonstrate a close bidirectional crosstalk between immune cells infiltration and myofibroblast activation.

## Summary and outlook

While fibroblasts have been historically mainly viewed as collagen-producing cells that induce cardiac fibrosis, technological improvements allowing for a detailed analysis of cell types and subsets of cells are giving novel insights into the crosstalk between the cells in the heart. In response to injury, stress or aging, fibroblasts become activated and change the microenvironment around them by secreting ECM, cytokines, growth factors, and other mediators of intercellular communication such as exosomes. These changes in the environment critically contribute to cardiovascular disease and dysfunction. Learning more about these communication pathways, but more specifically investigating subsets and states of fibroblasts, may lead to



**Fig. 4.** Fibroblast-immune cell crosstalk in diseased and aged heart. (A) Protective interaction. Monocyte-derived macrophages release CD74-MIF and thereby attenuate fibroblast activation [137]. (B) In the diseased heart, myofibroblasts secrete GM-CSF, which stimulates macrophages to recruit inflammatory cells [138,139]. In addition, NF- $\kappa$ B, MAPK8, and p38 activation induce fibroblasts to recruit further inflammatory cells [140,141]. (C) Ly6C<sup>high</sup> macrophages release profibrotic factors that activate cardiac fibroblasts and drive their conversion into myofibroblasts [142]. Neutrophil-derived ROS and eosinophil-derived TGF $\beta$  further activate fibroblast-to-myofibroblast differentiation [146,147]. Neutrophils can also act in an antifibrotic manner by releasing granules that contain matrix metalloproteases to degrade excessive collagen deposits [149].

the identification of novel possible interventional strategies to combat cardiovascular disease. Searching for secreted factors, for example, by using secretome mouse models allowing cell type-specific tracking of released proteins [152], may also lead to the identification of circulating biomarkers of fibroblast activation for identification of patients at risk of worsening cardiac function and chronic fibroblast activation.

## References

- Moore-Morris T, Guimarães-Camboa N, Yutzey KE, Pucéat M and Evans SM (2015) Cardiac fibroblasts: from development to heart failure. *J Mol Med* **93**, 823–830.
- Tallquist MD and Molkenin JD (2017) Redefining the identity of cardiac fibroblasts. *Nat Rev Cardiol* **14**, 484–491.
- Braitsch CM and Yutzey KE (2013) Transcriptional control of cell lineage development in epicardium-derived cells. *Journal of Developmental Biology* **1**, 92–111.
- Camelliti P, Borg TK and Kohl P (2005) Structural and functional characterisation of cardiac fibroblasts. *Cardiovasc Res* **65**, 40–51.
- Anderson DEJ and Hinds MT (2012) Extracellular matrix production and regulation in micropatterned endothelial cells. *Biochem Biophys Res Comm* **427**, 159–164.
- Su H, Cantrell AC, Zeng H, Zhu S-H and Chen J-X (2021) Emerging role of pericytes and their secretome in the heart. *Cells* **10**, 548.
- Nag AC (1980) Study of non-muscle cells of the adult mammalian heart: a fine structural analysis and distribution. *Cytobios* **28**, 41–61.
- Banerjee I, Fuseler JW, Price RL, Borg TK and Baudino TA (2007) Determination of cell types and numbers during cardiac development in the neonatal and adult rat and mouse. *Am J Physiol-Heart Circul Physiol* **293**, H1883–H1891.
- Pinto AR, Ilinykh A, Ivey MJ, Kuwabara JT, D'Antoni ML, Debuque R, Chandran A, Wang L, Arora K, Rosenthal NA *et al.* (2016) Revisiting cardiac cellular composition. *Circ Res* **118**, 400–409.
- Ali SR, Ranjbarvaziri S, Talkhabi M, Zhao P, Subat A, Hojjat A, Kamran P, Müller AMS, Volz KS, Tang Z *et al.* (2014) Developmental heterogeneity of cardiac fibroblasts does not predict pathological proliferation and activation. *Circ Res* **115**, 625–635.
- Gittenberger-de Groot AC, Vrancken Peeters M-PFM, Mentink MMT, Gourdie RG and Poelmann RE (1998) Epicardium-derived cells contribute a novel population to the myocardial wall and the atrioventricular cushions. *Circ Res* **82**, 1043–1052.
- Li Y, Urban A, Midura D, Simon H-G and Wang QT (2017) Proteomic characterization of epicardial-myocardial signaling reveals novel regulatory networks including a role for NF- $\kappa$ B in epicardial EMT. *PLoS One* **12**, e0174563.
- Smith CL, Baek ST, Sung CY and Tallquist MD (2011) Epicardial-derived cell epithelial-to-mesenchymal transition and fate specification require PDGF receptor signaling. *Circ Res* **108**, e15–e26.
- Acharya A, Baek ST, Huang G, Eskiocak B, Goetsch S, Sung CY, Banfi S, Sauer MF, Olsen GS, Duffield JS *et al.* (2012) The bHLH transcription factor Tcf21 is required for lineage-specific EMT of cardiac fibroblast progenitors. *Development* **139**, 2139–2149.
- Lu J, Richardson JA and Olson EN (1998) Capsulin: a novel bHLH transcription factor expressed in epicardial progenitors and mesenchyme of visceral organs. *Mech Dev* **73**, 23–32.
- Zhou B, Ma Q, Rajagopal S, Wu SM, Domian I, Rivera-Feliciano J, Jiang D, von Gise A, Ikeda S, Chien KR *et al.* (2008) Epicardial progenitors contribute to the cardiomyocyte lineage in the developing heart. *Nature* **454**, 109–113.
- Cai C-L, Martin JC, Sun Y, Cui L, Wang L, Ouyang K, Yang L, Bu L, Liang X, Zhang X *et al.* (2008) A myocardial lineage derives from Tbx18 epicardial cells. *Nature* **454**, 104–108.
- Mikawa T and Gourdie RG (1996) Pericardial mesoderm generates a population of coronary smooth muscle cells migrating into the heart along with ingrowth of the epicardial organ. *Dev Biol* **174**, 221–232.
- Le Douarin NM and Dupin E (2012) The neural crest in vertebrate evolution. *Curr Opin Genet Dev* **22**, 381–389.
- Bergwerff M, Verberne ME, DeRuiter MC, Poelmann RE and Gittenberger-de Groot AC (1998) Neural crest cell contribution to the developing circulatory system: implications for vascular morphology? *Circ Res* **82**, 221–231.
- Kirby ML (1993) Cellular and molecular contributions of the cardiac neural crest to cardiovascular development. *Trends Cardiovasc Med* **3**, 18–23.
- Lang D, Lu MM, Huang L, Engleka KA, Zhang M, Chu EY, Lipner S, Skoultschi A, Millar SE and Epstein JA (2005) Pax3 functions at a nodal point in melanocyte stem cell differentiation. *Nature* **433**, 884–887.
- Moore-Morris T, Guimarães-Camboa N, Banerjee I, Zambon AC, Kisseleva T, Velayoudon A, Stallcup WB, Gu Y, Dalton ND, Cedenilla M *et al.* (2014) Resident fibroblast lineages mediate pressure overload-induced cardiac fibrosis. *J Clin Invest* **124**, 2921–2934.
- Kanisisak O, Khalil H, Ivey MJ, Karch J, Maliken BD, Correll RN, Brody MJ, Lin J, Aronow BJ,

- Tallquist MD and Molkenin JD (2016) Genetic lineage tracing defines myofibroblast origin and function in the injured heart. *Nat Commun* **7**, 12260.
- 25 Kaur H, Takefuji M, Ngai CY, Carvalho J, Bayer J, Wietelmann A, Poetsch A, Hoelper S, Conway SJ, Möllmann H *et al.* (2016) Targeted ablation of periostin-expressing activated fibroblasts prevents adverse cardiac remodeling in mice. *Circ Res* **118**, 1906–1917.
- 26 Frangogiannis NG (2021) Cardiac fibrosis. *Cardiovasc Res* **117**, 1450–1488.
- 27 Serini G, Bochaton-Piallat ML, Ropraz P, Geinoz A, Borsi L, Zardi L and Gabbiani G (1998) The fibronectin domain ED-A is crucial for myofibroblastic phenotype induction by transforming growth factor-beta1. *J Cell Biol* **142**, 873–881.
- 28 Derynck R and Zhang YE (2003) Smad-dependent and Smad-independent pathways in TGF-beta family signalling. *Nature* **425**, 577–584.
- 29 Dobaczewski M, Gonzalez-Quesada C and Frangogiannis NG (2010) The extracellular matrix as a modulator of the inflammatory and reparative response following myocardial infarction. *J Mol Cell Cardiol* **48**, 504–511.
- 30 Nagalingam RS, Al-Hattab DS and Czubyrt MP (2019) What's in a name? On fibroblast phenotype and nomenclature 1. *Can J Physiol Pharmacol* **97**, 493–497.
- 31 Kong P, Christia P and Frangogiannis NG (2014) The pathogenesis of cardiac fibrosis. *Cell Mol Life Sci* **71**, 549–574.
- 32 Fu X, Khalil H, Kanisicak O, Boyer JG, Vagnozzi RJ, Maliken BD, Sargent MA, Prasad V, Valiente-Alandi I, Blaxall BC *et al.* (2018) Specialized fibroblast differentiated states underlie scar formation in the infarcted mouse heart. *J Clin Invest* **128**, 2127–2143.
- 33 Arslan F, Smeets MB, Riem Vis PW, Karper JC, Quax PH, Bongartz LG, Peters JH, Hoefler IE, Doevendans PA, Pasterkamp G *et al.* (2011) Lack of fibronectin-EDA promotes survival and prevents adverse remodeling and heart function deterioration after myocardial infarction. *Circ Res* **108**, 582–592.
- 34 Bretherton R, Bugg D, Olszewski E and Davis J (2020) Regulators of cardiac fibroblast cell state. *Matrix Biol* **91–92**, 117–135.
- 35 Desmoulière A, Geinoz A, Gabbiani F and Gabbiani G (1993) Transforming growth factor-beta 1 induces alpha-smooth muscle actin expression in granulation tissue myofibroblasts and in quiescent and growing cultured fibroblasts. *J Cell Biol* **122**, 103–111.
- 36 Tuuminen R, Nykänen AI, Krebs R, Soronen J, Pajusola K, Keränen MAI, Koskinen PK, Alitalo K and Lemström KB (2009) PDGF-A, -C, and -D but not PDGF-B increase TGF-beta1 and chronic rejection in rat cardiac allografts. *Arterioscler Thromb Vasc Biol* **29**, 691–698.
- 37 Yang F, Chung ACK, Huang XR and Lan HY (2009) Angiotensin II induces connective tissue growth factor and collagen I expression via transforming growth factor-beta-dependent and -independent Smad pathways: the role of Smad3. *Hypertension* **54**, 877–884.
- 38 Johar S, Cave AC, Narayanapanicker A, Grieve DJ and Shah AM (2006) Aldosterone mediates angiotensin II-induced interstitial cardiac fibrosis via a Nox2-containing NADPH oxidase. *FASEB J* **20**, 1546–1548.
- 39 Hayashi M, Tsutamoto T, Wada A, Tsutsui T, Ishii C, Ohno K, Fujii M, Taniguchi A, Hamatani T, Nozato Y *et al.* (2003) Immediate administration of mineralocorticoid receptor antagonist spironolactone prevents post-infarct left ventricular remodeling associated with suppression of a marker of myocardial collagen synthesis in patients with first anterior acute myocardial infarction. *Circulation* **107**, 2559–2565.
- 40 Shi-Wen X, Chen Y, Denton CP, Eastwood M, Renzoni EA, Bou-Gharios G, Pearson JD, Dashwood M, du Bois RM, Black CM *et al.* (2004) Endothelin-1 promotes myofibroblast induction through the ETA receptor via a rac/phosphoinositide 3-kinase/Akt-dependent pathway and is essential for the enhanced contractile phenotype of fibrotic fibroblasts. *Mol Biol Cell* **15**, 2707–2719.
- 41 Klingberg F, Hinz B and White ES (2013) The myofibroblast matrix: implications for tissue repair and fibrosis. *J Pathol* **229**, 298–309.
- 42 Kong P, Christia P, Saxena A, Su Y and Frangogiannis NG (2013) Lack of specificity of fibroblast-specific protein 1 in cardiac remodeling and fibrosis. *Am J Physiol Heart Circ Physiol* **305**, H1363–1372.
- 43 Tombor LS, John D, Glaser SF, Luxán G, Forte E, Furtado M, Rosenthal N, Baumgarten N, Schulz MH, Wittig J *et al.* (2021) Single cell sequencing reveals endothelial plasticity with transient mesenchymal activation after myocardial infarction. *Nat Commun* **12**, 681.
- 44 Litviňuková M, Talavera-López C, Maatz H, Reichart D, Worth CL, Lindberg EL, Kanda M, Polanski K, Heinig M, Lee M *et al.* (2020) Cells of the adult human heart. *Nature* **588**, 466–472.
- 45 Bergmann O, Zdunek S, Felker A, Salehpour M, Alkass K, Bernard S, Sjöstrom SL, Szewczykowska M, Jackowska T, Dos Remedios C *et al.* (2015) Dynamics of cell generation and turnover in the human heart. *Cell* **161**, 1566–1575.
- 46 Feinberg AW, Alford PW, Jin H, Ripplinger CM, Werdich AA, Sheehy SP, Grosberg A and Parker KK (2012) Controlling the contractile strength of engineered cardiac muscle by hierarchal tissue architecture. *Biomaterials* **33**, 5732–5741.

- 47 Schwan J, Kwaczala AT, Ryan TJ, Bartulos O, Ren Y, Sewanan LR, Morris AH, Jacoby DL, Qyang Y and Campbell SG (2016) Anisotropic engineered heart tissue made from laser-cut decellularized myocardium. *Sci Rep* **6**, 32068.
- 48 Ronaldson-Bouchard K, Ma SP, Yeager K, Chen T, Song L, Sirabella D, Morikawa K, Teles D, Yazawa M and Vunjak-Novakovic G (2018) Advanced maturation of human cardiac tissue grown from pluripotent stem cells. *Nature* **556**, 239–243.
- 49 Furtado MB, Nim HT, Boyd SE and Rosenthal NA (2016) View from the heart: cardiac fibroblasts in development, scarring and regeneration. *Development* **143**, 387–397.
- 50 Nicin L, Abplanalp WT, Schänzer A, Sprengel A, John D, Mellentin H, Tombor L, Keuper M, Ullrich E, Klingel K *et al.* (2021) Single nuclei sequencing reveals novel insights into the regulation of cellular signatures in children with dilated cardiomyopathy. *Circulation* **143**, 1704–1719.
- 51 Cartledge JE, Kane C, Dias P, Tesfom M, Clarke L, Mckee B, Al Ayoubi S, Chester A, Yacoub MH, Camelliti P *et al.* (2015) Functional crosstalk between cardiac fibroblasts and adult cardiomyocytes by soluble mediators. *Cardiovasc Res* **105**, 260–270.
- 52 Engler AJ, Carag-Krieger C, Johnson CP, Raab M, Tang H-Y, Speicher DW, Sanger JW, Sanger JM and Discher DE (2008) Embryonic cardiomyocytes beat best on a matrix with heart-like elasticity: scar-like rigidity inhibits beating. *J Cell Sci* **121**, 3794–3802.
- 53 Quinn TA, Camelliti P, Rog-Zielinska EA, Siedlecka U, Poggioli T, O'Toole ET, Knöpfel T and Kohl P (2016) Electrotonic coupling of excitable and nonexcitable cells in the heart revealed by optogenetics. *Proc Natl Acad Sci USA* **113**, 14852–14857.
- 54 Nguyen BY, Azam T and Wang X (2021) Cellular signaling cross-talk between different cardiac cell populations: an insight into the role of exosomes in the heart diseases and therapy. *Am J Physiol Heart Circ Physiol* **320**, H1213–H1234.
- 55 Vasquez C, Mohandas P, Louie KL, Benamer N, Bapat AC and Morley GE (2010) Enhanced fibroblast-myocyte interactions in response to cardiac injury. *Circ Res* **107**, 1011–1020.
- 56 LaFramboise WA, Scalise D, Stoodley P, Graner SR, Guthrie RD, Magovern JA and Becich MJ (2007) Cardiac fibroblasts influence cardiomyocyte phenotype *in vitro*. *Am J Physiol Cell Physiol* **292**, C1799–1808.
- 57 Dostal DE, Rothblum KN, Chernin MI, Cooper GR and Baker KM (1992) Intracardiac detection of angiotensinogen and renin: a localized renin-angiotensin system in neonatal rat heart. *Am J Physiol* **263**, C838–850.
- 58 Schieffer B, Wirger A, Meybrunn M, Seitz S, Holtz J, Riede UN and Drexler H (1994) Comparative effects of chronic angiotensin-converting enzyme inhibition and angiotensin II type 1 receptor blockade on cardiac remodeling after myocardial infarction in the rat. *Circulation* **89**, 2273–2282.
- 59 Sun Y, Zhang JQ, Zhang J and Ramires FJ (1998) Angiotensin II, transforming growth factor-beta1 and repair in the infarcted heart. *J Mol Cell Cardiol* **30**, 1559–1569.
- 60 Sadoshima J, Xu Y, Slayter HS and Izumo S (1993) Autocrine release of angiotensin II mediates stretch-induced hypertrophy of cardiac myocytes *in vitro*. *Cell* **75**, 977–984.
- 61 De Mello WC (2010) Cell swelling impairs dye coupling in adult rat ventricular myocytes. Cell volume as a regulator of cell communication. *Mol Cell Biochem* **343**, 107–113.
- 62 De Mello WC (1996) Renin-angiotensin system and cell communication in the failing heart. *Hypertension* **27**, 1267–1272.
- 63 De Mello WC and Specht P (2006) Chronic blockade of angiotensin II AT1-receptors increased cell-to-cell communication, reduced fibrosis and improved impulse propagation in the failing heart. *J Renin Angiotensin Aldosterone Syst* **7**, 201–205.
- 64 Khalil H, Kanisicak O, Prasad V, Correll RN, Fu X, Schips T, Vagnozzi RJ, Liu R, Huynh T, Lee S-J *et al.* (2017) Fibroblast-specific TGF- $\beta$ -Smad2/3 signaling underlies cardiac fibrosis. *J Clin Invest* **127**, 3770–3783.
- 65 Hanna A and Frangogiannis NG (2019) The role of the TGF- $\beta$  superfamily in myocardial infarction. *Front Cardiovasc Med* **6**, 140.
- 66 Kamimura D, Ishihara K and Hirano T (2003) IL-6 signal transduction and its physiological roles: the signal orchestration model. *Rev Physiol Biochem Pharmacol* **149**, 1–38.
- 67 Banerjee I, Fuseler JW, Intwala AR and Baudino TA (2009) IL-6 loss causes ventricular dysfunction, fibrosis, reduced capillary density, and dramatically alters the cell populations of the developing and adult heart. *Am J Physiol Heart Circ Physiol* **296**, H1694–1704.
- 68 Fontes JA, Rose NR and Ciháková D (2015) The varying faces of IL-6: From cardiac protection to cardiac failure. *Cytokine* **74**, 62–68.
- 69 Wang J-H, Zhao L, Pan X, Chen N-N, Chen J, Gong Q-L, Su F, Yan J, Zhang Y and Zhang S-H (2016) Hypoxia-stimulated cardiac fibroblast production of IL-6 promotes myocardial fibrosis via the TGF- $\beta$ 1 signaling pathway. *Lab Invest* **96**, 839–852.
- 70 Kumar S, Wang G, Zheng N, Cheng W, Ouyang K, Lin H, Liao Y and Liu J (2019) HIMF (Hypoxia-Induced Mitogenic Factor)-IL (Interleukin)-6 signaling mediates cardiomyocyte-fibroblast crosstalk to

- promote cardiac hypertrophy and fibrosis. *Hypertension* **73**, 1058–1070.
- 71 Sanada S, Hakuno D, Higgins LJ, Schreiter ER, McKenzie ANJ and Lee RT (2007) IL-33 and ST2 comprise a critical biomechanically induced and cardioprotective signaling system. *J Clin Invest* **117**, 1538–1549.
- 72 Saraswati S, Alfaro MP, Thorne CA, Atkinson J, Lee E and Young PP (2010) Pyrvinium, a potent small molecule Wnt inhibitor, promotes wound repair and post-MI cardiac remodeling. *PLoS One* **5**, e15521.
- 73 Matsushima K, Suyama T, Takenaka C, Nishishita N, Ikeda K, Ikada Y, Sawa Y, Jakt LM, Mori H and Kawamata S (2010) Secreted frizzled related protein 4 reduces fibrosis scar size and ameliorates cardiac function after ischemic injury. *Tissue Eng Part A* **16**, 3329–3341.
- 74 Barandon L, Dufourcq P, Costet P, Moreau C, Allières C, Daret D, Dos Santos P, Daniel Lamazière J-M, Couffignal T and Dupl a C (2005) Involvement of FrzA/sFRP-1 and the Wnt/frizzled pathway in ischemic preconditioning. *Circ Res* **96**, 1299–1306.
- 75 Wang F, Trial J, Diwan A, Gao F, Birdsall H, Entman M, Hornsby P, Sivasubramaniam N and Mann D (2002) Regulation of cardiac fibroblast cellular function by leukemia inhibitory factor. *J Mol Cell Cardiol* **34**, 1309–1316.
- 76 Tsuruda T, Jougasaki M, Boerrigter G, Huntley BK, Chen HH, D'Assoro AB, Lee SC, Larsen AM, Cataliotti A and Burnett JC (2002) Cardiotrophin-1 stimulation of cardiac fibroblast growth: roles for glycoprotein 130/leukemia inhibitory factor receptor and the endothelin type A receptor. *Circ Res* **90**, 128–134.
- 77 Moreira LM, Takawale A, Hulsurkar M, Menassa DA, Antanaviciute A, Lahiri SK, Mehta N, Evans N, Psarros C, Robinson P *et al.* (2020) Paracrine signalling by cardiac calcitonin controls atrial fibrogenesis and arrhythmia. *Nature* **587**, 460–465.
- 78 Spinale FG (2007) Myocardial matrix remodeling and the matrix metalloproteinases: influence on cardiac form and function. *Physiol Rev* **87**, 1285–1342.
- 79 Kn ll R, Hoshijima M and Chien K (2003) Cardiac mechanotransduction and implications for heart disease. *J Mol Med (Berl)* **81**, 750–756.
- 80 Lapidos KA, Kakkar R and McNally EM (2004) The dystrophin glycoprotein complex: signaling strength and integrity for the sarcolemma. *Circ Res* **94**, 1023–1031.
- 81 Michele DE, Kabaeva Z, Davis SL, Weiss RM and Campbell KP (2009) Dystroglycan matrix receptor function in cardiac myocytes is important for limiting activity-induced myocardial damage. *Circ Res* **105**, 984–993.
- 82 Zhuang J, Yamada KA, Saffitz JE and Kl ber AG (2000) Pulsatile stretch remodels cell-to-cell communication in cultured myocytes. *Circ Res* **87**, 316–322.
- 83 Yamada K, Green KG, Samarel AM and Saffitz JE (2005) Distinct pathways regulate expression of cardiac electrical and mechanical junction proteins in response to stretch. *Circ Res* **97**, 346–353.
- 84 Gomes RN, Manuel F and Nascimento DS (2021) The bright side of fibroblasts: molecular signature and regenerative cues in major organs. *NPJ Regen Med* **6**, 43.
- 85 Ieda M, Tsuchihashi T, Ivey KN, Ross RS, Hong T-T, Shaw RM and Srivastava D (2009) Cardiac fibroblasts regulate myocardial proliferation through beta1 integrin signaling. *Dev Cell* **16**, 233–244.
- 86 Russo I, Cavalera M, Huang S, Su Y, Hanna A, Chen B, Shinde AV, Conway SJ, Graff J and Frangogiannis NG (2019) Protective effects of activated myofibroblasts in the pressure-overloaded myocardium are mediated through smad-dependent activation of a matrix-preserving program. *Circ Res* **124**, 1214–1227.
- 87 Kakkar R and Lee RT (2010) Intramyocardial fibroblast myocyte communication. *Circ Res* **106**, 47–57.
- 88 Gaudesius G, Miragoli M, Thomas SP and Rohr S (2003) Coupling of cardiac electrical activity over extended distances by fibroblasts of cardiac origin. *Circ Res* **93**, 421–428.
- 89 Miragoli M, Salvarani N and Rohr S (2007) Myofibroblasts induce ectopic activity in cardiac tissue. *Circ Res* **101**, 755–758.
- 90 Salvarani N, Maguy A, De Simone SA, Miragoli M, Jousset F and Rohr S (2017) TGF- 1 (Transforming Growth Factor- 1) Plays a Pivotal Role in Cardiac Myofibroblast Arrhythmogenicity. *Circ Arrhythm Electrophysiol* **10**, e004567.
- 91 Vozzi C, Dupont E, Coppin SR, Yeh HI and Severs NJ (1999) Chamber-related differences in connexin expression in the human heart. *J Mol Cell Cardiol* **31**, 991–1003.
- 92 McArthur L, Chilton L, Smith GL and Nicklin SA (2015) Electrical consequences of cardiac myocyte: fibroblast coupling. *Biochem Soc Trans* **43**, 513–518.
- 93 Camelliti P, Devlin GP, Matthews KG, Kohl P and Green CR (2004) Spatially and temporally distinct expression of fibroblast connexins after sheep ventricular infarction. *Cardiovasc Res* **62**, 415–425.
- 94 Zhang Y, Kanter EM and Yamada KA (2010) Remodeling of cardiac fibroblasts following myocardial infarction results in increased gap junction intercellular communication. *Cardiovasc Pathol* **19**, e233–240.
- 95 Peters NS, Green CR, Poole-Wilson PA and Severs NJ (1993) Reduced content of connexin43 gap junctions in

- ventricular myocardium from hypertrophied and ischemic human hearts. *Circulation* **88**, 864–875.
- 96 Zhang Y, Hou M-C, Li J-J, Qi Y, Zhang Y, She G, Ren Y-J, Wu W, Pang Z-D, Xie W *et al.* (2020) Cardiac  $\beta$ -adrenergic receptor activation mediates distinct and cell type-dependent changes in the expression and distribution of connexin 43. *J Cell Mol Med* **24**, 8505–8517.
  - 97 Funken M, Bruegmann T and Sasse P (2020) Selective optogenetic stimulation of fibroblasts enables quantification of hetero-cellular coupling to cardiomyocytes in a three-dimensional model of heart tissue. *Europace* **22**, 1590–1599.
  - 98 Błyszczuk P, Zuppingner C, Costa A, Nurzynska D, Di Meglio FD, Stellato M, Agarkova I, Smith GL, Distler O and Kania G (2020) Activated cardiac fibroblasts control contraction of human fibrotic cardiac microtissues by a  $\beta$ -adrenoreceptor-dependent mechanism. *Cells* **9**, E1270.
  - 99 Bang C, Batkai S, Dangwal S, Gupta SK, Foinquinos A, Holzmann A, Just A, Remke J, Zimmer K, Zeug A *et al.* (2014) Cardiac fibroblast-derived microRNA passenger strand-enriched exosomes mediate cardiomyocyte hypertrophy. *J Clin Invest* **124**, 2136–2146.
  - 100 Lyu L, Wang H, Li B, Qin Q, Qi L, Nagarkatti M, Nagarkatti P, Janicki JS, Wang XL and Cui T (2015) A critical role of cardiac fibroblast-derived exosomes in activating renin angiotensin system in cardiomyocytes. *J Mol Cell Cardiol* **89**, 268–279.
  - 101 Yang J, Yu X, Xue F, Li Y, Liu W and Zhang S (2018) Exosomes derived from cardiomyocytes promote cardiac fibrosis via myocyte-fibroblast crosstalk. *Am J Transl Res* **10**, 4350–4366.
  - 102 Wang X, Gu H, Huang W, Peng J, Li Y, Yang L, Qin D, Essandoh K, Wang Y, Peng T *et al.* (2016) Hsp20-mediated activation of exosome biogenesis in cardiomyocytes improves cardiac function and angiogenesis in diabetic mice. *Diabetes* **65**, 3111–3128.
  - 103 Chaturvedi P, Kalani A, Medina I, Famil'tseva A and Tyagi SC (2015) Cardiosome mediated regulation of MMP9 in diabetic heart: role of mir29b and mir455 in exercise. *J Cell Mol Med* **19**, 2153–2161.
  - 104 Vidal R, Wagner JUG, Braeuning C, Fischer C, Patrick R, Tombor L, Muhly-Reinholz M, John D, Kliem M, Conrad T *et al.* (2019) Transcriptional heterogeneity of fibroblasts is a hallmark of the aging heart. *JCI Insight* **4**, 22.
  - 105 Chang ACY, Pardon G, Chang ACH, Wu H, Ong S-G, Eguchi A, Ancel S, Holbrook C, Ramunas J, Ribeiro AJS *et al.* (2021) Increased tissue stiffness triggers contractile dysfunction and telomere shortening in dystrophic cardiomyocytes. *Stem Cell Reports* **16**, 2169–2181.
  - 106 Watanabe M, Ichinose S and Sunamori M (2004) Age-related changes in gap junctional protein of the rat heart. *Exp Clin Cardiol* **9**, 130–132.
  - 107 Yan J, Thomson JK, Zhao W, Wu X, Gao X, DeMarco D, Kong W, Tong M, Sun J, Bakhos M *et al.* (2018) The stress kinase JNK regulates gap junction Cx43 gene expression and promotes atrial fibrillation in the aged heart. *J Mol Cell Cardiol* **114**, 105–115.
  - 108 Bowers SLK, Borg TK and Baudino TA (2010) The dynamics of fibroblast-myocyte-capillary interactions in the heart. *Ann N Y Acad Sci* **1188**, 143–152.
  - 109 Rafii S, Butler JM and Ding B-S (2016) Angiocrine functions of organ-specific endothelial cells. *Nature* **529**, 316–325.
  - 110 Psaltis PJ, Zannettino ACW, Worthley SG and Gronthos S (2008) Concise review: mesenchymal stromal cells: potential for cardiovascular repair. *Stem Cells* **26**, 2201–2210.
  - 111 Kastrup J, Mygind ND, Qayyum AA, Mathiasen AB, Haack-Sørensen M and Ekblond A (2016) Mesenchymal stromal cell therapy in ischemic heart disease. *Scand Cardiovasc J* **50**, 293–299.
  - 112 Zhao J, Li X, Hu J, Chen F, Qiao S, Sun X, Gao L, Xie J and Xu B (2019) Mesenchymal stromal cell-derived exosomes attenuate myocardial ischaemia-reperfusion injury through miR-182-regulated macrophage polarization. *Cardiovasc Res* **115**, 1205–1216.
  - 113 Farbehi N, Patrick R, Dorison A, Xaymardan M, Janbandhu V, Wystub-Lis K, Ho JW, Nordon RE and Harvey RP (2019) Single-cell expression profiling reveals dynamic flux of cardiac stromal, vascular and immune cells in health and injury. *Elife* **8**, e43882.
  - 114 Papadimitriou E, Pantazaka E, Castana P, Tsalios T, Polyzos A and Beis D (2016) Pleiotrophin and its receptor protein tyrosine phosphatase beta/zeta as regulators of angiogenesis and cancer. *Biochim Biophys Acta* **1866**, 252–265.
  - 115 *Matrix Biol* (2008) TIMP3, angiogenesis and macular degeneration. **27**, 379–380.
  - 116 Scholz B, Korn C, Wojtarowicz J, Mogler C, Augustin I, Boutros M, Niehrs C and Augustin HG (2016) Endothelial RSPO3 controls vascular stability and pruning through non-canonical WNT/Ca(2+)/NFAT signaling. *Dev Cell* **36**, 79–93.
  - 117 Xie L, Palmsten K, MacDonald B, Kieran MW, Potenta S, Vong S and Kalluri R (2008) Basement membrane derived fibulin-1 and fibulin-5 function as angiogenesis inhibitors and suppress tumor growth. *Exp Biol Med (Maywood)* **233**, 155–162.
  - 118 Hu J, Dong A, Fernandez-Ruiz V, Shan J, Kawa M, Martínez-Ansó E, Prieto J and Qian C (2009) Blockade of Wnt signaling inhibits angiogenesis and

- tumor growth in hepatocellular carcinoma. *Cancer Res* **69**, 6951–6959.
- 119 Ko YB, Kim B-R, Yoon K, Choi EK, Seo SH, Lee Y, Lee MA, Yang JB, Park MS and Rho SB (2014) WIF1 can effectively co-regulate pro-apoptotic activity through the combination with DKK1. *Cell Signal* **26**, 2562–2572.
- 120 Meyer IS, Jungmann A, Dieterich C, Zhang M, Lasitschka F, Werkmeister S, Haas J, Müller OJ, Boutros M, Nahrendorf M *et al.* (2017) The cardiac microenvironment uses non-canonical WNT signaling to activate monocytes after myocardial infarction. *EMBO Mol Med* **9**, 1279–1293.
- 121 Lu D, Dong W, Zhang X, Quan X, Bao D, Lu Y and Zhang L (2013) WIF1 causes dysfunction of heart in transgenic mice. *Transgenic Res* **22**, 1179–1189.
- 122 Ranjan P, Kumari R, Goswami SK, Li J, Pal H, Suleiman Z, Cheng Z, Krishnamurthy P, Kishore R and Verma SK (2021) Myofibroblast-derived exosome induce cardiac endothelial cell dysfunction. *Front Cardiovas Med* **8**, 319.
- 123 Gramley F, Lorenzen J, Knackstedt C, Rana OR, Saygili E, Frechen D, Stanzel S, Pezzella F, Koellensperger E, Weiss C *et al.* (2009) Age-related atrial fibrosis. *Age* **31**, 27–38.
- 124 Münzel T, Camici GG, Maack C, Bonetti NR, Fuster V and Kovacic JC (2017) Impact of oxidative stress on the heart and vasculature: Part 2 of a 3-part series. *J Am Coll Cardiol* **70**, 212–229.
- 125 Wagner JUG, Chavakis E, Rogg E-M, Muhly-Reinholz M, Glaser SF, Günther S, John D, Bonini F, Zeiher AM, Schaefer L *et al.* (2018) Switch in Laminin  $\beta$ 2 to Laminin  $\beta$ 1 isoforms during aging controls endothelial cell functions—brief report. *Arterioscler Thromb Vasc Biol* **38**, 1170–1177.
- 126 Goel A, Su B, Flavahan S, Lowenstein CJ, Berkowitz DE and Flavahan NA (2010) Increased endothelial exocytosis and generation of endothelin-1 contributes to constriction of aged arteries. *Circ Res* **107**, 242–251.
- 127 Wang M, Zhang J, Spinetti G, Jiang L-Q, Monticone R, Zhao D, Cheng L, Krawczyk M, Talan M, Pintus G *et al.* (2005) Angiotensin II activates matrix metalloproteinase type II and mimics age-associated carotid arterial remodeling in young rats. *Am J Pathol* **167**, 1429–1442.
- 128 Csiszar A, Ungvari Z, Koller A, Edwards JG and Kaley G (2003) Aging-induced proinflammatory shift in cytokine expression profile in coronary arteries. *FASEB J* **17**, 1183–1185.
- 129 Cieslik KA, Trial J and Entman ML (2011) Defective myofibroblast formation from mesenchymal stem cells in the aging murine heart rescue by activation of the AMPK pathway. *Am J Pathol* **179**, 1792–1806.
- 130 Shiraha H, Gupta K, Drabik K and Wells A (2000) Aging fibroblasts present reduced epidermal growth factor (EGF) responsiveness due to preferential loss of EGF receptors. *J Biol Chem* **275**, 19343–19351.
- 131 Xu M, Tchkonina T, Ding H, Ogrodnik M, Lubbers ER, Pirtskhalava T, White TA, Johnson KO, Stout MB, Mezera V *et al.* (2015) JAK inhibition alleviates the cellular senescence-associated secretory phenotype and frailty in old age. *Proc Natl Acad Sci USA* **112**, E6301–6310.
- 132 Lewis-McDougall FC, Ruchaya PJ, Domenjo-Vila E, Shin Teoh T, Prata L, Cottle BJ, Clark JE, Punjabi PP, Awad W, Torella D *et al.* (2019) Aged-senescent cells contribute to impaired heart regeneration. *Aging Cell* **18**, e12931.
- 133 Meyer K, Hodwin B, Ramanujam D, Engelhardt S and Sarikas A (2016) Essential role for premature senescence of myofibroblasts in myocardial fibrosis. *J Am Coll Cardiol* **67**, 2018–2028.
- 134 Feng T, Meng J, Kou S, Jiang Z, Huang X, Lu Z, Zhao H, Lau LF, Zhou B and Zhang H (2019) CCN1-induced cellular senescence promotes heart regeneration. *Circulation* **139**, 2495–2498.
- 135 Swirski FK and Nahrendorf M (2013) Leukocyte behavior in atherosclerosis, myocardial infarction, and heart failure. *Science* **339**, 161–166.
- 136 Wei L (2011) Immunological aspect of cardiac remodeling: T lymphocyte subsets in inflammation-mediated cardiac fibrosis. *Exp Mol Pathol* **90**, 74–78.
- 137 Heinrichs D, Knauel M, Offermanns C, Berres M-L, Nellen A, Leng L, Schmitz P, Bucala R, Trautwein C, Weber C *et al.* (2011) Macrophage migration inhibitory factor (MIF) exerts antifibrotic effects in experimental liver fibrosis via CD74. *Proc Natl Acad Sci USA* **108**, 17444–17449.
- 138 Anzai A, Choi JL, He S, Fenn AM, Nairz M, Rattik S, McAlpine CS, Mindur JE, Chan CT, Iwamoto Y *et al.* (2017) The infarcted myocardium solicits GM-CSF for the detrimental oversupply of inflammatory leukocytes. *J Exp Med* **214**, 3293–3310.
- 139 Wu L, Ong S, Talor MV, Barin JG, Baldeviano GC, Kass DA, Bedja D, Zhang H, Sheikh A, Margolick JB *et al.* (2014) Cardiac fibroblasts mediate IL-17A-driven inflammatory dilated cardiomyopathy. *J Exp Med* **211**, 1449–1464.
- 140 Turner NA (2016) Inflammatory and fibrotic responses of cardiac fibroblasts to myocardial damage associated molecular patterns (DAMPs). *J Mol Cell Cardiol* **94**, 189–200.
- 141 Zhang W, Lavine KJ, Epelman S, Evans SA, Weinheimer CJ, Barger PM and Mann DL (2015) Necrotic myocardial cells release damage-associated molecular patterns that provoke fibroblast activation *in vitro* and trigger myocardial inflammation and fibrosis *in vivo*. *J Am Heart Assoc* **4**, e001993.
- 142 Jung M, Ma Y, Iyer RP, DeLeon-Pennell KY, Yabluchanskiy A, Garrett MR and Lindsey ML



- (2017) IL-10 improves cardiac remodeling after myocardial infarction by stimulating M2 macrophage polarization and fibroblast activation. *Basic Res Cardiol* **112**, 33.
- 143 Wang C, Zhang C, Liu L, Chen B, Li Y and Du J (2017) Macrophage-Derived mir-155-containing exosomes suppress fibroblast proliferation and promote fibroblast inflammation during cardiac injury. *Mol Ther* **25**, 192–204.
- 144 Ramanujam D, Schön AP, Beck C, Vaccarello P, Felician G, Dueck A, Esfandyari D, Meister G, Meitinger T, Schulz C *et al.* (2021) MicroRNA-21-dependent macrophage-to-fibroblast signaling determines the cardiac response to pressure overload. *Circulation* **143**, 1513–1525.
- 145 Dick SA, Macklin JA, Nejat S, Momen A, Clemente-Casares X, Althagafi MG, Chen J, Kantores C, Hosseinzadeh S, Aronoff L *et al.* (2019) Self-renewing resident cardiac macrophages limit adverse remodeling following myocardial infarction. *Nat Immunol* **20**, 29–39.
- 146 Liu X-H, Pan L-L, Deng H-Y, Xiong Q-H, Wu D, Huang G-Y, Gong Q-H and Zhu Y-Z (2013) Leonurine (SCM-198) attenuates myocardial fibrotic response via inhibition of NADPH oxidase 4. *Free Radic Biol Med* **54**, 93–104.
- 147 Qin F, Simeone M and Patel R (2007) Inhibition of NADPH oxidase reduces myocardial oxidative stress and apoptosis and improves cardiac function in heart failure after myocardial infarction. *Free Radic Biol Med* **43**, 271–281.
- 148 Bendall JK, Cave AC, Heymes C, Gall N and Shah AM (2002) Pivotal role of a gp91(phox)-containing NADPH oxidase in angiotensin II-induced cardiac hypertrophy in mice. *Circulation* **105**, 293–296.
- 149 Ma Y, Yabluchanskiy A and Lindsey ML (2013) Neutrophil roles in left ventricular remodeling following myocardial infarction. *Fibrogen Tissue Repair* **6**, 11.
- 150 Liu J, Yang C, Liu T, Deng Z, Fang W, Zhang X, Li J, Huang Q, Liu C, Wang Y *et al.* (2020) Eosinophils improve cardiac function after myocardial infarction. *Nat Commun* **11**, 6396.
- 151 Diny NL, Hou X, Barin JG, Chen G, Talor MV, Schaub J, Russell SD, Klingel K, Rose NR and Čiháková D (2016) Macrophages and cardiac fibroblasts are the main producers of eotaxins and regulate eosinophil trafficking to the heart. *Eur J Immunol* **46**, 2749–2760.
- 152 Uezu A and Soderling S (2019) Identifying synaptic proteins by *in vivo* BioID from mouse brain. *Methods Mol Biol* **2008**, 107–119.