

1 **Effects of microplastics mixed with natural particles on *Daphnia magna* populations**

2

3 Christoph Schür^{a, b}, Joana Beck^b, Scott Lambert^b, Christian Scherer^{b, c}, Jörg Oehlmann^b,
4 Martin Wagner^{d*}

5

6 ^a Department of Environmental Toxicology, Eawag, Überlandstrasse 133, 8600 Dübendorf,
7 Switzerland

8 ^b Department Aquatic Ecotoxicology, Faculty of Biological Sciences, Goethe University,
9 Frankfurt am Main, Max-von-Laue-Str. 13, 60438 Frankfurt am Main, Germany

10 ^c Federal Institute of Hydrology, Department Biochemistry and Ecotoxicology, Am Mainzer
11 Tor 1, 56002, Koblenz, Germany

12 ^d Department of Biology, Norwegian University of Science and Technology, Høgskoleringen
13 5, 7491 Trondheim, Norway

14

15

16 * Corresponding author: Martin Wagner, martin.wagner@ntnu.no, +47 73 41 30 47

17 **Abstract**


18 The toxicity of microplastics on *Daphnia magna* as a key model for freshwater zooplankton is
19 well described. While several studies predict population-level effects based on short-term,
20 individual-level responses, only very few have validated these predictions experimentally.
21 Thus, we exposed *D. magna* populations to irregular polystyrene microplastics and diatomite
22 as natural particle (both $\leq 63 \mu\text{m}$) over 50 days. We used mixtures of both particle types at
23 fixed particle concentrations ($50,000 \text{ mL}^{-1}$) and recorded the effects on overall population size
24 and structure, the size of the individual animals, and resting egg production. Particle exposure
25 adversely affected the population density and structure, and induced resting egg production.
26 The terminal population size was 28–42% lower in exposed compared to control populations.
27 Interestingly, mixtures containing diatomite induced stronger effects than microplastics alone,
28 highlighting that natural particles are not *per se* less toxic than microplastics. Our results
29 demonstrate that an exposure to synthetic and natural particles has negative population-level
30 effects on zooplankton. Understanding the mixture toxicity of microplastics and natural
31 particles is important given that aquatic organisms will experience exposure to both. Just as
32 for chemical pollutants, better knowledge of such joint effects is essential to fully understand
33 the environmental impacts of complex particle mixtures.

34 Environmental Implications

35 While microplastics are commonly considered hazardous based on individual-level effects,
36 there is a dearth of information on how they affect populations. Since the latter is key for
37 understanding the environmental impacts of microplastics, we investigated how particle
38 exposures affect the population size and structure of *Daphnia magna*. In addition, we used
39 mixtures of microplastics and natural particles because neither occurs alone in nature and
40 joint effects can be expected in an environmentally realistic scenario. We show that such
41 mixtures adversely affect daphnid populations and highlight that population-level and
42 mixture-toxicity designs are one important step towards more environmental realism in
43 microplastics research.

44

45 Graphical Abstract



46

47 **Highlights**

- 48 ● *Daphnia* populations exposed to mixtures of microplastics and diatomite
- 49 ● Effects on population density, structure, and resting egg production
- 50 ● Diatomite as natural particle more toxic than microplastics
- 51 ● Particle mixtures induce negative population-level effects
- 52 ● Particle mixtures represent more realistic exposure scenario

53

54 **Keywords**

55 particulate matter, population dynamics, suspended matter

56 **Introduction**

57 Microplastics (MP) are a ubiquitous pollutant in the aquatic environment. They can interact
58 with and affect a broad range of species across all levels of biological organization, including
59 zooplankton such as the Cladoceran *Daphnia magna*. In the environment, MP are only one
60 type of non-food particles organisms interact with and MP as well as naturally occurring
61 particles have been shown to negatively affect daphnids, sometimes across generations (Kirk
62 1991b; 1991a; Robinson, Capper, and Klaine 2010; Ogonowski et al. 2016; Rist, Baun, and
63 Hartmann 2017; Martins and Guilhermino 2018; Schür et al. 2020). Nonetheless, as non-
64 selectively filter-feeding organisms, daphnids are well-adapted to non-food particles. This is
65 achieved through a number of behavioral and physiological mechanisms, including a
66 reduction in feeding rate, regurgitation of boluses, and the ability to remove adhering particles
67 from the filtering setae via the post-abdominal claw (Burns 1968a; 1968b; Kirk 1991a;
68 Ogonowski et al. 2016). Since exposure in the environment is never to a single type of
69 particle (synthetic or natural) and their effects in comparison to MP are often overlooked,
70 exposing animals to particle mixtures can be considered more environmentally relevant
71 (Gerdes et al. 2018; 2019). Additionally, the currently available literature is strongly biased
72 towards acute exposure scenarios, even though, due to their short generation time and the
73 environmental persistence of MP, daphnids are exposed continuously over generations and
74 not just intermittently (Rozman and Kalčíkova 2021; Yin et al. 2023). Thus, a long-term,
75 continuous exposure throughout an individual's lifetime, as well as following generations, is a
76 more realistic scenario (Schür et al. 2020; 2021). Daphnids as r-strategists form large, often
77 short-lived, populations. Population growth rates are high, but quickly reach a carrying
78 capacity limited by space and/or food. Such stressors are then often met with the formation of
79 resting eggs that can resurrect the population once conditions have returned to a more
80 favorable state (Smirnov 2017). In accordance with these considerations, we designed an
81 experiment in which *D. magna* populations with a defined age structure and size were
82 continuously exposed to mixtures of MP and the natural particle diatomite at constant particle
83 numbers and constant food levels. The aim of this study was to compare the effects of MP to
84 natural particles and their mixtures on the population level in a more realistic scenario.

85 **Materials and methods**

86 **Daphnia culture**

87 Ten *D. magna* individuals were cultured in 1 L of Elendt M4 medium (OECD 2012) at 20 °C
88 with a 16:8 h light:dark cycle. The daphnids were fed with the green algae *Desmodesmus*
89 *subspicatus* thrice a week at 0.2 mg carbon per individual per day (mgC daphnid⁻¹ d⁻¹). The
90 medium was fully renewed once a week.

91 **Particle preparation**

92 The irregularly shaped MP were produced from polystyrene coffee-to-go-cup lids obtained
93 from a local bakery. They were rinsed, cut into pieces using scissors, frozen in liquid nitrogen
94 and ground up in a ballmill (Retsch MM400, Retsch GmbH, Germany) at 30 Hz for 30 s.
95 Diatomite was purchased from Sigma Aldrich (CAS: 91053-39-3). Both particle types were
96 sieved to $\leq 63 \mu\text{m}$ using a sediment shaker (Retsch AS 200 basic, Retsch GmbH, Germany) to
97 achieve particles in a size range that is available for daphnids for ingestion (Scherer et al.
98 2018). Particle size distributions within the measuring margins of the Coulter counter
99 (Multisizer 3, Beckman Coulter, Germany; orifice tube with 100 μm aperture diameter for a
100 particle size range of 2.0–60 μm ; measurements in filtrated ($< 0.2 \mu\text{m}$) 0.98% NaCl solution)
101 are given in the supplementary material (Figure S1). Size distributions for the diatomite size
102 fraction $< 2 \mu\text{m}$ for a suspension prepared with the same method are given in the
103 supplementary materials of Scherer et al. (2019) but were not measured for this study.
104 Furthermore, Scanning electron microscope images of both particle types were taken using a
105 Hitachi S-4500 scanning electron microscope (supplementary material, Figure S2). For that,
106 20 μL of each suspension was transferred to the sample holder, dried under a heat lamp, and
107 sputtered with gold before imaging. Additional characterization of similar materials and
108 particle types can be found in Schür et al. (2021) and Scherer et al. (2019). Exposure
109 suspensions were prepared by dilution in Elendt M4 medium based on measured particle
110 concentrations of the stock solutions (Multisizer 3, Beckman Coulter) and used throughout
111 the experiment. Previous experiments, described in Schür et al. (2020), showed a good
112 correlation between nominal and measured particle concentrations. A new MP stock
113 suspension was prepared after day 37. Fourier transform infrared spectroscopy (ATR-FTIR)
114 spectra (FTIR Spectrum Two, PerkinElmer; LiTaO₃ detector, range: 4000–450 cm^{-1}) of the
115 raw plastic material before and after grinding and sieving are given in Figure S3 of the
116 supplementary material.

118 **Experimental design**

119 The initial daphnid populations consisted of 3 adults (2 weeks old), 5 juveniles (1 week old),
120 and 8 neonates (< 72 h old) held in 1 L glass vessels containing 900 mL Elendt M4 medium
121 (OECD 2012). Each population was kept for 50 d and fed a constant ration of 0.5 mgC d⁻¹ of
122 *D. subspicatus*. All treatment groups were exposed to a total of 50,000 particles mL⁻¹ of
123 varying ratios of MP and diatomite (n = 3, Table 1).

124 Populations were fed thrice per week, and the medium was exchanged on days 7, 14, 21, 28,
125 37, 42, and 50. During each feeding, vessels were covered with a lid and gently inverted to re-
126 suspend the particles. With each medium exchange, populations were sieved, transferred to an
127 hourglass, and photographed. ImageJ (Schindelin et al. 2012) was used to then quantify living
128 animals (total population size, Figure 1, Figure S4), the number of resting eggs (Figure 2,
129 Figure S5), body lengths (Figure 3, Figure S6). Resting eggs are seen as indicators of
130 population stress like insufficient food or high population density (Smirnov 2017). Individual
131 body lengths were measured from the center of the eye to the base of the apical spinus
132 (Ogonowski et al. 2016). Body lengths were categorized into three size/age classes in
133 accordance with Agatz et al. (2015), including neonates (≤ 1400 μm), juveniles (1400–2600
134 μm), and adults (> 2600 μm).

135

136 **Table 1: Ratios and absolute nominal particle concentrations of microplastics and**
137 **diatomite in the treatment groups of the population experiment.**

Treatment group	Microplastics		Diatomite	
	%	Particles mL ⁻¹	%	Particles mL ⁻¹
Control	0	0	0	0
MP100	100	50,000	0	0
MP80	80	40,000	20	10,000
MP60	60	30,000	40	20,000
MP50	50	25,000	50	25,000
MP40	40	20,000	60	30,000
MP20	20	10,000	80	40,000
MP0	0	0	100	50,000

138

139

140 **Statistical analysis**

141 The data was visualized using R (R Core Team 2021) with RStudio 2021.09.2+382 and the
142 tidyverse package (Wickham et al. 2019). The impact of exposure time and treatment on
143 population sizes (i.e., total number of animals at a given time) and structure was analyzed
144 using a one-way ANOVA with Holm-Šídák's multiple comparison test against the
145 corresponding control group for each time point in GraphPad Prism for Mac 9.3.1. The
146 number of resting eggs on day 50 of the experiment was compared against the control group
147 using a one-way ANOVA with Holm-Šídák's multiple comparisons test in GraphPad Prism
148 for Mac 9.3.1. The body length of individuals in each population was compared using
149 Kruskal-Wallis tests followed by Dunn's multiple comparison tests. Boxplots are created with
150 the `geom_boxplot()` function of the `ggplot2` package (Wickham 2016) in accordance with
151 McGill et al. (1978). Significance levels are indicated by asterisks as follows: * $p < 0.05$, ** p
152 < 0.01 , *** $p < 0.001$.

153 **Results**


154 Following the goal to investigate the population-level effects of mixtures of MP and natural
155 particles, the experiment included three main endpoints: population size (i.e., total number of
156 individuals per population at each time point), population structure based on the body lengths
157 of the individuals comprising each population, and the number of resting eggs (*ephippiae*) per
158 population.

159 All populations, both exposed to particles and of the control group, grew rapidly with regards
160 to the number of individuals during the first two weeks, with little variability between the
161 three replicates per treatment group (Figure 1). This is because the available food was
162 sufficient for such small populations coupled with low initial population densities acting as
163 triggers for rapid population growth. All population sizes peaked at day 14, declined from day
164 21 onwards, and reached their lowest size on day 50.

165 We observed a concentration-dependent effect in the populations exposed to particles in such
166 that in the phase of rapid decline (days 21–37), daphnid populations exposed to particle
167 mixtures that contained more diatomite had a lower population size (Figure 1, Figure S4). For
168 instance, populations exposed to 100% diatomite (MP0) consisted of significantly fewer
169 animals than the control populations on day 14 and 28 ($p < 0.05$, one-way ANOVA). At the
170 end of the experiment, on day 42 and 50, the size of all populations exposed to MP, diatomite
171 or mixtures thereof was significantly smaller than the control populations ($p < 0.05$). Notably,
172 the terminal population size in all treatments was 28–42% lower compared to control.

173 Resting egg formation occurred in all populations, including controls, after day 14 but to
174 varying degrees (Figure S5). Since the production of resting eggs is a stress response
175 (Smirnov 2017), this indicates a rapid onset of stress caused by increasing population
176 densities and/or decreasing food levels. The particle exposure had a significant effect on the
177 total number of resting eggs produced, with the populations in the MP60 ($p = 0.023$), MP40
178 ($p = 0.003$), MP20 ($p = 0.011$), and MP0 ($p = 0.008$) groups producing circa 100 ephippiae
179 compared to 70 in the control populations (Figure 2). Similar to the effect on intermediate
180 population sizes, this points towards a stronger effect of diatomite compared to MP.

181




182

183 **Figure 1: Boxplots of the population size of *Daphnia magna* exposed to polystyrene**
 184 **microplastics (MP100), diatomite (MP0), or their mixtures over 50 d (A). B–H represent**

185 the population sizes after 7, 14, 21, 28, 37, 42 and 50 d, respectively. Significant differences
186 compared to control populations are indicated by asterisks: * $p < 0.05$, ** $p < 0.01$, *** $p <$
187 0.001.

188




189

190 **Figure 2: Total number of resting eggs produced by *Daphnia magna* populations**
191 **exposed to polystyrene microplastics (MP100), diatomite (MP0), or their mixtures over**
192 **50 d (n = 3). Significant differences compared to control populations are indicated by**
193 **asterisks: * $p < 0.05$, ** $p < 0.01$.**

194

195 We measured the body length of each individual in a population weekly and used this to
196 describe the population structure by categorizing the daphnids into neonates, juveniles, and
197 adults. The initial population growth is largely driven by the production of neonates
198 (Figure 3). As a result of the lower reproduction from day 14 onwards, the population
199 structure shifts towards juveniles and adults. Particle exposure affected the number of
200 neonates and juveniles in the populations (Tables S2 and S4): At day 28, populations exposed
201 to diatomite or diatomite mixed with 20, 40, 60 and 80% MP consisted of significantly fewer
202 neonates. This effect translated into significantly fewer juveniles in all particle-exposed
203 populations on day 50. The number of adults was low compared to the other size classes and
204 increased over the course of the experiment without significant impact of the particle

205 exposure. In accordance with the loss of neonates and juveniles, individuals in particle-
206 exposed populations were in many cases significantly larger than in control populations
207 (Kruskal-Wallis tests, Table S6).



208

209 **Figure 3: Population structure of *Daphnia magna* exposed to polystyrene microplastics**
210 **(MP100), diatomite (MP0), or their mixtures over 50 d. Data presented as mean relative**

211 ratios of neonates (green), juveniles (blue), and adults (red) compared to the overall
212 population density ($n = 3$).

213 **Discussion**

214 We exposed *D. magna* populations to 50,000 particles mL⁻¹ of either polystyrene MP,
215 diatomite, or mixtures of both over the course of 50 d. Notwithstanding the ratio of MP to
216 diatomite, particle exposure significantly reduced the population size from day 43 onwards
217 and resulted in populations consisting of 28–42% less individuals than control populations at
218 the end of the experiment. This effect on population size is most likely due to particle
219 exposures having a negative impact on reproduction as previously shown by Ogonowski et al.
220 (2016) and Schür et al. (2020), especially after the population size has peaked. The
221 reproductive toxicity of particles is also reflected in the population structure with particle-
222 exposed populations consisting of larger and, thus, older individuals than control populations.
223 Taken together, this demonstrates that mixtures of synthetic and natural particles have
224 negative effects at the population level in *D. magna*.

225 The fact that MP as well as their mixtures with natural particles affected the population size
226 and structure highlights that the well-documented individual-level toxicity of MP and other
227 particles in daphnids translates into impacts at the population level. While multiple studies
228 predict effects of MP exposures on population growth rates based on individual level
229 responses (*e.g.*, Martins and Guilhermino (2018); Guilhermino et al. (2021)), population and
230 multigeneration effects were recently identified as severe data gaps in a review on the
231 ecotoxicology of MP in *Daphnia* (Yin et al. 2023). Bosker et al. (2019) reported that exposure
232 to polystyrene beads caused a significant decline in population size and biomass but did not
233 affect the size of individuals or *ephippiae* production. Besides using another type of MP, their
234 general approach was different from ours as they grew populations to holding capacity before
235 starting particle exposure at day 30. This probably reduced the overall stress level induced by
236 continuous particle exposures.

237 Zebrowski et al. (2022) investigated how the exposure to PS, high-density polyethylene (PE),
238 and the assumed biodegradable polyhydroxybutyrate (PHB) affected the growth (measured by
239 population density, *i.e.* individuals per L) of competing populations of *D. magna*, *Daphnia*
240 *pulex*, and *Daphnia galeata* under constant food levels (two species were paired in
241 competition). While the outcome of the competition experiments was not affected (the same
242 species outperformed their competitors as in the particle-free control treatments), two main
243 findings are worth mentioning: (I) The larger *D. magna* did not always outcompete the
244 smaller cladoceran species, but only the smallest *D. galeata*, while *D. pulex* consistently
245 persisted against both other species, and (II) *D. magna* and *D. pulex* populations were affected

246 by PS particle exposure, compared to the respective control groups, while *D. galeata*
247 populations grew very similarly to their control populations. Some of their results hint
248 towards a positive effect of the PHB exposure, about which the authors hypothesize that the
249 material is biodegraded and serves as nutrients for bacteria in the exposure vessel, which
250 subsequently improve the nutritional status of the cladocerans.

251 Al-Jaibachi et al. (2019) observed the initial decline but subsequent recovery of daphnid
252 populations in MP-exposed mesocosms, while no effect on other species was observed. Here,
253 high variability and unknown influencing factors from the mesocosm setup impede the
254 comparison between the two studies. Nonetheless, all three studies demonstrate that MP
255 effects also manifest on the population level, which is considered highly relevant for assessing
256 the environmental risks of these particles.

257 We used multiple mixtures of MP and diatomite at a fixed numerical concentration to explore
258 a more realistic exposure scenario (i.e., MP as part of a more diverse set of suspended solids)
259 and to investigate whether the mixtures' toxicity is driven by plastic or natural particles.
260 Indeed, our results show that diatomite is more toxic to daphnid populations than MP. With
261 regards to the intermediate population size, resting egg production, and population structure,
262 exposure to pure diatomite induced stronger effects than to pure MP (Figures 1–3). In the
263 treatments with particle mixtures, we often observed a concentration-dependent response with
264 mixtures containing more diatomite being more toxic. This is particularly obvious for the
265 population sizes at days 14–28 and the resting egg production. Accordingly, particle mixtures
266 consisting of more diatomite are more toxic.

267 The reason for the higher toxicity of diatomite compared to MP may be its porous and spiky
268 structure. Diatomite has biocidal properties (European Food Safety Authority (EFSA) et al.
269 2020) and its absorptive and abrasive capacities will damage insect cuticles (Korunic 1998)
270 and may injure the digestive system (Scherer et al. 2019). Diatomite has been used as a
271 natural reference material in previous MP studies. In the freshwater mollusks *Dreissena*
272 *polymorpha* and *Lymnea stagnalis*, diatomite was in general not more toxic than polystyrene
273 MP (Weber, von Randow, et al. 2021; Weber, Jeckel, et al. 2021) but induced a stronger
274 effect on the antioxidant capacity in the former species (Weber, Jeckel, and Wagner 2020) at
275 identical numerical concentrations. In *Chironomus riparius* larvae, diatomite was toxic but
276 less so than polyvinyl chloride MP at identical mass-based concentrations (Scherer et al.
277 2019). Since one of the main mechanisms of its toxicity appears to be the desorption of waxes

278 from the cuticle, arthropods, such as chironomids and daphnids, may be particularly sensitive
279 to diatomite exposures.

280 Our study shows that some natural particles can be more toxic than a mixture of natural
281 particles and MP or MP by themselves. Earlier work compared the effects of the natural
282 particle kaolin with polystyrene MP similar to those used in this study in a multigenerational
283 study with daphnids (Schür et al. 2020). There, we found that kaolin had no effect, while MP
284 affected all recorded endpoints in a concentration-dependent manner with effects increasing
285 over generations. This shows that transferring findings on one particle type to another is not
286 straightforward and MP may be more toxic than some but not all natural particles. Particle
287 shape may play an important role as diatomite is spiky and sharp compared to kaolin which is
288 rather round. Thus, the toxicity of natural particles will depend on their individual set of
289 physicochemical properties and cannot be easily generalized without a better mechanistic
290 understanding (see Scherer et al. (2019) for an in-depth discussion).

291 Finally, our study was not designed to mimic environmental concentrations of MP or natural
292 particles. Instead, our aim was to investigate the toxicity of mixtures of both, because this
293 exposure scenario is more realistic compared to the use of only MP in toxicity studies. Given
294 that in nature, aquatic organisms will most likely be exposed to natural and synthetic
295 particulate matter concurrently, a better understanding of the joint toxicity is needed to
296 develop realistic predictions of environmental impacts.

297

298 **Conclusions**

299 Our study demonstrates that an exposure to MP and diatomite alone as well as in mixture has
300 negative population-level effects in *D. magna*. This corroborates previous predictions based
301 on individual-level responses. Our findings are relevant because adverse impacts on
302 populations of a keystone zooplankton species will have ecological consequences. However,
303 the fact that we used one very high particle concentration only calls for follow-up studies to
304 generate concentration-response relationships. We used mixtures of plastic and the natural
305 particle diatomite because we deem this exposure scenario more realistic and found that
306 diatomite is more toxic than MP. This contradicts the common assumption that natural
307 particles are benign and highlights that – just as with MP – the toxicity of a particle type
308 depends on its individual set of physicochemical properties. This calls into question whether
309 general comparisons, such as MP are more or less toxic than something else, are meaningful.
310 It also highlights the challenge of finding an adequate reference particle when attempting to

311 perform such comparisons. Finally, we believe that investigating the mixture toxicity of
312 synthetic and natural particles is valuable given that aquatic organisms will experience
313 exposure to both. Similar to chemical pollutants, better knowledge of such joint effects is
314 essential to fully understand the environmental risks complex particle mixtures pose to
315 aquatic species.

316 **Author contributions**

317 Christoph Schür: Conceptualization, Data curation, Formal analysis, Investigation,
318 Methodology, Validation, Visualization, Project administration, Writing - original draft,
319 Writing - review & editing

320 Joana Beck: Data curation, Investigation, Writing - review & editing

321 Scott Lambert: Conceptualization, Methodology, Writing - review & editing

322 Christian Scherer: Conceptualization, Methodology, Investigation, Writing - review & editing

323 Jörg Oehlmann: Funding acquisition, Project administration, Resources, Writing - review &
324 editing

325 Martin Wagner: Conceptualization, Formal analysis, Funding acquisition, Resources, Project
326 administration, Visualization, Resources, Writing - review & editing

327 **Declaration of interest**

328 Martin Wagner is an unremunerated member of the Scientific Advisory Board of the Food
329 Packaging Forum (FPF). He has received travel funding from FPF to attend its annual board
330 meetings and from Hold Norge Rent (Keep Norway Beautiful) to speak at one of their
331 conferences. The other authors declare no conflict of interest.

332 **Acknowledgments**

333 This study was supported by the German Federal Ministry for Education and Research to CS,
334 JO, and MW (02WRS1378I, 03F0789D). The graphical abstract was created with BioRender.

335 **Supplementary Material**

336 **The supplemental data are available ###.**

337 References

- 338 Agatz, Annika et al. 2015. “Family-Portraits for Daphnids: Scanning Living Individuals and
339 Populations to Measure Body Length.” *Ecotoxicology* 24 (6): 1385–94.
340 <https://doi.org/10.1007/s10646-015-1490-0>.
- 341 Al-Jaibachi, Rana et al. 2019. “Impacts of Polystyrene Microplastics on *Daphnia Magna*: A
342 Laboratory and a Mesocosm Study.” *Science of The Total Environment*, 135800.
343 <https://doi.org/10.1016/j.scitotenv.2019.135800>.
- 344 Bosker, Thijs et al. 2019. “Significant Decline of *Daphnia Magna* Population Biomass Due to
345 Microplastic Exposure.” *Environmental Pollution*.
346 <https://doi.org/10.1016/j.envpol.2019.04.067>.
- 347 Burns, Carolyn W. 1968a. “Direct Observations of Mechanisms Regulating Feeding Behavior
348 of *Daphnia*, in Lakewater.” *Internationale Revue Der Gesamten Hydrobiologie Und*
349 *Hydrographie* 53 (1): 83–100. <https://doi.org/10.1002/iroh.19680530104>.
- 350 ———. 1968b. “The Relationship between Body Size of Filter-Feeding Cladocera and the
351 Maximum Size of Particle Ingested.” *Limnology and Oceanography* 13 (4): 675–78.
352 <https://doi.org/10.4319/lo.1968.13.4.0675>.
- 353 European Food Safety Authority (EFSA) et al. 2020. “Peer Review of the Pesticide Risk
354 Assessment of the Active Substance Kieselgur (Diatomaceous Earth).” *EFSA Journal*
355 18 (3). <https://doi.org/10.2903/j.efsa.2020.6054>.
- 356 Gerdes, Zandra et al. 2018. “A Serial Dilution Method for Assessment of Microplastic
357 Toxicity in Suspension.”
358 ———. 2019. “A Novel Method for Assessing Microplastic Effect in Suspension through
359 Mixing Test and Reference Materials.” *Scientific Reports* 9 (1).
360 <https://doi.org/10.1038/s41598-019-47160-1>.
- 361 Guilhermino, Lúcia et al. 2021. “Long-Term Adverse Effects of Microplastics on *Daphnia*
362 *Magna* Reproduction and Population Growth Rate at Increased Water Temperature
363 and Light Intensity: Combined Effects of Stressors and Interactions.” *Science of The*
364 *Total Environment* 784 (August): 147082.
365 <https://doi.org/10.1016/j.scitotenv.2021.147082>.
- 366 Kirk, Kevin L. 1991. “Suspended Clay Reduces *Daphnia* Feeding Rate: Behavioural
367 Mechanisms.” *Freshwater Biology* 25 (2): 357–65. <https://doi.org/10.1111/j.1365-2427.1991.tb00498.x>.
- 369 Korunic, Zlatko. 1998. “Diatomaceous Earths, a Group of Natural Insecticides.” *Journal of*
370 *Stored Products Research* 34 (2–3): 87–97. [https://doi.org/10.1016/S0022-474X\(97\)00039-8](https://doi.org/10.1016/S0022-474X(97)00039-8).
- 372 Martins, Alexandra, and Lúcia Guilhermino. 2018. “Transgenerational Effects and Recovery
373 of Microplastics Exposure in Model Populations of the Freshwater Cladoceran
374 *Daphnia Magna* Straus.” *Science of The Total Environment* 631–632: 421–28.
375 <https://doi.org/10.1016/j.scitotenv.2018.03.054>.
- 376 McGill, Robert, John W. Tukey, and Wayne A. Larsen. 1978. “Variations of Box Plots.” *The*
377 *American Statistician* 32 (1): 12–16.
378 <https://doi.org/10.1080/00031305.1978.10479236>.
- 379 OECD. 2012. *Test No. 211: Daphnia Magna Reproduction Test*. Paris: Organisation for
380 Economic Co-operation and Development. [http://www.oecd-](http://www.oecd-ilibrary.org/content/book/9789264185203-en)
381 [ilibrary.org/content/book/9789264185203-en](http://www.oecd-ilibrary.org/content/book/9789264185203-en).
- 382 Ogonowski, Martin et al. 2016. “The Effects of Natural and Anthropogenic Microparticles on
383 Individual Fitness in *Daphnia Magna*.” *PLOS ONE* 11 (5): e0155063.
384 <https://doi.org/10.1371/journal.pone.0155063>.
- 385 R Core Team. 2021. *R: A Language and Environment for Statistical Computing*. Vienna,
386 Austria: R Foundation for Statistical Computing. <https://www.R-project.org/>.

- 387 Rist, Sinja, Anders Baun, and Nanna B. Hartmann. 2017. “Ingestion of Micro- and
388 Nanoplastics in *Daphnia Magna* – Quantification of Body Burdens and Assessment of
389 Feeding Rates and Reproduction.” *Environmental Pollution* 228 (September): 398–
390 407. <https://doi.org/10.1016/j.envpol.2017.05.048>.
- 391 Robinson, Sarah E., Neil A. Capper, and Stephen J. Klaine. 2010. “The Effects of Continuous
392 and Pulsed Exposures of Suspended Clay on the Survival, Growth, and Reproduction
393 of *Daphnia Magna*.” *Environmental Toxicology and Chemistry* 29 (1): 168–75.
394 <https://doi.org/10.1002/etc.4>.
- 395 Rozman, Ula, and Gabriela Kalčíková. 2021. “Seeking for a Perfect (Non-Spherical)
396 Microplastic Particle – The Most Comprehensive Review on Microplastic Laboratory
397 Research.” *Journal of Hazardous Materials*, 127529.
398 <https://doi.org/10.1016/j.jhazmat.2021.127529>.
- 399 Scherer, Christian et al. 2018. “Interactions of Microplastics with Freshwater Biota.” In
400 *Freshwater Microplastics*, 153–80. The Handbook of Environmental Chemistry.
401 Springer, Cham. https://doi.org/10.1007/978-3-319-61615-5_8.
- 402 ———. 2019. “Toxicity of Microplastics and Natural Particles in the Freshwater Dipteran
403 *Chironomus Riparius*: Same Same but Different?” *Science of The Total Environment*
404 711: 134604. <https://doi.org/10.1016/j.scitotenv.2019.134604>.
- 405 Schneider, Caroline A., Wayne S. Rasband, and Kevin W. Eliceiri. 2012. “NIH Image to
406 ImageJ: 25 Years of Image Analysis.” *Nature Methods* 9 (7): 671–75.
407 <https://doi.org/10.1038/nmeth.2089>.
- 408 Schür, Christoph et al. 2020. “Microplastics but Not Natural Particles Induce
409 Multigenerational Effects in *Daphnia Magna*.” *Environmental Pollution* 260: 113904.
410 <https://doi.org/10.1016/j.envpol.2019.113904>.
- 411 ———. 2021. “Incubation in Wastewater Reduces the Multigenerational Effects of
412 Microplastics in *Daphnia Magna*.” *Environmental Science & Technology*,
413 [acs.est.0c07911](https://doi.org/10.1021/acs.est.0c07911). <https://doi.org/10.1021/acs.est.0c07911>.
- 414 Smirnov, N. N. 2017. *Physiology of the Cladocera*. Second edition. London, United
415 Kingdom; San Diego, CA, United States: Elsevier/AP, Academic Press, an imprint
416 of Elsevier.
- 417 Weber, Annkatrin, Nina Jeckel, and Martin Wagner. 2020. “Combined Effects of Polystyrene
418 Microplastics and Thermal Stress on the Freshwater Mussel *Dreissena Polymorpha*.”
419 *Science of The Total Environment* 718: 137253.
420 <https://doi.org/10.1016/j.scitotenv.2020.137253>.
- 421 Weber, Annkatrin, Marvin von Randow, et al. 2021. “Ingestion and Toxicity of Microplastics
422 in the Freshwater Gastropod *Lymnaea Stagnalis*: No Microplastic-Induced Effects
423 Alone or in Combination with Copper.” *Chemosphere* 263: 128040.
424 <https://doi.org/10.1016/j.chemosphere.2020.128040>.
- 425 Weber, Annkatrin, Nina Jeckel, et al. 2021. “Ingestion and Toxicity of Polystyrene
426 Microplastics in Freshwater Bivalves.” *Environmental Toxicology and Chemistry*,
427 etc.5076. <https://doi.org/10.1002/etc.5076>.
- 428 Wickham, Hadley. 2016. *Ggplot2: Elegant Graphics for Data Analysis*. Springer-Verlag New
429 York. <https://ggplot2.tidyverse.org>.
- 430 ———. 2019. “Welcome to the Tidyverse.” *Journal of Open Source Software* 4 (43): 1686.
431 <https://doi.org/10.21105/joss.01686>.
- 432