

Appendix 3: Reference list of included studies

*full text available in German language only

- *Abholz H.-H. (2006): Prevention in general practice - A proposal for a systematic approach. In: *Zeitschrift fur Allgemeinmedizin* 82 (2), S. 50–55. DOI: 10.1055/s-2006-921444.
- Achelrod D.; Welte T.; Schreyogg J.; Stargardt T. (2016): Costs and outcomes of the German disease management programme (DMP) for chronic obstructive pulmonary disease (COPD)-A large population-based cohort study. In: *Health Policy* 120 (9), S. 1029–1039. DOI: 10.1016/j.healthpol.2016.08.002.
- Adarkwah C.C.; Jegan N.; Heinzel-Gutenbrunner M.; Kuhne F.; Siebert U.; Popert U. et al. (2019): The Optimizing-Rrisk-Ccommunication (OptRisk) randomized trial - impact of decision-aid-based consultation on adherence and perception of cardiovascular risk. In: *Patient Preference and Adherence* 13 S. 441–452. DOI: 10.2147/PPA.S197545.
- Akmatov, Manas K.; Rubsamen, Nicole; Deyneko, Igor V.; Karch, Andre; Mikolajczyk, Rafael T. (2018): Poor knowledge of vaccination recommendations and negative attitudes towards vaccinations are independently associated with poor vaccination uptake among adults - Findings of a population-based panel study in Lower Saxony, Germany. In: *Vaccine* 36 (18), S. 2417–2426. DOI: 10.1016/j.vaccine.2018.03.050.
- Alber K.; Kuehlein T.; Schedlbauer A.; Schaffer S. (2017): Medical overuse and quaternary prevention in primary care - A qualitative study with general practitioners. In: *BMC family practice* 18 (1), S. 99. DOI: 10.1186/s12875-017-0667-4.
- *Allenberg J.-R.; Trampisch H.-J.; Darius H.; Lange S.; Haberl R.; Stritzky B.V. et al. (2004): The prevalence, comorbidity and intensity of treatment of peripheral arterial disease in primary care. Results of the getABI study. In: *Gefasschirurgie* 9 (3), S. 166–171. DOI: 10.1007/s00772-004-0360-5.
- *Althaus, D.; Niklewski, G.; Schmidtke, A.; Hegerl, U. (2007): [Changes in the frequency of suicidal behaviour after a 2-year intervention campaign]. In: *Der Nervenarzt* 78 (3), S. 272–282.
- *Althaus D.; Hegerl U. (2001): Evaluating suicide-preventive activities: The present state of research and the resulting implications. In: *Nervenarzt* 72 (9), S. 677–684. DOI: 10.1007/s001150170046.
- *Althaus D.; Niklewski G.; Schmidtke A.; Felber W.; Kunz J.; Lehfeld H.; Hegerl U. (2003): Changes in the number of suicide attempts: Results after nine months of intervention of the Nuremberg Alliance against Depression. In: *PsychoNeuro* 29 (1), S. 28–34. DOI: 10.1055/s-2003-37615.
- *Altiner A.; Abholz H.-H. (2000): Antithrombotic treatment of atrial fibrillation - Application of evidence. In: *Zeitschrift fur Allgemeinmedizin* 76 (6), S. 280–284.
- Altiner A.; Brockmann S.; Sielk M.; Wilm S.; Wegscheider K.; Abholz H.-H. (2007): Reducing antibiotic prescriptions for acute cough by motivating GPs to change their attitudes to communication and empowering patients: A cluster-randomized intervention study. In: *Journal of Antimicrobial Chemotherapy* 60 (3), S. 638–644. DOI: 10.1093/jac/dkm254.
- Altiner A.; Schafer I.; Mellert C.; Löffler C.; Mortsiefer A.; Ernst A. et al. (2012): Activating GENeral practitioners dialogue with patients on their Agenda (MultiCare AGENDA) study protocol for a cluster randomized controlled trial. In: *BMC family practice* 13 S. 118. DOI: 10.1186/1471-2296-13-118.
- *Altpeter B. (2017): E-health as a component of holistic therapy optimization. In: *Diabetologe* 13 (1), S. 29–37. DOI: 10.1007/s11428-016-0172-8.
- Amelang K.; Bauer S. (2019): Following the algorithm: How epidemiological risk-scores do accountability. In: *Social studies of science* 49 (4), S. 476–502. DOI: 10.1177/0306312719862049.
- Anders, Markus P.; Fengler, Susanne; Volkmer, Beate; Greinert, Rudiger; Breitbart, Eckhard W. (2017): Nationwide skin cancer screening in Germany: Evaluation of the training program. In: *International journal of dermatology* 56 (10), S. 1046–1051. DOI: 10.1111/ijd.13688.
- *Anders J.; Dapp U.; Laub S.; von Renteln-Kruse W. (2007): Impact of fall risk and fear of falling on mobility of independently living senior citizens transitioning to frailty: Screening results concerning fall prevention in the community. In: *Zeitschrift fur Gerontologie und Geriatrie* 40 (4), S. 255–267. DOI: 10.1007/s00391-007-0473-z.
- *Andrees V.; Girbig G.; Krensel M.; Augustin M.; Zander N. (2020): Access to statutory skin cancer screening: Organizational and regional factors in Germany. In: *Hautarzt* 71 (1), S. 39–45. DOI: 10.1007/s00105-019-04494-1.
- *Andres, Edith; Bleek, Julian; Stock, Johannes; Bader, Erwin; Gunter, Alexander; Wambach, Veit et al. (2018): [Measuring, assessing, acting: A practice test of quality indicators for coronary heart disease]. In: *Zeitschrift fur Evidenz, Fortbildung und Qualitat im Gesundheitswesen* 137-138 (101477604), S. 9–19. DOI: 10.1016/j.zefq.2018.08.003.

*Angenendt, J. (2014): [Interventions for mental health sequelae of accidents]. In: *Bundesgesundheitsblatt, Gesundheitsforschung, Gesundheitsschutz* 57 (6), S. 666–672. DOI: 10.1007/s00103-014-1967-7.

Arnou, Robert; Frank, Matthias; Hagel, Thomas; Prebet, Alice (2011): Willingness to vaccinate or get vaccinated with an intradermal seasonal influenza vaccine: a survey of general practitioners and the general public in France and Germany. In: *Advances in therapy* 28 (7), S. 555–565. DOI: 10.1007/s12325-011-0035-z.

*Arntz H.-R.; Tebbe U. (2004): Acute coronary syndrome - What can the family doctor do? In: *MMW-Fortschritte der Medizin* 146 (15), S. 30–35.

Atmann O.; Linde K.; Werner C.; Dorn U.; Schneider A. (2019): Participation factors for asthma education programs - A cross sectional survey. In: *BMC Pulmonary Medicine* 19 (1), S. 256. DOI: 10.1186/s12890-019-0979-3.

*Augustin, Matthias; Stadler, Rudolf; Reusch, Michael; Schafer, Ines; Kornek, Thomas; Luger, Thomas (2012): Skin cancer screening in Germany - perception by the public. In: *Journal der Deutschen Dermatologischen Gesellschaft = Journal of the German Society of Dermatology : JDDG* 10 (1), S. 42–49. DOI: 10.1111/j.1610-0387.2011.07761.x.

Bachmann, Viktoria; Volkner, Michael; Bosnerr, Stefan; Donner-Banzhoff, Norbert (2014): The experiences of Russian-speaking migrants in primary care consultations. In: *Deutsches Arzteblatt International* 111 (51-52), S. 871–876. DOI: 10.3238/artztebl.2014.0871.

*Bahrs O.; Heim S. (2005): Theme-progression in case orientated peer review groups. In: *Zeitschrift fur Allgemeinmedizin* 81 (3), S. 121–126. DOI: 10.1055/s-2005-836310.

*Banke, I. J.; Eisenhart-Rothe, R. von; Muhlhofer, H. M. L. (2015): [Epidemiology and prevention of prosthetic joint infection]. In: *Der Orthopade* 44 (12), 928–3. DOI: 10.1007/s00132-015-3187-8.

*Barlinn, J.; Barlinn, K.; Helbig, U.; Siepmann, T.; Pallesen, L-P; Urban, H. et al. (2016): [Organized Post-Stroke Care through Case Management on the Basis of a Standardized Treatment Pathway : Results of a Single-Centre Pilot Study]. In: *Der Nervenarzt* 87 (8), S. 860–869. DOI: 10.1007/s00115-016-0100-6.

Barth, Cordula; Neusser, Silke; Biermann, Janine; Wasem, Jurgen; Berg, Thomas; Wiegand, Johannes et al. (2018): Budget impact analysis on the introduction of a guideline based hepatitis B and C screening into a routine check-up in the German primary care setting. In: *Scandinavian journal of gastroenterology* 53 (1), S. 76–82. DOI: 10.1080/00365521.2017.1390599.

*Bastian H.; Kaiser T.; Matschewsky S. (2005): Promotion of general health and scientific literacy via consumer and patient information: The role of the IQWiG. In: *Zeitschrift fur Arztliche Fortbildung und Qualitatssicherung* 99 (6), S. 379–385.

Bauer, Antonia; Grunewald, Martin; Eberhardt, Hans; Schulz, Rieke; Martus, Peter; Bruggenjurgen, Bernd et al. (2020): Ambulatory screening and decontamination to prevent *Staphylococcus aureus* complications in patients undergoing elective surgery (STAUFrei): study protocol for a controlled intervention study. In: *BMC Infectious Diseases* 20 (1), S. 95. DOI: 10.1186/s12879-020-4804-7.

*Baum E. (2002): Female climacterium and hormonal replacement therapy. In: *Zeitschrift fur Allgemeinmedizin* 78 (6-7), S. 308–313.

*Baumgart D.; Egelhof T. (2007): Preventive whole-body screening encompassing modern imaging using magnetic resonance tomography. In: *Herz* 32 (5), S. 387–394. DOI: 10.1007/s00059-007-3020-1.

Bayrhuber M.; Anka N.; Camp J.; Glattacker M.; Farin E.; Rieg S. (2020): Prevention of post-splenectomy sepsis in patients with asplenia - A study protocol of a controlled trial. In: *BMC Infectious Diseases* 20 (1), S. 41. DOI: 10.1186/s12879-019-4752-2.

*Becker, C.; Rapp, K. (2011): [Falling in geriatrics. Diagnosis and treatment]. In: *Der Internist* 52 (8), S. 939–945. DOI: 10.1007/s00108-011-2805-8.

Becker A.; Herzberg D.; Marsden N.; Thomanek S.; Jung H.; Leonhardt C. (2011): A new computer-based counselling system for the promotion of physical activity in patients with chronic diseases-Results from a pilot study. In: *Patient Education and Counseling* 83 (2), S. 195–202. DOI: 10.1016/j.pec.2010.05.024.

*Beesdo-Baum, Katja; Knappe, Susanne; Einsle, Franziska; Knothe, Lisa; Wieder, Gesine; Venz, John et al. (2018): [How frequently are depressive disorders recognized in primary care patients? : A cross-sectional epidemiological study in Germany]. In: *Bundesgesundheitsblatt, Gesundheitsforschung, Gesundheitsschutz* 61 (1), S. 52–64. DOI: 10.1007/s00103-017-2662-2.

Beinhoff U.; Hilbert V.; Bittner D.; Gron G.; Riepe M.W. (2005): Screening for cognitive impairment: A triage for outpatient care. In: *Dementia and Geriatric Cognitive Disorders* 20 (5), S. 278–285. DOI: 10.1159/000088249.

*Beltermann, Esther; Krane, Sibylla; Kiesewetter, Jan; Fischer, Martin R.; Schelling, Jorg (2015): See your GP, see the world - An activating course concept for fostering students' competence in performing vaccine and travel consultations. In: *GMS Zeitschrift fur medizinische Ausbildung* 32 (3), Doc28. DOI: 10.3205/zma000970.

Bentele, Michael; Kriston, Levente; Clement, Hans-Willi; Harter, Martin; Mundt, Gotz; Berner, Michael M. (2007): The validity of the laboratory marker combinations DOVER and QUVER to detect physician's diagnosis of at-risk drinking. In: *Addiction biology* 12 (1), S. 85–92.

- Berg-Beckhoff, Gabriele; Heyer, Kristina; Kowall, Bernd; Breckenkamp, Jurgen; Razum, Oliver (2010): The views of primary care physicians on health risks from electromagnetic fields. In: *Deutsches Arzteblatt International* 107 (46), S. 817–823. DOI: 10.3238/ärztebl.2010.0817.
- Berner M.M.; Harter M.; Kriston L.; Lohmann M.; Ruf D.; Lorenz G.; Mundt G. (2007): Detection and management of alcohol use disorders in German primary care influenced by non-clinical factors. In: *Alcohol and Alcoholism* 42 (4), S. 308–316. DOI: 10.1093/alcalc/agm013.
- Berthold, Heiner K.; Bestehorn, Kurt P.; Jannowitz, Christina; Krone, Wilhelm; Gouni-Berthold, Ioanna (2011): Disease management programs in type 2 diabetes: quality of care. In: *The American journal of managed care* 17 (6), S. 393–403.
- Bickel H.; Ander K.-H.; Bronner M.; Etgen T.; Gnahn H.; Gotzler O. et al. (2012): Reduction of long-term care dependence after an 8-year primary care prevention program for stroke and dementia: The INVADE trial. In: *Journal of the American Heart Association* 1 (4), e000786. DOI: 10.1161/JAHA.112.000786.
- *Bickel H.; Block M.; Gotzler O.; Hartmann J.; Purner K.; Sander D.; Forstl H. (2020): Prevention of stroke and dementia in general practice: Evaluation of the project INVADE. In: *Deutsche Medizinische Wochenschrift* 145 (12), E61–E70. DOI: 10.1055/a-1124-9124.
- Bischof, G.; Freyer-Adam, J.; Meyer, C.; John, U.; Rumpf, H-J (2012): Changes in drinking behavior among control group participants in early intervention studies targeting unhealthy alcohol use recruited in general hospitals and general practices. In: *Drug and alcohol dependence* 125 (1-2), S. 81–88. DOI: 10.1016/j.drugalcdep.2012.03.018.
- Bischof, Gallus; Reinhardt, Susa; Grothues, Janina; Meyer, Christian; John, Ulrich; Rumpf, Hans-Jurgen (2007): Development and evaluation of a screening instrument for alcohol-use disorders and at-risk drinking: the brief alcohol screening instrument for medical care (BASIC). In: *Journal of studies on alcohol and drugs* 68 (4), S. 607–614.
- Bischoff B.; Silber S.; Richartz B.M.; Pieper L.; Klotsche J.; Wittchen H.-U. (2006): Inadequate medical treatment of patients with coronary artery disease by primary care physicians in Germany. In: *Clinical Research in Cardiology* 95 (8), S. 405–412. DOI: 10.1007/s00392-006-0399-2.
- *Bjerregaard, F.; Hull, M.; Stieglitz, R-D; Holzel, L. P. (2018): [Time for Change: What We can Learn from the USA about Primary Care of the Depressed Elderly]. In: *Gesundheitswesen (Bundesverband der Ärzte des Öffentlichen Gesundheitsdienstes (Germany))* 80 (1), S. 34–39. DOI: 10.1055/s-0042-107344.
- *Bleckwenn, M.; Theisel, N.; Mucke, M.; Steudel, H. (2018): [Cardiovascular Prevention: Acceptance of Enhanced Occupational Health Care]. In: *Gesundheitswesen (Bundesverband der Ärzte des Öffentlichen Gesundheitsdienstes (Germany))* 80 (5), S. 495–498. DOI: 10.1055/s-0042-107348.
- Bock C.; Diehm C.; Schneider S. (2012): Physical activity promotion in primary health care: Results from a German physician survey. In: *European Journal of General Practice* 18 (2), S. 86–91. DOI: 10.3109/13814788.2012.675504.
- *Bodeker, B.; Wichmann, O.; Mertens, B.; Seefeld, L.; Pott, E. (2014): [Implementation of the influenza vaccination recommendation in nursing homes in Germany : results of a survey as part of the national influenza immunization campaign]. In: *Bundesgesundheitsblatt, Gesundheitsforschung, Gesundheitsschutz* 57 (11), S. 1315–1321. DOI: 10.1007/s00103-014-2044-y.
- Bohlken J.; Kostev K. (2019): Coded Prevalence of Mild Cognitive Impairment in General and Neuropsychiatrists Practices in Germany between 2007 and 2017. In: *Journal of Alzheimer's Disease* 67 (4), S. 1313–1318. DOI: 10.3233/JAD-181180.
- Bozorgmehr, Kayvan; Szecsenyi, Joachim; Ose, Dominik; Besier, Werner; Mayer, Manfred; Krisam, Johannes et al. (2014): Practice network-based care management for patients with type 2 diabetes and multiple comorbidities (GEDIMAPplus): study protocol for a randomized controlled trial. In: *Trials* 15 (101263253), S. 243. DOI: 10.1186/1745-6215-15-243.
- *Bramesfeld, A.; Schwartz, F. W. (2006): [Possibilities for intervention in depression among adolescents and young adults from a public health point of view]. In: *Gesundheitswesen (Bundesverband der Ärzte des Öffentlichen Gesundheitsdienstes (Germany))* 68 (8-9), S. 508–516.
- Brandt, Henning Christian; Spiller, Inge; Song, In-Ho; Vahldiek, Janis L.; Rudwaleit, Martin; Sieper, Joachim (2007): Performance of referral recommendations in patients with chronic back pain and suspected axial spondyloarthritis. In: *Annals of the rheumatic diseases* 66 (11), S. 1479–1484.
- Braun, Sarah R.; Reiner, Katinka; Tegeler, Christina; Bucholtz, Nina; Boustani, Malaz A.; Steinhagen-Thiessen, Elisabeth (2014): Acceptance of and attitudes towards Alzheimer's disease screening in elderly German adults. In: *International psychogeriatrics* 26 (3), S. 425–434. DOI: 10.1017/S1041610213002044.
- *Brenner G.; Altenhofen L.; Kerek-Bodden H.; Koch H.; Lang A.; Weber I.; Willms S. (2000): Outpatient treatment of patients with cardiovascular diseases. In: *Herz* 25 (5), S. 502–514.
- *Brockmann S.; Sielk M.; Fiegen J.; Wilm S.; Altiner A. (2008): Inappropriate antibiotic prescriptions for acute cough by GPs - Concept and script of a successful persuasive intervention (peer-intervention). In: *Zeitschrift für Allgemeinmedizin* 84 (2), S. 51–57. DOI: 10.1055/s-2007-1022534.

*Bruni C.; Steinhauser J. (2014): Questions related to breastfeeding - What every family physician should know. In: *Zeitschrift fur Allgemeinmedizin* 90 (10), S. 424–427. DOI: 10.3238/zfa.2014.0424-0427.

*Bucker B.; Butzlaff M.; Isfort J.; Koneczny N.; Vollmar H.C.; Lange S. et al. (2010): [Effect of written patient information on knowledge and function of patients with acute uncomplicated back pain (PIK Study)]. In: *Gesundheitswesen (Bundesverband der Arzte des Offentlichen Gesundheitsdienstes (Germany))* 72 (12), e78-88.

Buhse S.; Kuniss N.; Liethmann K.; Muller U.A.; Lehmann T.; Muhlhauser I. (2018): Informed shared decision-making programme for patients with type 2 diabetes in primary care: Cluster randomised controlled trial. In: *BMJ Open* 8 (12), e024004. DOI: 10.1136/bmjopen-2018-024004.

Buhse S.; Muhlhauser I.; Kuniss N.; Muller U.A.; Lehmann T.; Liethmann K.; Lenz M. (2015): An informed shared decision making programme on the prevention of myocardial infarction for patients with type 2 diabetes in primary care: protocol of a cluster randomised, controlled trial. In: *BMC family practice* 16 S. 43. DOI: 10.1186/s12875-015-0257-2.

*Bungartz, Jessica; Szecsenyi, Joachim; Joos, Stefanie (2011): [He that knows nothing doubts nothing: availability of foreign language patient education material for immigrant patients in Germany - a survey]. In: *Zeitschrift fur Evidenz, Fortbildung und Qualitat im Gesundheitswesen* 105 (10), S. 743–750. DOI: 10.1016/j.zefq.2011.09.028.

Buntrock C.; Ebert D.D.; Lehr D.; Smit F.; Riper H.; Berking M.; Cuijpers P. (2016): Effect of a web-based guided self-help intervention for prevention of major depression in adults with subthreshold depression a randomized clinical trial. In: *JAMA - Journal of the American Medical Association* 315 (17), S. 1854–1863. DOI: 10.1001/jama.2016.4326.

Burgdorf F.; Sundmacher L. (2014): Potentially avoidable hospital admissions in Germany: An analysis of factors influencing rates of ambulatory care sensitive hospitalizations. In: *Deutsches Arzteblatt International* 111 (13), S. 215–223. DOI: 10.3238/arztebl.2014.0215.

Campbell S.M.; Ludt S.; Van Lieshout J.; Boffin N.; Wensing M.; Petek D. et al. (2008): Quality indicators for the prevention and management of cardiovascular disease in primary care in nine European countries. In: *European Journal of Cardiovascular Prevention and Rehabilitation* 15 (5), S. 509–515. DOI: 10.1097/HJR.0b013e328302f44d.

Chehab, Gamal; Richter, Jutta G.; Brinks, Ralph; Fischer-Betz, Rebecca; Winkler-Rohlfing, Borgi; Schneider, Matthias (2018): Vaccination coverage in systemic lupus erythematosus-a cross-sectional analysis of the German long-term study (LuLa cohort). In: *Rheumatology (Oxford, England)* 57 (8), S. 1439–1447. DOI: 10.1093/rheumatology/key120.

Chenot, Jean-Francois; Scherer, Martin; Becker, Annette; Donner-Banzhoff, Norbert; Baum, Erika; Leonhardt, Corinna et al. (2008): Acceptance and perceived barriers of implementing a guideline for managing low back in general practice. In: *Implementation science : IS* 3 (101258411), S. 7. DOI: 10.1186/1748-5908-3-7.

*Chenot J.-F.; Egidi G. (2015): Skin cancer screening recommendation in the skin cancer prevention guideline - Critic of the DEGAM Part 2. In: *Zeitschrift fur Allgemeinmedizin* 91 (3), S. 121–125. DOI: 10.3238/zfa.2015.0121-0125.

Chenot J.-F.; Hua T.D.; Abu Abed M.; Schneider-Rudt H.; Friede T.; Schneider S.; Vormfelde S.V. (2014): Safety relevant knowledge of orally anticoagulated patients without self-monitoring: a baseline survey in primary care. In: *BMC family practice* 15 S. 104. DOI: 10.1186/1471-2296-15-104.

*Chenot J.-F.; Keller S. (2004): Smoking Cessation - What can Help? In: *Zeitschrift fur Allgemeinmedizin* 80 (3), S. 113–118.

Chenot J.-F.; Kochen M.M.; Himmel W. (2009): Student evaluation of a primary care clerkship: Quality assurance and identification of potential for improvement. In: *BMC Medical Education* 9 (1), S. 17. DOI: 10.1186/1472-6920-9-17.

*Chenot J.-F.; Pfingsten M.; Marnitz U.; Pfeifer K.; Kohlmann T.; Lindena G.; Schmidt C.O. (2019): Effectiveness of a risk-tailored short intervention to prevent chronic low back pain: A cluster-randomized study in general practice. In: *Schmerz* 33 (3), S. 226–235. DOI: 10.1007/s00482-019-0362-6.

*Claus, D.; Fleischer, S.; Mattern, E.; Ayerle, G. (2016): [A survey of Local Physicians and Psychotherapists on Cooperation in Regional Networks for Early Child Interventions in Saxony-Anhalt]. In: *Klinische Padiatrie* 228 (4), S. 202–207. DOI: 10.1055/s-0042-101244.

*Conrad, S.; Huppe, A.; Raspe, H. (2012): [Preference of patients with inflammatory bowel disease regarding information and shared decision-making: results from a cross-sectional survey in Germany]. In: *Zeitschrift fur Gastroenterologie* 50 (4), S. 364–372. DOI: 10.1055/s-0031-1281949.

*Constantinidis A.; Beyer M.; Karl S.; Gerlach F.M.; Erler A. (2013): Family practices in Germany: A Medical Home for patients with chronic illness? In: *Zeitschrift fur Allgemeinmedizin* 89 (7-8), S. 317–324. DOI: 10.3238/zfa.2013.0317-0324.

Cuddihy R.M.; Philis-Tsimikas A.; Nazeri A. (2011): Type 2 diabetes care and insulin intensification: Is a more multidisciplinary approach needed? Results from the modify survey. In: *Diabetes Educator* 37 (1), S. 111–123. DOI: 10.1177/0145721710388426.

- *Dapp U.; Anders J.; Meier-Baumgartner H.P.; von Renteln-Kruse W. (2007): Geriatric health promotion and prevention for independently living senior citizens: Programmes and target groups. In: *Zeitschrift fur Gerontologie und Geriatrie* 40 (4), S. 226–240. DOI: 10.1007/s00391-007-0469-8.
- Dapp U.; Anders J.; von Renteln-Kruse W.; Golgert S.; Meier-Baumgartner H.P.; Minder C.E. (2012): The Longitudinal Urban Cohort Ageing Study (LUCAS): study protocol and participation in the first decade. In: *BMC geriatrics* 12 ((Dapp) Albertinen-Haus Geriatrics Centre, Scientific Department at the University of Hamburg, Sellhopsweg 18-22, D-22459, Hamburg, Germany), S. 35.
- Dapp U.; Anders J.A.; von Renteln-Kruse W.; Minder C.E.; Meier-Baumgartner H.P.; Swift C.G. et al. (2011): A randomized trial of effects of health risk appraisal combined with group sessions or home visits on preventive behaviors in older adults. In: *The journals of gerontology. Series A, Biological sciences and medical sciences* 66 (5), S. 591–598.
- *Deinzer, A.; Babel, H.; Veelken, R.; Kohnen, R.; Schmieder, R. E. (2006): [Shared decision-making with hypertensive patients. Results of an implementation in Germany]. In: *Deutsche medizinische Wochenschrift* (1946) 131 (46), S. 2592–2596.
- Deinzer A.; Veelken R.; Kohnen R.; Schmieder R.E. (2009): Is a shared decision-making approach effective in improving hypertension management? In: *Journal of Clinical Hypertension* 11 (5), S. 266–270. DOI: 10.1111/j.1751-7176.2009.00112.x.
- *Demmel R.; Rist F.; Hagen J.; Aulhorn I.; Scheuren B.; Scherbaum N. et al. (2003): Secondary prevention beyond screening and brief advice. In: *Suchtmedizin in Forschung und Praxis* 5 (1), S. 33–36.
- *Dhein Y.; Munks-Lederer C.; Worth H. (2003): Evaluation of a Structured Education Programme for Patients with COPD under Outpatient Conditions - A Pilot Study. In: *Pneumologie* 57 (10), S. 591–597. DOI: 10.1055/s-2003-43019.
- Diederichs C.; Jordan S.; Domanska O.; Neuhauser H. (2018): Health literacy in men and women with cardiovascular diseases and its association with the use of health care services - Results from the population-based GEDA2014/2015-EHIS survey in Germany. In: *PLoS ONE* 13 (12), e0208303. DOI: 10.1371/journal.pone.0208303.
- Diehl K.; Ganseforth D.; Herr R.M.; Gorig T.; Bock C.; Mayer M.; Schneider S. (2015): Physician gender and lifestyle counselling to prevent cardiovascular disease: A nationwide representative study. In: *Journal of Public Health Research* 4 (2), S. 113–119. DOI: 10.4081/jphr.2015.534.
- *Diehm C.; Allenberg J.R.; Pittrow D.; Darius H. (2007): Importance of the Ankle-Brachial Index (ABI) in the prevention of cardiovascular diseases. Ten questions and answers. In: *Herz* 32 (5), S. 404–409. DOI: 10.1007/s00059-007-3010-3.
- Diener, Annette; Celemin-Heinrich, Salome; Wegscheider, Karl; Kolpatzik, Kai; Tomaschko, Katrin; Altiner, Attila et al. (2013): In-vivo-validation of a cardiovascular risk prediction tool: the arriba-pro study. In: *BMC family practice* 14 (100967792), S. 13. DOI: 10.1186/1471-2296-14-13.
- *Doering T.J.; Vahlbruch A.; Steuernagel B.; Fischer G.C. (2002): Prevention in general practitioner's daily work. In: *Pravention und Rehabilitation* 14 (1), S. 28–41.
- *Drabik A.; Graf C.; Buscher G.; Stock S. (2012): Evaluating the effectiveness of a disease management program diabetes in the German Statutory Health Insurance - First results and methodological considerations. In: *Zeitschrift fur Evidenz, Fortbildung und Qualitat im Gesundheitswesen* 106 (9), S. 649–655. DOI: 10.1016/j.zefq.2011.12.007.
- Drewelow E.; Wollny A.; Pentzek M.; Immecke J.; Lambrecht S.; Wilm S. et al. (2012): Improvement of primary health care of patients with poorly regulated diabetes mellitus type 2 using shared decision-making--the DEBATE trial. In: *BMC family practice* 13 ((Drewelow) Rostock University Medical Center, Rostock, Germany), S. 88. DOI: 10.1186/1471-2296-13-88.
- Dudouet P.; Gautret P.; Larsen C.S.; Diaz-Menendez M.; Trigo E.; von Sonnenburg F. et al. (2020): Chikungunya resurgence in the Maldives and risk for importation via tourists to Europe in 2019-2020: A GeoSentinel case series. In: *Travel Medicine and Infectious Disease* 36 S. 101814. DOI: 10.1016/j.tmaid.2020.101814.
- Dusch, Martin; Narcis, Elisabeth; Strohmer, Renate; Schuttpelz-Brauns, Katrin (2018): Competency-based learning in an ambulatory care setting: Implementation of simulation training in the Ambulatory Care Rotation during the final year of the MaReCuM model curriculum. In: *GMS journal for medical education* 35 (1), Doc6. DOI: 10.3205/zma001153.
- Dybek, Inga; Bischof, Gallus; Grothues, Janina; Reinhardt, Susa; Meyer, Christian; Hapke, Ulfert et al. (2006): The reliability and validity of the Alcohol Use Disorders Identification Test (AUDIT) in a German general practice population sample. In: *Journal of studies on alcohol* 67 (3), S. 473–481.
- *Eckle V.-S.; Hucklenbruch C. (2010): General practitioner's medical records for preoperative anesthetic visit. In: *MMW-Fortschritte der Medizin* 152 (4), S. 43–46. DOI: 10.1007/bf03365975.
- *Egidi G. (2015): Controversies over guidelines - Exemplified in the S3 guideline "secondary prevention of ischemic stroke and transient ischemic attack". In: *Zeitschrift fur Allgemeinmedizin* 91 (7-8), S. 304–309. DOI: 10.3238/zfa.2015.0304-0309.
- *Egidi G.; Chenot J.-F. (2015): Skin cancer screening recommendation in the skin cancer prevention guideline - Critique of the DEGAM Part 1. In: *Zeitschrift fur Allgemeinmedizin* 91 (3), S. 117–120. DOI: 10.3238/zfa.2015.0117-0120.

*Eich, W.; Hauser, W.; Arnold, B.; Bernardy, K.; Brucke, W.; Eidmann, U. et al. (2012): [Fibromyalgia syndrome. General principles and coordination of clinical care and patient education]. In: *Schmerz (Berlin, Germany)* 26 (3), S. 268–275. DOI: 10.1007/s00482-012-1167-z.

*Eickhoff A.; Riemann J.F. (2003): The prevention of colorectal cancer - What can the family doctor do? In: *MMW-Fortschritte der Medizin* 145 (16), S. 35–41.

*Ellsaser G.; Trost-Brinkhues G. (2013): [Vaccination among students in grades 6-10, 2011 - a comparison of German states: need for action for a targeted nationwide immunisation strategy]. In: *Gesundheitswesen (Bundesverband der Ärzte des Öffentlichen Gesundheitsdienstes (Germany))* 75 (11), S. 705–713.

Engler, Jennifer; Dahlhaus, Anne; Guthlin, Corina (2017): The readiness of German GPs to recommend and conduct cancer screening is associated with patient-physician gender concordance. Results of a survey. In: *The European journal of general practice* 23 (1), S. 11–19. DOI: 10.1080/13814788.2016.1240166.

Ernsting, Clemens; Stuhmann, Lena Mareike; Dombrowski, Stephan U.; Voigt-Antons, Jan-Niklas; Kuhlmeier, Adelheid; Gellert, Paul (2019): Associations of Health App Use and Perceived Effectiveness in People With Cardiovascular Diseases and Diabetes: Population-Based Survey. In: *JMIR mHealth and uHealth* 7 (3), e12179. DOI: 10.2196/12179.

*Eschweiler G.W. (2010): New developments in the diagnosis of dementia. In: *Deutsches Arzteblatt* 107 (39), S. 677–683. DOI: 10.3238/arztebl.2010.0677.

*Etzel, M.; Mons, U.; Schmitt, S.; Lang, P.; Potschke-Langer, M. (2008): [Smoking cessation in Germany in 2007. Patterns of outpatient smoking cessation counselling and treatment]. In: *Bundesgesundheitsblatt, Gesundheitsforschung, Gesundheitsschutz* 51 (12), S. 1453–1461. DOI: 10.1007/s00103-008-0719-y.

Fankhaenel, Thomas; Samos, Franziska; Luck-Sikorski, Claudia; Thiel, Carolin; Klement, Andreas; Frese, Thomas (2018): Patient Satisfaction as a Moderator of Risky Alcohol Consumers' Attitude Towards Screening and Brief Intervention: A Cross Sectional Survey. In: *Alcohol and alcoholism (Oxford, Oxfordshire)* 53 (4), S. 403–407. DOI: 10.1093/alcalc/agx001.

*Fankhanel, Thomas; Julia, Lenz; Stephanie, Papert; Karen, Voigt; Andreas, Klement (2014): [Screening and brief intervention in primary care: barriers against an early intervention for patients with alcohol misuse]. In: *Psychotherapie, Psychosomatik, medizinische Psychologie* 64 (9-10), S. 373–377. DOI: 10.1055/s-0034-1374595.

*Fendler, C.; Saracbasi, E.; Dybowski, F.; Heldmann, F.; Braun, J. (2012): [Practical problems by implementation of vaccination recommendations]. In: *Zeitschrift für Rheumatologie* 71 (2), 147–5. DOI: 10.1007/s00393-011-0946-8.

Fiederling, Jonas; Ulrich, Cornelia M.; Hemminki, Kari; Haug, Ulrike (2014): Consideration of family history of cancer in medical routine: a survey in the primary care setting in Germany. In: *European journal of cancer prevention : the official journal of the European Cancer Prevention Organisation (ECP)* 23 (3), S. 199–205. DOI: 10.1097/CEJ.0b013e328364743d.

Filipas, D.; Spix, C.; Schulz-Lampel, D.; Michaelis, J.; Hohenfellner, R.; Roth, S.; Thuroff, J. W. (2003): Screening for renal cell carcinoma using ultrasonography: a feasibility study. In: *BJU international* 91 (7), S. 595–599.

Fink A.; Fach E.-M.; Schroder S.L. (2019): 'Learning to shape life' - A qualitative study on the challenges posed by a diagnosis of diabetes mellitus type 2. In: *International Journal for Equity in Health* 18 (1), S. 924. DOI: 10.1186/s12939-019-0924-3.

*Fischer, G. C.; Beyer, M.; Gerlach, F. M.; Rohde-Kampmann, R. (2001): [Importance and possibilities of general guidelines in family practice strategies--do we need 'basic guidelines' in general practice?]. In: *Zeitschrift für Ärztliche Fortbildung und Qualitätssicherung* 95 (6), S. 435–442.

Fissen G.; Lewis D.K.; Abholz H.-H. (2008): Understanding the concept of medical risk reduction: A comparison between the UK and Germany. In: *European Journal of General Practice* 14 (3-4), S. 109–116. DOI: 10.1080/13814780802580247.

*Floer B.; Schnee M.; Bocken J.; Streich W.; Kunstrmann W.; Isfort J.; Butzlaff M. (2004): Shared decision making. The perspective of practicing physicians. In: *Medizinische Klinik* 99 (8), S. 435–440. DOI: 10.1007/s00063-004-1066-x.

*Frank, Luca; Hueber, Susann; Kuhlein, Thomas; Schedlbauer, Angela; Roos, Marco (2018): [Evidence and practice: How do primary care physicians evaluate screening interventions? A questionnaire study]. In: *Zeitschrift für Evidenz, Fortbildung und Qualität im Gesundheitswesen* 135-136 (101477604), S. 1–9. DOI: 10.1016/j.zefq.2018.06.003.

Freier C.; Heintze C.; Herrmann W.J. (2020): Prescribing and medical non-adherence after myocardial infarction: qualitative interviews with general practitioners in Germany. In: *BMC family practice* 21 (1), S. 81. DOI: 10.1186/s12875-020-01145-6.

Frese, Thomas; Deutsch, Tobias; Keyser, Melanie; Sandholzer, Hagen (2012): In-home preventive comprehensive geriatric assessment (CGA) reduces mortality--a randomized controlled trial. In: *Archives of gerontology and geriatrics* 55 (3), S. 639–644. DOI: 10.1016/j.archger.2012.06.012.

Freund T.; Freund M.; Mahler C.; Schwantes U.; Ose D.; Joos S.; Szecsenyi J. (2010): Family practice web sites: new perspectives for patient care and health information? In: *Family medicine* 42 (7), S. 501–506.

*Freund T.; Lekutat C.; Schwantes U.; Szecsenyi J.; Joos S. (2010): [Health screening and vaccination status--results from a patient survey in German general practices]. In: *Gesundheitswesen (Bundesverband der Ärzte des Öffentlichen Gesundheitsdienstes (Germany))* 72 (11), S. 804–807.

Freund T.; Mahler C.; Erler A.; Gensichen J.; Ose D.; Szecsenyi J.; Peters-Klimm F. (2011): Identification of patients likely to benefit from care management programs. In: *American Journal of Managed Care* 17 (5), S. 345–352. Online verfügbar unter <http://www.ajmc.com/media/pdf/824f49680d95f8d8323703f5fcdec279.pdf>.

Freund T.; Wensing M.; Geissler S.; Peters-Klimm F.; Mahler C.; Boyd C.M.; Szecsenyi J. (2012): Primary care physicians' experiences with case finding for practice-based care management. In: *American Journal of Managed Care* 18 (4), e155-e161. Online verfügbar unter <http://www.ajmc.com/articles/Primary-Care-Physicians-Experiences-With-Case-Finding-for-Practice-Based-Care-Management>.

Freyer J.; Kasprick L.; Sultzer R.; Schiek S.; Bertsche T. (2018): A dual intervention in geriatric patients to prevent drug-related problems and improve discharge management. In: *International Journal of Clinical Pharmacy* 40 (5), S. 1189–1198. DOI: 10.1007/s11096-018-0643-7.

Führer A.; Niedermann A.; Kalfa V.; Mikolajczyk R.; Wienke A. (2020): Serious shortcomings in assessment and treatment of asylum seekers' mental health needs. In: *PLoS ONE* 15 (10), e0239211. DOI: 10.1371/journal.pone.0239211.

Fuller, N. R.; Colagiuri, S.; Schofield, D.; Olson, A. D.; Shrestha, R.; Holzapfel, C. et al. (2013): A within-trial cost-effectiveness analysis of primary care referral to a commercial provider for weight loss treatment, relative to standard care--an international randomised controlled trial. In: *International journal of obesity* (2005) 37 (6), S. 828–834. DOI: 10.1038/ijo.2012.139.

*Fulling, Tim; Bula, Philipp; Defer, Alexander; Bonnaire, Felix Alois (2020): Dresdener Network Osteoporosis. In: *Zeitschrift für Orthopädie und Unfallchirurgie* (101308227). DOI: 10.1055/a-1149-9588.

Gabrys L.; Jordan S.; Schlaud M. (2015): Prevalence and temporal trends of physical activity counselling in primary health care in Germany from 1997-1999 to 2008-2011. In: *International Journal of Behavioral Nutrition and Physical Activity* 12 (1), S. 136. DOI: 10.1186/s12966-015-0299-9.

*Gaede L.; Blumenstein J.; Liebetrau C.; Dorr O.; Kim W.-K.; Hamm C. et al. (2019): Heart valve diseases: How sufficient is the knowledge of the German population? In: *Herz* 44 (2), S. 161–169. DOI: 10.1007/s00059-017-4632-8.

*Gallois, K.; Strube, H.; Hassel, H.; Pigeot, I.; Zeeb, H. (2015): [OptimaHI 60plus--Development, Implementation and Transfer of a Counselling Aid for Elderly People through Participation]. In: *Gesundheitswesen (Bundesverband der Ärzte des Öffentlichen Gesundheitsdienstes (Germany))* 77 Suppl 1 (bfd, 9204210), S89-90. DOI: 10.1055/s-0032-1331247.

*Gandjour A.; Lauterbach K.W. (2005): When is it worth reducing overuse of health care services? The example of prescribing expensive antihypertensives. In: *Medizinische Klinik* 100 (9), S. 535–541. DOI: 10.1007/s00063-005-1073-6.

*Gardemann J. (2001): Towards a new line in public school health service; the medical profession in the fields of social pediatrics, occupational health and multiprofessional health promotion. In: *Gesundheitswesen* 63 (11), S. 658–666. DOI: 10.1055/s-2001-18408.

Geest T.A.; Wetzel R.; Raposo V.; Ferreira P.L.; Baker R.; Wensing M.; Olesen F. (2005): Elderly patients' and GPs' views on different methods for patient involvement: An international qualitative interview study. In: *Family Practice* 22 (2), S. 184–191. DOI: 10.1093/fampra/cmh733.

Gehrke-Beck S.; Banfer M.; Schilling N.; Schmidt K.; Gensichen J.; Heintze C. (2017): The specific needs of patients following sepsis: A nested qualitative interview study. In: *BJGP Open* 1 (1). DOI: 10.3399/bjgpopen17X100725.

*Gensichen J.; Hensler S.; Sonnichsen A. (2008): Monitoring in secondary prevention: Cardio-vascular prevention as an example DEGAM series: Comprehensive care for patient with chronic illness. In: *Zeitschrift für Allgemeinmedizin* 84 (5), S. 191–194. DOI: 10.1055/s-2008-1062710.

*Gerber, B.; Bolz, M.; Kringel, U.; Janni, W. (2006): [Gynecological and obstetric aspects of general practice]. In: *MMW Fortschritte der Medizin* 148 (6), S. 23–27.

*Gerlach F.M.; Szecsenyi J. (2013): Family doctor-centred care in Baden-Württemberg: Concept and results of a controlled evaluation study. In: *Zeitschrift für Evidenz, Fortbildung und Qualität im Gesundheitswesen* 107 (6), S. 365–371. DOI: 10.1016/j.zefq.2013.07.002.

Gerlach N.; Michiels-Corsten M.; Viniol A.; Schleef T.; Junius-Walker U.; Krause O.; Donner-Banzhoff N. (2020): Professional roles of general practitioners, community pharmacists and specialist providers in collaborative medication deprescribing - a qualitative study. In: *BMC family practice* 21 (1), S. 183. DOI: 10.1186/s12875-020-01255-1.

Gitt A.K.; Sonntag F.; Jannowitz C.; Weizel A.; Karmann B.; Schaefer J.R. et al. (2016): Better lipid target achievement for secondary prevention through disease management programs for diabetes mellitus and coronary heart disease in clinical practice in Germany. In: *Current Medical Research and Opinion* 32 (3), S. 417–426. DOI: 10.1185/03007995.2015.1120715.

Glaab T.; Banik N.; Rutschmann O.; Wencker M. (2006): National survey of guideline-compliant COPD management among pneumologists and primary care physicians. In: *COPD: Journal of Chronic Obstructive Pulmonary Disease* 3 (3), S. 141–148. DOI: 10.1080/15412550600829299.

- *Gogol, Manfred; Siebenhofer, Andrea (2016): [Choosing wisely--against overuse in healthcare systems--activities in Germany and Austria in geriatric medicine]. In: *Wiener medizinische Wochenschrift* (1946) 166 (5-6), S. 155–160. DOI: 10.1007/s10354-015-0424-z.
- *Gohlke, H.; Jarmatz, H.; Zaumseil, J.; Bestehorn, K.; Jansen, C.; Hasford, J. (2000): [Effect of optimized integrated management on long-term effectiveness of cardiologic rehabilitation]. In: *Deutsche medizinische Wochenschrift* (1946) 125 (48), S. 1452–1456.
- *Gohlke, H.; Schirmer, A.; Theobald, K-H (2004): [Cardiovascular risk stratification in general practice]. In: *Zeitschrift fur Kardiologie* 93 Suppl 2 (xw7, 0360430), II26-32.
- Goldbeck L; Schmid K. (2003): Effectiveness of autogenic relaxation training on children and adolescents with behavioral and emotional problems. In: *Journal of the American Academy of Child and Adolescent Psychiatry* 42 (9), S. 1046–1054. DOI: 10.1097/01.CHI.0000070244.24125.F.
- *Goltz L; Bodechtel U.; Siepmann T. (2014): Statins and ASS for primary prevention of cardiovascular and cerebrovascular disease. In: *Deutsche Medizinische Wochenschrift* 139 (6), S. 283–286. DOI: 10.1055/s-0033-1359905.
- *Golubnitschaja O. (2018): The Keyrole of Multiomics in the Predictive, Preventive and Personalised Medical Approach towards Glaucoma Management. In: *Klinische Monatsblatter fur Augenheilkunde* 235 (2), S. 146–150. DOI: 10.1055/s-0044-101164.
- Goriga T.; Mayer M.; Bock C.; Diehl K.; Hilger J.; Herr R.M.; Schneider S. (2014): Dietary counselling for cardiovascular disease prevention in primary care settings: Results from a German physician survey. In: *Family Practice* 31 (3), S. 325–332. DOI: 10.1093/fampra/cmu007.
- Graf, Christine; Schlepper, Stefanie; Bauer, Carina; Ferrari, Nina; Frank, Stefan; Gartner, Lena et al. (2016): Feasibility and acceptance of exercise recommendations (10,000 steps a day) within routine German health check (Check-Up 35/GOA29)-study protocol. In: *Pilot and feasibility studies* 2 (101676536), S. 52.
- Grammer T.B.; Dressel A.; Gergei I.; Kleber M.E.; Laufs U.; Scharnagl H. et al. (2019): Cardiovascular risk algorithms in primary care: Results from the DETECT study. In: *Scientific reports* 9 (1), S. 1101. DOI: 10.1038/s41598-018-37092-7.
- *Grewe, H. A.; Heckenhahn, S.; Blattner, B. (2014): [Health protection during heat waves: European recommendations and experience in Hesse]. In: *Zeitschrift fur Gerontologie und Geriatrie* 47 (6), S. 483–489. DOI: 10.1007/s00391-014-0676-z.
- Grochtdreis, Thomas; Brettschneider, Christian; Bjerregaard, Frederike; Bleich, Christiane; Boczor, Sigrid; Harter, Martin et al. (2019): Cost-effectiveness analysis of collaborative treatment of late-life depression in primary care (GermanIMPACT). In: *European psychiatry : the journal of the Association of European Psychiatrists* 57 (djf, 9111820), S. 10–18. DOI: 10.1016/j.eurpsy.2018.12.007.
- *Grulke, Norbert; Bailer, Harald; Blaser, Gerd; Geyer, Michael; Hinz, Andreas; Schmutzler, Gabriele et al. (2005): [Screening for depression -- two questions practically oriented]. In: *Wiener medizinische Wochenschrift* (1946) 155 (13-14), S. 297–302.
- *Grusser M.; Hoffstadt K.; Jorgens V. (2000): Preventive care for early detection of diabetes mellitus complications: a model project in Wolfsburg. In: *Zeitschrift fur arztliche Fortbildung und Qualitatssicherung : in Zusammenarbeit mit der Kaiserin-Friedrich-Stiftung fur das arztliche Fortbildungswesen* 94 (5), S. 411–416.
- *Gstettner, A.; Holzapfel, C.; Stoll, J.; Hauner, H. (2013): [Weight reduction: evaluation of the possibilities in primary care and patient satisfaction. Results from a weight reduction trial]. In: *Deutsche medizinische Wochenschrift* (1946) 138 (19), S. 989–994. DOI: 10.1055/s-0033-1343164.
- *Gube M.; Ell W.; Schiefer J.; Kraus T. (2010): Medicolegal assessment of the ability to drive a motor vehicle in persons with epilepsy. In: *Deutsches Arzteblatt* 107 (13), S. 217–223. DOI: 10.3238/arztebl.2010.0217.
- Gumz A.; Weigel A.; Wegscheider K.; Romer G.; Lowe B. (2018): The psychenet public health intervention for anorexia nervosa: a pre-post-evaluation study in a female patient sample. In: *Primary health care research & development* 19 (1), S. 42–52. DOI: 10.1017/S1463423617000524.
- *Gutz S.; Gillissen A. (2006): Prevention of lung cancer. In: *MMW-Fortschritte der Medizin* 148 (22), S. 26–27.
- Guttmann H. (2013): The health care situation of the mentally ill elderly in Germany. In: *Geriatric Mental Health Care* 1 (1), S. 20–25. DOI: 10.1016/j.gmhc.2012.11.001.
- *Gysan, D. B.; Latsch, J.; Bjarnason-Wehrens, B.; Albus, C.; Falkowski, G.; Herold, G. et al. (2004): [The PreFord Study. A prospective cohort study to evaluate the risk of a cardiovascular event (overall-collective) as well as a prospective, randomized, controlled, multicentre clinical intervention study (high-risk-collective) on primary prevention of ca. In: *Zeitschrift fur Kardiologie* 93 (2), S. 131–136.
- Haeusler K.G.; Gerth A.; Limbourg T.; Tebbe U.; Oeff M.; Wegscheider K. et al. (2015): Use of vitamin K antagonists for secondary stroke prevention depends on the treating healthcare provider in Germany - results from the German AFNET registry. In: *BMC Neurology* 15 (1), S. 129. DOI: 10.1186/s12883-015-0371-8.
- Hagen, Bernd; Strauch, Stefan (2011): The J1 adolescent health check-up: analysis of data from the German KiGGS survey. In: *Deutsches Arzteblatt International* 108 (11), S. 180–186. DOI: 10.3238/arztebl.2011.0180.

*Haisch J.; Remmele W. (2000): Effectiveness and efficiency of out-patient instruction of diabetics: A comparison of specialist and general practitioner practices. In: *Deutsche Medizinische Wochenschrift* 125 (7), S. 171–176. DOI: 10.1055/s-2007-1024014.

Hannover W.; Kopke D.; Hannich H.-J. (2010): Perceived Barriers to Prostate Cancer Screenings Among Middle-Aged Men in North-Eastern Germany. In: *Public Health Nursing* 27 (6), S. 504–512. DOI: 10.1111/j.1525-1446.2010.00889.x.

Harder, Sebastian; Fischer, Philipp; Krause-Schafer, Markus; Ostermann, Klaus; Helms, Gottfried; Prinz, Helge et al. (2005): Structure and markers of appropriateness, quality and performance of drug treatment over a 1-year period after hospital discharge in a cohort of elderly patients with cardiovascular diseases from Germany. In: *European journal of clinical pharmacology* 60 (11), S. 797–805.

Harter, Martin; Bartsch, Anna-Lena; Egger, Nina; Konig, Hans-Helmut; Kriston, Levente; Schulz, Holger et al. (2015): Evaluating a collaborative smoking cessation intervention in primary care (ENTER): study protocol for a cluster-randomized controlled trial. In: *Trials* 16 (101263253), S. 447. DOI: 10.1186/s13063-015-0977-6.

Hartmann, Uwe; Burkart, Martin (2007): Erectile dysfunctions in patient-physician communication: optimized strategies for addressing sexual issues and the benefit of using a patient questionnaire. In: *The journal of sexual medicine* 4 (1), S. 38–46. DOI: 10.1111/j.1743-6109.2006.00385.x.

*Haumann H.; Thielmann A.; Sikora M.; Schnell U.; Weltermann B. (2016): How do you explain vaccine effects to patients? A representative, web-based survey among primary care physicians from North-Rhine Westphalia. In: *Deutsche Medizinische Wochenschrift* 141 (12), e115–e120. DOI: 10.1055/s-0042-101463.

*Hauner H.; Beyer-Reiners E.; Bischoff G.; Breidenassel C.; Ferschke M.; Gebhardt A. et al. (2019): Manual of Nutritional Therapy in Patient Care. In: *Aktuelle Ernährungsmedizin* 44 (6), S. 384–419. DOI: 10.1055/a-1030-5207.

Haussmann, Alexander; Ungar, Nadine; Gabrian, Martina; Tsioris, Angeliki; Sieverding, Monika; Wiskemann, Joachim; Steindorf, Karen (2018): Are healthcare professionals being left in the lurch? The role of structural barriers and information resources to promote physical activity to cancer patients. In: *Supportive care in cancer : official journal of the Multinational Association of Supportive Care in Cancer* 26 (12), S. 4087–4096. DOI: 10.1007/s00520-018-4279-x.

*Haustein K.-O. (2003): Juveniles, smoking and the role of the general practitioner. In: *Tagliche Praxis* 44 (1), S. 123–132.

*Hauswaldt, J.; Kersting, M.; Hummers-Pradier, E. (2010): [Influenza vaccination by Lower Saxonian general practitioners - a secondary analysis of physicians' data from 1995/1996, 2002/2003, and 2005/2006]. In: *Gesundheitswesen (Bundesverband der Ärzte des Öffentlichen Gesundheitsdienstes (Germany))* 72 (6), S. 332–339. DOI: 10.1055/s-0030-1249690.

*Hauswaldt J.; Junius-Walker U.; Kersting M.; Hummers-Pradier E. (2009): Preventive health examination in German general practice - A secondary analysis of data from 1996 to 2006. In: *Zeitschrift für Allgemeinmedizin* 85 (10), S. 411–417. DOI: 10.3238/zfa.2009.0416.

*Hebebrand J.; Hamelmann E.; Hartmann A.; Holtmann M.; Jockel K.-H.; Kremer U. et al. (2017): Current practice of pre-and postnatal screening and future developments for evidence based guidelines. In: *Klinische Pädiatrie* 229 (1), S. 2–13. DOI: 10.1055/s-0042-118187.

*Heberlein, I.; Demary, W.; Bloching, H.; Braun, J.; Buttgereit, F.; Dreher, R. et al. (2011): [Diagnosis and treatment of osteoporosis and rheumatoid arthritis in accordance with German guidelines. Results of a survey of patients, primary care physicians and rheumatologists]. In: *Zeitschrift für Rheumatologie* 70 (7), S. 592–601. DOI: 10.1007/s00393-011-0821-7.

Hecht M.; Heuss D.; Hilz M.J. (2003): Diagnosis and treatment of polyneuropathy. What can the family doctor do? In: *MMW-Fortschritte der Medizin* 145 (SUPPL. 2), S. 81–85.

Heddaeus, Daniela; Dirmaier, Jorg; Daubmann, Anne; Grochtdreis, Thomas; Konig, Hans-Helmut; Lowe, Bernd et al. (2019): [Clinical trial of a stepped and collaborative care model for mental illnesses and comorbidities in the Hamburg Network for Health Services Research]. In: *Bundesgesundheitsblatt, Gesundheitsforschung, Gesundheitsschutz* 62 (2), S. 205–213. DOI: 10.1007/s00103-018-2865-1.

Heddaeus, Daniela; Steinmann, Maya; Liebherz, Sarah; Harter, Martin; Watzke, Birgit (2015): [psychenet - The Hamburg Network for Mental Health: Evaluation of the Health Network Depression from the Perspective of Participating General Practitioners, Psychotherapists and Psychiatrists]. In: *Psychiatrische Praxis* 42 Suppl 1 (qck, 0423204), S54-9. DOI: 10.1055/s-0034-1387688.

Hegerl U.; Althaus D. (2003): Elements for an Effective Disease Management for Patients with Depressive Disorders at the Primary Care Level. In: *MMW-Fortschritte der Medizin* 145 (41), S. 24–27.

Hegerl U.; Althaus D.; Schmidtko A.; Niklewski G. (2006): The alliance against depression: 2-year evaluation of a community-based intervention to reduce suicidality. In: *Psychological Medicine* 36 (9), S. 1225–1233. DOI: 10.1017/S003329170600780X.

Hegerl U.; Maxwell M.; Harris F.; Koburger N.; Mergl R.; Szekely A. et al. (2019): Prevention of suicidal behaviour: Results of a controlled community-based intervention study in four European countries. In: *PLoS ONE* 14 (11), e0224602. DOI: 10.1371/journal.pone.0224602.

Hegerl U.; Wittenburg L.; Arensman E.; Van Audenhove C.; Coyne J.C.; McDaid D. et al. (2009): Optimizing Suicide Prevention Programs and Their Implementation in Europe (OSPI Europe): An evidence-based multi-level approach. In: *BMC Public Health* 9

((Hegerl, Wittenburg, Bramesfeld) University of Leipzig, Department of Psychiatry, Germany(Varnik) Estonian-Swedish Mental Health and Suicidology Institute, Estonia(McDaid) London School of Economics, Personal Social Services Research Unit, LSE Health and), S. 428. DOI: 10.1186/1471-2458-9-428.

Heidrich, Benjamin; Cetindere, Ahmet; Beyaz, Mustafa; Stahmeyer, Jona T.; Basaran, Michael M.; Braynis, Boris et al. (2014): High prevalence of hepatitis markers in immigrant populations: a prospective screening approach in a real-world setting. In: *European journal of gastroenterology & hepatology* 26 (10), S. 1090–1097. DOI: 10.1097/MEG.00000000000000164.

Heidrich J.; Behrens T.; Raspe F.; Keil U. (2005): Knowledge and perception of guidelines and secondary prevention of coronary heart disease among general practitioners and internists. Results from a physician survey in Germany. In: *European journal of cardiovascular prevention and rehabilitation : official journal of the European Society of Cardiology, Working Groups on Epidemiology & Prevention and Cardiac Rehabilitation and Exercise Physiology* 12 (6), S. 521–529.

Heigl F.; Pfleiderer T.; Schettler V.; Grutzmacher P.; Julius U.; Ringel J. et al. (2017): Lipidological competence centres and networks: Future perspectives to improve healthcare of patients with disorders of lipid metabolism. In: *Atherosclerosis Supplements* 30 ((Heigl, Pfleiderer, Hettich, Lotz, Reeg, Graf) Medical Care Centre Kempten-Allgau, Kempten, Germany(Schettler) Centre of Nephrology Gottingen GbR, Gottingen, Germany(Grutzmacher) Medical Clinic II, Agaplesion Markus Hospital, Frankfurt, Germany(Julius) Dep), S. 63–71. DOI: 10.1016/j.atherosclerosissup.2017.05.014.

Heinemann S.; Weiss V.; Straube K.; Nau R.; Grimmssmann T.; Himmel W.; Hummers-Pradier E. (2016): Understanding and reducing the prescription of hypnotics and sedatives at the interface of hospital care and general practice: A protocol for a mixed-methods study. In: *BMJ Open* 6 (8), e011908. DOI: 10.1136/bmjopen-2016-011908.

*Heintze, Christoph (2014): [Prevention in general practice between evidence- and narrative-based medicine: squaring the circle?]. In: *Zeitschrift fur Evidenz, Fortbildung und Qualitat im Gesundheitswesen* 108 (4), S. 203–207. DOI: 10.1016/j.zefq.2014.03.016.

Helmut I.G.; Poulsen A.; Suppli C.H.; Molbak K. (2015): Varicella in Europe-A review of the epidemiology and experience with vaccination. In: *Vaccine* 33 (21), S. 2406–2413. DOI: 10.1016/j.vaccine.2015.03.055.

*Hensler S.; Hoidn S.; Jork K. (2006): DEGAM practice guideline for stroke. In: *Zeitschrift fur Allgemeinmedizin* 82 (9), S. 404–408. DOI: 10.1055/s-2006-942213.

*Hensler S.; Hoidn S.; Jork K. (2007): Is ASS + Dipyridamole better than ASS for preventing recurrent stroke? In: *Zeitschrift fur Allgemeinmedizin* 83 (8), S. 324–327. DOI: 10.1055/s-2007-984364.

Herber O.R.; Gies V.; Schwappach D.; Thurmann P.; Wilm S. (2014): Patient information leaflets: informing or frightening? A focus group study exploring patients' emotional reactions and subsequent behavior towards package leaflets of commonly prescribed medications in family practices. In: *BMC family practice* 15 ((Herber, Gies, Schwappach, Thurmann, Wilm) Institute of General Practice, Medical Faculty of the Heinrich Heine University Dusseldorf, Moorenstr, 5, Building 14,97, 40225 Dusseldorf, Germany. Oliver.Herber@med.uni-duesseldorf.de), S. 163. DOI: 10.1186/1471-2296-15-163.

*Hermann K.; Szecsenyi J.; Ludt S. (2008): Comparative evaluation of 2 tools for counselling patients in general practice about cardiovascular risk: CARRISMA and e-ARRIBA. Effects on motivation for change and shared decision making. In: *Zeitschrift fur Allgemeinmedizin* 84 (5), S. 207–213. DOI: 10.1055/s-2008-1077070.

*Hildebrandt H.; Schmitt G.; Roth M.; Stunder B. (2011): Implementing population-based integrated care for a region: A work-in-progress report on the project "gesundes Kinzigtal". In: *Zeitschrift fur Evidenz, Fortbildung und Qualitat im Gesundheitswesen* 105 (8), S. 585–589. DOI: 10.1016/j.zefq.2011.09.003.

*Hinrichs T.; Brach M.; Bucchi C.; Moschny A.; Wilm S.; Thiem U.; Platen P. (2013): An exercise programme for community-dwelling, mobility-restricted and chronically ill older adults with structured support by the general practitioner's practice (HOMEfit): From feasibility to evaluation. In: *Zeitschrift fur Gerontologie und Geriatrie* 46 (1), S. 56–63. DOI: 10.1007/s00391-012-0329-z.

Hinrichs T.; Moschny A.; Klaassen-Mielke R.; Trampisch U.; Thiem U.; Platen P. (2011): General practitioner advice on physical activity: analyses in a cohort of older primary health care patients (getABI). In: *BMC family practice* 12 ((Hinrichs) Department of Sports Medicine and Sports Nutrition, Ruhr-University Bochum, 44780 Bochum, Germany), S. 26. DOI: 10.1186/1471-2296-12-26.

Hobbs, F. D. Richard; Erhardt, Leif (2002): Acceptance of guideline recommendations and perceived implementation of coronary heart disease prevention among primary care physicians in five European countries: the Reassessing European Attitudes about Cardiovascular Treatment (REACT) survey. In: *Family Practice* 19 (6), S. 596–604.

Hoch, Eva; Muehlig, Stephan; Hofler, Michael; Lieb, Roselind; Wittchen, Hans-Ulrich (2004): How prevalent is smoking and nicotine dependence in primary care in Germany? In: *Addiction (Abingdon, England)* 99 (12), S. 1586–1598.

*Hofreuter-Gatgens K.; Mnich E.; Thomas D.; Salomon T.; Von Dem Knesebeck O. (2011): Active health promotion among the aged in a rural region: Participants, acceptance, and implementation. In: *Bundesgesundheitsblatt - Gesundheitsforschung - Gesundheitsschutz* 54 (8), S. 933–941. DOI: 10.1007/s00103-011-1319-9.

Hohmann C.; Radziwill R.; Klotz J.M.; Jacobs A.H. (2010): Health-Related Quality of Life after Ischemic Stroke: The Impact of Pharmaceutical Interventions on Drug Therapy (Pharmaceutical Care Concept). In: *Health and Quality of Life Outcomes* 8 ((Hohmann, Klotz) Klinikum Fulda gAG, Department of Neurology, Pacelliallee 4, 36043 Fulda, Germany(Hohmann, Radziwill)

Klinikum Fulda gAG, Department of Pharmacy and Patient Counselling, Pacelliallee 4, 36043 Fulda, Germany(Jacobs) European Institute for), S. 59. DOI: 10.1186/1477-7525-8-59.

*Holmberg C.; Muckelbauer R.; Sarganas G.; Braun V.; Heintze C.; Dini L.; Muller-Nordhorn J. (2018): Primary Prevention in General Medical Practice: A Survey. In: *Gesundheitswesen (Bundesverband der Arzte des Offentlichen Gesundheitsdienstes (Germany))* 80 (5), S. 465–470. DOI: 10.1055/s-0042-113601.

Holmberg C.; Sarganas G.; Mittring N.; Braun V.; Dini L.; Heintze C. et al. (2014): Primary prevention in general practice - views of German general practitioners: a mixed-methods study. In: *BMC family practice* 15 ((Holmberg, Sarganas, Mittring, Braun, Dini, Heintze, Rieckmann, Muckelbauer, Muller-Nordhorn) Berlin School of Public Health, Charite Universitätsmedizin Berlin, Seestr, 73, 13347 Berlin, Germany), S. 103. DOI: 10.1186/1471-2296-15-103.

Holzapfel, C.; Cresswell, L.; Ahern, A. L.; Fuller, N. R.; Eberhard, M.; Stoll, J. et al. (2014): The challenge of a 2-year follow-up after intervention for weight loss in primary care. In: *International journal of obesity (2005)* 38 (6), S. 806–811. DOI: 10.1038/ijo.2013.180.

Holzel, Lars P.; Ries, Zivile; Kriston, Levente; Dirmaier, Jorg; Zill, Jordis M.; Rummel-Kluge, Christine et al. (2016): Effects of culture-sensitive adaptation of patient information material on usefulness in migrants: a multicentre, blinded randomised controlled trial. In: *BMJ Open* 6 (11), e012008. DOI: 10.1136/bmjopen-2016-012008.

Holzel, Lars P.; Ries, Zivile; Zill, Jordis M.; Kriston, Levente; Dirmaier, Jorg; Harter, Martin; Bermejo, Isaac (2014): Development and testing of culturally sensitive patient information material for Turkish, Polish, Russian and Italian migrants with depression or chronic low back pain (KULTINFO): study protocol for a double-blind randomized controlled trial. In: *Trials* 15 (101263253), S. 265. DOI: 10.1186/1745-6215-15-265.

*Holzel L.P.; Vollmer M.; Kriston L.; Siegel A.; Harter M. (2012): Patient participation in medical decision making within an integrated health care system in Germany: Results of a controlled cohort study. In: *Bundesgesundheitsblatt - Gesundheitsforschung - Gesundheitsschutz* 55 (11-12), S. 1524–1533. DOI: 10.1007/s00103-012-1567-3.

*Hua T.D.; Vormfelde S.V.; Abed M.A.; Schneider-Rudt H.; Sobotta P.; Chenot J.-F. (2010): Oral anticoagulation therapy in family medicine. In: *Zeitschrift fur Allgemeinmedizin* 86 (10), S. 382–389. DOI: 10.3238/zfa.2010.0382.

Hua T.D.; Vormfelde S.V.; Abed M.A.; Schneider-Rudt H.; Sobotta P.; Friede T.; Chenot J.F. (2011): Practice nurse-based, individual and video-assisted patient education in oral anticoagulation--protocol of a cluster-randomized controlled trial. In: *BMC family practice* 12 ((Hua) Department of General Practice, University Medical Centre Gottingen, Humboldtallee 38, Gottingen, 37073, Germany), S. 17. DOI: 10.1186/1471-2296-12-17.

Hubner-Liebermann B.; Neuner T.; Hegerl U.; Hajak G.; Spiessl H. (2010): Reducing suicides through an alliance against depression? In: *General Hospital Psychiatry* 32 (5), S. 514–518. DOI: 10.1016/j.genhosppsych.2010.06.008.

*Hutt H.J.; Bennerscheidt P.; Thiel B.; Arand M. (2010): Immunosenescence and vaccinations in the elderly. In: *Medizinische Klinik* 105 (11), S. 802–807. DOI: 10.1007/s00063-010-1137-0.

*Huy C.; Diehm C.; Schneider S. (2012): Cardiovascular prevention at the general practitioner? First results of a study on attitudes, services, success and barriers in practice. In: *Deutsche Medizinische Wochenschrift* 137 (1-2), S. 17–22. DOI: 10.1055/s-0031-1292884.

Jacob, Louis; Kostev, Karel (2017): Compliance with vaccination against hepatitis A virus in Germany: A retrospective analysis. In: *International Journal of Clinical Pharmacology and Therapeutics* 55 (9), S. 740–745. DOI: 10.5414/CP202990.

Jacobs, Esther; Tamayo, Miguel; Rosenbauer, Joachim; Schulze, Matthias B.; Kuss, Oliver; Rathmann, Wolfgang (2018): Protocol of a cluster randomized trial to investigate the impact of a type 2 diabetes risk prediction model on change in physical activity in primary care. In: *BMC endocrine disorders* 18 (1), S. 72. DOI: 10.1186/s12902-018-0299-2.

Jager C.; Szecsenyi J.; Steinhauser J. (2015): Design and delivery of a tailored intervention to implement recommendations for multimorbid patients receiving polypharmacy into primary care practices. In: *BioMed Research International* 2015 ((Jager, Szecsenyi, Steinhauser) Department of General Practice and Health Services Research, University Hospital Heidelberg, Vosstrasse 2, Heidelberg 69115, Germany), S. 938069. DOI: 10.1155/2015/938069.

*Jahne, A.; Ruther, T.; Deest, H.; Gehrig, H.; Zeeuw, J. de; Alberti, A.; Mulzer, K. (2014): [Evaluation of a structured smoking cessation program for primary care medicine]. In: *Deutsche medizinische Wochenschrift (1946)* 139 (4), S. 127–133. DOI: 10.1055/s-0033-1349646.

*Jauss M.; Grond M. (2005): Progress in ischemic brain infarction: Acute treatment of stroke by the general practitioner, the emergency physician and in the hospital. In: *Notfall und Hausarztmedizin* 31 (5), S. 214–221. DOI: 10.1055/s-2005-871840.

Jung, Julia; Neumann, Melanie; Wirtz, Markus; Ernstmann, Nicole; Staratschek-Jox, Andrea; Wolf, Jurgen; Pfaff, Holger (2010): Validation of the "SmoCess-GP" instrument - a short patient questionnaire for assessing the smoking cessation activities of general practitioners: a cross-sectional study. In: *BMC family practice* 11 (100967792), S. 9. DOI: 10.1186/1471-2296-11-9.

*Junius U.; Fischer G. (2002): Geriatric assessment for use in primary care - Results of a concerted action involving seven European countries. In: *Zeitschrift fur Gerontologie und Geriatrie* 35 (3), S. 210–223. DOI: 10.1007/s00391-002-0069-6.

*Junius U.; Schmidt C.; Fischer G.; Breull A.; Langner D. (2003): The European geriatric assessment in general practice: Results of the German feasibility and acceptance study. In: *Zeitschrift für Allgemeinmedizin* 79 (12), S. 620–623.

*Junius U.; Schultz C.; Fischer G.; Breull A.; Langner D. (2003): Evidence based preventative assessment for older patients: The implementation of the results of a European study in a concrete medical check-up in Germany. In: *Zeitschrift für Allgemeinmedizin* 79 (3), S. 143–148.

Jurges H.; Pohl V. (2012): Medical guidelines, physician density, and quality of care: Evidence from German SHARE data. In: *European Journal of Health Economics* 13 (5), S. 635–649. DOI: 10.1007/s10198-011-0372-5.

*Kaiser P.; Noack A.; Donner-Banzhoff N.; Keller S.; Baum E. (2005): Comparing physicians' and lawyers' health behaviour. In: *Zeitschrift für Allgemeinmedizin* 81 (10), S. 419–422. DOI: 10.1055/s-2005-836914.

*Kamtsiuris, P.; Bergmann, E.; Rattay, P.; Schlaud, M. (2007): [Use of medical services. Results of the German Health Interview and Examination Survey for Children and Adolescents (KiGGS)]. In: *Bundesgesundheitsblatt, Gesundheitsforschung, Gesundheitsschutz* 50 (5-6), S. 836–850.

Katsarava, Zaza; Mania, Maka; Lampl, Christian; Herberhold, Johanna; Steiner, Timothy J. (2018): Poor medical care for people with migraine in Europe - evidence from the Eurolight study. In: *The journal of headache and pain* 19 (1), S. 10. DOI: 10.1186/s10194-018-0839-1.

*Kaufmann D. (2013): Diabetic retinopathy. In: *Medizinische Welt* 64 (6), S. 349–355. Online verfügbar unter <http://www.schattauer.de/de/magazine/uebersicht/zeitschriften-a-z/die-medizinische-welt/issue/special/manuscript/20577/download.html>.

*Keil, Jan; Brendler, Verena; Sachse, Catherina; Zulke, Andrea; Zeynalova, Samira; Engel, Christoph et al. (2020): [Gender-Specific Differences in the Utilization of Health Care Services in an Urban Population Sample]. In: *Gesundheitswesen (Bundesverband der Ärzte des Öffentlichen Gesundheitsdienstes (Germany))* 82 (3), e17-e23. DOI: 10.1055/a-0820-3584.

Keller, Heidemarie; Krones, Tanja; Becker, Annette; Hirsch, Oliver; Sonnichsen, Andreas C.; Popert, Uwe et al. (2012): Arriba: effects of an educational intervention on prescribing behaviour in prevention of CVD in general practice. In: *European journal of preventive cardiology* 19 (3), S. 322–329. DOI: 10.1177/1741826711404502.

Keller H.; Hirsch O.; Kaufmann-Kolle P.; Krones T.; Becker A.; Sonnichsen A.C. et al. (2013): Evaluating an implementation strategy in cardiovascular prevention to improve prescribing of statins in Germany: an intention to treat analysis. In: *BMC Public Health* 13 ((Hirsch, Kaufmann-Kolle, Krones, Becker, Sonnichsen, Baum, Donner-Banzhoff) Department of General Practice/Family Medicine, Philipps University of Marburg, Karl-von-Frisch-Strasse 4, Marburg, 35043, Germany), S. 623. DOI: 10.1186/1471-2458-13-623.

*Keller H.; Krones T.; Sonnichsen A.; Sadowski E.; Popert U.; Rochon J. et al. (2007): Medical prevention of cardiovascular disease in general practice: Is medical treatment appropriate to global risk? In: *Zeitschrift für Allgemeinmedizin* 83 (9), S. 359–364. DOI: 10.1055/s-2007-985833.

*Keller M.; Grau A. (2018): Recent developments in secondary stroke prevention and aftercare. In: *Internistische Praxis* 59 (3), S. 386–395. Online verfügbar unter http://cme.marseille-verlag.com/Site/Content/ejournals/proxy.php?file=ip-059-03_online.pdf&rid=418.

*Kellner, Christiane; Kuniss, Nadine; Kloos, Christof; Muller, Ulrich Alfons; Muller, Nicolle (2018): No selection, but higher satisfaction of people participating in the disease management programme diabetes type 2 in Germany. In: *Acta diabetologica* 55 (4), S. 363–367. DOI: 10.1007/s00592-018-1106-2.

*Kiel, Simone; Zimak, Carolin; Schmidt, Carsten Oliver; Chenot, Jean-Francois (2020): [Effect of ambulatory geriatric rehabilitation on polypharmacy and potentially inappropriate medication-an observational study]. In: *Zeitschrift für Gerontologie und Geriatrie* 53 (5), S. 416–422. DOI: 10.1007/s00391-019-01589-1.

Kiel S.; Zimak C.; Chenot J.-F.; Schmidt C.O. (2020): Evaluation of an ambulatory geriatric rehabilitation program - results of a matched cohort study based on claims data. In: *BMC geriatrics* 20 (1), S. 30. DOI: 10.1186/s12877-020-1415-5.

Kircher T.T.J.; Wormstall H.; Muller P.H.; Schwarzler F.; Buchkremer G.; Wild K. et al. (2007): A randomised trial of a geriatric evaluation and management consultation services in frail hospitalised patients. In: *Age and Ageing* 36 (1), S. 36–42. DOI: 10.1093/ageing/afl102.

Klein G.; Ulbricht S.; Haug S.; Gross B.; Rumpf H.-J.; John U.; Meyer C. (2010): Effects of practitioner-delivered brief counseling and computer-generated tailored letters on cigarettes per day among smokers who do not quit-A quasi-randomized controlled trial. In: *Drug and alcohol dependence* 112 (1-2), S. 81–89. DOI: 10.1016/j.drugalcdep.2010.05.016.

*Klement, Andreas; Oemler, Matthias; Wienke, Andreas; Richter, Matthias; Wolfradt, Uwe (2015): [(Expected) Consultation length, mental (co-)morbidity and patient satisfaction in the family practice encounter]. In: *Zeitschrift für Evidenz, Fortbildung und Qualität im Gesundheitswesen* 109 (8), S. 560–569. DOI: 10.1016/j.zefq.2015.08.005.

*Klement A.; Oemler M.; Braun K.; Unverzagt S. (2015): Strategies for the implementation of guidelines: Synopsis of a systematic review. In: *Zeitschrift für Allgemeinmedizin* 91 (2), S. 57–63. DOI: 10.3238/zfa.2015.0057-0063.

Klett-Tammen, Carolina Judith; Krause, Gerard; Lengerke, Thomas von; Castell, Stefanie (2016): Advising vaccinations for the elderly: a cross-sectional survey on differences between general practitioners and physician assistants in Germany. In: *BMC family practice* 17 (100967792), S. 98. DOI: 10.1186/s12875-016-0502-3.

*Kliche T.; Mann R. (2008): Prevention and care for obese children and adolescents in Germany: Quality and efficiencies of programs and interventions. In: *Bundesgesundheitsblatt - Gesundheitsforschung - Gesundheitsschutz* 51 (6), S. 646–656. DOI: 10.1007/s00103-008-0541-6.

*Kling C.; Hansmann-Wiest J.; Kabelitz D. (2017): Immunosuppression and vaccination in family practices. In: *Zeitschrift fur Allgemeinmedizin* 93 (2), S. 56–63. DOI: 10.3238/zfa.2017.0056-0063.

*Klingenber A.; Brand T.; Andres E.; Kaufmann-Kolle P.; Wambach V.; Szecsenyi J. (2019): What do patients know about antibiotics and how often do they expect an antibiotic prescription? In: *Zeitschrift fur Allgemeinmedizin* 2019 (5), S. 198–202. DOI: 10.3238/zfa.2019.0198-0202.

Kloos, C.; Muller, N.; Wolf, G.; Hartmann, P.; Lehmann, T.; Muller, U. A. (2011): Better HbA1c and blood pressure control in patients with diabetes mellitus treated at a primary health care level 10 years after initiation of a diabetes quality improvement program. In: *Experimental and clinical endocrinology & diabetes : official journal, German Society of Endocrinology [and] German Diabetes Association* 119 (8), S. 459–462. DOI: 10.1055/s-0031-1273746.

*Klosterhalfen S.; Funke C.; Schneider A.; Kotz D. (2020): The assessment of burden of COPD (ABC) tool" in family medicine results of a feasibility study. In: *Zeitschrift fur Allgemeinmedizin* 96 (5), S. 205–210. DOI: 10.3238/zfa.2020.0205-0210.

*Knesebeck O.V.D.; Mielck A. (2009): Social inequality and health care among the aged. In: *Zeitschrift fur Gerontologie und Geriatrie* 42 (1), S. 39–46. DOI: 10.1007/s00391-008-0522-2.

*Knobe, M.; Rasche, P.; Rentemeister, L.; Bliemel, C.; Bucking, B.; Bollheimer, L. C.; Pape, H-C (2018): [Evaluation of a simple screening tool for ambulant fall prevention]. In: *Der Unfallchirurg* 121 (11), S. 901–910. DOI: 10.1007/s00113-018-0462-2.

*Knoll T.; Bach T.; Humke U.; Neisius A.; Stein R.; Schonthaler M.; Wendt-Nordahl G. (2016): S2k guidelines on diagnostics, therapy and metaphylaxis of urolithiasis (AWMF 043/025): Compendium. In: *Urologe* 55 (7), S. 904–922. DOI: 10.1007/s00120-016-0133-2.

Koch, Judith; Hellenbrand, Wiebke; Schink, Susanne; Wichmann, Ole; Carganico, Andreas; Drewes, Jochen et al. (2016): Evaluation of a temporary vaccination recommendation in response to an outbreak of invasive meningococcal serogroup C disease in men who have sex with men in Berlin, 2013-2014. In: *Euro surveillance : bulletin European sur les maladies transmissibles = European communicable disease bulletin* 21 (5), S. 12–22. DOI: 10.2807/1560-7917.ES.2016.21.5.30122.

Kogler, Harald; Metzdorf, Norbert; Glaab, Thomas; Welte, Tobias (2010): Preselection of patients at risk for COPD by two simple screening questions. In: *Respiratory medicine* 104 (7), S. 1012–1019. DOI: 10.1016/j.rmed.2010.01.005.

*Kohn D.; Pukrop R.; Niedersteberg A.; Schultze-Lutter F.; Ruhrmann S.; Bechdolf A. et al. (2004): Pathways to care: Help-seeking behavior in first-episode psychosis. In: *Fortschritte der Neurologie Psychiatrie* 72 (11), S. 635–642. DOI: 10.1055/s-2004-818418.

*Kotter T. (2016): "Family practitioners' counseling regarding PSA screening": practice recommendation of the German college of general practitioners and family physicians (DEGAM). In: *Zeitschrift fur Allgemeinmedizin* 92 (12), S. 495–499. DOI: 10.3238/zfa.2016.0495-0499.

*Kowall B.; Rathmann W.; Landgraf R. (2014): Use of risk scores for type 2 diabetes in clinical practice. In: *Diabetologe* 10 (7), S. 547–553. DOI: 10.1007/s11428-014-1212-x.

*Kranz, J.; Schmidt, S.; Lebert, C.; Schneidewind, L.; Vahlensieck, W.; Sester, U. et al. (2017): [Epidemiology, diagnostics, therapy, prevention and management of uncomplicated bacterial outpatient acquired urinary tract infections in adult patients : Update 2017 of the interdisciplinary AWMF S3 guideline]. In: *Der Urologe. Ausg. A* 56 (6), S. 746–758. DOI: 10.1007/s00120-017-0389-1.

*Krasselt M.; Baerwald C.; Seifert O. (2018): Insufficient vaccination rates in patients with systemic lupus erythematosus in a German outpatient clinic. In: *Zeitschrift fur Rheumatologie* 77 (8), S. 727–734. DOI: 10.1007/s00393-017-0410-5.

Krause O.; Wiese B.; Doyle I.-M.; Kirsch C.; Thurmann P.; Wilm S. et al. (2019): Multidisciplinary intervention to improve medication safety in nursing home residents: protocol of a cluster randomised controlled trial (HIOPP-3-iTBX study). In: *BMC geriatrics* 19 (1), S. 24. DOI: 10.1186/s12877-019-1027-0.

Krobot K.J.; Wagner A.; Siebert U. (2012): Risk factor levels, risk factor combinations, and residual coronary risk: Population-based estimates for secondary prevention patients using statins. In: *European Journal of Cardiovascular Prevention and Rehabilitation* 19 (1), S. 109–117. DOI: 10.1177/1741826710392668.

*Kroger, Knut (2009): [Prophylaxis of deep vein thrombosis with enoxaparin 40 mg in outpatients compared to hospitalized medically ill patients]. In: *Medizinische Klinik (Munich, Germany : 1983)* 104 (8), S. 608–613. DOI: 10.1007/s00063-009-1132-5.

Kruger K.; Leppkes N.; Gehrke-Beck S.; Herrmann W.; Algharably E.A.; Kreutz R. et al. (2018): Improving long-term adherence to statin therapy: A qualitative study of GPS' experiences in primary care. In: *British Journal of General Practice* 68 (671), e401-e407. DOI: 10.3399/bjgp18X696173.

Kuchenbecker, Ulrike; Chase, Daniela; Reichert, Anika; Schiffner-Rohe, Julia; Atwood, Mark (2018): Estimating the cost-effectiveness of a sequential pneumococcal vaccination program for adults in Germany. In: *PLoS ONE* 13 (5), e0197905. DOI: 10.1371/journal.pone.0197905.

*Kunstmann W.; Wollgarten D.; Vollenbroich R.; Hildenbrand F.; Grunert J.-F. (2002): Integrated learning in doctor's practices from 1st year on - An innovative curriculum in family medicine at Witten/Herdecke University. In: *Medizinische Ausbildung* 19 (2), S. 213–217.

*Kupper, B.; Krause, P.; Glaesmer, H.; Wittchen, H-U (2004): [How do risk patients with overweight/obesity differ in their health knowledge and behaviour from patients with normal weight? A study in primary medical care]. In: *Gesundheitswesen (Bundesverband der Ärzte des Öffentlichen Gesundheitsdienstes (Germany))* 66 (6), S. 361–369.

*Kurch-Bek, Diana; Gallowitz, Christian; Tenckhoff, Bernhard; Sundmacher, Leonie (2019): [Influenza vaccination rates of patients with statutory ambulatory medical care in protective facilities and in nursing and special care homes]. In: *Bundesgesundheitsblatt, Gesundheitsforschung, Gesundheitsschutz* 62 (1), S. 84–93. DOI: 10.1007/s00103-018-2854-4.

*Kuver C.; Becker A.; Ludt S. (2008): Counseling and education for people with chronic illness. DEGAM series: Comprehensive care for patients with chronic illness. In: *Zeitschrift für Allgemeinmedizin* 84 (11), S. 471–476. DOI: 10.1055/s-0028-1093327.

*Labenz, Christian; Adarkwah, Charles C.; Worns, Marcus-Alexander; Miehlke, Stephan; Hofmann, Wolf P.; Buggisch, Peter et al. (2020): Management of hepatic encephalopathy in Germany: a survey among physicians. In: *Zeitschrift für Gastroenterologie* 58 (1), S. 49–56. DOI: 10.1055/a-1010-6974.

Lahiri K.; Rettig-Ewen V.; Bohm M.; Laufs U. (2007): Perceived psychosocial stress and cardiovascular risk factors in obese and non-obese patients. In: *Clinical Research in Cardiology* 96 (6), S. 365–374. DOI: 10.1007/s00392-007-0512-1.

*Landgraf M.N.; Heinen F. (2017): Diagnostics of fetal alcohol spectrum disorders in pediatrics: Recommendations for the practice according to the S3 guidelines. In: *Monatsschrift für Kinderheilkunde* 165 (9), S. 786–793. DOI: 10.1007/s00112-016-0191-y.

*Lang E.; Kastner S.; Liebig K.; Neundorfer B. (2002): Interventions for improvement of primary care in patients with low back pain: How effective are advice to primary care physicians on therapies and a multimodal therapy program arising out of cooperation of outpatient health care structures? In: *Schmerz* 16 (1), S. 22–23. DOI: 10.1007/s004820100091.

Lang E.; Liebig K.; Kastner S.; Neundorfer B.; Heuschmann P. (2003): Multidisciplinary rehabilitation versus usual care for chronic low back pain in the community: Effects on quality of life. In: *Spine Journal* 3 (4), S. 270–276. DOI: 10.1016/S1529-9430(03)00028-7.

*Lau D.; Klein R.; Steinkohl M. (2003): Prevention through focused nutritional advice in general practice. In: *Zeitschrift für Allgemeinmedizin* 79 (5), S. 234–237.

*Lehmkuhl, Dieter (2019): [Climate change and its significance in the healthcare community: history, landmarks, and major players]. In: *Bundesgesundheitsblatt, Gesundheitsforschung, Gesundheitsschutz* 62 (5), S. 546–555. DOI: 10.1007/s00103-019-02935-9.

Lehrke, P.; Nuebling, M.; Hofmann, F.; Stoessel, U. (2001): Attitudes of homoeopathic physicians towards vaccination. In: *Vaccine* 19 (32), S. 4859–4864.

Leiter U.; Hepp M.V.; Steeb T.; Amaral T.; Bauer A.; Becker J.C. et al. (2020): S3 guideline for actinic keratosis and cutaneous squamous cell carcinoma (cSCC) - short version, part 2: epidemiology, surgical and systemic treatment of cSCC, follow-up, prevention and occupational disease. In: *JDDG - Journal of the German Society of Dermatology* 18 (4), S. 400–413. DOI: 10.1111/ddg.14072.

Lempp T.; Heinzel-Gutenbrunner M.; Bachmann C. (2016): Child and adolescent psychiatry: which knowledge and skills do primary care physicians need to have? A survey in general practitioners and paediatricians. In: *European Child and Adolescent Psychiatry* 25 (4), S. 443–451. DOI: 10.1007/s00787-015-0757-6.

Leonhardt C.; Keller S.; Chenot J.-F.; Luckmann J.; Basler H.-D.; Wegscheider K. et al. (2008): TTM-based motivational counselling does not increase physical activity of low back pain patients in a primary care setting-A cluster-randomized controlled trial. In: *Patient Education and Counseling* 70 (1), S. 50–60. DOI: 10.1016/j.pec.2007.09.018.

Linden, Michael; Christof, Tatjana; Rentzsch, Christiane (2008): Contents of general practitioner-patient consultations in the treatment of depression. In: *Journal of general internal medicine* 23 (10), S. 1567–1570. DOI: 10.1007/s11606-008-0648-3.

Loeffert S.; Ommen O.; Ernstmann N.; Pfaff H. (2009): Changes in secondary pharmacological prevention of acute coronary syndromes and stroke after hospital discharge: A 6-month follow-up study of German primary care patients. In: *Journal of Public Health* 17 (1), S. 3–7. DOI: 10.1007/s10389-008-0210-0.

Loerch, B.; Szegedi, A.; Kohnen, R.; Benkert, O. (2000): The primary care evaluation of mental disorders (PRIME-MD), German version: a comparison with the CIDI. In: *Journal of psychiatric research* 34 (3), S. 211–220.

*Lohler, Jan; Lehmann, Martin; Segler, Veronika; Volkenstein, Stefan; Battmer, Rolf-Dieter; Ernst, Arneborg et al. (2019): [The Mini-Audio-Test (MAT) - a screening method on hearing impairment to be used by general practitioners and specialized physicians]. In: *Laryngo- rhino- otologie* 98 (1), S. 27–34. DOI: 10.1055/a-0805-5741.

*Lollgen H.; Volker K.; Bockenhoff A.; Lollgen D. (2006): Physical activity and prevention of cardiovascular diseases. In: *Herz* 31 (6), S. 519–523. DOI: 10.1007/s00059-006-2859-x.

*Lorenz H.-J.; Reeg J.; Welke J.; Schwantes U.; Herrmann M. (2006): Comparison of patients' subjective assessment of and objective cardiovascular risk dependent on socio-economic status. In: *Zeitschrift für Allgemeinmedizin* 82 (10), S. 435–440. DOI: 10.1055/s-2006-942192.

*Low M.; Stegmaier C.; Ziegler H.; Rothenbacher D.; Brenner H. (2004): Epidemiological investigations of the chances of preventing, recognizing early and optimally treating chronic diseases in an elderly population (ESTHER study). In: *Deutsche Medizinische Wochenschrift* 129 (49), S. 2643–2647. DOI: 10.1055/s-2004-836089.

Luck T.; Riedel-Heller S.G.; Kaduszkiewicz H.; Bickel H.; Jessen F.; Pentzek M. et al. (2007): Mild cognitive impairment in general practice: Age-specific prevalence and correlate results from the German study on ageing, cognition and dementia in primary care patients (AgeCoDe). In: *Dementia and Geriatric Cognitive Disorders* 24 (4), S. 307–316. DOI: 10.1159/000108099.

*Ludt S.; Campbell S.M.; Wensing M.; Van Lieshout J.; Boffin N.; Grol R.; Szecsenyi J. (2008): Prevention and risk-management of cardiovascular disease in European primary care - EPA-CARDIO - Development of quality indicators: Results from Germany. In: *Zeitschrift für Allgemeinmedizin* 84 (11), S. 477–488. DOI: 10.1055/s-0028-1086033.

Lullau, Anja; Petroff, David; Batz, Olaf; Kramer, Jan; Jedrysiak, Katrin; Tenckhoff, Hannelore et al. (2018): Linkage to care of HbsAg-positive and anti-HCV-positive patients after a systematic screening approach in the German primary care setting. In: *European journal of gastroenterology & hepatology* 30 (3), S. 280–283. DOI: 10.1097/MEG.00000000000001052.

*Lutgehetmann, M.; Meyer, F.; Volz, T.; Lohse, A. W.; Fischer, C.; Dandri, M.; Petersen, Jorg (2010): [Knowledge about HBV, prevention behaviour and treatment adherence of patients with chronic hepatitis B in a large referral centre in Germany]. In: *Zeitschrift für Gastroenterologie* 48 (9), S. 1126–1132. DOI: 10.1055/s-0029-1245304.

*Luttje, D.; Varwig, D.; Teigel, B.; Gilhaus, B. (2011): [Geriatric assessment. Development, status quo and perspectives]. In: *Der Internist* 52 (8), S. 925–933. DOI: 10.1007/s00108-011-2803-x.

Maikranz V.; Siebenhofer A.; Ulrich L.-R.; Mergenthal K.; Schulz-Rothe S.; Kemperdick B. et al. (2017): Does a complex intervention increase patient knowledge about oral anticoagulation? - a cluster-randomised controlled trial. In: *BMC family practice* 18 (1), S. 15. DOI: 10.1186/s12875-017-0588-2.

*Manthey, Jakob; Lindemann, Christina; Verthein, Uwe; Frischknecht, Ulrich; Kraus, Ludwig; Reimer, Jens et al. (2020): [Provision of healthcare for people with risky alcohol use and severe alcohol use disorders in the state of Bremen, Germany: demand and guideline concordance?]. In: *Bundesgesundheitsblatt, Gesundheitsforschung, Gesundheitsschutz* 63 (1), S. 122–130. DOI: 10.1007/s00103-019-03072-z.

Marcus, Ulrich; Gassowski, Martyna; Kruspe, Martin; Drewes, Jochen (2015): Recency and frequency of HIV testing among men who have sex with men in Germany and socio-demographic factors associated with testing behaviour. In: *BMC Public Health* 15 (100968562), S. 727. DOI: 10.1186/s12889-015-1945-5.

Martin S.; Schramm W.; Schneider B.; Neeser K.; Weber C.; Lodwig V. et al. (2007): Epidemiology of complications and total treatment costs from diagnosis of type 2 diabetes in Germany (ROSSO 4). In: *Experimental and Clinical Endocrinology and Diabetes* 115 (8), S. 495–501. DOI: 10.1055/s-2007-981470.

Matrisch, M.; Trampisch, U.; Klaassen-Mielke, R.; Pientka, L.; Trampisch, H. J.; Thiem, U. (2012): [Screening for dementia using telephone interviews. An evaluation and reliability study of the Telephone Interview for Cognitive Status (TICS) in its modified German version]. In: *Zeitschrift für Gerontologie und Geriatrie* 45 (3), S. 218–223. DOI: 10.1007/s00391-011-0220-3.

*Mau W.; Reuter S. (2011): Health promotion, prevention, and rehabilitation in the elderly. In: *Deutsche Medizinische Wochenschrift* 136 (43), S. 2199–2204. DOI: 10.1055/s-0031-1292033.

*Mayer, Hermann (2006): [Diagnosis and treatment of overweight in children and adolescents]. In: *MMW Fortschritte der Medizin* 148 (40), S. 38–40.

Mayer J.E.; Swetter S.M.; Fu T.; Geller A.C. (2014): Screening, early detection, education, and trends for melanoma: Current status (2007–2013) and future directions Part II. Screening, education, and future directions. In: *Journal of the American Academy of Dermatology* 71 (4), S. 611. DOI: 10.1016/j.jaad.2014.05.045.

*Meffert C.; Mittag O.; Jackel W.H. (2013): Work place health promotion programmes of the statutory German pension insurance. In: *Rehabilitation (Germany)* 52 (6), S. 391–398. DOI: 10.1055/s-0032-1333220.

Mehring, Michael; Donnachie, Ewan; Mutschler, Robert; Hofmann, Frank; Keller, Manfred; Schneider, Antonius (2013): Disease management programs for patients with asthma in Germany: a longitudinal population-based study. In: *Respiratory care* 58 (7), S. 1170–1177.

Mehring M.; Haag M.; Linde K.; Wagenpfeil S.; Frensch F.; Blome J.; Schneider A. (2013): Effects of a general practice guided web-based weight reduction program--results of a cluster-randomized controlled trial. In: *BMC family practice* 14 ((Mehring) Institute of General Practice, Technische Universität München, Orleansstr. 47, Munich 81667, Germany), S. 76.

*Meinck M.; Lubke N.; Lauterberg J.; Robra B.-P. (2004): Preventive home visits to the elderly: Systematic review of available evidence. In: *Gesundheitswesen* 66 (11), S. 732–738. DOI: 10.1055/s-2004-813779.

Meyer, Christian; Ulbricht, Sabina; Baumeister, Sebastian E.; Schumann, Anja; Ruge, Jeannette; Bischof, Gallus et al. (2008): Proactive interventions for smoking cessation in general medical practice: a quasi-randomized controlled trial to examine the efficacy of computer-tailored letters and physician-delivered brief advice. In: *Addiction (Abingdon, England)* 103 (2), S. 294–304.

*Meyer F. (2004): Is it useful to establish a "Do-it-yourself" blood-pressure measurement station in a general practice? In: *Zeitschrift fur Allgemeinmedizin* 80 (10), S. 421–424. DOI: 10.1055/s-2004-820423.

Michaelis M.; Balint E.M.; Junne F.; Zipfel S.; Gundel H.; Lange R. et al. (2019): Who should play a key role in preventing common mental disorders that affect employees in the workplace? Results of a survey with occupational health physicians, primary care physicians, psychotherapists, and human resource managers. In: *International Journal of Environmental Research and Public Health* 16 (8), S. 1383. DOI: 10.3390/ijerph16081383.

Michels I.I.; Stover H.; Gerlach R. (2007): Substitution treatment for opioid addicts in Germany. In: *Harm Reduction Journal* 4 ((Michels) Office of the Federal Drug Commissioner, Federal Ministry of Health, Berlin, Germany(Stover) Bremen Institute of Drug Research, University of Bremen, Bremen, Germany(Gerlach) Institute for the Advancement of Qualitative Drug Research (INDRO), Mu), S. 5. DOI: 10.1186/1477-7517-4-5.

*Micke O.; Vormann J.; Classen H.-G.; Kisters K. (2020): Magnesium: Relevance for general practitioners - A position paper of the Society for Magnesium Research e V. In: *Deutsche Medizinische Wochenschrift* 145 (22), S. 1628–1634. DOI: 10.1055/a-1166-7229.

*Mildner C.; Kramer I. (2013): Close the gap - Reduction of supply and information gaps at the interface of inpatient and outpatient treatment through participation of the hospital pharmacist. In: *Krankenhauspharmazie* 34 (4), S. 187–196.

*Moerchel C.; Kroeger K. (2007): Thromboprophylaxis in Germany. In: *Deutsches Arzteblatt* 104 (42), S. 2886–2893.

*Moller J.; Kuver C.; Beyer M.; Gerlach F.M. (2004): Patient Education in General Practice as a Contribution to Disease Management: A Qualitative Study of the Motivation and Obstacles in the Implementation of Patient Education for Type-2 Diabetics. In: *Zeitschrift fur Allgemeinmedizin* 80 (4), S. 146–149.

*Mols V.; Jahn H.; Hetzel A.; Luckner A.; Kampmann M.; Niebling W. (2005): Quality circles in the secondary prevention of stroke - A controlled interventional study. In: *Zeitschrift fur Allgemeinmedizin* 81 (10), S. 435–441. DOI: 10.1055/s-2005-836913.

*Mortsiefer, A.; Ludt, S.; Pentzek, M.; Wilm, S.; Brotons, C. (2014): [Patients' needs and expectations concerning prevention and health advice in primary care. Results from the EUROPREVIEW survey]. In: *Gesundheitswesen (Bundesverband der Arzte des Offentlichen Gesundheitsdienstes (Germany))* 76 (7), S. 417–422. DOI: 10.1055/s-0034-1381985.

*Mortsiefer A.; Abholz H.-H.; Schumacher M.; Wegscheider K.; In Der Schmitten J. (2013): Treatment of patients with hypertension in consideration of their global cardiovascular risk: Cross-sectional study in 89 German family practices. In: *Zeitschrift fur Allgemeinmedizin* 89 (7-8), S. 311–316. DOI: 10.3238/zfa.2013.0311-0316.

Mortsiefer A.; Meysen T.; Schumacher M.; Lintges C.; Stamer M.; Schmacke N. et al. (2008): CRISTOPH - A cluster-randomised intervention study to optimise the treatment of patients with hypertension in General Practice. In: *BMC family practice* 9 ((Mortsiefer, Meysen, Schumacher, Lintges, Abholz, In Der Schmitten) Department of General Practice, University Hospital, P.O. Box 101001, 40225 Dusseldorf, Germany(Stamer, Schmacke) University of Bremen, Bibliothekstr. 1, 28359 Bremen, Germany(Wegscheider), S. 33. DOI: 10.1186/1471-2296-9-33.

Mortsiefer A.; Wilm S.; Santos S.; Loscher S.; Wollny A.; Drewelow E. et al. (2020): Family conferences and shared prioritisation to improve patient safety in the frail elderly (COFRAIL): Study protocol of a cluster randomised intervention trial in primary care. In: *Trials* 21 (1), S. 285. DOI: 10.1186/s13063-020-4182-x.

Moschny A.; Platen P.; Klaassen-Mielke R.; Trampisch U.; Hinrichs T. (2011): Physical activity patterns in older men and women in Germany: a cross-sectional study. In: *BMC Public Health* 11 ((Moschny) Department of Sports Medicine and Sports Nutrition, Ruhr-University Bochum, Germany), S. 559.

*Moshammer, Dirk; Natanzon, Iris; Manske, Ira; Grutschkowski, Philipp; Rieger, Monika A. (2012): [Deficiencies and barriers of the cooperation between German general practitioners and occupational health physicians? A qualitative content analysis of focus groups]. In: *Zeitschrift fur Evidenz, Fortbildung und Qualitat im Gesundheitswesen* 106 (9), S. 639–648. DOI: 10.1016/j.zefq.2011.09.027.

Moshammer, Dirk; Natanzon, Iris; Manske, Ira; Grutschkowski, Philipp; Rieger, Monika A. (2014): Cooperation between general practitioners and occupational health physicians in Germany: how can it be optimised? A qualitative study. In: *International archives of occupational and environmental health* 87 (2), S. 137–146. DOI: 10.1007/s00420-013-0845-1.

Moshammer D.; Michaelis M.; Mehne J.; Wilm S.; Rieger M.A. (2016): General practitioners' and occupational health physicians' views on their cooperation: a cross-sectional postal survey. In: *International archives of occupational and environmental health* 89 (3), S. 449–459. DOI: 10.1007/s00420-015-1084-4.

*Moshammer D.; Muche R.; Hermes J.; Zollner I.; Lorenz G. (2009): Factors associated with influenza vaccination information - A cross-sectional study in elderly primary care patients. In: *Zeitschrift fur Evidenz, Fortbildung und Qualitat im Gesundheitswesen* 103 (7), S. 445–451. DOI: 10.1016/j.zefq.2009.03.001.

Muche-Borowski, Cathleen; Abiry, Dorit; Wagner, Hans-Otto; Barzel, Anne; Luhmann, Dagmar; Egidi, Gunther et al. (2018): Protection against the overuse and underuse of health care - methodological considerations for establishing prioritization criteria and recommendations in general practice. In: *BMC health services research* 18 (1), S. 768. DOI: 10.1186/s12913-018-3569-9.

Muehlig, Stephan; Hoch, Eva; Hoefler, Michael; Pittrow, David; Wittchen, Hans-Ulrich (2003): Aims, design and methods of the Smoking and Nicotine Dependence Awareness and Screening (SNICAS) study. In: *International journal of methods in psychiatric research* 12 (4), S. 208–228.

Mueller G.; Schumacher P.; Wetzlmaier J.; Pallauf M. (2016): Screening questionnaires to identify problem drinking in the primary care setting: a systematic review. In: *Journal of Public Health (Germany)* 24 (1), S. 9–19. DOI: 10.1007/s10389-015-0694-3.

*Muhlig S.; De Vries U.; Waldmann H.-C.; Petermann F. (2005): Evaluation of asthma patient education from the subjective perspective. In: *Pravention und Rehabilitation* 17 (1), S. 1–15.

Muller, D.; Szucs, T. D. (2007): Influenza vaccination coverage rates in 5 European countries: a population-based cross-sectional analysis of the seasons 02/03, 03/04 and 04/05. In: *Infection* 35 (5), S. 308–319.

*Muller, M.; Kogler, H.; Glaab, T.; Welte, T. (2012): [Use of a lung function screening device for identifying patients at risk for COPD in general practice]. In: *Pneumologie (Stuttgart, Germany)* 66 (11), S. 645–649. DOI: 10.1055/s-0032-1325791.

Muller-Buhl, Uwe; Laux, Gunter; Szecsenyi, Joachim (2011): Secondary Pharmacotherapeutic Prevention among German Primary Care Patients with Peripheral Arterial Disease. In: *International journal of vascular medicine* 2011 (101542608), S. 316496. DOI: 10.1155/2011/316496.

Muller-Riemenschneider F.; Holmberg C.; Rieckmann N.; Kliems H.; Rufer V.; Muller-Nordhorn J.; Willich S.N. (2010): Barriers to routine risk-score use for healthy primary care patients: Survey and qualitative study. In: *Archives of Internal Medicine* 170 (8), S. 719–724. DOI: 10.1001/archinternmed.2010.66.

Muth C.; van den Akker M.; Blom J.W.; Mallen C.D.; Rochon J.; Schellevis F.G. et al. (2014): The Ariadne principles: How to handle multimorbidity in primary care consultations. In: *BMC Medicine* 12 (1), S. 223. DOI: 10.1186/s12916-014-0223-1.

*Nass-Griegoleit I. (2002): Medical care in breast health - The expectations of healthy women. In: *Gynakologe* 35 (11), S. 1117–1120.

Netzer N.; Strohl K.; Faulhaber M.; Gatterer H.; Burtscher M. (2013): Hypoxia-related altitude illnesses. In: *Journal of Travel Medicine* 20 (4), S. 247–255. DOI: 10.1111/jtm.12017.

Neufeind J.; Betsch C.; Habersaat K.B.; Eckardt M.; Schmid P.; Wichmann O. (2020): Barriers and drivers to adult vaccination among family physicians - Insights for tailoring the immunization program in Germany. In: *Vaccine* 38 (27), S. 4252–4262. DOI: 10.1016/j.vaccine.2020.04.052.

*Nobis H.-G.; Pielsticker A. (2013): Medical education and communication in primary pain treatment: Clinical relevance and pedagogic challenge. In: *Schmerz* 27 (3), S. 317–324. DOI: 10.1007/s00482-012-1289-3.

Nouwens E.; Van Lieshout J.; van den Hombergh P.; Laurant M.; Wensing M. (2014): Shifting cardiovascular care to nurses results in structured chronic care. In: *The American journal of managed care* 20 (7), e278–e284.

Oedekoven M.; Herrmann W.J.; Ernsting C.; Schnitzer S.; Kanzler M.; Kuhlmeijer A.; Gellert P. (2019): Patients' health literacy in relation to the preference for a general practitioner as the source of health information. In: *BMC family practice* 20 (1), S. 94. DOI: 10.1186/s12875-019-0975-y.

Oertelt-Prigione S.; Ruecke M.; Seeland U.; Regitz-Zagrosek V. (2017): Gynecological Practice Represents the Ideal Setting for Early Opportunistic Cardiovascular Prevention: A Cross-Sectional Survey in an Urban Female Population. In: *Journal of Women's Health* 26 (1), S. 29–35. DOI: 10.1089/jwh.2015.5729.

Ong J.P.; Oehler G.; Kruger-Jansen C.; Lambert-Baumann J.; Younossi Z.M. (2011): Oral L-ornithine-L-aspartate improves health-related quality of life in cirrhotic patients with hepatic encephalopathy: An open-label, prospective, multicentre observational study. In: *Clinical Drug Investigation* 31 (4), S. 213–220. DOI: 10.2165/11586700-000000000-00000.

Oriol-Zerbe C.; Abholz H.-H. (2007): Primary prevention of cardiovascular diseases by lipid-lowering treatment in German general practice: Results from GPs ignoring guidelines and risk calculators. In: *European Journal of General Practice* 13 (1), S. 27–34. DOI: 10.1080/13814780601050541.

*Otto F.; de Wall S. (2019): Inpatient treatment for informal caregivers of relatives suffering from dementia. In: *Zeitschrift für Allgemeinmedizin* 2019 (11), S. 462–467. DOI: 10.3238/zfa.2019.0462-0467.

*Patzelt, C.; Deitermann, B.; Heim, S.; Lupp, K.; Stahmeyer, J.; Krauth, C. et al. (2015): [The Preventive Home Visit as a Preventive Intervention to Reach the Elderly--Effectiveness and Cost-Effectiveness of Different Ways of Approach]. In: *Gesundheitswesen (Bundesverband der Ärzte des Öffentlichen Gesundheitsdienstes (Germany))* 77 Suppl 1 (bfd, 9204210), S45-6. DOI: 10.1055/s-0033-1337914.

Pausch M.; Schedlbauer A.; Weiss M.; Kuehlein T.; Hueber S. (2020): Is it really always only the others who are to blame? GP's view on medical overuse. A questionnaire study. In: *PLoS ONE* 15 (1), e0227457. DOI: 10.1371/journal.pone.0227457.

Perleth M.; Busse R.; Gibis B.; Brand A. (2001): Evaluation of preventive technologies in Germany: Case studies of mammography, prostate cancer screening, and fetal ultrasound. In: *International Journal of Technology Assessment in Health Care* 17 (3), S. 329–337. DOI: 10.1017/S0266462301106057.

Peters-Klimm F.; Campbell S.; Hermann K.; Kunz C.U.; Muller-Tasch T.; Szecsenyi J. (2010): Case management for patients with chronic systolic heart failure in primary care: The HICMan exploratory randomised controlled trial. In: *Trials* 11 ((Peters-Klimm, Campbell, Hermann, Szecsenyi) Department of General Practice and Health Services Research, University Hospital Heidelberg, Heidelberg, Germany(Campbell) National Primary Care Research and Development Centre, University of Manchester, Manche), S. 56. DOI: 10.1186/1745-6215-11-56.

Pfister, Roman; Ihle, Peter; Mews, Birgit; Kohnen, Elisabeth; Wahner, Marcus; Karbach, Ute et al. (2017): Feasibility of including patients with migration background in a structured heart failure management programme: A prospective case-control study exemplarily on Turkish migrants. In: *PLoS ONE* 12 (11), e0187358. DOI: 10.1371/journal.pone.0187358.

*Pilars de Pilar, Marieke; Abholz, Heinz-Harald; Becker, Nicole; Sielk, Martin (2012): [How do general practitioners deal with patients they do not consider to be depressed but who are classified as such according the PHQ-9?]. In: *Psychiatrische Praxis* 39 (2), S. 71–78. DOI: 10.1055/s-0031-1276897.

Plass, Anne Marie C.; Baars, Marieke J. H.; Cornel, Martina C.; Julian-Reynier, Claire; Nippert, Irmgard; Harris, Hillary et al. (2009): Testing the children: do non-genetic health-care providers differ in their decision to advise genetic presymptomatic testing on minors? A cross-sectional study in five countries in the European Union. In: *Genetic testing and molecular biomarkers* 13 (3), S. 367–376. DOI: 10.1089/gtmb.2008.0119.

*Plath, Jasper; Siebenhofer, Andrea; Schulz-Rothe, Sylvia; Guthlin, Corina (2018): [Who Should Provide Persons with Familial Risk of Colorectal Cancer with Information on Early Detection? - The View of those Affected]. In: *Gesundheitswesen (Bundesverband der Ärzte des Öffentlichen Gesundheitsdienstes (Germany))* 80 (10), S. 878–881. DOI: 10.1055/s-0042-124671.

Pohontsch N.J.; Heser K.; Loffler A.; Haenisch B.; Parker D.; Luck T. et al. (2017): General practitioners' views on (long-term) prescription and use of problematic and potentially inappropriate medication for oldest-old patients-A qualitative interview study with GPs (CIM-TRIAD study). In: *BMC family practice* 18 (1), S. 22. DOI: 10.1186/s12875-017-0595-3.

Poinhos R.; Oliveira B.M.P.M.; Van Der Lans I.A.; Fischer A.R.H.; Berezowska A.; Rankin A. et al. (2017): Providing Personalised Nutrition: Consumers' Trust and Preferences Regarding Sources of Information, Service Providers and Regulators, and Communication Channels. In: *Public Health Genomics* 20 (4), S. 218–228. DOI: 10.1159/000481357.

Poss-Doering R.; Kuehn L.; Kamradt M.; Glassen K.; Wensing M. (2020): Applying digital information delivery to convert habits of antibiotic use in primary care in Germany: Mixed-methods study. In: *Journal of Medical Internet Research* 22 (10), e18200. DOI: 10.2196/18200.

Probst C.; Manthey J.; Martinez A.; Rehm J. (2015): Alcohol use disorder severity and reported reasons not to seek treatment: a cross-sectional study in European primary care practices. In: *Substance abuse treatment, prevention, and policy* 10 ((Probst, Rehm) Social and Epidemiological Research Department, Centre for Addiction and Mental Health, 33 Russell Street, Toronto, ON, M5S 2S1, Canada(Probst, Manthey, Rehm) Institute for Clinical Psychology and Psychotherapy, Technische Universität Dresden), S. 32. DOI: 10.1186/s13011-015-0028-z.

Prufer F.; Joos S.; Miksch A. (2013): What do resource-oriented approaches mean to general practitioners and how can they be facilitated in primary care? A qualitative study. In: *Evidence-based Complementary and Alternative Medicine* 2013 ((Prufer, Joos, Miksch) Department of General Practice and Health Services Research, University Hospital Heidelberg, Vosstrasse 2, Geb.37, 69115 Heidelberg, Germany(Prufer, Joos, Miksch) Competence Centre for General Practice, Baden-Württemberg, Germany), S. 187641. DOI: 10.1155/2013/187641.

Pullen R.; Weltermann B.; Homann J.; Fusgen I. (2001): How much do geriatric high-risk patients know about stroke? In: *European Journal of Geriatrics* 3 (3), S. 131–134.

*Radtke, Marc Alexander; Rustenbach, Stephan Jeff; Reusch, Michael; Stromer, Klaus; Augustin, Matthias (2013): Influenza vaccination rate among patients with moderate to severe psoriasis. In: *Journal der Deutschen Dermatologischen Gesellschaft = Journal of the German Society of Dermatology : JDDG* 11 (9), S. 837–844. DOI: 10.1111/ddg.12010.

*Rapke C.; Hofmann G.; Stedtfeld H.-W.; Scherbel G. (2004): Cooperation between physician and pharmacist to determine and resolve errors in concomitant medication previously prescribed for trauma patients. Quality assurance study. In: *Unfallchirurg* 107 (11), S. 1041–1049. DOI: 10.1007/s00113-004-0826-7.

*Rattay, P.; Starker, A.; Domanska, O.; Butschalowsky, H.; Gutsche, J.; Kamtsiuris, P.; KiGGS Study Group (2014): [Trends in the utilization of outpatient medical care in childhood and adolescence: results of the KiGGS study - a comparison of baseline and first follow up (KiGGS Wave 1)]. In: *Bundesgesundheitsblatt, Gesundheitsforschung, Gesundheitsschutz* 57 (7), S. 878–891. DOI: 10.1007/s00103-014-1989-1.

*Rau R.; Mensing M.; Brand H. (2006): Community knowledge about stroke. A survey in the District of Wesel, Germany (2002). In: *Bundesgesundheitsblatt - Gesundheitsforschung - Gesundheitsschutz* 49 (5), S. 450–458. DOI: 10.1007/s00103-006-1247-2.

Raupach T.; Merker J.; Hasenfuss G.; Andreas S.; Pipe A. (2011): Knowledge gaps about smoking cessation in hospitalized patients and their doctors. In: *European Journal of Cardiovascular Prevention and Rehabilitation* 18 (2), S. 334–341. DOI: 10.1177/1741826710389370.

*Regus S.; Sonntag U.; Bockelbrink A.; Welke J.; Braun V.; Heintze C. (2011): Check up 35: What benefits do general practitioners in Brandenburg see? In: *Zeitschrift für Evidenz, Fortbildung und Qualität im Gesundheitswesen* 105 (6), S. 421–426. DOI: 10.1016/j.zefq.2010.07.012.

*Rehm J.; Gmel G.; Kiefer F.; Kreutz R.; Kugler J.; Muller-Walther M. et al. (2014): Improvement of management of hypertension by implementation of alcohol screening and subsequent interventions in primary practice. In: *Deutsche Medizinische Wochenschrift* 139 (48), S. 2457–2462. DOI: 10.1055/s-0034-1387417.

*Reiter, S.; Haas, W. (2005): [Influenza pandemic: A real threat ?]. In: *MMW Fortschritte der Medizin* 147 (9), S. 35–38.

*Renz, Jorg-Christian; Meinck, Matthias (2020): [Preventive Home Visits for Elderly People: A Systematic Appraisal of Current Application in Germany]. In: *Gesundheitswesen (Bundesverband der Ärzte des Öffentlichen Gesundheitsdienstes (Germany))* 82 (4), S. 339–344. DOI: 10.1055/a-0658-5880.

*Rexroth, U.; Buda, S. (2014): [Occupational health and practice management of primary care practitioners during influenza pandemic 2009/10 in Germany - a survey of 1150 physicians participating in syndromic influenza surveillance at Robert Koch Institute]. In: *Gesundheitswesen (Bundesverband der Ärzte des Öffentlichen Gesundheitsdienstes (Germany))* 76 (10), S. 670–675. DOI: 10.1055/s-0033-1355402.

Richter, Christin; Berg, Almuth; Fleischer, Steffen; Kopke, Sascha; Balzer, Katrin; Fick, Eva-Maria et al. (2015): Effect of person-centred care on antipsychotic drug use in nursing homes (EPCentCare): study protocol for a cluster-randomised controlled trial. In: *Implementation science : IS* 10 (101258411), S. 82. DOI: 10.1186/s13012-015-0268-3.

Richter, Christin; Berg, Almuth; Langner, Henriette; Meyer, Gabriele; Kopke, Sascha; Balzer, Katrin et al. (2019): Effect of person-centred care on antipsychotic drug use in nursing homes (EPCentCare): a cluster-randomised controlled trial. In: *Age and Ageing* 48 (3), S. 419–425. DOI: 10.1093/ageing/afz016.

*Richter, Jutta G.; Wessel, Ewa; Klimt, Ralf; Willers, Reinhardt; Schneider, Matthias (2008): [RheumaCheck: development and evaluation of a German language screening instrument for rheumatic diseases]. In: *Wiener klinische Wochenschrift* 120 (3-4), S. 103–111. DOI: 10.1007/s00508-008-0929-1.

*Richter J.G.; Becker A.; Specker C.; Monser R.; Schneider M. (2004): Disease-oriented Internet use in outpatients with inflammatory rheumatic diseases. In: *Zeitschrift für Rheumatologie* 63 (3), S. 216–222.

*Richter K.; Acker J.G.; Geiss L.; Niklewski G. (2017): Sleep disorders in the general practitioners office. In: *Internistische Praxis* 57 (3), S. 479–486. Online verfügbar unter <http://www.marseille-verlag.com>.

Rieckert A.; Sommerauer C.; Krumeich A.; Sonnichsen A. (2018): Reduction of inappropriate medication in older populations by electronic decision support (the PRIMA-eDS study): a qualitative study of practical implementation in primary care. In: *BMC family practice* 19 (1), S. 110. DOI: 10.1186/s12875-018-0789-3.

Rieckert A.; Teichmann A.-L.; Drewelow E.; Kriechmayr C.; Piccoliori G.; Woodham A.; Sonnichsen A. (2019): Reduction of inappropriate medication in older populations by electronic decision support (the PRIMA-eDS project): A survey of general practitioners' experiences. In: *Journal of the American Medical Informatics Association* 26 (11), S. 1323–1332. DOI: 10.1093/jamia/ocz104.

*Riedl B.; Linde K. (2018): Recommending screening colonoscopy in a family physician's practice: Retrospective evaluation of routine data between 2002 and 2017. In: *Zeitschrift für Allgemeinmedizin* 94 (11), S. 443–448. Online verfügbar unter https://www.online-zfa.com/fileadmin/user_upload/Riedl-Koloskopie_MMK.pdf.

*Riegel, W.; Hahn, K.; Kreutz, R.; Weber, M.; Zidek, W.; Schmieder, R. (2005): [BENEFIT Kidney--significance of a nephrology screening at intervention outset and therapy success]. In: *Deutsche medizinische Wochenschrift* (1946) 130 (13), S. 792–796.

*Rock D.; Schnellbacher E.; Fritsche A.; Ziemssen T.; Ziemssen F.; Marahrens L. (2018): Knowledge of Patients and Health Professionals about Diabetes-related Eye Diseases (Risk Factors, Screening, Treatment). In: *Diabetologie und Stoffwechsel* 13 (3), S. 263–269. DOI: 10.1055/a-0578-8112.

Roggendorf, Hedwig; Santibanez, Sabine; Mankertz, Annette; van Treeck, Ulrich; Roggendorf, Michael (2012): Two consecutive measles outbreaks with genotypes D8 and D4 in two mainly unvaccinated communities in Germany. In: *Medical microbiology and immunology* 201 (3), S. 349–355. DOI: 10.1007/s00430-012-0240-7.

Ropers, Gwendolin; Krause, Gerard; Tiemann, Friedrich; Du Ry van Beest Holle, Mirna; Stark, Klaus (2004): Nationwide survey of the role of travel medicine in primary care in Germany. In: *Journal of Travel Medicine* 11 (5), S. 287–294.

Rosano A.; Peschel P.; Kugler J.; Ricciardi W.; Guasticchi G.; Van Der Zee J. (2013): Preventable hospitalization and the role of primary care: A comparison between Italy and Germany. In: *Journal of Public Health (Germany)* 21 (5), S. 445–454. DOI: 10.1007/s10389-013-0563-x.

Rosemann T.; Hermann K.; Miksch A.; Engeser P.; Szecsenyi J. (2007): The PAMINO-project: Evaluating a primary care-based educational program to improve the quality of life of palliative patients. In: *BMC Palliative Care* 6 ((Rosemann, Hermann, Miksch, Engeser, Szecsenyi) Department of General Practice and Health Services Research, University Hospital of Heidelberg, Vosstr. 2, 69115 Heidelberg, Germany), S. 5. DOI: 10.1186/1472-684X-6-5.

Roske K.; Coder B.; John U.; Hannich H.-J.; Ulbricht S. (2006): The role of general practitioners in the care for patients with addiction problems in a rural area of Germany. In: *Family Medicine and Primary Care Review* 8 (4), S. 1223–1227.

Rosta J. (2003): Determinants of physicians' prevention-related practice: A comparison between Aarhus, Denmark, and Mainz, Germany. In: *Journal of Substance Use* 8 (1), S. 42–46. DOI: 10.1080/1465989031000067245.

*Rothenbacher D.; Ruter G.; Saam S.; Brenner H. (2002): Management of patients with type 2 diabetes: Results in 12 practices of general practitioners. In: *Deutsche Medizinische Wochenschrift* 127 (22), S. 1183–1187. DOI: 10.1055/s-2002-31940.

Rothermund E.; Michaelis M.; Jarczok M.N.; Balint E.M.; Lange R.; Zipfel S. et al. (2018): Prevention of common mental disorders in employees. Perspectives on collaboration from three health care professions. In: *International Journal of Environmental Research and Public Health* 15 (2), S. 278. DOI: 10.3390/ijerph15020278.

*Rubsam M.-L.; Esch M.; Baum E.; Bosner S. (2015): Dermatology in primary care: Coordination, prevention, therapy and educational needs. In: *Zeitschrift fur Allgemeinmedizin* 91 (5), S. 227–232. Online verfügbar unter https://www.online-zfa.de/media/article/2015/05/1EE4823A-B8FD-4F77-A1D7-42472F4E88FF/1EE4823AB8FD4F77A1D742472F4E88FF_boesner_dermatologie_in_der_primaerversorgung_1_original.pdf.

Rumpf, H. J.; Bohlmann, J.; Hill, A.; Hapke, U.; John, U. (2001): Physicians' low detection rates of alcohol dependence or abuse: a matter of methodological shortcomings? In: *General Hospital Psychiatry* 23 (3), S. 133–137.

Ruof J.; Klein G.; Marz W.; Wollschlager H.; Neiss A.; Wehling M. (2002): Lipid-lowering medication for secondary prevention of coronary heart disease in a german outpatient population: The gap between treatment guidelines and real life treatment patterns. In: *Preventive Medicine* 35 (1), S. 48–53. DOI: 10.1006/pmed.2002.1050.

*Ruter G. (2003): Individuality and salutogenesis in general medicine: A paradigm shift for care, research, training and policy. In: *Zeitschrift fur Allgemeinmedizin* 79 (6), S. 278–282.

Sahlan S.; Wollny A.; Brockmann S.; Fuchs A.; Altiner A. (2008): Reducing unnecessary prescriptions of antibiotics for acute cough: Adaptation of a leaflet aimed at Turkish immigrants in Germany. In: *BMC family practice* 9 ((Sahlan, Wollny, Brockmann, Fuchs, Altiner) Department of General Practice, University Hospital, Heinrich-Heine-University, 40001 Duesseldorf, Germany), S. 57. DOI: 10.1186/1471-2296-9-57.

Salize H.J.; Merkel S.; Reinhard I.; Twardella D.; Mann K.; Brenner H. (2009): Cost-effective primary care-based strategies to improve smoking cessation. In: *Archives of Internal Medicine* 169 (3), S. 230–235. DOI: 10.1001/archinternmed.2008.556.

Samann A.; Tajiyeva O.; Muller N.; Tschauner T.; Hoyer H.; Wolf G.; Muller U.A. (2008): Prevalence of the diabetic foot syndrome at the primary care level in Germany: A cross-sectional study. In: *Diabetic Medicine* 25 (5), S. 557–563. DOI: 10.1111/j.1464-5491.2008.02435.x.

*Sander D.; Schwertfeger M.; Kofuncu E.; Diehm C.; Pittrow D. (2008): Secondary prevention with clopidogrel or acetylsalicylic acid after cerebrovascular event - Prospective cross-sectional study in primary care. In: *Deutsche Medizinische Wochenschrift* 133 (36), S. 1773–1778. DOI: 10.1055/s-0028-1082810.

*Sandholzer H.; Hellenbrand W.; Renteln-Kruse W.; Van Weel C.; Walker P. (2004): STEP - Standardized assessment of elderly people in primary care. In: *Deutsche Medizinische Wochenschrift* 129 (SUPPL. 4), S183-S226. DOI: 10.1055/s-2004-836107.

Sanftenberg, Linda; Brombacher, Felix; Schelling, Jorg; Klug, Stefanie J.; Gensichen, Jochen (2019): Increasing Influenza Vaccination Rates in People With Chronic Illness. In: *Deutsches Arzteblatt International* 116 (39), S. 645–652. DOI: 10.3238/arztebl.2019.0645.

*Sanftenberg, Linda; Salavati, Puria; Schelling, Jorg (2017): [Implementing vaccination recommendations in discharge letters]. In: *Zeitschrift fur Evidenz, Fortbildung und Qualitat im Gesundheitswesen* 127-128 (101477604), S. 30–35. DOI: 10.1016/j.zefq.2017.09.011.

Sarnighausen H.-E. (2010): The role of the general practitioner in diabetes prevention. In: *Austrian Journal of Clinical Endocrinology and Metabolism* 3 (2), S. 26–29. Online verfügbar unter <http://www.kup.at/kup/pdf/8977.pdf>.

*Schacher C.; Dhein Y.; Munks-Lederer C.; Vollmer T.; Worth H. (2006): Evaluation of a structured outpatient education program for adult asthmatics. In: *Deutsche Medizinische Wochenschrift* 131 (12), S. 606–610. DOI: 10.1055/s-2006-933702.

Schaefer J.R.; Gitt A.K.; Sonntag F.; Weizel A.; Jannowitz C.; Karmann B. et al. (2013): Lipid management in 13,000 high risk cardiovascular patients treated under daily practice conditions: LIMA registry. In: *Vascular Health and Risk Management* 9 (1), S. 71–80. DOI: 10.2147/VHRM.S37143.

Schafer I.; Kaduszkiewicz H.; Mellert C.; Loffler C.; Mortsiefer A.; Ernst A. et al. (2018): Narrative medicine-based intervention in primary care to reduce polypharmacy: Results from the cluster-randomised controlled trial MultiCare AGENDA. In: *BMJ Open* 8 (1), e017653. DOI: 10.1136/bmjopen-2017-017653.

- Schafer I.; Pawels M.; Kuver C.; Pohontsch N.J.; Scherer M.; van den Bussche H.; Kaduszkiewicz H. (2014): Strategies for improving participation in diabetes education. A qualitative study. In: *PLoS ONE* 9 (4), e95035. DOI: 10.1371/journal.pone.0095035.
- *Schang, Laura; Sundmacher, Leonie; Grill, Eva (2020): [New Forms of Cooperation Between The Outpatient and Inpatient Sectors: An Innovative Funding Concept]. In: *Gesundheitswesen (Bundesverband der Ärzte des Öffentlichen Gesundheitsdienstes (Germany))* 82 (6), S. 514–519. DOI: 10.1055/a-0829-6465.
- *Schaper, M.; Berndt, M.; Schrimpf, C.; Wilhelm, M.; Elff, M.; Haverich, A. (2016): [Fax Survey to Elucidate the Information Needs of General Practitioners in Lower Saxony Regarding the Topic of Medical Implants]. In: *Zentralblatt für Chirurgie* 141 (6), S. 677–681.
- Scharf J.; Vu-Eickmann P.; Li J.; Muller A.; Wilm S.; Angerer P.; Loerbroks A. (2019): Desired improvements of working conditions among medical assistants in Germany: A cross-sectional study. In: *Journal of Occupational Medicine and Toxicology* 14 (1), S. 18. DOI: 10.1186/s12995-019-0237-x.
- *Scharl A.; Salterberg A.; Papathemeis T.; Scharl S. (2017): The German breast cancer screening program. In: *Internistische Praxis* 58 (1), S. 75–84. Online verfügbar unter <http://www.marseille-verlag.com>.
- *Schelling J.; Weinberger S.; Kunisch R.; Taylor D.; Weiss N. (2013): Peripheral arterial disease: Implementation of secondary prevention measures in primary care according to guidelines. In: *Zeitschrift für Allgemeinmedizin* 89 (2), S. 61–66. DOI: 10.3238/zfa.2013.0061-0066.
- Schenck, U.; Karsa, L. von (2000): Cervical cancer screening in Germany. In: *European journal of cancer (Oxford, England : 1990)* 36 (17), S. 2221–2226.
- *Scherbaum, W. A.; Dicken, H-D; Verheyen, F.; Baehring, T. (2006): [Detection of a previously unrecognized type 2 diabetes using a risk questionnaire. Efficiency for general population screening]. In: *Deutsche medizinische Wochenschrift (1946)* 131 (40), S. 2208–2212.
- Scheuermann W.; Razum O.; Scheidt R.; Wiesemann A.; Von Frankenberg H.; Topf G.; Nussel E. (2000): Effectiveness of a decentralized, community-related approach to reduce cardiovascular disease risk factor levels in Germany. In: *European Heart Journal* 21 (19), S. 1591–1597. DOI: 10.1053/euhj.1999.2037.
- *Schiel R.; Muller U.A. (2003): Structured treatment and teaching programs and an improvement in private health care lead to a better quality of diabetes care. JEVIN, a population-based trial 1989/90 up to 1999/2000. In: *Medizinische Klinik* 98 (6), S. 303–312. DOI: 10.1007/s00063-003-1264-y.
- Schiel R.; Voigt U.; Ross I.S.; Braun A.; Rillig A.; Hunger-Dathe W. et al. (2006): Structured diabetes therapy and education improves the outcome of patients with insulin treated diabetes mellitus: The 10 year follow-up of a prospective, population-based survey on the quality of diabetes care (The JEVIN Trial). In: *Experimental and Clinical Endocrinology and Diabetes* 114 (1), S. 18–27. DOI: 10.1055/s-2005-873079.
- Schmedt N.; Schiffner-Rohe J.; Sprenger R.; Walker J.; Von Eiff C.; Hackl D. (2019): Pneumococcal vaccination rates in immunocompromised patients-A cohort study based on claims data from more than 200,000 patients in Germany. In: *PLoS ONE* 14 (8), e0220848. DOI: 10.1371/journal.pone.0220848.
- *Schmidt, C. O.; Lindena, G.; Pfingsten, M.; Kohlmann, T.; Chenot, J-F (2014): [Comparison of two screening questionnaires for patients with low back pain. Collation of risk factors for chronicification]. In: *Schmerz (Berlin, Germany)* 28 (4), S. 365–373. DOI: 10.1007/s00482-014-1431-5.
- Schmidt, Carsten Oliver; Chenot, Jean-Francois; Pfingsten, Michael; Fahland, Ruth Anja; Lindena, Gabriele; Marnitz, Ulf et al. (2010): Assessing a risk tailored intervention to prevent disabling low back pain--protocol of a cluster randomized controlled trial. In: *BMC musculoskeletal disorders* 11 (100968565), S. 5. DOI: 10.1186/1471-2474-11-5.
- Schmidt, L. G. (2004): [Helping smokers kick the habit]. In: *MMW Fortschritte der Medizin* 146 (17), S. 34–39.
- Schmidt, Silke; Herrmann-Garitz, Carsten; Bomba, Franziska; Thyen, Ute (2016): A multicenter prospective quasi-experimental study on the impact of a transition-oriented generic patient education program on health service participation and quality of life in adolescents and young adults. In: *Patient Education and Counseling* 99 (3), S. 421–428. DOI: 10.1016/j.pec.2015.10.024.
- *Schmidt-Petruschkat, S. (2007): [HPV-infection]. In: *MMW Fortschritte der Medizin* 149 (12), S. 29–31.
- Schmieder R.E.; Goebel M.; Bramlage P. (2012): Barriers to cardiovascular risk prevention and management in Germany - An analysis of the EURIKA study. In: *Vascular Health and Risk Management* 8 (1), S. 177–186. DOI: 10.2147/VHRM.S29915.
- *Schmiemann, Guido; Biesewig-Siebenmorgen, Jurgen; Gebhardt, Klaus; Egidi, Gunther (2014): [The Bremen periodic health exam - feasibility of a new concept]. In: *Zeitschrift für Evidenz, Fortbildung und Qualität im Gesundheitswesen* 108 (4), S. 196–202. DOI: 10.1016/j.zefq.2013.12.002.
- *Schmiemann G. (2011): Injury by a bird of prey. Antibiotic prophylaxis and rabies vaccination. In: *Internistische Praxis* 51 (3), S. 549–553.
- *Schmiemann G.; Biesewig-Siebenmorgen J.; Egidi G. (2013): The new bremen preventive medical check-up - Development and concept of an age adjusted health examination. In: *Zeitschrift für Allgemeinmedizin* 89 (6), S. 261–265. DOI: 10.3238/zfa.2013.0261-0265.

*Schmitz, Astrid (2011): [Information needs of people suffering from venous leg ulcer. Expertise of people concerned as a basis of patient-centered information]. In: *Pflege* 24 (3), S. 161–170. DOI: 10.1024/1012-5302/a000117.

Schnabel K.; Binting S.; Witt C.M.; Teut M. (2014): Use of complementary and alternative medicine by older adults--a cross-sectional survey. In: *BMC geriatrics* 14 ((Teut) Institute for Social Medicine, Epidemiology and Health Economics, Charite Universitätsmedizin, D-10098 Berlin, Germany), S. 38. DOI: 10.1186/1471-2318-14-38.

*Schnakenberg, R.; Rothe, M.; Bleckwenn, M. (2017): [Early Selection of At-risk Patients for Type 2 Diabetes in the GP Surgery by Medical Assistants - A Feasibility Study]. In: *Gesundheitswesen (Bundesverband der Ärzte des Öffentlichen Gesundheitsdienstes (Germany))* 79 (8-09), S. 633–637. DOI: 10.1055/s-0035-1548858.

Schneider, Antonius; Wensing, Michel; Quinzler, Renate; Bieber, Christiane; Szecsenyi, Joachim (2007): Higher preference for participation in treatment decisions is associated with lower medication adherence in asthma patients. In: *Patient Education and Counseling* 67 (1-2), S. 57–62.

Schneider C.; Joos S.; Bozorgmehr K. (2015): Disparities in health and access to healthcare between asylum seekers and residents in Germany: A population-based cross-sectional feasibility study. In: *BMJ Open* 5 (11), e008784. DOI: 10.1136/bmjopen-2015-008784.

Schneider S.; Diehl K.; Bock C.; Herr R.M.; Mayer M.; Gorig T. (2014): Modifying health behavior to prevent cardiovascular diseases: A nationwide survey among German primary care physicians. In: *International Journal of Environmental Research and Public Health* 11 (4), S. 4218–4232. DOI: 10.3390/ijerph110404218.

*Schoierer, Julia; Mertes, Hanna; Wershofen, Birgit; Bose-O'Reilly, Stephan (2019): [Training modules on climate change, heat, and health for medical assistants and nurses in outpatient care]. In: *Bundesgesundheitsblatt, Gesundheitsforschung, Gesundheitsschutz* 62 (5), S. 620–628. DOI: 10.1007/s00103-019-02942-w.

Schopf A.C.; von Hirschhausen M.; Farin E.; Maun A. (2018): Elderly patients' and GPs' perspectives of patient-GP communication concerning polypharmacy: a qualitative interview study. In: *Primary health care research & development* 19 (4), S. 355–364. DOI: 10.1017/S1463423617000883.

Schoppmeyer, Konrad; Spieker, Henning; Mossner, Joachim (2009): Failure of screening or failure to screen? The screening behavior of patients with colorectal cancer in the Leipzig area. In: *Deutsches Arzteblatt International* 106 (12), S. 195–201. DOI: 10.3238/arztebl.2009.0195.

Schray, D.; Neuerburg, C.; Stein, J.; Gosch, M.; Schieker, M.; Bocker, W.; Kammerlander, C. (2016): Value of a coordinated management of osteoporosis via Fracture Liaison Service for the treatment of orthogeriatric patients. In: *European journal of trauma and emergency surgery : official publication of the European Trauma Society* 42 (5), S. 559–564.

Schray, Deborah; Stumpf, Ulla; Kammerlander, Christian; Bocker, Wolfgang; Neuerburg, Carl (2016): Diagnosis and therapy of osteoporosis in geriatric trauma patients: an update. In: *Innovative surgical sciences* 1 (2), S. 65–71. DOI: 10.1515/iss-2016-0028.

Schubert I.; Koster I.; Ihle P.; Von Ferber L. (2001): Development of indicators for assessing the quality of prescribing of lipid-lowering drugs: Data from the pharmacotherapeutic quality circles in Hesse, Germany. In: *International Journal of Clinical Pharmacology and Therapeutics* 39 (11), S. 492–498.

*Schubmann R.M. (2000): Integration of prevention groups in an existing outpatient cardiac group. In: *Pravention und Rehabilitation* 12 (3), S. 114–118.

*Schultze, A.; Akmatov, M. K.; Castell, S.; Karch, A.; Ahrens, W.; Gunther, K. et al. (2014): Collection of vaccination data in the German National Cohort : Findings of a feasibility study in three study centers. In: *Bundesgesundheitsblatt, Gesundheitsforschung, Gesundheitsschutz* 57 (11), S. 1292–1299. DOI: 10.1007/s00103-014-2050-0.

*Schulze Horn, C.; Sander, K.; Ilg, R.; Bickel, H.; Briesenick, C.; Sander, D. (2007): [Modification of vascular risk factors by intervention on the primary care level. Results of the INVADE project]. In: *Der Nervenarzt* 78 (12), S. 1413–1419.

*Schuttrumpf S.; Binder C. (2003): Infections in patients with malignant diseases - Prophylaxis and therapy. In: *Zeitschrift für Allgemeinmedizin* 79 (2), S. 77–83.

*Schwarz P.E.H. (2015): Screening and prevention of diabetes. In: *Internist* 56 (10), S. 1124–1133. DOI: 10.1007/s00108-015-3737-5.

*Schwarz P.E.H.; Muller U.; Schulz M.; Hauner H.; Landgraf R. (2007): Prevention of typ 2 diabetes in Germany - A challenge for health professionals. In: *Medizinische Monatsschrift für Pharmazeuten* 30 (8), S. 289–296.

*Seebauer L.; Simon D.; Bermejo I.; Herbarth L.; Siegmund-Schultze E.; Temmert D. et al. (2011): [Attitudes of chronically-ill patients to telephone health coaching]. In: *Gesundheitswesen (Bundesverband der Ärzte des Öffentlichen Gesundheitsdienstes (Germany))* 73 (7), S. 430–437.

*Seibt, K.; Schulz, M.; Hensel, F. J. (2000): [Opinions and attitudes on the topic of vaccination of established physicians, pharmacists and their personnel and current vaccination status of these groups]. In: *Gesundheitswesen (Bundesverband der Ärzte des Öffentlichen Gesundheitsdienstes (Germany))* 62 (7), S. 376–382.

Seidel, Hans Joachim (2002): Environmental medicine in Germany--a review. In: *Environmental health perspectives* 110 Suppl 1 (ei0, 0330411), S. 113–118.

Send A.F.J.; Peters-Klimm F.; Bruckner T.; Haefeli W.E.; Seidling H.M. (2017): A randomized controlled trial to assess the effect of a medication plan containing drug administration recommendations on patients' drug knowledge after 2 months. In: *Journal of Clinical Pharmacy and Therapeutics* 42 (1), S. 69–74. DOI: 10.1111/jcpt.12476.

*Shedden-Mora, Meike; Lau, Katharina; Kuby, Amina; Gros, Beatrice; Gladigau, Maria; Fabisch, Alexandra; Lowe, Bernd (2015): [Improving Health Care for Patients with Somatoform and Functional Disorders: A Collaborative Stepped Care Network (Sofu-Net)]. In: *Psychiatrische Praxis* 42 Suppl 1 (qck, 0423204), S60-4. DOI: 10.1055/s-0034-1387690.

Shellenberger R.; Nabhan M.; Kakaraparthi S. (2016): Melanoma screening: A plan for improving early detection. In: *Annals of Medicine* 48 (3), S. 142–148. DOI: 10.3109/07853890.2016.1145795.

*Siebenhofer, Andrea; Horvath, Karl; Semlitsch, Thomas; Plath, Jasper; Jeitler, Klaus (2014): [General practitioner relevant recommendations in German cancer guidelines: a systematic review]. In: *Zeitschrift fur Evidenz, Fortbildung und Qualitat im Gesundheitswesen* 108 (5-6), S. 283–292. DOI: 10.1016/j.zefq.2014.04.017.

Siebenhofer, Andrea; Plath, Jasper; Taubenroth, Maja; Singer, Susanne; Hechtner, Marlene; Dahlhaus, Anne et al. (2015): Positive family history of colorectal cancer in a general practice setting [FRIDA.Frankfurt]: study protocol of a cross-sectional study. In: *BMC cancer* 15 (100967800), S. 605. DOI: 10.1186/s12885-015-1600-7.

Siebenhofer, Andrea; Ulrich, Lisa-Rebekka; Mergenthal, Karola; Berghold, Andrea; Pregartner, Gudrun; Kemperdick, Birgit et al. (2019): Primary care management for patients receiving long-term antithrombotic treatment: A cluster-randomized controlled trial. In: *PLoS ONE* 14 (1), e0209366. DOI: 10.1371/journal.pone.0209366.

Siedler A.; Arndt U. (2010): Impact of the routine varicella vaccination programme on varicella epidemiology in Germany. In: *Euro surveillance : bulletin European sur les maladies transmissibles = European communicable disease bulletin* 15 (13).

Siegrist, Monika; Freiberger, Ellen; Geilhof, Barbara; Salb, Johannes; Hentschke, Christian; Landendoerfer, Peter et al. (2016): Fall Prevention in a Primary Care Setting. In: *Deutsches Arzteblatt International* 113 (21), S. 365–372. DOI: 10.3238/arztebl.2016.0365.

*Siepmann, M. (2012): [Prevention of psychic disorders]. In: *Deutsche medizinische Wochenschrift* (1946) 137 (23), S. 1259–1263. DOI: 10.1055/s-0032-1305036.

*Simbrich, A.; Semjonow, A.; Donner-Banzhoff, N.; Hense, H-W (2018): [Practice of early detection of prostate cancer : Descriptive survey in preparation for the PSAInForm study]. In: *Der Urologe. Ausg. A* 57 (6), S. 702–708. DOI: 10.1007/s00120-018-0644-0.

*Skoeries B.A.; Ulbricht S.; Koepsell S.; Rumpf H.J.; John U.; Meyer C. (2010): Readiness to provide smoking cessation counselling--results of a survey among general practitioners in Brandenburg. In: *Gesundheitswesen (Bundesverband der Ärzte des Öffentlichen Gesundheitsdienstes (Germany))* 72 (4), S. 228–232. DOI: 10.1055/s-0029-1215561.

*Smolka, M. N.; Kiefer, F.; Mann, K. (2003): [Advances in the treatment of alcohol dependence: pharmacological relapse prevention]. In: *MMW Fortschritte der Medizin* 145 Suppl 3 (dhp, 100893959), S. 65–69.

*Sonnichsen A. (2014): Shall all adults be screened for HIV? In: *Zeitschrift für Allgemeinmedizin* 90 (11), S. 435–437. Online verfügbar unter https://www.online-zfa.de/media/article/2014/11/5829F563-B0D2-4D73-93DC-8AB73F0D0E4F/5829F563B0D24D7393DC8AB73F0D0E4F_soennichsen_ebm_hiv_1_original.pdf.

Sonnichsen A.; Trampisch U.S.; Rieckert A.; Piccoliori G.; Vogele A.; Flamm M. et al. (2016): Polypharmacy in chronic diseases--Reduction of Inappropriate Medication and Adverse drug events in older populations by electronic Decision Support (PRIMA-eDS): Study protocol for a randomized controlled trial. In: *Trials* 17 (1), S. 57. DOI: 10.1186/s13063-016-1177-8.

*Sonnichsen A.C.; Rambeck M.; Donner-Banzhoff N.; Baum E. (2006): Cholesterol screening at the "chek-up 35": Does it make a difference? In: *Zeitschrift für Allgemeinmedizin* 82 (10), S. 431–434. DOI: 10.1055/s-2006-942198.

*Sonntag H.; Hoch E.; Jahn B.; Spiegel B.; Pfisterl H.; Wittchen H.-U. (2003): Smoking cessation in primary care: Implementation effectiveness and optimized allocation. In: *Suchtmedizin in Forschung und Praxis* 5 (2), S. 137–141.

*Sonntag H.-G.; Harke H.-P. (2006): Contribution of hygiene for infection prevention: Report on the working meeting of the Rudolf Schulke Foundation, November 24-25, 2005 in Hamburg. In: *Hygiene + Medizin* 31 (7-8), S. 332–339.

Steckelings, U. Muscha; Stoppelhaar, Martina; Sharma, Arya M.; Wittchen, Hans-Ulrich; Krause, Petra; Kupper, Beate et al. (2004): HYDRA: possible determinants of unsatisfactory hypertension control in German primary care patients. In: *Blood pressure* 13 (2), S. 80–88.

Stein, Janine; Luppa, Melanie; Kaduszkiewicz, Hanna; Eisele, Marion; Weyerer, Siegfried; Werle, Jochen et al. (2015): Is the Short Form of the Mini-Mental State Examination (MMSE) a better screening instrument for dementia in older primary care patients than the original MMSE? Results of the German study on ageing, cognition, and dementia in primary care patients (AgeCoD). In: *Psychological assessment* 27 (3), S. 895–904. DOI: 10.1037/pas0000076.

*Steinkohl, Catarina; Donner-Banzhoff, Norbert (2014): [Periodic health examinations - an international comparison]. In: *Zeitschrift für Evidenz, Fortbildung und Qualität im Gesundheitswesen* 108 (4), S. 185–191. DOI: 10.1016/j.zefq.2014.04.003.

*Sterdt, E.; Brandes, I.; Dathe, R.; Walter, U. (2010): [Post-rehabilitation care in the context of cardiac rehabilitation--aspects of prevention and health promotion]. In: *Die Rehabilitation* 49 (2), S. 87–94. DOI: 10.1055/s-0030-1249027.

- *Steuernagel B.; Doering T. (2003): Health prevention in children and adolescents in general practices. In: *Pravention und Rehabilitation* 15 (1), S. 17–23. DOI: 10.5414/prp15017.
- *Stiel, S.; Joppich, R.; Korb, K.; Hahnen, M.; Elsner, F.; Rossaint, R.; Radbruch, L. (2009): [Problems and deficits in the transition from inpatient and outpatient care of cancer patients. A qualitative analysis]. In: *Schmerz (Berlin, Germany)* 23 (5), S. 510–517. DOI: 10.1007/s00482-009-0828-z.
- *Stock K.; Wollny A.; Brockmann S.; Abholz H.-H.; Altiner A. (2008): A qualitative glance into the blackbox: Educational intervention to reduce unnecessary antibiotic prescriptions. In: *Zeitschrift fur Allgemeinmedizin* 84 (10), S. 444–450. DOI: 10.1055/s-0028-1087183.
- Stockbrugger R. (2012): Isolated colorectal cancer screening or integrated cancer prevention? A provocative suggestion! In: *Digestive Diseases* 30 (3), S. 316–319. DOI: 10.1159/000337005.
- Storr, Constanze; Sanftenberg, Linda; Schelling, Joerg; Heininger, Ulrich; Schneider, Antonius (2018): Measles Status–Barriers to Vaccination and Strategies for Overcoming Them. In: *Deutsches Arzteblatt International* 115 (43), S. 723–730. DOI: 10.3238/arztebl.2018.0723.
- Strassner C.; Steinhauser J.; Freund T.; Szecsenyi J.; Wensing M. (2018): German healthcare professionals' perspective on implementing recommendations about polypharmacy in general practice: A qualitative study. In: *Family Practice* 35 (4), S. 503–510. DOI: 10.1093/fampra/cmx127.
- *Strech D. (2013): Would you mind a little bit less? Methodological and ethical challenges posed by the reduction of overtreatment. In: *Zeitschrift fur Evidenz, Fortbildung und Qualitat im Gesundheitswesen* 107 (2), S. 156–163. DOI: 10.1016/j.zefq.2013.02.009.
- Stuhrmann, Manfred; Hoy, Ludwig; Nippert, Irmgard; Schmidtke, Jorg (2005): Genotype-based screening for hereditary hemochromatosis: II. Attitudes toward genetic testing and psychosocial impact--a report from a German pilot study. In: *Genetic testing* 9 (3), S. 242–254.
- Sturm H.; Wildermuth R.; Stolz R.; Bertram L.; Eschweiler G.W.; Thomas C. et al. (2019): Diverging awareness of postoperative delirium and cognitive dysfunction in German health care providers. In: *Clinical Interventions in Aging* 14 ((Sturm, Wildermuth, Stolz, Bertram, Joos) Institute for General Practice and Interprofessional Care, University Hospital Tubingen, Tubingen, Germany(Eschweiler) Geriatric Center, University Hospital Tubingen, Tubingen 72076, Germany(Thomas) Department of), S. 2125–2135. DOI: 10.2147/CIA.S230800.
- Szecsenyi, Joachim; Rosemann, Thomas; Joos, Stefanie; Peters-Klimm, Frank; Miksch, Antje (2008): German diabetes disease management programs are appropriate for restructuring care according to the chronic care model: an evaluation with the patient assessment of chronic illness care instrument. In: *Diabetes care* 31 (6), S. 1150–1154. DOI: 10.2337/dc07-2104.
- Tamayo, T.; Tamayo, M.; Rathmann, W.; Potthoff, P. (2016): Prevalence of gestational diabetes and risk of complications before and after initiation of a general systematic two-step screening strategy in Germany (2012–2014). In: *Diabetes research and clinical practice* 115 (ebi, 8508335), S. 1–8. DOI: 10.1016/j.diabres.2016.03.001.
- Tanislav C.; Kostev K. (2019): Late Detection of Atrial Fibrillation after Stroke: Implications for the Secondary Prevention. In: *European Neurology* 81 (5–6), S. 262–269. DOI: 10.1159/000503562.
- *Thelen M.; Scheidt-Nave C.; Schaeffer D.; Bluher S.; Nitsche-Neumann L.; Sass A.-C.; Herweck R. (2012): National health target "healthy ageing". Medical, psychosocial and nursing care for elderly people. In: *Bundesgesundheitsblatt - Gesundheitsforschung - Gesundheitsschutz* 55 (8), S. 991–997. DOI: 10.1007/s00103-012-1517-0.
- Thielmann A.; Viehmann A.; Weltermann B.M. (2015): Effectiveness of a web-based education program to improve vaccine storage conditions in primary care (Keep Cool): study protocol for a randomized controlled trial. In: *Trials* 16 (1), S. 301. DOI: 10.1186/s13063-015-0824-9.
- Thiem U.; Wilm S.; Greiner W.; Rudolf H.; Trampisch H.; Muller C. et al. (2020): Reduction of potentially inappropriate medication in the elderly: design of a cluster-randomised controlled trial in German primary care practices (RIME). In: *Therapeutic Advances in Drug Safety* 12 ((Thiem, Trampisch) Department of Medical Informatics, Biometry and Epidemiology, University of Bochum, Bochum, Germany(Wilm) Institute of General Practice, University of Dusseldorf, Dusseldorf, Germany(Greiner) Department of Health Economics and Health Ca). DOI: 10.1177/2042098620918459.
- *Thomasius F.; Baum E.; Bernecker P.; Bocker W.; Brabant T.; Clarenz P. et al. (2018): S-3 DVO Guidelines 2017 in prophylaxis, diagnosis and therapy of osteoporosis in postmenopausal women und men. In: *Osteologie* 27 (3), S. 154–160. DOI: 10.1055/s-0038-1673537.
- *Tillmanns H.; Erdogan A.; Sedding D. (2009): Treatment of chronic CAD--do the guidelines (ESC, AHA) reflect daily practice? In: *Herz* 34 (1), S. 39–54.
- Tinsel, Iris; Buchholz, Anika; Vach, Werner; Siegel, Achim; Durk, Thorsten; Buchholz, Angela et al. (2013): Shared decision-making in antihypertensive therapy: a cluster randomised controlled trial. In: *BMC family practice* 14 (100967792), S. 135. DOI: 10.1186/1471-2296-14-135.

Tinsel, Iris; Buchholz, Anika; Vach, Werner; Siegel, Achim; Durk, Thorsten; Loh, Andreas et al. (2012): Implementation of shared decision making by physician training to optimise hypertension treatment. Study protocol of a cluster-RCT. In: *BMC cardiovascular disorders* 12 (100968539), S. 73. DOI: 10.1186/1471-2261-12-73.

Tinsel, Iris; Siegel, Achim; Schmoor, Claudia; Buchholz, Anika; Niebling, Wilhelm (2017): DECADE-pilot: decision aid, action planning, and follow-up support for patients to reduce the 10-year risk of cardiovascular diseases-a protocol of a randomized controlled pilot trial. In: *Pilot and feasibility studies* 3 (101676536), S. 32. DOI: 10.1186/s40814-017-0172-5.

*Trojan, J. (2011): [Colorectal cancer--current screening, treatment and follow-up standards]. In: *Versicherungsmedizin* 63 (3), S. 132–136.

*Twardella D.; Brenner H. (2003): New strategies for the promotion of smoking cessation in general practice: A cluster-randomised trial. In: *Suchtmedizin in Forschung und Praxis* 5 (2), S. 142–144.

Twardella D.; Brenner H. (2005): Lack of training as a central barrier to the promotion of smoking cessation: A survey among general practitioners in Germany. In: *European Journal of Public Health* 15 (2), S. 140–145. DOI: 10.1093/eurpub/cki123.

Tzaribachev N.; Benseler S.M.; Tyrrell P.N.; Meyer A.; Kuemmerle-Deschner J.B. (2009): Predictors of delayed referral to a pediatric rheumatology center. In: *Arthritis Care and Research* 61 (10), S. 1367–1372. DOI: 10.1002/art.24671.

Uhl M.C.; Muth C.; Gerlach F.M.; Schoch G.-G.; Muller B.S. (2018): Patient-perceived barriers and facilitators to the implementation of a medication review in primary care: a qualitative thematic analysis. In: *BMC family practice* 19 (1), S. 3. DOI: 10.1186/s12875-017-0707-0.

*Ulbricht, S.; Meyer, C.; Schumann, A.; Rumpf, H-J; Bischof, G.; Hapke, U.; John, U. (2004): [Promoting the intention to quit smoking among patients in family medicine practice]. In: *Gesundheitswesen (Bundesverband der Ärzte des Öffentlichen Gesundheitsdienstes (Germany))* 66 (8-9), S. 518–521.

Ulbricht, Sabina; Baumeister, Sebastian E.; Meyer, Christian; Schmidt, Carsten Oliver; Schumann, Anja; Rumpf, Hans-Jurgen; John, Ulrich (2009): Does the smoking status of general practitioners affect the efficacy of smoking cessation counselling? In: *Patient Education and Counseling* 74 (1), S. 23–28. DOI: 10.1016/j.pec.2008.07.047.

Ulbricht, Sabina; Meyer, Christian; Schumann, Anja; Rumpf, Hans-Jurgen; Hapke, Ulfert; John, Ulrich (2006): Provision of smoking cessation counseling by general practitioners assisted by training and screening procedure. In: *Patient Education and Counseling* 63 (1-2), S. 232–238.

*Ulbricht S.; Koepsell S.; Rumpf H.J.; John U.; Meyer C. (2010): [The readiness of staff in general medical practices to participate in a smoking cessation programme]. In: *Gesundheitswesen (Bundesverband der Ärzte des Öffentlichen Gesundheitsdienstes (Germany))* 72 (8-9), S. 492–495.

Ulrich, Lisa-R; Mergenthal, Karola; Petersen, Juliana J.; Roehl, Ina; Rauck, Sandra; Kemperdick, Birgit et al. (2014): Anticoagulant treatment in German family practices - screening results from a cluster randomized controlled trial. In: *BMC family practice* 15 (100967792), S. 170. DOI: 10.1186/s12875-014-0170-0.

Van Eerd E.A.M.; Risor M.B.; Spigt M.; Godycki-Cwirko M.; Andreeva E.; Francis N. et al. (2017): Why do physicians lack engagement with smoking cessation treatment in their COPD patients? A multinational qualitative study. In: *npj Primary Care Respiratory Medicine* 27 (1), S. 41. DOI: 10.1038/s41533-017-0038-6.

Van Lieshout J.; Wensing M.; Campbell S.M.; Grol R. (2009): Primary care strength linked to prevention programs for cardiovascular disease. In: *American Journal of Managed Care* 15 (4), S. 255–262.

*Vass, M.; Hendriksen, C. (2005): Polypharmacy and older people--the GP perspective. In: *Zeitschrift für Gerontologie und Geriatrie* 38 Suppl 1 (cc3, 9506215), I14-7.

*Voigt K.; Kuhne F.; Twork S.; Gobel A.; Kugler J.; Bergmann A. (2008): Current vaccination status of health-care personnel in Brandenburg, saxony and saxony-anhalt. In: *Gesundheitswesen* 70 (7), S. 408–414. DOI: 10.1055/s-0028-1082077.

Voigt-Radloff, Sebastian; Schopf, Andrea C.; Boeker, Martin; Frank, Luca; Farin, Erik; Kaier, Klaus et al. (2019): Well informed physician-patient communication in consultations on back pain - study protocol of the cluster randomized GAP trial. In: *BMC family practice* 20 (1), S. 33. DOI: 10.1186/s12875-019-0925-8.

Vollbracht C.; Schneider B.; Leendert V.; Weiss G.; Auerbach L.; Beuth J. (2011): Intravenous vitamin C administration improves quality of life in breast cancer patients during chemo-/radiotherapy and aftercare: Results of a retrospective, multicentre, epidemiological cohort study in Germany. In: *In Vivo* 25 (6), S. 983–990.

Von Dem Knesebeck O.; Hoehne A.; Link C.; Marceau L.; Adams A.; Roland M. et al. (2012): Talking about smoking in primary care medical practice-Results of experimental studies from the US, UK and Germany. In: *Patient Education and Counseling* 89 (1), S. 51–56. DOI: 10.1016/j.pec.2012.04.011.

Von Lengerke T.; Wichmann H.-E.; Lowel H.; Meisinger C.; Illig T.; Holle R.; John J. (2007): General practitioners' opportunities for preventing ill health in healthy vs morbid obese adults: A general population study on consultations. In: *Journal of Public Health* 15 (2), S. 71–80. DOI: 10.1007/s10389-006-0086-9.

*von Renteln-Kruse W.; Anders J.; Dapp U.; Meier-Baumgartner H.P. (2003): Preventative home visits by a specially trained nurse for 60-year olds and elderly in Hamburg. In: *Zeitschrift fur Gerontologie und Geriatrie* 36 (5), S. 378–391. DOI: 10.1007/s00391-003-0179-9.

*Von Voshaar T.; Behr J.; Bruggenjorgen B.; Deimling A.; Kruger M.; Virchow C.; Wiens C. (2012): [On general practitioners' care of patients with asthma]. In: *MMW Fortschritte der Medizin* 154 Suppl 1 ((Von Voshaar) Med. Klinik III, Pneumologie, Allergologie, Zentrum für Schlaf und Beatmungsmedizin), S. 1–7.

Vormfelde S.V.; Abu Abed M.; Hua T.D.; Schneider S.; Friede T.; Chenot J.-F. (2014): Educating orally anticoagulated patients in drug safety: A cluster-randomized study in general practice. In: *Deutsches Arzteblatt International* 111 (37), S. 607–614. DOI: 10.3238/arztebl.2014.0607.

*Wagner A.; Becker A. (2008): The role of sports as medicine in general practice exemplified by the coronary artery disease. In: *Zeitschrift fur Allgemeinmedizin* 84 (3), S. 125–140. DOI: 10.1055/s-2008-1062718.

*Wahle K. (2009): Responsibilities of the general practitioner in vaccination control. In: *Deutsche medizinische Wochenschrift* (1946) 134 Suppl 2 ((Wahle) Arzt für Allgemeinmedizin, Am Pastorenbusch 2, Munster), S65–70. DOI: 10.1055/s-0029-1220212.

Walter U.; Flick U.; Neuber A.; Fischer C.; Hussein R.J.; Schwartz F.W. (2010): Putting prevention into practice: qualitative study of factors that inhibit and promote preventive care by general practitioners, with a focus on elderly patients. In: *BMC family practice* 11 ((Walter) Institute for Epidemiology, Social Medicine and Health System Research, Hannover Medical School, Carl-Neuberg Strasse 1, OE 5410, 30623 Hannover, Germany), S. 68.

*Wangler J.; Jansky M. (2018): What benefits do health apps provide for family medicine?: A survey among family practitioners in hesse and thuringia. In: *Zeitschrift fur Allgemeinmedizin* 94 (6), S. 259–264. DOI: 10.3238/zfa.2018.0259-0264.

*Webendorfer S.; Messerer P.; Eberle F.; Zober A. (2004): Screening for colorectal cancer: An initiative for secondary prevention at the BASF Aktiengesellschaft. In: *Deutsche Medizinische Wochenschrift* 129 (6), S. 239–243. DOI: 10.1055/s-2004-817998.

*Weber, Annemarie; Schelling, Jorg; Kohls, Niko; van Dyck, Marcus; Poggenburg, Stephanie; Vajda, Christian et al. (2018): [Current Research Activities on Person-Centered Medicine in Academic Institutes of General Practice in Germany and Austria]. In: *Gesundheitswesen (Bundesverband der Ärzte des Öffentlichen Gesundheitsdienstes (Germany))* 80 (11), S. 1006–1012. DOI: 10.1055/s-0043-117734.

Weimar, Christian; Diener, Hans-Christoph; Alberts, Mark J.; Steg, P. Gabriel; Bhatt, Deepak L.; Wilson, Peter W. F. et al. (2009): The Essen stroke risk score predicts recurrent cardiovascular events: a validation within the REduction of Atherothrombosis for Continued Health (REACH) registry. In: *Stroke* 40 (2), S. 350–354. DOI: 10.1161/STROKEAHA.108.521419.

*Weiss H. (2019): Nutritional Counselling at the Edge between Clinic and Outpatient Care. In: *Aktuelle Ernährungsmedizin* 44 (4), S. 261–268. DOI: 10.1055/a-0894-1224.

*Weltermann, Birgitta; Reinders, Sabrina; Bettin, Markus; Gesenhues, Stefan; Hermann, Martin (2013): [Screening for diabetic retinopathy in type 2 diabetes: a critical review of an annual routine]. In: *Zeitschrift für Evidenz, Fortbildung und Qualität im Gesundheitswesen* 107 (6), S. 403–409. DOI: 10.1016/j.zefq.2012.11.019.

*Weltermann, Birgitta M.; Doost, Sahar Mousa; Kersting, Christine; Gesenhues, Stefan (2014): Hypertension management in primary care: how effective is a telephone recall for patients with low appointment adherence in a practice setting? In: *Wiener klinische Wochenschrift* 126 (19–20), S. 613–618. DOI: 10.1007/s00508-014-0590-9.

Weltermann, Birgitta M.; Markic, Marta; Thielmann, Anika; Gesenhues, Stefan; Hermann, Martin (2014): Vaccination management and vaccination errors: a representative online-survey among primary care physicians. In: *PLoS ONE* 9 (8), e105119. DOI: 10.1371/journal.pone.0105119.

*Weltermann B.; Kempis P.; Reinders S.; Gesenhues S. (2013): Patient-centered health care management: A concept developed in a teaching practice. In: *Zeitschrift für Allgemeinmedizin* 89 (3), S. 127–132.

Weltermann B.; Viehmann A.; Kersting C. (2015): Hypertension management in primary care: Study protocol for a cluster randomized controlled trial. In: *Trials* 16 (1), S. 105. DOI: 10.1186/s13063-015-0627-z.

Weltermann B.M.; Driouach-Bleckmann Y.; Reinders S.; Berndt P.; Gesenhues S. (2013): Stroke knowledge among diabetics: A cross-sectional study on the influence of age, gender, education, and migration status. In: *BMC Neurology* 13 ((Weltermann, Driouach-Bleckmann, Reinders, Berndt, Gesenhues) Institute for General Medicine, University of Duisburg-Essen, Hufelandstr 55, 45122 Essen, Germany), S. 202. DOI: 10.1186/1471-2377-13-202.

Weltermann B.M.; Schroeder V.; Rieger M.A.; Rind E.; Siegel A.; Wagner A. et al. (2020): IMPROVEjob - Participatory intervention to improve job satisfaction of general practice teams: A model for structural and behavioural prevention in small and medium-sized enterprises - A study protocol of a cluster-randomised controlled trial. In: *Trials* 21 (1), S. 532. DOI: 10.1186/s13063-020-04427-7.

Welzel, Franziska D.; Stein, Janine; Pabst, Alexander; Lappa, Melanie; Kersting, Anette; Bluher, Matthias et al. (2018): Five A's counseling in weight management of obese patients in primary care: a cluster-randomized controlled trial (INTERACT). In: *BMC family practice* 19 (1), S. 97. DOI: 10.1186/s12875-018-0785-7.

Wennehorst K.; Mildenstein K.; Saliger B.; Tigges C.; Diehl H.; Keil T.; Englert H. (2016): A Comprehensive Lifestyle Intervention to Prevent Type 2 Diabetes and Cardiovascular Diseases: the German CHIP Trial. In: *Prevention science : the official journal of the Society for Prevention Research* 17 (3), S. 386–397. DOI: 10.1007/s11121-015-0623-2.

*Westhofen, Johannes; Puth, Marie-Therese; Steudel, Henrike; Munster, Eva; Weckbecker, Klaus (2020): [Health Expertise of Seasonal Influenza Vaccination among Midwife Trainees in North Rhine-Westphalia: Results of a Cross-Sectional Study]. In: *Gesundheitswesen (Bundesverband der Arzte des Offentlichen Gesundheitsdienstes (Germany))* 82 (3), S. 253–259. DOI: 10.1055/a-0839-4810.

*Wiesemann A.; Barlet J.; Engeser P.; Kuth N.; Muller-Buhl U. (2006): Obesity behavior change promotion in primary care. In: *Zeitschrift fur Allgemeinmedizin* 82 (3), S. 103–107. DOI: 10.1055/s-2006-921505.

*Wiesemann A.; Engeser P.; Reichert K.; Mollien P.; Schach S.; Scheidt R. (2004): Quality of Primary Medical Care after In-patient Stroke Rehabilitation - (HANS-Study of GP). In: *Gesundheitswesen* 66 (1), S. 29–36. DOI: 10.1055/s-2004-812829.

Wiesemann A.; Ludt S.; Szecsenyi J.; Scheuermann W.; Scheidt R. (2004): Cardiovascular risk factors and motivation for a healthy life-style in a German community - Results of the GP-based Oestringen study. In: *Patient Education and Counseling* 55 (1), S. 40–47. DOI: 10.1016/j.pec.2003.07.002.

*Wiesemann A.; Mueller-Buehl U.; Scheidt R.; Boehme W.; Scheuermann W. (2004): Patient willingness to pay for preventive measures in primary care: A study of five GPs in a German community. In: *Sozial- und Praventivmedizin* 49 (4), S. 254–260. DOI: 10.1007/s00038-004-3039-5.

*Wittchen, H. U.; Krause, P.; Hofler, M.; Pfister, H.; Kupper, B.; Pittrow, D. et al. (2003): [Aim, design and methods of the "Hypertension and diabetes screening and awareness" -- (HYDRA) study]. In: *Fortschritte der Medizin. Originalien* 121 Suppl 1 (101120496), S. 2–11.

*Wittmund B.; Wilms H.-U.; Bull N.; Angermeyer M.C. (2003): Cooperation of a specialized outpatient unit, general practitioners and psychiatrists in the treatment of anxiety disorders. In: *Verhaltenstherapie* 13 (4), S. 284–290. DOI: 10.1159/000075844.

Wolffram, Ingmar; Petroff, David; Batz, Olaf; Jedrysiak, Katrin; Kramer, Jan; Tenckhoff, Hannelore et al. (2015): Prevalence of elevated ALT values, HBsAg, and anti-HCV in the primary care setting and evaluation of guideline defined hepatitis risk scenarios. In: *Journal of hepatology* 62 (6), S. 1256–1264. DOI: 10.1016/j.jhep.2015.01.011.

Wollny A.; Altiner A.; Brand T.; Garbe K.; Kamradt M.; Kaufmann-Kolle P. et al. (2019): Converting habits of antibiotic use for respiratory tract infections in German primary care - Study protocol of the cluster-randomized controlled CHANGE-3 trial. In: *Trials* 20 (1), S. 103. DOI: 10.1186/s13063-019-3209-7.

*Wortberg, S.; Walter, D. (2010): [Recallsystems in primary care practices to increase vaccination rates against seasonal influenza]. In: *Deutsche medizinische Wochenschrift* (1946) 135 (22), S. 1113–1117. DOI: 10.1055/s-0030-1247867.

*Wortberg S.; Walter D.; Knesebeck M.; Reiter S. (2009): Physicians as key communicators of the influenza vaccination for the elderly, patients with chronic conditions, and health care workers. Results of a nationwide survey in the context of the national influenza vaccination campaign. In: *Bundesgesundheitsblatt, Gesundheitsforschung, Gesundheitsschutz* 52 (10), S. 945–952. DOI: 10.1007/s00103-009-0871-z.

Wutzler, Alexander; Krogias, Christos; Grau, Anna; Veltkamp, Roland; Heuschmann, Peter U.; Haeusler, Karl Georg (2019): Stroke prevention in patients with acute ischemic stroke and atrial fibrillation in Germany - a cross sectional survey. In: *BMC Neurology* 19 (1), S. 25. DOI: 10.1186/s12883-019-1249-y.

Zelenak C.; Radenovic S.; Musial-Bright L.; Tahirovic E.; Sacirovic M.; Lee C.-B. et al. (2017): Heart failure awareness survey in Germany: general knowledge on heart failure remains poor. In: *ESC heart failure* 4 (3), S. 224–231. DOI: 10.1002/ehf2.12144.

*Zeymer U. (2007): Secondary prevention in outpatients with coronary artery disease: Adherence with recommendations within 4 weeks after hospital discharge. In: *Deutsche Medizinische Wochenschrift* 132 (45), S. 2367–2370. DOI: 10.1055/s-2007-991659.

*Zhan, Tianzuo; Hielscher, Thomas; Bilge, Asme; Giese, Thomas; Schafer, Christoph; Riemann, Jurgen F. et al. (2017): Invitation letters increase participation in colorectal cancer screening - results from an observational study. In: *Zeitschrift fur Gastroenterologie* 55 (12), S. 1307–1312. DOI: 10.1055/s-0043-121347.

*Ziegenhagen D.J.; Frye C.; Kottmair S. (2005): Patient-oriented healthcare programs: Concepts and practical experience in the field of chronic heart failure. In: *Zeitschrift fur Arztliche Fortbildung und Qualitatssicherung* 99 (3), S. 209–215.

Ziegler, Dan; Strom, Alexander; Lobmann, Ralf; Reiners, Karlheinz; Rett, Kristian; Schnell, Oliver (2015): High prevalence of diagnosed and undiagnosed polyneuropathy in subjects with and without diabetes participating in a nationwide educational initiative (PROTECT study). In: *Journal of diabetes and its complications* 29 (8), S. 998–1002. DOI: 10.1016/j.jdiacomp.2015.09.008.

*Ziegler, M.; Krause, S.; Kolligs, F. T. (2011): [How physicians inform about colorectal cancer and screening]. In: *Zeitschrift fur Gastroenterologie* 49 (5), S. 584–590. DOI: 10.1055/s-0029-1246026.

*Zieren J.; Neuss H.; Maecker F.; Muller J.M. (2002): What is the value of the internet as a source of information for patients with inguinal hernias? First results of an observational study. In: *Der Chirurg; Zeitschrift fur alle Gebiete der operativen Medizinen* 73 (5).

*Zimmermann, Thomas; Puschmann, Egina; Porzelt, Sarah; Ebersbach, Martin; Ernst, Annette; Thomsen, Patricia; Scherer, Martin (2015): [Promoting Self-Management in Primary Care - the Association of Motivation for Change, Self-Efficacy and Psychological Distress Prior to the Onset of Intervention]. In: *Psychiatrische Praxis* 42 Suppl 1 (qck, 0423204), S44-8. DOI: 10.1055/s-0034-1387686.

Zimmermann T.; Puschmann E.; Ebersbach M.; Daubmann A.; Steinmann S.; Scherer M. (2014): Effectiveness of a primary care based complex intervention to promote self-management in patients presenting psychiatric symptoms: Study protocol of a cluster-randomized controlled trial. In: *BMC Psychiatry* 14 (1), S. 2. DOI: 10.1186/1471-244X-14-2.

Zimmermann T.; Puschmann E.; van den Bussche H.; Wiese B.; Ernst A.; Porzelt S. et al. (2016): Collaborative nurse-led self-management support for primary care patients with anxiety, depressive or somatic symptoms: Cluster-randomised controlled trial (findings of the SMADS study). In: *International journal of nursing studies* 63 ((Zimmermann, Puschmann, Ernst, Porzelt, Scherer) Department of General Practice / Primary Care, Center for Psychosocial Medicine, University Medical Center Hamburg-Eppendorf, Martinstr. 52, 20246 Hamburg, Germany(van den Bussche) Department of General Pr), S. 101–111. DOI: 10.1016/j.ijnurstu.2016.08.007.

Zulke A.; Luck T.; Pabst A.; Hoffmann W.; Thyrian J.R.; Gensichen J. et al. (2019): AgeWell.de - study protocol of a pragmatic multi-center cluster-randomized controlled prevention trial against cognitive decline in older primary care patients. In: *BMC geriatrics* 19 (1), S. 203. DOI: 10.1186/s12877-019-1212-1.