i

Physics Letters B 704 (2011) 442-455



Contents lists available at SciVerse ScienceDirect

Physics Letters B

www.elsevier.com/locate/physletb



Rapidity and transverse momentum dependence of inclusive J/ ψ production in pp collisions at $\sqrt{s}=7$ TeV $^{\diamondsuit}$

ALICE Collaboration

ARTICLE INFO

Article history:
Received 3 May 2011
Received in revised form 9 September 2011
Accepted 14 September 2011
Available online 19 September 2011
Editor: L. Rolandi

ABSTRACT

The ALICE experiment at the LHC has studied inclusive J/ψ production at central and forward rapidities in pp collisions at $\sqrt{s}=7$ TeV. In this Letter, we report on the first results obtained detecting the J/ψ through the dilepton decay into e^+e^- and $\mu^+\mu^-$ pairs in the rapidity ranges |y|<0.9 and 2.5< y<4, respectively, and with acceptance down to zero p_T . In the dielectron channel the analysis was carried out on a data sample corresponding to an integrated luminosity $L_{\rm int}=5.6$ nb $^{-1}$ and the number of signal events is $N_{\rm J/\psi}=352\pm32$ (stat.) ±28 (syst.); the corresponding figures in the dimuon channel are $L_{\rm int}=15.6$ nb $^{-1}$ and $N_{\rm J/\psi}=1924\pm77$ (stat.) ±144 (syst.). The measured production cross sections are $\sigma_{\rm J/\psi}(|y|<0.9)=10.7\pm1.0$ (stat.) ±1.6 (syst.) $_{-2.3}^{+1.6}$ (syst.pol.) μ b and $\sigma_{\rm J/\psi}(2.5< y<4)=6.31\pm0.25$ (stat.) ±0.76 (syst.) $_{-1.96}^{+0.95}$ (syst.pol.) μ b. The differential cross sections, in transverse momentum and rapidity, of the $\rm J/\psi$ were also measured.

 $\ensuremath{\text{@}}$ 2011 CERN. Published by Elsevier B.V. All rights reserved.

1. Introduction

The hadroproduction of heavy quarkonium states is governed by both perturbative and non-perturbative aspects of Quantum Chromodynamics (QCD) and was extensively studied at the Tevatron [1-4] and RHIC [5] hadron colliders. Various theoretical approaches, recently reviewed in [6,7], were proposed to describe the data. They mainly differ in the details of the non-perturbative evolution of the heavy quark pair towards a bound state. The models are not able to consistently reproduce the production cross section, the transverse momentum (p_T) distributions and the polarization. Recently, theoretical studies focused on the calculation of NLO and NNLO contributions, finding that their impact on the results is quantitatively important [8-12]. Measurements in the new energy domain of the LHC are crucial for a deeper understanding of the physics involved in hadroproduction processes. Furthermore, the range of Bjorken-x values accessible at LHC energies is unique. Low- p_T charmonium measurements, in particular at forward rapidity, are sensitive to an unexplored region ($x < 10^{-5}$ at $Q^2 = m_{1/y_t}^2$) of the gluon distribution function of the proton.

Heavy quarkonia are measured in the ALICE experiment [13] through their e⁺e⁻ and $\mu^+\mu^-$ decays. In this Letter we present the results on inclusive J/ ψ production in pp collisions at $\sqrt{s}=7$ TeV, measured in the rapidity regions |y|<0.9 for the dielectron channel and 2.5 < y < 4.0 for the dimuon decay channel.¹

First, a description of the ALICE experimental apparatus is given, mainly focusing on the muon detection. Track reconstruction in the central rapidity region was discussed previously [14]. Details are provided concerning the data analysis, the reconstruction algorithm, the event selection criteria and the techniques used for the extraction of the signal. After describing the determination of the acceptance and efficiency corrections, and the methods used for the evaluation of the luminosity, the values of the integrated, y-differential and $p_{\rm T}$ -differential J/ ψ cross sections are presented and compared with the results obtained by the other LHC experiments [15–17].

2. Experimental apparatus and data taking conditions

The ALICE experiment [13] consists of two main parts: a central barrel and a muon spectrometer. The central barrel detectors ($|\eta| < 0.9$) are embedded in a large solenoidal magnet, providing a magnetic field of 0.5 T. Various detector systems track particles down to $p_{\rm T}$ of about 100 MeV/c, and can provide particle identification over a wide momentum range. The muon spectrometer covers the pseudo-rapidity range $-4 < \eta < -2.5$ and detects muons having p > 4 GeV/c. Finally, various sets of forward detectors further extend the charged particle pseudo-rapidity coverage up to $\eta = 5.1$ and can be used for triggering purposes.

The barrel detectors used in this analysis are the Inner Tracking System (ITS) and the Time Projection Chamber (TPC). The ITS

 $^{^1}$ The muon spectrometer covers, in the ALICE official reference frame, a negative η range and, consequently, a negative y range. However, since in pp the physics is

symmetric with respect to y = 0, we have dropped the negative sign when quoting rapidity values.

[13,18] is a cylindrically-shaped silicon tracker that surrounds the central beam pipe. It consists of six layers, with radii between 3.9 cm and 43.0 cm, covering the pseudo-rapidity range $|\eta| < 0.9$. The two innermost layers are equipped with Silicon Pixel Detectors (SPD), the two intermediate layers contain Silicon Drift Detectors (SDD), and Silicon Strip Detectors (SSD) are used on the two outermost layers. The main task of the ITS is to provide precise track and vertex reconstruction close to the interaction point, to improve the overall momentum resolution and to extend tracking down to very low $p_{\rm T}$. The SPD can also deliver a signal for the first level trigger (L0), which is based on a hit pattern recognition system at the level of individual readout chips.

The TPC [19] is a large cylindrical drift detector with a central high voltage membrane maintained at $-100~\rm kV$ and two readout planes at the end-caps. The active volume extends over the ranges $85 < r < 247~\rm cm$ and $-250 < z < 250~\rm cm$ in the radial and longitudinal (beam) directions, respectively. Besides being the main tracking detector in the central barrel, the TPC also provides charged hadron identification with very good purity up to a total momentum of about $3~\rm GeV/c$ [20] and electrons up to about $10~\rm GeV/c$, via specific energy loss (d $E/\rm dx$) measurement, with a resolution of $\sigma = 5.5\%$ for minimum ionizing particles.

Other central barrel detectors with full azimuthal coverage, in particular the Time-Of-Flight (TOF) [21], the Transition Radiation Detector (TRD) [22] and the Electromagnetic Calorimeter (EMCAL) [23], are not used in this analysis, but are expected to significantly improve the electron identification and triggering capabilities of the experiment in the future.

The muon spectrometer consists of a front absorber followed by a 3 Tm dipole magnet, coupled to tracking and triggering devices. Muons emitted in the forward rapidity region are filtered by means of a 10 interaction length (λ_I) thick front absorber made of carbon, concrete and steel, and placed between 0.9 and 5.0 m from the nominal position of the interaction point (IP). Muon tracking is performed by means of 5 tracking stations, positioned between 5.2 and 14.4 m from the IP, each one based on two planes of Cathode Pad Chambers. The total number of electronic channels is close to 1.1×10^6 , and the intrinsic spatial resolution for these detectors is of the order of 70 µm in the bending direction. Stations 1 and 2 (4 and 5) are located upstream (downstream) of the dipole magnet, while station 3 is embedded inside its gap. A muon triggering system is placed downstream of a 1.2 m thick iron wall $(7.2\lambda_I)$, which absorbs secondary hadrons escaping the front absorber and lowmomentum muons (having p < 1.5 GeV/c at the exit of the front absorber). It consists of two stations positioned at 16.1 and 17.1 m from the IP, equipped with two planes of Resistive Plate Chambers (RPC) each. The spatial resolution achieved is better than 1 cm, while the time resolution is of the order of 2 ns. Throughout its full length, a conical absorber ($\theta < 2^{\circ}$) made of tungsten, lead and steel protects the muon spectrometer against secondary particles produced by the interaction of large- η primaries in the beam pipe.

Finally, the VZERO detector consists of two scintillator arrays covering the range $2.8 < \eta < 5.1$ and $-3.7 < \eta < -1.7$, and positioned, respectively, at z=340 and z=-90 cm from the IP. It provides timing information to the L0 trigger with a resolution better than 1 ns. This feature proves to be useful in the offline rejection of beam-halo and beam-gas events.

The results presented in this Letter were obtained by analyzing data collected in the first year of operation of the LHC, corresponding to pp collisions at $\sqrt{s}=7$ TeV. During this period, the LHC reached its goal of delivering more than 10^{32} cm $^{-2}$ s $^{-1}$ instantaneous luminosity. In ALICE, the instantaneous luminosity was kept to $0.6-1.2\times10^{29}$ cm $^{-2}$ s $^{-1}$ in order to have a collision pile-up rate in the same bunch crossing below 5%. In order to increase the statistics for low cross-section processes, ALICE ran, during short

periods, at luminosities about 10 times higher, therefore with a much larger pile-up rate in the same bunch crossing. For the analysis presented in this Letter, the collected data were divided into three sub-periods. Each one corresponds to similar average luminosity and pile-up rates, and is characterized by reasonably stable tracking and trigger detector configuration and performance.

The event sample used in this analysis corresponds to minimum bias events (MB trigger) and, for the muon analysis, to events where the detection of at least one muon in the angular acceptance of the muon spectrometer (μ -MB trigger) is additionally required. The MB trigger is defined as the logical OR between the requirement of at least one fired readout chip in the SPD, and a signal in at least one of the two VZERO detectors [24]. It also requires a coincidence with signals from two beam pick-up counters, one on each side of the interaction region, indicating the passage of proton bunches. The μ -MB trigger allows the selection of events where at least one particle was detected in the trigger chambers of the muon spectrometer. The trigger logic is based on the requirement of having at least 3 hits (out of 4) in the trigger stations, both in the bending and non-bending directions [25]. In this way one can define a "trigger track", and compute its deviation with respect to a track with infinite momentum. By requiring such a deviation to be smaller than a certain value one can select muon candidate tracks having a transverse momentum larger than a predefined value. Such a $p_{\rm T}^{\rm trig}$ cut can be used to reject soft muons, dominated by π and K decays, and is able to limit the muon trigger rate when the machine luminosity is high. The instantaneous luminosity at ALICE allowed for the choice of the lowest $p_{\mathrm{T}}^{\mathrm{trig}}$ threshold (0.5 GeV/c), leading to a μ -MB trigger rate between 30 and 500 Hz. With this $p_{\rm T}^{\rm trig}$, the effect of the trigger response function on the J/ ψ detection efficiency is negligible. Note that the effect of such a cut is not sharp and that the selection efficiency reaches the plateau value only at $p_T \sim 1.5$ GeV/c. Finally, in order to limit the systematic uncertainties related to non-uniformities in the detector response, data were collected by periodically varying the polarities of the solenoidal and dipole magnets.

3. Data analysis

For the dielectron analysis, 3.5×10^8 minimum bias events $(N_{\rm MB})$ are analyzed. An event with a reconstructed vertex position $z_{\rm V}$ is accepted if $|z_{\rm V}| < 10$ cm. The tracks are required to have a minimum $p_{\rm T}$ of 1.0 GeV/c, a minimum number of 70 TPC clusters per track (out of a maximum of 159), a χ^2 per space point of the momentum fit lower than 4, and to point back to the interaction vertex within 1 cm in the transverse plane. A hit in at least one of the two innermost layers of the ITS is required in order to reduce the contribution of electrons from γ conversions. For full-length tracks, the geometrical coverage of the central barrel detectors is $|\eta| < 0.9$.

The particle identification performance of the TPC is essential for the J/ ψ measurement. In Fig. 1 the specific energy loss in the TPC is shown as a function of momentum in the region of interest for the present measurement. A $\pm 3\sigma$ inclusion cut for electrons and a $\pm 3.5\sigma$ ($\pm 3\sigma$) exclusion cuts for pions (protons) were employed. As seen in Fig. 1, with our current identification strategy, the electron identification is performed with an efficiency better than 50% for momenta below 7–8 GeV/c.

Electron candidates compatible, together with a positron candidate, with being products of γ conversions were removed, in order to reduce the combinatorial background. It was verified, using a Monte Carlo simulation, that this procedure does not affect the J/ ψ signal.




Fig. 1. Specific energy loss in the TPC as a function of momentum with superimposed Bethe–Bloch lines for various particle species. The dashed lines show the pion and proton exclusion bands. The dotted line corresponds to the $+3\sigma$ cut for electrons (see text).




Fig. 2. Top panel: invariant mass distributions for opposite-sign (OS) and like-sign (LS) electron pairs (|y| < 0.9, all p_T), as well as for pairs obtained with one track randomly rotated (TrkRot, see text). Bottom panel: the difference of the OS and LS distributions with the fit to the Monte Carlo (MC) signal superimposed.

The invariant mass distribution for the opposite-sign (OS) electron pairs is shown in Fig. 2. In the same figure we also show the background contribution, obtained as the sum of the like-sign (LS) pairs, $N^{++} + N^{--}$, scaled to match the integral of the OS distribution in the mass interval 3.2–5.0 GeV/ c^2 . The scale factor, 1.23, originates from the presence of correlated background (mostly from semi-leptonic charm decays) in the OS distribution, but is also influenced by misidentified electrons and by electrons

from conversions. In the top panel of Fig. 2 we also show the background estimated using a track rotation method (TrkRot),² used later in the estimate of the systematic uncertainties related to signal extraction. The signal, obtained by subtracting the scaled LS distribution from the OS, is shown in the bottom panel of Fig. 2 in comparison with the signal from Monte Carlo (MC) simulations (described below). A good agreement between data and MC is observed, both for the bulk of the signal and for the bremsstrahlung tail. Integration of the signal in the mass range 2.92–3.16 GeV/ c^2 yields $N_{\mathrm{I/\psi}} = 352 \pm 32 \; (\mathrm{stat.}) \pm 28 \; (\mathrm{syst.})$ counts (the systematic uncertainty on this quantity is described below); the signal to background ratio is $S/B = 1.2 \pm 0.1$ and the significance is $S/\sqrt{S+B} = 13.9 \pm 0.6$. The tagging and corresponding rejection of γ conversions is found to improve S/B by ~30%. The MC simulations show that $(73.4 \pm 2.0)\%$ of the signal is within the integration range. The error on this quantity was obtained by analyzing MC samples where the detector material budget was varied by $\pm 6\%$ [26] with respect to the nominal value, and by varying the track-related cuts (p_T and required number of TPC clusters) around their nominal values. A fit to the invariant mass distribution after background subtraction with a Crystal Ball function [27] gives a mass resolution of 28.3 \pm 1.8 MeV/ c^2 .

For the dimuon channel, the total data sample available for physics analysis amounts to 1.9×10^8 MB events, of which 1.0×10^7 satisfy the μ -MB condition.

An accurate alignment of the tracking chambers of the muon spectrometer is an essential pre-requisite to identify resonances in the $\mu^+\mu^-$ invariant mass spectrum. This was carried out using a modified version of the MILLEPEDE package [28,29], starting from a sample of 3×10^5 tracks, taken with no magnetic field in the dipole and in the solenoid. The resulting alignment precision is $\sim\!750~\mu\mathrm{m}$ in the bending and non-bending directions.

Track reconstruction is based on a Kalman filter algorithm [29, 30]. The procedure starts from the most downstream tracking stations (4 and 5), which are less subject to the background due to soft particles that escape the front absorber. Straight line segments are formed by joining clusters on the two planes of each station and a first estimate of the track parameters (position, slope and inverse bending momentum) and corresponding errors is made. The momentum is first estimated assuming that the track originates from the vertex and is bent by a constant magnetic field in the dipole. In a second step, track candidates on station 4 are extrapolated to station 5 (or vice versa) and paired with at least one cluster on the basis of a χ^2 cut. If several clusters are found, the track is duplicated to consider all the possible combinations. After this association the track parameters and errors are recalculated using the Kalman filter.

The same procedure is repeated iteratively for the upstream stations, rejecting, at each step, the candidates for which no cluster is found or those whose parameters indicate that they will exit the geometrical acceptance of the spectrometer in the next steps. At the end of the procedure, additional algorithms are applied to improve the track quality by adding/removing clusters based on a χ^2 cut, and removing fake tracks sharing clusters with others. Finally, the remaining tracks are extrapolated to the primary vertex position as given by the SPD [24], and their parameters are recomputed taking into account the energy loss and multiple Coulomb scattering in the absorber. With the alignment precision obtained for the analyzed data sample the relative momentum resolution of

 $^{^2}$ The method consists in rotating, around the z axis, one of the tracks of the OS pair by a random azimuthal angle. More pairs can be obtained by applying the method several times to the same pair.




Fig. 3. Invariant mass distribution for opposite-sign muon pairs (2.5 < y < 4, all $p_{\rm T}$), in the mass region $1.5 < m_{\mu\mu} < 5~{\rm GeV}/c^2$ with the result of the fit. The plot refers to the sub-period with the largest statistics $(N_{\rm J/\psi} = 957 \pm 56$, corresponding to $L_{\rm int} = 7.9~{\rm nb}^{-1}$). The fitted ${\rm J/\psi}$ and $\psi(2S)$ contributions, as well as the background, are also shown.

the reconstructed tracks ranges between 2% at 10 ${\rm GeV}/c$ and 10% at 100 ${\rm GeV}/c$.

After reconstruction, 4.1×10^5 events having at least two muon candidates are found, out of which only 6% have three or more muons. Various selection cuts are then applied to this data sample. First, events are required to have at least one interaction vertex reconstructed by the SPD. This cut rejects 0.5% of the statistics. Then, it is required that at least one of the two muon candidates matches the corresponding hits in the trigger chambers. In this way hadrons produced in the absorber, which are stopped by the iron wall positioned upstream of the trigger chambers, are efficiently rejected. This cut rejects ~24% of the muon pairs, and its effect is important only for $m_{\mu\mu} < 1 \text{ GeV}/c^2$. In fact, since in 99% of the cases at least one of the two J/ ψ decay muons has a transverse momentum larger than the trigger p_T threshold, the signal loss induced by this cut is negligible. Requiring both candidate muon tracks to be matched with the corresponding "trigger tracks" would increase the purity of the muon sample, but it was checked that this cut would lead to a loss of \sim 20% of the J/ ψ events without decisively increasing the signal to background ratio at the I/ψ invariant mass. Furthermore, the cut $R_{\rm abs} > 17.5$ cm, where $R_{\rm abs}$ is the radial coordinate of the track at the end of the front absorber, was applied. In this way, muons emitted at small angles, that have crossed a significant fraction of the thick beam shield, can be rejected. Finally, to remove events very close to the edge of the spectrometer acceptance, the cut 2.5 < y < 4 on the pair rapidity was applied. These quality cuts reject 10.3% of the muon pairs.

After selection, the dimuon sample consists of 1.75×10^5 OS muon pairs. In Fig. 3 we present the OS invariant mass spectrum for the mass region $1.5 < m_{\mu\mu} < 5$ GeV/ c^2 , corresponding to the sub-period having the largest statistics. A peak corresponding to the J/ $\psi \rightarrow \mu^+\mu^-$ decay is clearly visible in the spectrum, on top of a large continuum. A weaker signal, corresponding to the $\psi(2S)$ decay, is also visible, in spite of the poor signal to background ratio.

The number of signal events $N_{\rm J/\psi}$ was extracted by fitting the mass range $1.5 < m_{\mu\mu} < 5~{\rm GeV/c^2}$. The ${\rm J/\psi}$ and $\psi(2S)$ line shapes are described with Crystal Ball functions [27], while the underlying continuum was parameterized using the sum of two exponentials. The functions representing the resonances were obtained by fitting the expected mass distribution of a pure MC signal sample. Such a sample was obtained by generating, for each sub-period, ${\rm J/\psi}$ and $\psi(2S)$ events with realistic differential distributions (see below for details). In order to account for small uncertainties in the MC description of the set-up, the position of the ${\rm J/\psi}$ mass pole, as well as the width of the Crystal Ball function, were kept as free parameters in the invariant mass fit. Due to the small statistics, the $\psi(2S)$ parameters were tied to those of the ${\rm J/\psi}$, imposing the mass difference between the two states to be equal to the one given by the Particle Data Group (PDG) [31], and the ratio of the resonance widths to be equal to the one obtained in the MC.

This choice of parameters leads to a satisfactory fit of the invariant mass spectrum ($\chi^2/ndf=1.14$), as shown in Fig. 3. The Crystal Ball function describing the J/ ψ is peaked at $m_{\mathrm{J/\psi}}=3.118\pm0.005~\mathrm{GeV/c^2}$. Such a value is larger than the one quoted by the PDG group by only 0.6%, showing that the accuracy of the magnetic field mapping and of the energy loss correction is reasonably under control. The measured width of the Crystal Ball function is $\sigma_{\mathrm{J/\psi}}=94\pm8~\mathrm{MeV/c^2}$, in agreement within less than 2% with the MC, and its FWHM is 221 MeV/ c^2 .

The same fitting procedure, applied to the other sub-periods, gives consistent results in terms of both $m_{J/\psi}$ (within 0.2%) and $\sigma_{J/\psi}$ (within 4%). The signal to background ratio, in the mass range $2.9 < m_{\mu\mu} < 3.3 \text{ GeV/}c^2$, varies between 2.3 and 2.9 in the various sub-periods. The total number of J/ψ signal events, obtained by integrating the Crystall Ball function over the full mass range, is $N_{J/\psi} = 1924 \pm 77$ (stat.) ± 144 (syst.). The determination of the systematic uncertainty on $N_{J/\psi}$ is described later in Section 5.

4. Acceptance and efficiency corrections, luminosity normalization

In order to extract the J/ψ yield, the number of signal events must be corrected, with a MC procedure, for the acceptance of the apparatus and for reconstruction and triggering efficiencies. This procedure is based on the generation of a large sample of signal events, with a p_T distribution extrapolated from CDF measurements [1] and a y distribution parameterized from Color Evaporation Model (CEM) calculations [32]. To avoid the loss of events due to smearing effects at the edge of the angular acceptance, the generation was performed over y ranges wider than those covered by the two detector systems. It was also assumed that J/ψ production is unpolarized. The acceptance factors are obtained with respect to the J/ψ rapidity ranges |y| < 0.9 and 2.5 < y < 4.0 for the central barrel and muon detectors, respectively.

For the central barrel detectors, the acceptance times efficiency value $(A \times \epsilon)$ is 9.8% and is the product of four contributions: (i) a kinematic factor, namely the requirement of having both e^+ and e^- within the acceptance $(|\eta^{e^+,e^-}| < 0.9)$, satisfying a transverse momentum cut $p_T^{e^+,e^-} > 1$ GeV/c. This factor amounts to 32.8%; (ii) the reconstruction efficiency for the e^+e^- pair, which is 50.3%; (iii) the identification efficiency, which is 81.0%; (iv) the fraction of the signal within the mass range 2.92–3.16 GeV/ c^2 , which is 73.4%.

For the muon spectrometer, the tracking efficiency is calculated from MC simulations, including a realistic map of dead channels and the residual misalignment of the detection elements. This efficiency, obtained from a sample of generated tracks which match the hit pattern on the various chambers required by the reconstruction algorithm, is $(97.1 \pm 0.8)\%$.

The efficiencies of the muon trigger chambers are obtained from the analysis of the "trigger tracks" collected in the measured data sample. The "trigger tracks", as explained in Section 2, are defined by the presence of a hit in at least 3 (out of 4) trigger planes. The efficiency for a chamber belonging to a certain trigger plane is calculated starting from a sample of "trigger tracks" where the corresponding chambers on the other 3 planes have recorded a hit, and then looking for the presence or absence of a hit in the chamber under study. Such a requirement on the sample does not introduce a bias in the efficiency calculation because the response of the detector planes are independent. Typical efficiency values are around 96%, with 90% of the detector surface having an efficiency larger than 91%. The obtained efficiencies are then plugged in the simulations in order to provide a realistic description of the detector. The time variation of the tracking and trigger detector efficiencies was accounted for in the simulation. Internally to each sub-period, the response of the tracking chamber channels is further weighted run by run.

For each sub-period i, the ratio between the total number of reconstructed events, satisfying the analysis cuts, and the generated events in the range 2.5 < y < 4 gives the product $A \times \varepsilon_i$ for the J/ ψ . The differences between the $A \times \varepsilon_i$ for the various sub-periods do not exceed 8%, and their average value is $\langle A \times \varepsilon \rangle = 32.9\%$. It is worth noting that $A \times \varepsilon$ exhibits, for both the central barrel detectors and the muon spectrometer, a rather small variation as a function of the J/ ψ $p_{\rm T}$, down to zero $p_{\rm T}$.

To get the production cross section value, the ratio $N_{{\rm J/\psi}}^{\rm cor}=N_{{\rm J/\psi}}/\langle A \times \varepsilon \rangle$ must be normalized to the integrated luminosity, or to the measured cross section for a chosen reference process. For this analysis, the adopted reference is the occurrence of the MB condition itself. One has simply

$$\sigma_{\mathrm{J/\psi}} = \frac{N_{\mathrm{J/\psi}}^{\mathrm{cor}}}{\mathrm{BR}(\mathrm{J/\psi} \to \ell^{+}\ell^{-})} \times \frac{\sigma_{\mathrm{MB}}}{N_{\mathrm{MB}}} \tag{1}$$

where BR(J/ $\psi \to \ell^+ \ell^-$) = (5.94 \pm 0.06)% [31], $N_{\rm MB}$ is the number of minimum bias collisions and $\sigma_{\rm MB}$ is the measured cross section for such events. $N_{\rm MB}$ was corrected, run by run, for the probability of having multiple interactions in a single bunch crossing.

In the muon channel, the J/ ψ signal was collected using the μ -MB trigger condition. Therefore, Eq. (1) has to include a multiplicative factor R that links the occurrence of a reference process in the μ -MB and MB event samples. We have chosen as a reference process the yield N_{μ} of single muons, detected in the region $-4 < \eta < -2.5$, and with $p_{\rm T} > 1$ GeV/c. The R factor is then defined as the ratio $R = N_{\mu}^{\rm MB}/N_{\mu}^{\mu-{\rm MB}}$ of the single muon yields for the two event samples. The numerical values of the R factor strongly depend on the relative bandwidth assigned by the data acquisition to the two trigger samples in each sub-period, and vary between 0.10 and 0.42. However, the choice of the $p_{\rm T}$ cut has no significant influence on these values (<1% for cut values between 0 and 3 GeV/c). This is due to the fact that both the μ -MB and MB muon samples are subject to the same set of cuts, including the requirement of matching of the reconstructed track with the corresponding trigger track.

The $\sigma_{\rm MB}$ value is 62.3 mb, and is affected by a 4% systematic uncertainty. It was obtained relative to the cross section $\sigma_{\rm VOAND}$ [33], measured in a van der Meer scan [34], of the coincidence VOAND between signals in the two VZERO detectors. The relative factor $\sigma_{\rm VOAND}/\sigma_{\rm MB}$ was obtained as the fraction of MB events where the LO trigger input corresponding to the VOAND condition has fired. Its value is 0.87, and is stable within 0.5% over the analyzed data sample. The integrated luminosity $L_{\rm int} = N_{\rm MB}/\sigma_{\rm MB}$

is 5.6 nb⁻¹ for the dielectron sample. For the dimuon sample $L_{\rm int} = (N_{\rm MB}/\sigma_{\rm MB})/R = 15.6 \ {\rm nb}^{-1}$.

5. Systematic uncertainties

The systematic uncertainty on the inclusive J/ψ cross section measurement was obtained considering the following sources:

- The uncertainty on the signal extraction procedure, for the electron channel, was estimated using the track rotation method as an alternative background calculation procedure, see Fig. 2, and by fitting the invariant mass distributions with a convolution of a polynomial and a Crystal Ball function (with parameters constrained via Monte Carlo). We have also varied the invariant mass ranges for the signal extraction and for background normalization. The value obtained is 8%. Including the contribution from the uncertainty on the material budget leads to a value of 8.5%. For the muon channel, various tests were performed. In particular, we tried to release in the fit the values of the parameters governing the asymmetric left tail of the Crystal Ball line shape, which were fixed to their MC values in the default fitting procedure. Alternative functions for the description of the signal and background shapes were also used. In particular, for the I/ψ a variable-width Gaussian function, adopted in the past by the NA50 and NA60 Collaborations [35], was used. For the background, a different shape was tested, based on a Gaussian having a width continuously increasing with the mass. The estimated overall systematic uncertainty on the signal extraction is 7.5%.
- The acceptance calculation depends on the y and p_T input distributions. For the electrons, the uncertainty is mainly determined by the choice of the p_T spectrum. By varying the $\langle p_{\rm T} \rangle$ of the input distribution within a factor 2, a 1.5% variation in the acceptance was obtained. Such a small value is indeed a consequence of the weak p_T dependence of the acceptance for the bulk of the spectrum. For the muons, both y and p_T were varied, using as alternative distributions those expected for pp collisions at $\sqrt{s} = 4$ and 10 TeV [36]. As a further test, the measured J/ψ differential distributions obtained from the analysis described later in Section 6, were used as an input in the calculation of the acceptance. In this way, a 5% systematic uncertainty on this quantity was determined. The larger systematic uncertainty for the muon channel is due to the larger influence, for this channel, of the choice of the shape of the rapidity distribution.
- The uncertainty on the muon trigger efficiency calculation was estimated comparing N^{cor}_{J/ψ} for the sample where only one of the two decay muons is required to match the trigger condition, with the same quantity for the sample where both muons satisfy that condition. The 4% discrepancy between the two quantities is taken as the systematic uncertainty on the evaluation of the trigger efficiency.
- The uncertainty on the reconstruction efficiency, for the central barrel analysis, is due to the track quality (4%) and particle identification (10%) cuts and originates from residual mismatches between data and MC simulations.
- For the muon analysis, the systematic uncertainty can be estimated by comparing determinations of the tracking efficiency based on real data and on a MC approach. In the first case, the tracking efficiency can be evaluated starting from the determination of the efficiency per chamber, computed using the redundancy of the tracking information in each station. The values thus obtained are in the range from 91.8 to 99.8%. The tracking efficiency evaluated starting from these chamber efficiencies is $(98.8 \pm 0.8)\%$. The very same procedure, in

-10

Table 1Systematic uncertainties (in percent) on the quantities associated to the integrated V_{t} / V_{t} cross section measurement.

Channel	e^+e^-		$\mu^+\mu^-$	
Signal extraction	8.5		7.5	
Acceptance input	1.5		5	
Trigger efficiency	0		4	
Reconstruction efficiency	11		3	
R factor	-		3	
Luminosity	4		5.5	
B.R.			1	
Polarization	$\lambda = -1$	$\lambda = 1$	$\lambda = -1$	$\lambda = 1$
CS	+19	-13	+31	-15

-15

+21

HE

- a MC approach, gives a $(99.8^{+0.2}_{-0.8})\%$ tracking efficiency. These two quantities differ by 1%, which is taken as an estimate of the systematic uncertainty. However, this method is not able to detect losses of efficiency due to the presence of correlated dead-areas in the same region of two chambers belonging to the same station. Such correlated dead-areas were singled out by studying, on data, the cluster maps of each station, and the corresponding loss of efficiency was estimated to be $(2.8 \pm 0.4)\%$. Taking into account this effect, the resulting tracking efficiency is in good agreement with the value previously quoted (see Section 4) from realistic MC simulations, $(97.1 \pm 0.8)\%$. Nevertheless, a 1% additional systematic error (30% of the efficiency loss discussed above) was assumed, to take into account possible small-area correlations that could be missed in the present approach. Combining this error with the one previously mentioned, the overall systematic uncertainty on the muon tracking efficiency is 1.5%, which gives 3% for muon pair detection.
- The error on the luminosity measurement is dominated by the 4% systematic uncertainty on the determination of σ_{VOAND} , which is due to the uncertainties on the beam intensities [37] and on the analysis procedure related to the van der Meer scan of the VOAND signal. Other effects, such as the oscillation in the ratio between the MB and VOAND counts, contribute to less than 1%. The cross section $\sigma_{\mu\text{-MB}}$, relative to the occurrence of the μ -MB trigger, was also measured in a van der Meer scan [33,34] and was used as an alternative reference for the luminosity determination in the muon analysis [38]. Using $\sigma_{\mu\text{-MB}}$ as a reference cross section, a 4% difference has been found with respect to the integrated luminosity based on $\sigma_{\rm MR}$. For safety, this 4% has been quadratically added to the luminosity systematic error in the muon analysis. In addition, for the muons, the calculation of the integrated luminosity, as described above, is also connected with the estimate of the R factor. This quantity was evaluated in an alternative way, using the information from the trigger scalers and taking into account the dead-time of the triggers. By comparing the two results, a 3% systematic uncertainty on the R factor was estimated.
- The branching ratio of the J/ψ decay to lepton pairs is known with a 1% accuracy.
- The acceptance values significantly depend on the degree of polarization assumed in the J/ ψ distributions. They were calculated in the two cases of fully transverse ($\lambda=1$) or longitudinal ($\lambda=-1$) polarization,³ in the Collins–Soper (CS) and helicity (HE) reference frames.

The systematic uncertainties are summarized in Table 1. The systematic uncertainty on the inclusive J/ψ cross section is obtained by quadratically combining the errors from the sources described above, except polarization, and is 12.1% for the dimuon channel and 14.5% for the dielectron one. The systematic uncertainty due to the unknown J/ψ polarization will be quoted separately.

6. Integrated and differential J/ψ cross sections

The inclusive J/ψ production cross sections in pp collisions at $\sqrt{s} = 7$ TeV are:

$$\begin{split} \sigma_{J/\psi} \left(|y| < 0.9 \right) \\ &= 10.7 \pm 1.0 \text{ (stat.)} \pm 1.6 \text{ (syst.)} + 1.6 \text{ } (\lambda_{HE} = 1) \\ &- 2.3 \text{ } (\lambda_{HE} = -1) \text{ } \mu\text{b} \quad \text{and} \\ \sigma_{J/\psi} \left(2.5 < y < 4 \right) \\ &= 6.31 \pm 0.25 \text{ (stat.)} \pm 0.76 \text{ (syst.)} + 0.95 \text{ } (\lambda_{CS} = 1) \\ &- 1.96 \text{ } (\lambda_{CS} = -1) \text{ } \mu\text{b}. \end{split}$$

The systematic uncertainties related to the unknown polarization are quoted for the reference frame where they are larger.

In the dielectron channel, the ${\rm d}\sigma_{{\rm J/\psi}}/{\rm d}p_{\rm T}$ differential cross section was measured in five $p_{\rm T}$ bins, between 0 and 7 GeV/c. In each bin, the signal was extracted with the same approach used for the integrated invariant mass spectrum. In Fig. 4 the OS invariant mass spectra are shown, together with the LS and the TrkRot backgrounds. The corrections for acceptance and reconstruction efficiency and the systematic errors are given in Table 2. Some of the contributions to the systematic uncertainty do not depend on $p_{\rm T}$, thus affecting only the overall normalization, and they are separately quoted in Table 2. The contributions which depend on $p_{\rm T}$, even when they are correlated bin by bin, were included among the non-correlated systematic errors.

For the analysis in the dimuon channel, a differential study of J/ψ production was performed in the two kinematic variables y and p_T separately. In particular, $d\sigma_{J/\psi}/dp_T$ was studied in seven bins between 0 and 8 GeV/c, and $d\sigma_{J/\psi}/dy$ in five bins between 2.5 and 4. The event sample used for the determination of the differential cross sections is slightly smaller (by about 15%, corresponding to $L_{\rm int}=13.3~{\rm nb}^{-1}$) than the one analyzed for the integrated cross section. This is due to the fact that the statistics in one of the three sub-periods of the data taking is too small to allow a satisfactory fit of the differential invariant mass spectra.

The J/ψ signal was extracted, for each y or p_T bin, with the same fitting technique used for the integrated invariant mass spectra. Since the $\psi(2S)$ yield is rather small and cannot be safely constrained by the data themselves, its contribution was fixed in such a way as to have the same $\psi(2S)/(J/\psi)$ ratio extracted from the integrated spectrum. Anyway, the results of the fit, for what concerns $N_{J/\psi}$, are quite insensitive to the precise level of the $\psi(2S)$ contribution. It has been verified, for example, that fixing for each p_T bin the ratios $\psi(2S)/(J/\psi)$ to the values measured (in the range $p_T > 2$ GeV/c) by CDF [39], $N_{J/\psi}$ varies by less than 1%. In Fig. 5 the OS invariant mass spectra are shown, together with the result of the fits.

The acceptance times efficiency was calculated differentially in y and p_T and the values are reported in Table 2. It can be noted that as a function of p_T , the $A \times \varepsilon$ coverage of the muon spectrometer for J/ψ production extends down to zero p_T , and that the values vary by less than a factor 1.6 in the analyzed p_T range. $A \times \varepsilon$ has a stronger y dependence, but its values are larger than 10% everywhere.

³ The polar angle distribution of the J/ ψ decay leptons is given by dN/d cos θ = 1 + λ cos² θ .




Fig. 4. Invariant mass spectra for OS electron pairs (|y| < 0.9), in bins of p_T . The background calculated using LS and TrkRot approaches are also shown.

Table 2 Summary of the results on the ${\rm J}/\psi$ differential cross sections.

p_{T} (GeV/ c)	$N_{ m J/\psi}$	$A \times \varepsilon$	$\mathrm{d}^2\sigma_{\mathrm{J/\psi}}/\mathrm{d}p_\mathrm{T}\mathrm{d}y \ (\mu\mathrm{b}/(\mathrm{GeV}/c))$	Systematic errors				
				Correl. (μb/(GeV/c))	Non-correl. (μb/(GeV/c))	Polariz., CS (μb/(GeV/c))	Polariz., HE (μb/(GeV/c))	
y < 0.9								
[0; 1]	50 ± 17	0.141	0.59 ± 0.21	0.02	0.18	+0.14, -0.16	+0.07, -0.10	
[1; 2]	86 ± 17	0.088	1.62 ± 0.32	0.06	0.27	+0.36, -0.43	+0.24, -0.34	
[2; 3]	79 ± 13	0.080	1.64 ± 0.27	0.07	0.20	+0.29, -0.37	+0.30, -0.38	
[3; 5]	75 ± 13	0.099	0.62 ± 0.11	0.02	0.08	+0.05, -0.07	+0.14, -0.11	
[5; 7]	50 ± 9	0.120	$\boldsymbol{0.35 \pm 0.06}$	0.01	0.04	+0.001, -0.004	+0.05, -0.07	
2.5 < y < 4								
[0; 1]	229 ± 29	0.280	0.67 ± 0.08	0.06	0.05	+0.14, -0.21	+0.13, -0.19	
[1; 2]	453 ± 40	0.287	$\boldsymbol{1.30 \pm 0.11}$	0.12	0.10	+0.31, -0.35	+0.19, -0.29	
[2; 3]	324 ± 26	0.289	0.92 ± 0.07	0.09	0.07	+0.17, -0.26	+0.09, -0.19	
[3; 4]	253 ± 21	0.312	0.67 ± 0.06	0.06	0.05	+0.12, -0.18	+0.07, -0.11	
[4; 5]	120 ± 17	0.359	$\boldsymbol{0.28 \pm 0.04}$	0.03	0.02	+0.04, -0.05	+0.02, -0.03	
[5; 6]	86 ± 12	0.392	$\boldsymbol{0.18 \pm 0.02}$	0.02	0.01	+0.01, -0.03	+0.01, -0.02	
[6; 8]	80 ± 12	0.452	$\boldsymbol{0.07 \pm 0.01}$	0.01	0.01	+0.007, -0.003	+0.007, -0.008	
y			$d\sigma_{I/\psi}/dy$ (µb)	(µb)	(μb)	(μb)	(μb)	
[-0.9; 0.9]	352 ± 32	0.098	5.97 ± 0.54	0.24	0.83	+0.8, -1.1	+0.9, -1.3	
[2.5; 2.8]	272 ± 28	0.117	5.12 ± 0.77	0.49	0.38	+1.29, -1.62	+0.94, -1.29	
[2.8; 3.1]	326 ± 30	0.383	4.34 ± 0.34	0.41	0.32	+0.97, -1.06	+0.85, -0.99	
[3.1; 3.4]	409 ± 32	0.469	4.64 ± 0.35	0.44	0.35	+0.53, -0.92	+0.46, -0.87	
[3.4; 3.7]	271 ± 26	0.417	3.59 ± 0.30	0.34	0.27	+0.57, -0.77	+0.22, -0.53	
[3.7; 4.0]	172 ± 23	0.215	3.05 ± 0.40	0.29	0.23	+0.67, -1.01	+0.09, -0.46	

pp √s=7 TeV




Fig. 5. Invariant mass spectra for OS muon pairs (2.5 < y < 4), in bins of p_T . The results of the fits are also shown.

dσ_{J/ψ} /dy (μb)

8





Fig. 6. $d^2\sigma_{J/\psi}/dp_T\,dy$ for the midrapidity range and for the forward rapidity data, compared with results from the other LHC experiments [15–17], obtained in similar rapidity ranges. The error bars represent the quadratic sum of the statistical and systematic errors, while the systematic uncertainties on luminosity are shown as boxes. The symbols are plotted at the center of each bin.



ALICE, e⁺e

ALICE, μ⁺μ⁻ CMS

LHCb

Fig. 7. $d\sigma_{J/\psi}/dy$, compared with results from the other LHC experiments [15–17]. The error bars represent the quadratic sum of the statistical and systematic errors, while the systematic uncertainties on luminosity are shown as boxes. The symbols are plotted at the center of each bin.

The differential cross sections are then calculated with the same approach used for the integrated cross section, normalizing $N_{J/\psi}^{\rm cor}(y)$ and $N_{J/\psi}^{\rm cor}(p_{\rm T})$ to the integrated luminosity. The differential cross sections are affected by the same systematic error sources discussed in the previous section. All except the one related to the signal extraction can be considered as common, or strongly correlated. Table 2 gives a summary of the results, including the various

sources of systematic uncertainties (correlated, uncorrelated and polarization-related).

The results are presented in Fig. 6 and Fig. 7 for the $p_{\rm T}$ -differential cross section ${\rm d}^2\sigma_{{\rm J/\psi}}/{\rm d}p_{\rm T}\,{\rm d}y$ and ${\rm d}\sigma_{{\rm J/\psi}}/{\rm d}y$ ($p_{\rm T}>0$), respectively. For the rapidity distribution, the values obtained in the forward region were also reflected with respect to y=0. In both figures, the symbols are plotted at the center of each bin. The statistical and systematic errors were added in quadrature, apart

from the 4 (5.5)% systematic uncertainties on luminosity for the dielectron (dimuon) channels, shown as boxes. The differential cross sections shown in Figs. 6 and 7 assume unpolarized J/ψ production. Systematic uncertainties due to the unknown I/ψ polarization are not shown. Our results are compared with those by the CMS [15], LHCb [16] and ATLAS [17] Collaborations. Also for these data the uncertainty due to luminosity, which is 11% for CMS, 10% for LHCb and 3.4% for ATLAS, is shown separately (boxes), while the error bars contain the statistical and the other sources of systematic errors added in quadrature. Our measurement at central rapidity reaches $p_T = 0$ and is therefore complementary to the data of CMS, available at |y| < 1.2 for $p_T > 6.5$ GeV/c, and AT-LAS, which covers the region |y| < 0.75, $p_T > 7$ GeV/c. In order to compare our $d^2\sigma_{I/\psi}/dp_T dy$ in the forward rapidity range with that of LHCb, we added the LHCb data for prompt and non-prompt production and integrated in the range 2.5 < y < 4 to match our measurement. The agreement between the two data sets is good.

In Fig. 7 our results are compared with the corresponding values from the CMS and LHCb experiments, for the rapidity bins where the $p_{\rm T}$ coverage extends down to zero (ATLAS has no coverage down to $p_{\rm T}=0$ in any rapidity range). For CMS, the value for 1.6 < |y| < 2.4 was obtained by integrating the published ${\rm d}^2\sigma_{{\rm J/\psi}}/{\rm d}p_{\rm T}{\rm d}y$ data [15], while for LHCb the published ${\rm d}\sigma_{{\rm J/\psi}}/{\rm d}y$ for prompt and non-prompt production [16] were added. Our data, together with that of the other LHC experiments, constitute a comprehensive measurement of inclusive ${\rm J/\psi}$ production cross section as a function of rapidity. At the LHC, the inclusive ${\rm J/\psi}$ production cross section at central rapidity is almost twice larger than at Tevatron ($\sqrt{s}=1.96$ TeV) [1] and about ten times larger than at RHIC ($\sqrt{s}=0.2$ TeV) [5]. The width (FWHM) of the rapidity distribution derived from our data is about twice larger than at RHIC [5].

We stress that the results described in this Letter refer to inclusive I/ψ production. Therefore the measured yield is a superposition of a direct component and of I/ψ coming from the decay of higher-mass charmonium states, in particular the χ_{c1} , χ_{c2} and $\psi(2S)$ states. These contributions were measured in lower-energy experiments and were found to be $\sim 25\%$ ($\chi_{c1} + \chi_{c2}$) and $\sim 8\%$ $(\psi(2S))$ of the total measured J/ ψ yield [40,41]. The χ_{c0} contribution is negligible since its B.R. into J/ψ is of the order of 1%. In addition to this "prompt" production, decays of beauty hadrons are also known to give a sizeable contribution (of the order of 10-15% in the p_T range accessed by ALICE [16]) to the observed J/ψ yield. With future high-statistics data samples, the ALICE experiment will identify, at central rapidities, J/ψ from b-decays, via the measurement of the pseudo-proper decay time distributions [42], and will also reconstruct the $\chi_c \rightarrow J/\psi + \gamma$ decay [43]. At forward rapidity, the contribution from b-decays will be estimated from the beauty cross section measurement carried out in the semi-leptonic decay channel [44].

7. Conclusions

The ALICE experiment has measured inclusive J/ ψ production in the rapidity ranges |y|<0.9 and 2.5< y<4, through the decays J/ $\psi\to e^+e^-$ and J/ $\psi\to \mu^+\mu^-$, respectively. The p_T -integrated cross sections, based on data samples corresponding to integrated luminosities $L_{\rm int}=5.6~{\rm nb}^{-1}$ (for the J/ $\psi\to e^+e^-$ channel) and $L_{\rm int}=15.6~{\rm nb}^{-1}$ (for J/ $\psi\to \mu^+\mu^-$) are $\sigma_{\rm J/\psi}(|y|<0.9)=10.7\pm1.0$ (stat.) ±1.6 (syst.) ±1.6 (syst.) ±1.6 ($\lambda_{\rm HE}=1$) ±1.6 (syst.) \pm

LHC, for its coverage of both central and forward rapidities and for the lowest p_T reach at y = 0.

Acknowledgements

The ALICE Collaboration would like to thank all its engineers and technicians for their invaluable contributions to the construction of the experiment and the CERN accelerator teams for the outstanding performance of the LHC complex. The ALICE Collaboration acknowledges the following funding agencies for their support in building and running the ALICE detector: Calouste Gulbenkian Foundation from Lisbon and Swiss Fonds Kidagan, Armenia; Conselho Nacional de Desenvolvimento Científico e Tecnológico (CNPq), Financiadora de Estudos e Projetos (FINEP), Fundação de Amparo à Pesquisa do Estado de São Paulo (FAPESP); National Natural Science Foundation of China (NSFC), the Chinese Ministry of Education (CMOE) and the Ministry of Science and Technology of China (MSTC); Ministry of Education and Youth of the Czech Republic; Danish Natural Science Research Council, the Carlsberg Foundation and the Danish National Research Foundation; The European Research Council under the European Community's Seventh Framework Programme; Helsinki Institute of Physics and the Academy of Finland; French CNRS-IN2P3, the 'Region Pays de Loire', 'Region Alsace', 'Region Auvergne' and CEA, France; German BMBF and the Helmholtz Association; Greek Ministry of Research and Technology; Hungarian OTKA and National Office for Research and Technology (NKTH); Department of Atomic Energy and Department of Science and Technology of the Government of India; Istituto Nazionale di Fisica Nucleare (INFN) of Italy; MEXT Grant-in-Aid for Specially Promoted Research, Japan; Joint Institute for Nuclear Research, Dubna; National Research Foundation of Korea (NRF); CONACYT, DGAPA, México, ALFA-EC and the HELEN Program (High-Energy physics Latin-American-European Network); Stichting voor Fundamenteel Onderzoek der Materie (FOM) and the Nederlandse Organisatie voor Wetenschappelijk Onderzoek (NWO), Netherlands; Research Council of Norway (NFR); Polish Ministry of Science and Higher Education: National Authority for Scientific Research - NASR (Autoritatea Națională pentru Cercetare Științifică -ANCS); Federal Agency of Science of the Ministry of Education and Science of Russian Federation, International Science and Technology Center, Russian Academy of Sciences, Russian Federal Agency of Atomic Energy, Russian Federal Agency for Science and Innovations and CERN-INTAS; Ministry of Education of Slovakia; CIEMAT, EELA, Ministerio de Educación y Ciencia of Spain, Xunta de Galicia (Consellería de Educación), CEADEN, Cubaenergía, Cuba, and IAEA (International Atomic Energy Agency); The Ministry of Science and Technology and the National Research Foundation (NRF), South Africa; Swedish Research Council (VR) and Knut & Alice Wallenberg Foundation (KAW); Ukraine Ministry of Education and Science; United Kingdom Science and Technology Facilities Council (STFC); The United States Department of Energy, the United States National Science Foundation, the State of Texas, and the State of Ohio.

Open access

This article is published Open Access at sciencedirect.com. It is distributed under the terms of the Creative Commons Attribution License 3.0, which permits unrestricted use, distribution, and reproduction in any medium, provided the original authors and source are credited.

References

[1] D. Acosta, et al., CDF Collaboration, Phys. Rev. D 71 (2005) 032001.

- [2] A. Abulencia, et al., CDF Collaboration, Phys. Rev. Lett. 99 (2007) 132001.
- [3] S. Abachi, et al., D0 Collaboration, Phys. Lett. B 370 (1996) 239.
- [4] B. Abbott, et al., D0 Collaboration, Phys. Rev. Lett. 82 (1999) 35.
- [5] A. Adare, et al., PHENIX Collaboration, Phys. Rev. Lett. 98 (2007) 232002.
- [6] N. Brambilla, et al., Eur. Phys. J. C 71 (2011) 1534.
- [7] J.P. Lansberg, Eur. Phys. J. C 61 (2009) 693.
- [8] J. Campbell, F. Maltoni, F. Tramontano, Phys. Rev. Lett. 98 (2007) 252002.
- [9] B. Gong, J.-X. Wang, Phys. Rev. Lett. 100 (2008) 232001.
- [10] B. Gong, X.Q. Li, J.-X. Wang, Phys. Lett. B 673 (2009) 197.
- [11] S. Brodsky, J.P. Lansberg, Phys. Rev. D 81 (2010) 051502.
- [12] M. Butenschön, B.A. Kniehl, Phys. Rev. Lett. 106 (2011) 022003.
- [13] K. Aamodt, et al., ALICE Collaboration, JINST 3 (2008) S08002.[14] K. Aamodt, et al., ALICE Collaboration, Phys. Lett. B 693 (2010) 53.
- [15] V. Khachatryan, et al., CMS Collaboration, Eur. Phys. J. C 71 (2011)
- [16] R. Aaij, et al., LHCb Collaboration, Eur. Phys. J. C 71 (2011) 1645.
- [17] G. Aad, et al., ATLAS Collaboration, Nucl. Phys. B 850 (2011) 387.
- [18] K. Aamodt, et al., ALICE Collaboration, JINST 5 (2010) P03003.
- [19] J. Alme, et al., ALICE Collaboration, Nucl. Inst. Meth. A 622 (2010) 316.
- [20] K. Aamodt, et al., ALICE Collaboration, Eur. Phys. J. C 71 (2011) 1655.
- [21] A. Akindinov, et al., Nucl. Inst. Meth. A 615 (2010) 37.
- [22] M.J. Kweon, et al., ALICE Collaboration, Nucl. Phys. A 830 (2009) 535c.
- [23] U. Abeysekara, et al., ALICE EMCAL Physics Performance Report arXiv:1008.0413.
- [24] K. Aamodt, et al., ALICE Collaboration, Eur. Phys. J. C 65 (2010) 111.
- [25] F. Guerin, et al., ALICE Internal Note ALICE-INT-2006-002, https://edms.cern.ch/document/713307/1.
- [26] K. Koch, et al., ALICE Collaboration, Nucl. Phys. A 855 (2011) 281.

- [27] I.E. Gaiser, Ph.D. thesis, SLAC-R-255, 1982.
- [28] V. Blobel, C. Kleinwort, arXiv:hep-ex/0208022.
- [29] L. Aphecetche, et al., ALICE Internal Note ALICE-INT-2009-044, https://edms. cern.ch/document/1054937/1.
- [30] G. Chabratova, et al., ALICE Internal Note ALICE-INT-2003-002, https://edms. cern.ch/document/371480/1.
- [31] K. Nakamura, et al., Particle Data Group, J. Phys. G 37 (2010) 075021.
- [32] D. Stocco, et al., ALICE Internal Note ALICE-INT-2006-029, https://edms.cern. ch/document/803009/1.
- [33] M. Gagliardi, et al., ALICE Collaboration, in: Proceedings of the International Workshop on Early Physics with Heavy-Ion Collisions at LHC, July 6th-8th, 2011, Bari (Italy);
 - K. Aamodt, et al., ALICE Collaboration, Measurement of inelastic, single and double diffraction cross sections in proton–proton collisions at LHC with ALICE, in preparation.
- [34] S. van der Meer, ISR-PO/68-31, KEK68-64.
- [35] R. Shahoyan, Ph.D. thesis, IST, Lisbon, 2001.
- [36] S. Grigoryan, A. DeFalco, ALICE Internal Note ALICE-INT-2008-016, https://edms.cern.ch/document/938944/1.
- [37] G. Anders, et al. CERN-ATS-Note-2011-016 PERF.
- [38] L. Aphecetche, et al., ALICE Scientific Note 2011-001.
- [39] T. Aaltonen, et al., CDF Collaboration, Phys. Rev. D 80 (2009) 031103.
- [40] P. Faccioli, et al., JHEP 0810 (2008) 004.
- [41] A. Adare, et al., PHENIX Collaboration, arXiv:1105.1966.
- [42] G.E. Bruno, et al. ALICE Collaboration, in: Proceedings of the 35th International Conference on High Energy Physics, PoS (ICHEP2010), p. 194.
- [43] P. Gonzalez, et al., ALICE Collaboration, Eur. Phys. J. C 61 (2009) 899.
- [44] D. Stocco, et al., ALICE Collaboration, Nucl. Phys. A 855 (2011) 323.

ALICE Collaboration

A.G. Agocs ⁷, A. Agostinelli ⁸, S. Aguilar Salazar ⁹, Z. Ahammed ¹⁰, N. Ahmad ¹¹, A. Ahmad Masoodi ¹¹, S.U. Ahn ^{12,1}, A. Akindinov ¹³, D. Aleksandrov ¹⁴, B. Alessandro ¹⁵, R. Alfaro Molina ⁹, A. Alici ¹⁶, A. Alkin ¹⁷, E. Almaráz Aviña ⁹, J. Alme ¹⁸, T. Alt ¹⁹, V. Altini ^{20,ii}, I. Altsybeev ²¹, C. Andrei ²², A. Andronic ²³, V. Anguelov ^{19,iii}, C. Anson ²⁴, T. Antičić ²⁵, F. Antinori ²⁶, P. Antonioli ²⁷, L. Aphecetche ²⁸, H. Appelshäuser ²⁹, N. Arbor ³⁰, S. Arcelli ⁸, A. Arend ²⁹, N. Armesto ³¹, R. Arnaldi ¹⁵, T. Aronsson ⁴, I.C. Arsene ²³, A. Asryan ²¹, A. Augustinus ⁶, R. Averbeck ²³, T.C. Awes ³², J. Äystö ³³, M.D. Azmi ¹¹, M. Bach ¹⁹, A. Badalà ³⁴, Y.W. Baek ^{12,i}, R. Bailhache ²⁹, R. Bala ¹⁵, R. Baldini Ferroli ¹⁶, A. Baldisseri ³⁵, A. Baldit ³⁶, J. Bán ³⁷, R. Barbera ³⁸, F. Barile ²⁰, G.G. Barnaföldi ⁷, L.S. Barnby ³⁹, V. Barret ³⁶, J. Barrtke ⁴⁰, M. Basile ⁸, N. Bastid ³⁶, B. Bathen ⁴¹, G. Batigne ²⁸, B. Batyunya ⁴², C. Baumann ²⁹, I.G. Bearden ⁴³, H. Beck ²⁹, I. Belikov ⁴⁴, F. Bellini ⁸, R. Bellwied ⁴⁵, E. Belmont-Moreno ⁹, S. Beole ⁴⁶, I. Berceanu ²², A. Bercuci ²², E. Berdermann ²³, Y. Berdnikov ⁴⁷, C. Bergmann ⁴¹, L. Betev ⁶, A. Bhasin ⁴⁸, A.K. Bhati ⁵, L. Bianchi ⁴⁶, N. Bianchi ⁴⁶, N. Bianchi ⁴⁶, N. Bianchi ⁴⁶, N. Bonchi ⁴⁷, D. Blau ⁴, C. Blume ²⁹, N. Bock ²⁴, A. Bogdanov ⁵⁴, H. Bøggild ⁴³, M. Bogolyubsky ⁵⁵, L. Boldizsár ⁷, M. Bombara ⁵⁶, C. Bombonati ⁵⁰, J. Book ²⁹, H. Borel ³⁵, A. Borissov ⁵⁷, C. Bortolin ^{50,iv}, S. Bose ⁵⁸, F. Bossú ^{6,v}, M. Botje ⁵², S. Böttger ⁵⁹, B. Boyer ⁶⁰, P. Braun-Munzinger ²³, L. Bravina ⁶¹, M. Bregant ²⁸, T. Breitner ⁵⁹, M. Broz ⁶², R. Brun ⁶, E. Bruna ⁴, G.E. Bruno ²⁰, D. Budnikov ⁶³, H. Buesching ²⁹, S. Bufalino ⁴⁶, O. Busch ⁶⁴, Z. Buthelezi ⁶⁵, D. Caffarrii ⁵⁰, X. Cai ⁶⁶, H. Caines ⁴, E. Calvo Villar ⁵⁷, P. Camerini ⁶⁸, V. Canoa Roman ^{69,ii}

K. Aamodt¹, A. Abrahantes Quintana², D. Adamová³, A.M. Adare⁴, M.M. Aggarwal⁵, G. Aglieri Rinella⁶,

Sanchez ⁶, H. Delagrange ²⁸, Y. Delgado Mercado ⁶⁷, G. Dellacasa ^{78,vii}, A. Deloff ⁸⁶, V. Demanov ⁶³, E. Dénes ⁷, A. Deppman ⁸², D. Di Bari ²⁰, C. Di Giglio ²⁰, S. Di Liberto ⁸⁷, A. Di Mauro ⁶, P. Di Nezza ⁴⁹, E. Dénes ⁷, A. Deppman ⁸², D. Di Bari ²⁰, C. Di Giglio ²⁰, S. Di Liberto ⁸⁷, A. Di Mauro ⁶, P. Di Nezza ⁴⁹, T. Dietel ⁴¹, R. Divià ⁶, Ø. Djuvsland ¹, A. Dobrin ⁵⁷, T. Dobrowolski ⁸⁶, I. Domínguez ⁷⁹, B. Dönigus ²³, O. Dordic ⁶¹, O. Driga ²⁸, A.K. Dubey ¹⁰, L. Ducroux ⁷¹, P. Dupieux ³⁶, A.K. Dutta Majumdar ⁵⁸, M.R. Dutta Majumdar ¹⁰, D. Elia ⁸⁴, D. Emschermann ⁴¹, H. Engel ⁵⁹, H.A. Erdal ¹⁸, B. Espagnon ⁶⁰, M. Estienne ²⁸, S. Esumi ⁷⁵, D. Evans ³⁹, S. Evrard ⁶, G. Eyyubova ⁶¹, D. Fabris ²⁶, J. Faivre ³⁰, D. Falchieri ⁸, A. Fantoni ⁴⁹, M. Fasel ²³, R. Fearick ⁶⁵, A. Fedunov ⁴², D. Fehlker ¹, V. Fekete ⁶², D. Felea ⁸⁰, G. Feofilov ²¹, A. Fernández Téllez ⁷⁷, E.G. Ferreiro ³¹, A. Ferretti ⁴⁶, R. Ferretti ⁷⁸, M.A.S. Figueredo ⁸², S. Filchagin ⁶³, R. Fini ⁸⁴, D. Finogeev ⁸⁸, F.M. Fionda ²⁰, E.M. Fiore ²⁰, M. Floris ⁶, S. Foertsch ⁶⁵, P. Foka ²³, S. Fokin ¹⁴, E. Fragiacomo ⁸⁹, M. Fragkiadakis ⁹⁰, U. Frankenfeld ²³, U. Fuchs ⁶, F. Furano ⁶, C. Furget ³⁰, M. Fusco Girard ⁸³, J.J. Gaardhøje ⁴³, S. Gadrat ³⁰, M. Gagliardi ⁴⁶, A. Gago ⁶⁷, M. Gallio ⁴⁶, P. Ganoti ³², C. Garabatos ²³, F. Garcia-Solis ⁹¹, R. Gemme ⁷⁸, I. Gerhard ¹⁹, M. Germain ²⁸, C. Geuna ³⁵, A. Gheata ⁶ C. Garabatos ²³, E. Garcia-Solis ⁹¹, R. Gemme ⁷⁸, J. Gerhard ¹⁹, M. Germain ²⁸, C. Geuna ³⁵, A. Gheata ⁶, M. Gheata ⁶, B. Ghidini ²⁰, P. Ghosh ¹⁰, P. Gianotti ⁴⁹, M.R. Girard ⁹², P. Giubellino ⁴⁶, ^{viii}, E. Gladysz-Dziadus ⁴⁰, P. Glässel ⁶⁴, R. Gomez ⁹³, L.H. González-Trueba ⁹, P. González-Zamora ⁵³, E. Gladysz-Dziadus ⁴⁰, P. Glässel ⁶⁴, R. Gomez ⁹³, L.H. González-Trueba ⁹, P. González-Zamora ³³, S. Gorbunov ¹⁹, S. Gotovac ⁹⁴, V. Grabski ⁹, L.K. Graczykowski ⁹², R. Grajcarek ⁶⁴, A. Grelli ⁷³, A. Grigoras ⁶, C. Grigoras ⁶, V. Grigoriev ⁵⁴, A. Grigoryan ⁹⁵, S. Grigoryan ⁴², B. Grinyov ¹⁷, N. Grion ⁸⁹, P. Gros ⁷⁴, J.F. Grosse-Oetringhaus ⁶, J.-Y. Grossiord ⁷¹, F. Guber ⁸⁸, R. Guernane ³⁰, C. Guerra Gutierrez ⁶⁷, B. Guerzoni ⁸, K. Gulbrandsen ⁴³, H. Gulkanyan ⁹⁵, T. Gunji ⁹⁶, A. Gupta ⁴⁸, R. Gupta ⁴⁸, H. Gutbrod ²³, Ø. Haaland ¹, C. Hadjidakis ⁶⁰, M. Haiduc ⁸⁰, H. Hamagaki ⁹⁶, G. Hamar ⁷, L.D. Hanratty ³⁹, Z. Harmanova ⁵⁶, J.W. Harris ⁴, M. Hartig ²⁹, D. Hasegan ⁸⁰, D. Hatzifotiadou ²⁷, A. Hayrapetyan ^{95,ii}, M. Heide ⁴¹, M. Heinz ⁴, H. Helstrup ¹⁸, A. Herghelegiu ²², G. Herrera Corral ⁶⁹, N. Herrmann ⁶⁴, K.F. Hetland ¹⁸, B. Hicks ⁴, P.T. Hille ⁴, B. Hippolyte ⁴⁴, T. Horaguchi ⁷⁵, Y. Hori ⁹⁶, P. Hristov ⁶, L. Hřivnáčová ⁶⁰, M. Huang ¹, S. Huber ²³, T.I. Humanic ²⁴, D.S. Hwang ⁹⁷, R. Ilkaev ⁶³, L. Ilkiv ⁸⁶ M. Heide ⁴¹, M. Heinz ⁴, H. Helstrup ¹⁸, Å. Herghelegiu ²², G. Herrera Corral ⁶⁹, N. Hermann ⁶⁴, K.F. Hetland ¹⁸, B. Hicks ⁴, P.T. Hille ⁴, B. Hippolyte ⁴⁴, T. Horaguchi ⁷⁵, Y. Hori ⁹⁶, P. Hristov ⁶, I. Hřivnáčová ⁶⁰, M. Huang ¹, S. Huber ²³, T.J. Humanic ²⁴, D.S. Hwang ⁹⁷, R. Ilkaev ⁶³, I. Ilkiv ⁸⁶, M. Inaba ⁷⁵, E. Incani ⁸⁵, G.M. Innocenti ⁴⁶, M. Ippolitov ¹⁴, M. Irfan ¹¹, C. Ivan ²³, A. Ivanov ²¹, M. Ivanov ²³, V. Ivanov ⁴⁷, A. Jachołkowski ⁶, P.M. Jacobs ⁹⁸, L. Jancurová ⁴², S. Jangal ⁴⁴, M.A. Janik ⁹², R. Janik ⁶², P.H.S.Y. Jayarathna ⁴⁵, ½, S. Jena ⁹⁹, L. Jirden ⁶, G.T. Jones ³⁹, P.G. Jones ³⁹, P. Jovanović ³⁹, H. Jung ¹², W. Jung ¹², A. Jusko ³⁹, S. Kalcher ¹⁹, P. Kaliiñák ³⁷, M. Kalisky ⁴¹, T. Kalliokoski ³³, A. Kalweit ¹⁰⁰, R. Kamermans ^{73, vii}, K. Kanaki ¹, E. Kang ¹², J.H. Kang ¹⁰¹, V. Kaplin ⁵⁴, A. Karasu Uysal ⁶, O. Karavichev ⁸⁸, T. Karavicheva ⁸⁸, E. Karpechev ⁸⁸, A. Kazantsev ¹⁴, U. Kebschull ⁵⁹, R. Keidel ¹⁰², M.M. Khan ¹¹, P. Khan ¹⁸, A. Khanzadeev ⁴⁷, Y. Kharlov ⁵⁵, B. Kileng ¹⁸, D.J. Kim ³³, D.S. Kim ¹², D.W. Kim ¹², J.H. Kim ⁹⁷, J.S. Kim ¹², N. Kim ¹⁰¹, S. Kiisel ⁹, S. Kiselev ¹³, A. Kisiel ⁵, J.L. Klay ¹⁰³, J. Klein ⁶⁴, C. Klein-Bösing ⁴¹, M. Kliemant ²⁹, A. Kluge ⁶, M.L. Knichel ²³, K. Koch ⁶⁴, M.K. Köhler ²³, A. Kolojvari ²¹, V. Kondratiev ²¹, N. Kondratyeva ⁵⁴, A. Konevskih ⁸⁸, E. Kornaś ⁴⁰, C. Kottachchi Kankanamge Don ⁵⁷, R. Kramer ²⁹, I. Kraus ²³, T. Krawutschke ^{64,xi}, M. Kretz ¹⁹, M. Krivda ^{39,xii}, F. Krizek ³³, M. Krus ⁵¹, F. Kramer ²⁹, I. Kraus ²³, T. Krawutschke ^{64,xi}, M. Kretz ¹⁹, M. Krivda ^{39,xii}, F. Krizek ³³, M. Krus ⁵¹, P. Ladrón de Guevara ⁷⁹, V. Lafage ⁶⁰, I. Lakomov ²¹, C. Kuhn ⁴⁴, P. G. Kuijer ⁵², P. Kurashvili ⁸⁶, A. Kurepin ⁸⁸, A. B. Kurepin ⁸⁸, A. Kuryakin ⁶³, S. Kushpil ³, V. Kushpil ³, M.J. Kweon ⁶⁴, Y. Kwon ¹⁰¹, P. La Rocca ³⁸, P. Ladr D. Miśkowiec ^{6,xv}, C. Mitu ⁸⁰, J. Mlynarz ⁵⁷, B. Mohanty ¹⁰, L. Molnar ⁶, L. Montaño Zetina ⁶⁹,

M. Monteno ¹⁵, E. Montes ⁵³, M. Morando ⁵⁰, D.A. Moreira De Godoy ⁸², S. Moretto ⁵⁰, A. Morsch ⁶, V. Muccifora ⁴⁹, E. Mudnic ⁹⁴, H. Müller ⁶, S. Muhuri ¹⁰, M.G. Munhoz ⁸², L. Musa ⁶, A. Musso ¹⁵, B.K. Nandi ⁹⁹, R. Nania ²⁷, E. Nappi ⁸⁴, C. Nattrass ¹⁰⁷, F. Navach ²⁰, S. Navin ³⁹, T.K. Nayak ¹⁰, S. Nazarenko ⁶³, G. Nazarov ⁶³, A. Nedosekin ¹³, F. Nendaz ⁷¹, M. Nicassio ²⁰, B.S. Nielsen ⁴³, S. Nikolaev ¹⁴, V. Nikolic ²⁵, S. Nikulin ¹⁴, V. Nikulin ⁴⁷, B.S. Nilsen ⁷⁰, M.S. Nilsson ⁶¹, F. Noferini ²⁷, G. Nooren ⁷³, N. Novitzky ³³, A. Nyanin ¹⁴, A. Nyatha ⁹⁹, C. Nygaard ⁴³, J. Nystrand ¹, H. Obayashi ¹⁰⁹, A. Ochirov ²¹, H. Oeschler ¹⁰⁰, S.K. Oh ¹², J. Oleniacz ⁹², C. Oppedisano ¹⁵, A. Ortiz Velasquez ⁷⁹, G. Ortona ^{6,v}, H. Oeschler ¹⁰⁰, S.K. Oh ¹², J. Oleniacz ⁹², C. Oppedisano ¹⁵, A. Ortiz Velasquez ⁷⁹, G. Ortona ^{6,v}, A. Oskarsson ⁷⁴, P. Ostrowski ⁹², I. Otterlund ⁷⁴, J. Otwinowski ²³, G. Øvrebekk ¹, K. Oyama ⁶⁴, K. Ozawa ⁹⁶, Y. Pachmayer ⁶⁴, M. Pachr ⁵¹, F. Padilla ⁴⁶, P. Pagano ⁸³, G. Paić ⁷⁹, F. Painke ¹⁹, C. Pajares ³¹, S. Pal ³⁵, S.K. Pal ¹⁰, A. Palaha ³⁹, A. Palmeri ³⁴, G.S. Pappalardo ³⁴, W.J. Park ²³, V. Paticchio ⁸⁴, A. Pavlinov ⁵⁷, T. Pawlak ⁹², T. Peitzmann ⁷³, D. Peresunko ¹⁴, C.E. Pérez Lara ⁵², D. Perini ⁶, D. Perrino ²⁰, W. Peryt ⁹², A. Pesci ²⁷, V. Peskov ^{6,xvi}, Y. Pestov ¹¹⁰, A.J. Peters ⁶, V. Petráček ⁵¹, M. Petran ⁵¹, M. Petris ²², P. Petrov ³⁹, M. Petrovici ²², C. Petta ³⁸, S. Piano ⁸⁹, A. Piccotti ¹⁵, M. Pikna ⁶², P. Pillot ²⁸, O. Pinazza ⁶, L. Pinsky ⁴⁵, N. Pitz ²⁹, F. Piuz ⁶, D.B. Piyarathna ^{57,xvii}, R. Platt ³⁹, M. Płoskoń ⁹⁸, J. Pluta ⁹², T. Pocheptsov ^{42,xviii}, S. Pochybova ⁷, P.L.M. Podesta-Lerma ⁹³, M.G. Poghosyan ⁴⁶, B. Polichtchouk ⁵⁵, A. Pop ²², V. Pospíšil ⁵¹, B. Potukuchi ⁴⁸, S.K. Prasad ⁵⁷, R. Preghenella ¹⁶, F. Prino ¹⁵, C.A. Pruneau ⁵⁷, I. Pshenichnov ⁸⁸, G. Puddu ⁸⁵, A. Pulvirenti ^{38,ii}, V. Punin ⁶³, M. Putiš ⁵⁶, L. Putschke ⁴, F. Ouercigh ⁶, H. Ovigstad ⁶¹ B. Ротикиспі то, S.K. Prasad то, K. Pregnenella то, F. Prino то, C.A. Pruneau то, I. Pshenichnov то, G. Puddu 85, A. Pulvirenti 38,ii, V. Punin 63, M. Putiš 56, J. Putschke 4, E. Quercigh 6, H. Qvigstad 61, A. Rachevski 89, A. Rademakers 6, S. Radomski 64, T.S. Räihä 33, J. Rak 33, A. Rakotozafindrabe 35, L. Ramello 78, A. Ramírez Reyes 69, M. Rammler 41, R. Raniwala 111, S. Raniwala 111, S.S. Räsänen 33, D. Rathee 5, K.F. Read 107, J.S. Real 30, K. Redlich 86,xix, R. Renfordt 29, A.R. Reolon 49, A. Reshetin 88, F. Rettig 19, J.-P. Revol 6, K. Reygers 64, H. Ricaud 100, L. Riccati 15, R.A. Ricci 112, M. Richter 1,xx, P. Riedler 6, W. Riegler 6, F. Riggi 38, M. Rodríguez Cahuantzi 77, D. Rohr 19, D. Röhrich 1, R. Romita 23, F. Ronchetti 49, P. Rosinský 6, P. Rosnet 36, S. Rossegger 6, A. Rossi 50, F. Roukoutakis 90, S. Rousseau 60, C. Roy 44, P. Roy 58, A. L. Rubio Montero 53, R. Rui 68, F. Ryabinkin 14, A. Rubio Montero 53, R. Rui 68, Rui 6 P. Rosinsky F. Rosnet F. Rosnet F. Roukoutakis F. R K. Schweda 64, G. Scioli 8, E. Scomparin 15,*, P.A. Scott 39, R. Scott 107, G. Segato 50, S. Senyukov 78, J. Seo 12, S. Serci 85, E. Serradilla 53, A. Sevcenco 80, I. Sgura 84, G. Shabratova 42, R. Shahoyan 6, N. Sharma 5, S. Sharma 48, K. Shigaki 109, M. Shimomura 75, K. Shtejer 2, Y. Sibiriak 14, M. Siciliano 46, E. Sicking 6, T. Siemiarczuk 86, D. Silvermyr 32, G. Simonetti 6, R. Singaraju 10, R. Singh 48, S. Singha 10, B.C. Sinha 10, T. Sinha 58, B. Sitar 62, M. Sitta 78, T.B. Skaali 61, K. Skjerdal 1, R. Smakal 51, N. Smirnov 4, R. Snellings 73, C. Søgaard 43, R. Soltz 114, H. Son 97, J. Song 115, M. Song 101, C. Soos 6, F. Soramel 50, M. Spyropoulou-Stassinaki 90, B.K. Srivastava 113, J. Stachel 64, I. Stan 80, G. Stefanek 86, G. Stefanini 6, T. Steinbeck 19, M. Steinpreis 24, E. Stenlund 74, G. Steyn 65, D. Stocco 28, R. Stock 29, C.H. Stokkevag 1, M. Stolpovskiy 55, P. Strmen 62, A.A.P. Suaide 82, M.A. Subieta Vásquez 46, T. Sugitate 109, C. Suire 60, M. Sukhorukov 63, M. Šumbera 3, T. Susa 25, D. Swoboda 6, T.J.M. Symons 98, A. Szanto de Toledo 82, L. Szarka 62, A. Szostak 1, C. Tagridis 90, L. Takahashi 72, L.D. Tapia Takaki 60, A. Tauro 6, G. Tejeda Muños I. Szarka ⁶², A. Szostak ¹, C. Tagridis ⁹⁰, J. Takahashi ⁷², J.D. Tapia Takaki ⁶⁰, A. Tauro ⁶, G. Tejeda Muñoz ⁷⁷, A. Telesca ⁶, C. Terrevoli ²⁰, J. Thäder ²³, D. Thomas ⁷³, J.H. Thomas ²³, R. Tieulent ⁷¹, A.R. Timmins ⁴⁵, A. Telesca o, C. Terrevoli 20, J. Thader 23, D. Thomas 73, J.H. Thomas 23, R. Tieulent 71, A.R. Timmins 43, D. Tlusty 51, A. Toia 6, H. Torii 109, F. Tosello 15, T. Traczyk 92, D. Truesdale 24, W.H. Trzaska 33, A. Tumkin 63, R. Turrisi 26, A.J. Turvey 70, T.S. Tveter 61, J. Ulery 29, K. Ullaland 1, A. Uras 85, J. Urbán 56, G.M. Urciuoli 87, G.L. Usai 85, M. Vajzer 51, M. Vala 42,xii, L. Valencia Palomo 60, S. Vallero 64, N. van der Kolk 52, M. van Leeuwen 73, P. Vande Vyvre 6, L. Vannucci 112, A. Vargas 77, R. Varma 99, M. Vasileiou 90, A. Vasiliev 14, V. Vechernin 21, M. Veldhoen 73, M. Venaruzzo 68, E. Vercellin 46, S. Vergara 77, D.C. Vernekohl 41, R. Vernet 116, M. Verweij 73, L. Vickovic 94, G. Viesti 50, O. Vikhlyantsev 63, Z. Vilakazi 65, O. Villalobos Baillie 39, A. Vinogradov 14, L. Vinogradov 21, Y. Vinogradov 63, T. Virgili 83, Y.P. Viyogi 10, A. Vodopvanov 42, K. Valoship 13, S. Veloship 57, G. Velosh A. Vodopyanov ⁴², K. Voloshin ¹³, S. Voloshin ⁵⁷, G. Volpe ²⁰, B. von Haller ⁶, D. Vranic ²³, J. Vrláková ⁵⁶, B. Vulpescu ³⁶, A. Vyushin ⁶³, B. Wagner ¹, V. Wagner ⁵¹, R. Wan ^{44,xxii}, D. Wang ⁶⁶, M. Wang ⁶⁶, Y. Wang ⁶⁴, Y. Wang ⁶⁶, K. Watanabe ⁷⁵, J.P. Wessels ^{41,viii}, U. Westerhoff ⁴¹, J. Wiechula ^{64,xxiii}, J. Wikne ⁶¹, M. Wilde ⁴¹, A. Wilk ⁴¹, G. Wilk ⁸⁶, M.C.S. Williams ²⁷, B. Windelband ⁶⁴, H. Yang ³⁵, S. Yasnopolskiy ¹⁴, J. Yi ¹¹⁵, Z. Yin ⁶⁶, H. Yokoyama ⁷⁵, I.-K. Yoo ¹¹⁵, X. Yuan ⁶⁶, I. Yushmanov ¹⁴, E. Zabrodin ⁶¹, C. Zach ⁵¹,

C. Zampolli ⁶, S. Zaporozhets ⁴², A. Zarochentsev ²¹, P. Závada ¹⁰⁵, N. Zaviyalov ⁶³, H. Zbroszczyk ⁹², P. Zelnicek ^{59,ii}, A. Zenin ⁵⁵, I. Zgura ⁸⁰, M. Zhalov ⁴⁷, X. Zhang ^{66,i}, D. Zhou ⁶⁶, F. Zhou ⁶⁶, Y. Zhou ⁷³, X. Zhu ⁶⁶, A. Zichichi ^{8,xxiv}, G. Zinovjev ¹⁷, Y. Zoccarato ⁷¹, M. Zynovyev ¹⁷

- ¹ Department of Physics and Technology, University of Bergen, Bergen, Norway
- ² Centro de Aplicaciones Tecnológicas y Desarrollo Nuclear (CEADEN), Havana, Cuba
- ³ Nuclear Physics Institute, Academy of Sciences of the Czech Republic, Řež u Prahy, Czech Republic
- ⁴ Yale University, New Haven, CT, United States
- ⁵ Physics Department, Panjab University, Chandigarh, India
- ⁶ European Organization for Nuclear Research (CERN), Geneva, Switzerland
- ⁷ KFKI Research Institute for Particle and Nuclear Physics, Hungarian Academy of Sciences, Budapest, Hungary
- ⁸ Dipartimento di Fisica dell'Università and Sezione INFN, Bologna, Italy
- ⁹ Instituto de Física, Universidad Nacional Autónoma de México, Mexico City, Mexico
- ¹⁰ Variable Energy Cyclotron Centre, Kolkata, India
- ¹¹ Department of Physics, Aligarh Muslim University, Aligarh, India
- ¹² Gangneung-Wonju National University, Gangneung, South Korea
- 13 Institute for Theoretical and Experimental Physics, Moscow, Russia
- ¹⁴ Russian Research Centre Kurchatov Institute, Moscow, Russia
- 15 Sezione INFN, Turin, Italy
- ¹⁶ Centro Fermi Centro Studi e Ricerche e Museo Storico della Fisica "Enrico Fermi", Rome, Italy
- ¹⁷ Bogolyubov Institute for Theoretical Physics, Kiev, Ukraine
- ¹⁸ Faculty of Engineering, Bergen University College, Bergen, Norway
- ¹⁹ Frankfurt Institute for Advanced Studies, Johann Wolfgang Goethe-Universität Frankfurt, Frankfurt, Germany
- ²⁰ Dipartimento Interateneo di Fisica 'M. Merlin' and Sezione INFN, Bari, Italy
- ²¹ V. Fock Institute for Physics, St. Petersburg State University, St. Petersburg, Russia
- ²² National Institute for Physics and Nuclear Engineering, Bucharest, Romania
- ²³ Research Division and ExtreMe Matter Institute EMMI, GSI Helmholtzzentrum für Schwerionenforschung, Darmstadt, Germany
- ²⁴ Department of Physics, Ohio State University, Columbus, OH, United States
- ²⁵ Rudjer Bošković Institute, Zagreb, Croatia
- ²⁶ Sezione INFN, Padova, Italy
- ²⁷ Sezione INFN, Bologna, Italy
- ²⁸ SUBATECH, Ecole des Mines de Nantes, Université de Nantes, CNRS-IN2P3, Nantes, France
- ²⁹ Institut für Kernphysik, Johann Wolfgang Goethe-Universität Frankfurt, Frankfurt, Germany
- 30 Laboratoire de Physique Subatomique et de Cosmologie (LPSC), Université Joseph Fourier, CNRS-IN2P3, Institut Polytechnique de Grenoble, Grenoble, France
- ³¹ Departamento de Física de Partículas and IGFAE, Universidad de Santiago de Compostela, Santiago de Compostela, Spain
- ³² Oak Ridge National Laboratory, Oak Ridge, TN, United States
- ³³ Helsinki Institute of Physics (HIP) and University of Jyväskylä, Jyväskylä, Finland
- ³⁴ Sezione INFN, Catania, Italy
- 35 Commissariat à l'Energie Atomique, IRFU, Saclay, France
- 36 Laboratoire de Physique Corpusculaire (LPC), Clermont Université, Université Blaise Pascal, CNRS-IN2P3, Clermont-Ferrand, France
- ³⁷ Institute of Experimental Physics, Slovak Academy of Sciences, Košice, Slovakia
- ³⁸ Dipartimento di Fisica e Astronomia dell'Università and Sezione INFN, Catania, Italy
- ³⁹ School of Physics and Astronomy, University of Birmingham, Birmingham, United Kingdom
- ⁴⁰ The Henryk Niewodniczanski Institute of Nuclear Physics, Polish Academy of Sciences, Cracow, Poland
- ⁴¹ Institut für Kernphysik, Westfälische Wilhelms-Universität Münster, Münster, Germany
- ⁴² Joint Institute for Nuclear Research (JINR), Dubna, Russia
- ⁴³ Niels Bohr Institute, University of Copenhagen, Copenhagen, Denmark
- 44 Institut Pluridisciplinaire Hubert Curien (IPHC), Université de Strasbourg, CNRS-IN2P3, Strasbourg, France
- ⁴⁵ University of Houston, Houston, TX, United States
- ⁴⁶ Dipartimento di Fisica Sperimentale dell'Università and Sezione INFN, Turin, Italy
- ⁴⁷ Petersburg Nuclear Physics Institute, Gatchina, Russia
- ⁴⁸ Physics Department, University of Jammu, Jammu, India
- ⁴⁹ Laboratori Nazionali di Frascati, INFN, Frascati, Italy
- ⁵⁰ Dipartimento di Fisica dell'Università and Sezione INFN, Padova, Italy
- ⁵¹ Faculty of Nuclear Sciences and Physical Engineering, Czech Technical University in Prague, Prague, Czech Republic
- ⁵² Nikhef, National Institute for Subatomic Physics, Amsterdam, Netherlands
- ⁵³ Centro de Investigaciones Energéticas Medioambientales y Tecnológicas (CIEMAT), Madrid, Spain
- 54 Moscow Engineering Physics Institute, Moscow, Russia
- ⁵⁵ Institute for High Energy Physics, Protvino, Russia
- ⁵⁶ Faculty of Science, P.J. Šafárik University, Košice, Slovakia
- ⁵⁷ Wayne State University, Detroit, MI, United States
- ⁵⁸ Saha Institute of Nuclear Physics, Kolkata, India
- ⁵⁹ Kirchhoff-Institut für Physik, Ruprecht-Karls-Universität Heidelberg, Heidelberg, Germany
- 60 Institut de Physique Nucléaire d'Orsay (IPNO), Université Paris-Sud, CNRS-IN2P3, Orsay, France
- 61 Department of Physics, University of Oslo, Oslo, Norway
- ⁶² Faculty of Mathematics, Physics and Informatics, Comenius University, Bratislava, Slovakia
- 63 Russian Federal Nuclear Center (VNIIEF), Sarov, Russia
- ⁶⁴ Physikalisches Institut, Ruprecht-Karls-Universität Heidelberg, Heidelberg, Germany
- ⁶⁵ Physics Department, University of Cape Town, iThemba LABS, Cape Town, South Africa
- ⁶⁶ Hua-Zhong Normal University, Wuhan, China
- 67 Sección Física, Departamento de Ciencias, Pontificia Universidad Católica del Perú, Lima, Peru
- 68 Dipartimento di Fisica dell'Università and Sezione INFN, Trieste, Italy
- ⁶⁹ Centro de Investigación y de Estudios Avanzados (CINVESTAV), Mexico City and Mérida, Mexico
- 70 Physics Department, Creighton University, Omaha, NE, United States
- ⁷¹ Université de Lyon, Université Lyon 1, CNRS/IN2P3, IPN-Lyon, Villeurbanne, France
- ⁷² Universidade Estadual de Campinas (UNICAMP), Campinas, Brazil
- 73 Nikhef, National Institute for Subatomic Physics and Institute for Subatomic Physics of Utrecht University, Utrecht, Netherlands
- ⁷⁴ Division of Experimental High Energy Physics, University of Lund, Lund, Sweden

- ⁷⁵ University of Tsukuba, Tsukuba, Japan
- ⁷⁶ Sezione INFN, Cagliari, Italy
- 77 Benemérita Universidad Autónoma de Puebla, Puebla, Mexico
- ⁷⁸ Dipartimento di Scienze e Tecnologie Avanzate dell'Università del Piemonte Orientale and Gruppo Collegato INFN, Alessandria, Italy
- ⁷⁹ Instituto de Ciencias Nucleares, Universidad Nacional Autónoma de México, Mexico City, Mexico
- ⁸⁰ Institute of Space Sciences (ISS), Bucharest, Romania
- ⁸¹ Institute of Physics, Bhubaneswar, India
- 82 Universidade de São Paulo (USP), São Paulo, Brazil
- 83 Dipartimento di Fisica 'E.R. Caianiello' dell'Università and Gruppo Collegato INFN, Salerno, Italy
- 84 Sezione INFN. Bari. Italy
- ⁸⁵ Dipartimento di Fisica dell'Università and Sezione INFN, Cagliari, Italy
- 86 Soltan Institute for Nuclear Studies, Warsaw, Poland
- 87 Sezione INFN, Rome, Italy
- 88 Institute for Nuclear Research, Academy of Sciences, Moscow, Russia
- 89 Sezione INFN, Trieste, Italy
- ⁹⁰ Physics Department, University of Athens, Athens, Greece
- ⁹¹ Chicago State University, Chicago, IL, United States
- 92 Warsaw University of Technology, Warsaw, Poland
- ⁹³ Universidad Autónoma de Sinaloa, Culiacán, Mexico
- ⁹⁴ Technical University of Split FESB, Split, Croatia
- ⁹⁵ Yerevan Physics Institute, Yerevan, Armenia
- ⁹⁶ University of Tokyo, Tokyo, Japan
- ⁹⁷ Department of Physics, Sejong University, Seoul, South Korea
- ⁹⁸ Lawrence Berkeley National Laboratory, Berkeley, CA, United States
- 99 Indian Institute of Technology, Mumbai, India
- ¹⁰⁰ Institut für Kernphysik, Technische Universität Darmstadt, Darmstadt, Germany
- ¹⁰¹ Yonsei University, Seoul, South Korea
- ¹⁰² Zentrum für Technologietransfer und Telekommunikation (ZTT), Fachhochschule Worms, Worms, Germany
- 103 California Polytechnic State University, San Luis Obispo, CA, United States
- ¹⁰⁴ China Institute of Atomic Energy, Beijing, China
- 105 Institute of Physics, Academy of Sciences of the Czech Republic, Prague, Czech Republic
- 106 The University of Texas at Austin, Physics Department, Austin, TX, United States
- ¹⁰⁷ University of Tennessee, Knoxville, TN, United States
- ¹⁰⁸ Dipartimento di Fisica dell'Università 'La Sapienza' and Sezione INFN, Rome, Italy
- ¹⁰⁹ Hiroshima University, Hiroshima, Japan
- ¹¹⁰ Budker Institute for Nuclear Physics, Novosibirsk, Russia
- 111 Physics Department, University of Rajasthan, Jaipur, India
- 112 Laboratori Nazionali di Legnaro, INFN, Legnaro, Italy
- ¹¹³ Purdue University, West Lafayette, IN, United States
- ¹¹⁴ Lawrence Livermore National Laboratory, Livermore, CA, United States
- ¹¹⁵ Pusan National University, Pusan, South Korea
- 116 Centre de Calcul de l'IN2P3, Villeurbanne, France
- * Corresponding author.
- E-mail address: scomparin@to.infn.it (E. Scomparin).
- i Also at Laboratoire de Physique Corpusculaire (LPC), Clermont Université, Université Blaise Pascal, CNRS-IN2P3, Clermont-Ferrand, France.
- ii Also at European Organization for Nuclear Research (CERN), Geneva, Switzerland.
- iii Now at Physikalisches Institut, Ruprecht-Karls-Universität Heidelberg, Heidelberg, Germany.
- iv Also at Dipartimento di Fisica dell'Università, Udine, Italy.
- $^{\nu}\,$ Also at Dipartimento di Fisica Sperimentale dell'Università and Sezione INFN, Turin, Italy.
- $^{\it vi}$ Also at Benemérita Universidad Autónoma de Puebla, Puebla, Mexico.
- vii Deceased
- viii Now at European Organization for Nuclear Research (CERN), Geneva, Switzerland.
- ix Also at Wayne State University, Detroit, Michigan, United States.
- x Also at Frankfurt Institute for Advanced Studies, Johann Wolfgang Goethe-Universität Frankfurt, Frankfurt, Germany.
- xi Also at Fachhochschule Köln, Köln, Germany.
- xii Also at Institute of Experimental Physics, Slovak Academy of Sciences, Košice, Slovakia.
- xiii Also at Laboratoire de Physique Subatomique et de Cosmologie (LPSC), Université Joseph Fourier, CNRS-IN2P3, Institut Polytechnique de Grenoble, Grenoble, France.
- xiv Also at "Vinča" Institute of Nuclear Sciences, Belgrade, Serbia.
- xv Also at Research Division and ExtreMe Matter Institute EMMI, GSI Helmholtzzentrum für Schwerionenforschung, Darmstadt, Germany.
- xvi Also at Instituto de Ciencias Nucleares, Universidad Nacional Autónoma de México, Mexico City, Mexico.
- xvii Also at University of Houston, Houston, Texas, United States.
- xviii Also at Department of Physics, University of Oslo, Oslo, Norway.
- xix Also at Institute of Theoretical Physics, University of Wroclaw, Wroclaw, Poland.
- xx Now at Department of Physics, University of Oslo, Oslo, Norway.
- xxi Also at Eberhard Karls Universität Tübingen, Tübingen, Germany.
- xxii Also at Hua-Zhong Normal University, Wuhan, China.
- xxiii Now at Eberhard Karls Universität Tübingen, Tübingen, Germany.
- xxiv Also at Centro Fermi Centro Studi e Ricerche e Museo Storico della Fisica "Enrico Fermi", Rome, Italy.

<u>Update</u>

Physics Letters B

Volume 718, Issue 2, 5 December 2012, Page 692–698

DOI: https://doi.org/10.1016/j.physletb.2012.10.060



Contents lists available at SciVerse ScienceDirect

Physics Letters B

www.elsevier.com/locate/physletb



Erratum

Erratum to "Rapidity and transverse momentum dependence of inclusive J/ ψ production in pp collisions at $\sqrt{s}=7$ TeV" [Phys. Lett. B 704 (5) (2011) 442] $^{\frac{1}{2}}$

ALICE Collaboration

ARTICLE INFO

Article history:
Received 18 October 2012
Accepted 22 October 2012
Available online 2 November 2012
Editor: L. Rolandi

We have identified a bias in the calculation of the cross section of J/ψ production measured in the e^+e^- channel at central rapidity (|y|<0.9) [1]. The acceptance and efficiency corrections were evaluated using a Monte Carlo simulation, based on PYTHIA [2,3], which did not include the radiative decays of J/ψ ($J/\psi \rightarrow e^+e^-\gamma$). In these decays the reconstructed dielectron invariant mass, $m_{e^+e^-}$, is biased towards smaller values than the nominal J/ψ mass, since the photon contribution is neglected. As a consequence, the fraction of signal events in the $m_{e^+e^-}$ range 2.92–3.16 GeV/ c^2 was overestimated by about 10%. Moreover, the requirement of the transverse momentum of the daughter electrons being larger than 1 GeV/c is more selective for radiative than non-radiative J/ψ decays. Therefore, the J/ψ acceptance was also overestimated by about 5%.

We have now evaluated the acceptance and efficiency corrections with a simulation where the decay of the J/ ψ particles is handled by the EvtGen package [4], and where the final state radiation is described using PHOTOS [5,6]. The new acceptance times efficiency value ($A \times \epsilon$) after all analysis cuts is a factor 1.155 smaller than that previously evaluated, independently of $p_{\rm T}$. Neglecting the effect of radiative decays therefore results in underestimating both the $p_{\rm T}$ -integrated and the differential cross sections by 15.5%.

For the dimuon channel, where no invariant mass cut is applied and the occurrence of final state radiation is reduced (by about a factor of three [7]), the differences in the $A \times \epsilon$ values obtained with the new and previous simulations are about 1–2%, well within the systematic uncertainty associated with the signal extraction.

We have further verified that in the dielectron channel the procedure used to derive the $p_{\rm T}$ -differential cross section, which is based on the computation of the $A \times \epsilon$ values, produces a result fully compatible with that obtained by applying an unfolding cor-




Fig. 1. Top panel: invariant mass distributions for opposite-sign (OS) and like-sign (LS) electron pairs (|y| < 0.9, all p_T), as well as for pairs obtained with one track randomly rotated (TrkRot). Bottom panel: the difference of the OS and LS distributions with a fit to the Monte Carlo (MC) signal superimposed.

rection procedure (see, e.g., review [8]), even when considering the radiative decays.

In Fig. 1 the invariant mass distributions of electron pairs are shown. In particular, in the bottom panel the new Monte Carlo line shape is superimposed on the difference of the opposite and like sign distributions. The fraction of the signal within the invariant mass range $2.92-3.16 \text{ GeV}/c^2$ estimated using this Monte Carlo

DOI of original article: http://dx.doi.org/10.1016/j.physletb.2011.09.054.

[©] CERN for the benefit of the ALICE Collaboration.

Table 1 Summary of the results on the J/ψ differential cross sections.

p _T (GeV/c)	$N_{ m J/\psi}$	$A imes \epsilon$	$\mathrm{d}^2\sigma_{\mathrm{J/\psi}}/\mathrm{d}p_{\mathrm{T}}\mathrm{d}y \ (\mu\mathrm{b}/(\mathrm{GeV}/c))$	Systematic uncertainties			
				Correl. (μb/(GeV/c))	Non-correl. (μb/(GeV/c))	Polariz., CS (μb/(GeV/c))	Polariz., HE (μb/(GeV/c))
y < 0.9							
[0; 1]	50 ± 17	0.122	0.68 ± 0.24	0.02	0.21	+0.16, -0.18	+0.08, -0.12
[1; 2]	86 ± 17	0.076	1.87 ± 0.37	0.07	0.31	+0.42, -0.50	+0.28, -0.39
[2; 3]	79 ± 13	0.069	1.89 ± 0.31	0.08	0.23	+0.33, -0.43	+0.35, -0.44
[3; 5]	75 ± 13	0.086	0.72 ± 0.13	0.02	0.09	+0.06, -0.08	+0.16, -0.13
[5; 7]	50 ± 9	0.104	$\textbf{0.40} \pm \textbf{0.07}$	0.01	0.05	+0.001, -0.005	+0.06, -0.08
y			$\mathrm{d}\sigma_{\mathrm{J/\psi}}/\mathrm{d}y$ ($\mu\mathrm{b}$)	(µb)	(µb)	(µb)	(µb)
[-0.9; 0.9]	352 ± 32	0.085	6.90 ± 0.62	0.28	0.96	+0.9, -1.3	+1.0, -1.5




Fig. 2. Double differential J/ψ production cross section as a function of p_T for the midrapidity range and for the forward rapidity data, compared with results from the other LHC experiments [10–12], obtained in similar rapidity ranges. The error bars represent the quadratic sum of the statistical and systematic errors, while the systematic uncertainties on luminosity are shown as boxes. The symbols are plotted at the center of each bin.

is $66.8 \pm 1.9\%$. The main contribution to the uncertainty on this quantity comes from the accuracy of the description of the detector material, as discussed in [1]. A smaller contribution (1%, in terms of the relative error) is attributed to the small discrepancies between the invariant mass distribution as provided by QED at next to leading order [9] and by the event generator (EvtGen + PHOTOS); the latter contribution remains even after taking into account the detector resolution.

The corrected value of the production cross section is $\sigma_{J/\psi}(|y| < 0.9) = 12.4 \pm 1.1 \; (\text{stat.}) \pm 1.8 \; (\text{syst.}) + 1.8 \; (\lambda_{\text{HE}} = 1) - 2.7 \; (\lambda_{\text{HE}} = -1) \; \mu\text{b.}$ In Table 1 the resulting differential cross sections are summarized. Finally, in Figs. 2 and 3 we have updated accordingly the ALICE data points at central rapidity.

Open access

This article is published Open Access at sciencedirect.com. It is distributed under the terms of the Creative Commons Attribu-




Fig. 3. J/ ψ cross section as a function of rapidity, compared with results from the other LHC experiments [10–12]. The error bars represent the quadratic sum of the statistical and systematic errors, while the systematic uncertainties on luminosity are shown as boxes. The symbols are plotted at the center of each bin.

tion License 3.0, which permits unrestricted use, distribution, and reproduction in any medium, provided the original authors and source are credited.

References

- [1] K. Aamodt, et al., ALICE Collaboration, Phys. Lett. B 704 (2011) 442.
- [2] T. Sjöstrand, Comput. Phys. Commun. 82 (1994) 74.
- [3] T. Sjöstrand, S. Mrenna, P. Skands, JHEP 0605 (2006) 026.
- [4] D.J. Lange, Nucl. Instrum. Meth. A 462 (2001) 152.
- [5] E. Barberio, B. van Eijk, Z. Was, Comput. Phys. Commun. 66 (1991) 115.
- [6] E. Barberio, Z. Was, Comput. Phys. Commun. 79 (1994) 291.
- [7] A. Spiridonov, Bremsstrahlung in leptonic onia decays: effects on mass spectra, arXiv:hep-ex/0510076v1, 2005.
- [8] V. Blobel, in: Proceedings of the 1984 CERN School of Computing, CERN 85-09, 1985, pp. 88-127.
- [9] J. Fleischer, F. Jegerlehner, Z. Phys. C 26 (1985) 629.
- [10] V. Khachatryan, et al., CMS Collaboration, Eur. Phys. J. C 71 (2011) 1575.
- [11] R. Aaij, et al., LHCb Collaboration, Eur. Phys. J. C 71 (2011) 1645.
- [12] G. Aad, et al., ATLAS Collaboration, Nucl. Phys. B 850 (2011) 387.

ALICE Collaboration

K. Aamodt ¹, A. Abrahantes Quintana ², D. Adamová ³, A.M. Adare ⁴, M.M. Aggarwal ⁵, G. Aglieri Rinella ⁶, A.G. Agocs ⁷, A. Agostinelli ⁸, S. Aguilar Salazar ⁹, Z. Ahammed ¹⁰, N. Ahmad ¹¹, A. Ahmad Masoodi ¹¹, S.U. Ahn ^{12,i}, A. Akindinov ¹³, D. Aleksandrov ¹⁴, B. Alessandro ¹⁵, R. Alfaro Molina ⁹, A. Alici ¹⁶,

A. Alkin ¹⁷, E. Almaráz Aviña ⁹, J. Alme ¹⁸, T. Alt ¹⁹, V. Altini ^{20,ii}, I. Altsybeev ²¹, C. Andrei ²², A. Andronic ²³, V. Anguelov ^{19,iii}, C. Anson ²⁴, T. Antičić ²⁵, F. Antinori ²⁶, P. Antonioli ²⁷, L. Aphecetche ²⁸, H. Appelshäuser ²⁹, N. Arbor ³⁰, S. Arcelli ⁸, A. Arend ²⁹, N. Armesto ³¹, R. Arnaldi ¹⁵, T. Aronsson ⁴, I.C. Arsene ²³, A. Asryan ²¹, A. Augustinus ⁶, R. Averbeck ²³, T.C. Awes ³², J. Äystö ³³, M.D. Azmi ¹¹, M. Bach ¹⁹, A. Badalà ³⁴, Y.W. Baek ^{12,i}, R. Bailhache ²⁹, R. Bala ¹⁵, R. Baldini Ferroli ¹⁶, A. Baldisseri ³⁵, A. Baldit ³⁶, J. Bán ³⁷, R. Barbera ³⁸, F. Barile ²⁰, G.G. Barnaföldi ⁷, L.S. Barnby ³⁹, V. Barret ³⁶, J. Bartke ⁴⁰, M. Basile ⁸, N. Bastid ³⁶, B. Bathen ⁴¹, G. Batigne ²⁸, B. Batyunya ⁴², C. Baumann ²⁹, I.G. Bearden ⁴³, H. Beck ²⁹, I. Belikov ⁴⁴, F. Bellini ⁸, R. Bellwied ⁴⁵, E. Belmont-Moreno ⁹, S. Beole ⁴⁶, I. Berceanu ²², A. Bercuci ²², E. Berdermann ²³, Y. Berdnikov ⁴⁷, C. Bergmann ⁴¹, L. Betev ⁶, A. Bhasin ⁴⁸, A.K. Bhati ⁵, L. Bianchi ⁴⁶, N. Bianchi ⁴⁹, C. Bianchin ⁵⁰, J. Bielčík ⁵¹, J. Bielčíková ³, A. Bilandzic ⁵², E. Biolcati ⁴⁶, A. Blaganov ⁵⁴, H. Borgild ⁴³, M. Bogolyubsky ⁵⁵, L. Boldizsár ⁷, M. Bombara ⁵⁶, C. Bombonati ⁵⁰, J. Book ²⁹, H. Borel ³⁵, A. Borissov ⁵⁷, C. Bortolin ^{50,iv}, S. Bose ⁵⁸, F. Bossú ⁶⁴, M. Borje ⁵², S. Böttger ⁵⁹, B. Boyer ⁶⁰, P. Braun-Munzinger ²³, L. Bravina ⁶¹, M. Bregant ²⁸, T. Breitner ⁵⁹, M. Broz ⁶², R. Brun ⁶, E. Bruna ⁴, G.E. Bruno ^{20,**}, D. Budnikov ⁶³, H. Buesching ²⁹, S. Bufalino ⁴⁶, O. Busch ⁶⁴, Z. Buthelezi ⁶⁵, D. Caffarri ⁵⁰, X. Cai ⁶⁶, H. Caines ⁴, E. Calvo Villar ⁶⁷, P. Camerini ⁶⁸, V. Canoa Roman ^{69,vi}, G. Cara Romeo ²⁷, F. Carena ⁶, W. Carena ⁶, V. Canoa Roman ^{69,vi}, G. Cara Romeo ²⁷, F. Carena ⁶, W. Carena ⁶, P. Christiansen ⁷⁴, T. Chujo ⁷⁵, C. Cicalo ⁷⁶, L. Cifarelli ^{8,ii}, F. Cindolo ²⁷, J. Cleymans S. De ¹⁰, A. De Azevedo Moregula ⁴⁹, G.O.V. de Barros ⁸², A. De Caro ⁸³, G. de Cataldo ⁸⁴, J. de Cuveland ¹⁹. A. De Falco ⁸⁵, D. De Gruttola ⁸³, N. De Marco ¹⁵, S. De Pasquale ⁸³, R. de Rooij ⁷³, E. Del Castillo Sanchez ⁶, H. Delagrange ²⁸, Y. Delgado Mercado ⁶⁷, G. Dellacasa ⁷⁸, vii, A. Deloff ⁸⁶, V. Demanov ⁶³, E. Dénes ⁷, A. Deppman ⁸², D. Di Bari ²⁰, C. Di Giglio ²⁰, S. Di Liberto ⁸⁷, A. Di Mauro ⁶, P. Di Nezza ⁴⁹, T. Dietel ⁴¹, R. Divià ⁶, Ø. Djuvsland ¹, A. Dobrin ⁵⁷, T. Dobrowolski ⁸⁶, I. Domínguez ⁷⁹, B. Dönigus ²³, O. Dordic ⁶¹, O. Driga ²⁸, A.K. Dubey ¹⁰, L. Ducroux ⁷¹, P. Dupieux ³⁶, A.K. Dutta Majumdar ⁵⁸, M.R. Dutta Majumdar ¹⁰, D. Elia ⁸⁴, D. Emschermann ⁴¹, H. Engel ⁵⁹, H.A. Erdal ¹⁸, B. Espagnon ⁶⁰, M. Estienne ²⁸, S. Esumi ⁷⁵, D. Evans ³⁹, S. Evrard ⁶, G. Eyyubova ⁶¹, D. Fabris ²⁶, J. Faivre ³⁰, D. Falchieri ⁸, A. Fantoni ⁴⁹, M. Fasel ²³, R. Fearick ⁶⁵, A. Fedunov ⁴², D. Fehlker ¹, V. Fekete ⁶², D. Felea ⁸⁰, G. Feofilov ²¹, A. Fernández Téllez ⁷⁷, E.G. Ferreiro ³¹, A. Ferretti ⁴⁶, R. Ferretti ⁷⁸, M.A.S. Figueredo ⁸², S. Filchagin ⁶³, R. Fini ⁸⁴, D. Finogeev ⁸⁸, F.M. Fionda ²⁰, E.M. Fiore ²⁰, M. Floris ⁶, S. Foertsch ⁶⁵, P. Foka ²³, S. Fokin ¹⁴, E. Fragiacomo ⁸⁹, M. Fragkiadakis ⁹⁰, U. Frankenfeld ²³, U. Fuchs ⁶, F. Furano ⁶, C. Furget ³⁰, M. Fusco Girard ⁸³, J.J. Gaardhøje ⁴³, S. Gadrat ³⁰, M. Gagliardi ⁴⁶, A. Gago ⁶⁷, M. Gallio ⁴⁶, P. Ganoti ³², C. Garabatos ²³, E. Garcia-Solis ⁹¹, R. Gemme ⁷⁸, J. Gerhard ¹⁹, M. Germain ²⁸, C. Geuna ³⁵, A. Gheata ⁶, Girard 63, J.J. Gaardhøje 43, S. Gadrat 30, M. Gagliardi 40, A. Gago 7, M. Gallio 40, P. Ganoti 32, C. Garabatos 23, E. Garcia-Solis 91, R. Gemme 78, J. Gerhard 19, M. Germain 28, C. Geuna 35, A. Gheata 6, M. Gheata 6, B. Ghidini 20, P. Ghosh 10, P. Gianotti 49, M.R. Girard 92, P. Giubellino 46,viii, E. Gladysz-Dziadus 40, P. Glässel 64, R. Gomez 93, L.H. González-Trueba 9, P. González-Zamora 53, S. Gorbunov 19, S. Gotovac 94, V. Grabski 9, L.K. Graczykowski 92, R. Grajcarek 64, A. Grelli 73, A. Grigoras 6, C. Grigoras 6, V. Grigoriev 54, A. Grigoryan 95, S. Grigoryan 42, B. Grinyov 17, N. Grion 89, P. Gros 74, J.F. Grosse-Oetringhaus 6, J.-Y. Grossiord 71, F. Guber 88, R. Guernane 30, C. Guerra Gutierrez 67, B. Guerzoni 8, K. Gulbrandsen 43, H. Gulkanyan 95, T. Gunji 96, A. Gupta 48, R. Gupta 48, H. Gutbrod 23, Ø. Haaland 1, C. Hadjidakis 60, M. Haiduc 80, H. Hamagaki 96, G. Hamar 7, L.D. Hanratty 39, T. Harmanova 56, L.W. Harris 4, M. Hartig 29, D. Hasegan 80, D. Hatzifotiadou 27, A. Hayranetyan 95, ii Z. Harmanova ⁵⁶, J.W. Harris ⁴, M. Hartig ²⁹, D. Hasegan ⁸⁰, D. Hatzifotiadou ²⁷, A. Hayrapetyan ^{95,ii}, M. Heide ⁴¹, M. Heinz ⁴, H. Helstrup ¹⁸, A. Herghelegiu ²², G. Herrera Corral ⁶⁹, N. Herrmann ⁶⁴, K.F. Hetland ¹⁸, B. Hicks ⁴, P.T. Hille ⁴, B. Hippolyte ⁴⁴, T. Horaguchi ⁷⁵, Y. Hori ⁹⁶, P. Hristov ⁶, I. Hřivnáčová ⁶⁰, M. Huang ¹, S. Huber ²³, T.J. Humanic ²⁴, D.S. Hwang ⁹⁷, R. Ilkaev ⁶³, I. Ilkiv ⁸⁶, M. Inaba ⁷⁵, E. Incani ⁸⁵, G.M. Innocenti ⁴⁶, M. Ippolitov ¹⁴, M. Irfan ¹¹, C. Ivan ²³, A. Ivanov ²¹, M. Ivanov ²³, V. Ivanov ⁴⁷, A. Jachołkowski ⁶, P.M. Jacobs ⁹⁸, L. Jancurová ⁴², S. Jangal ⁴⁴, M.A. Janik ⁹²,

R. Janik ⁶², P.H.S.Y. Jayarathna ^{45,ix}, S. Jena ⁹⁹, L. Jirden ⁶, G.T. Jones ³⁹, P.G. Jones ³⁹, P. Jovanović ³⁹, H. Jung ¹², W. Jung ¹², A. Jusko ³⁹, S. Kalcher ¹⁹, P. Kaliňák ³⁷, M. Kalisky ⁴¹, T. Kalliokoski ³³, A. Kalweit ¹⁰⁰, R. Kamermans ^{73,vii}, K. Kanaki ¹, E. Kang ¹², J.H. Kang ¹⁰¹, V. Kaplin ⁵⁴, A. Karasu Uysal ⁶, O. Karavichev ⁸⁸, T. Karavicheva ⁸⁸, E. Karpechev ⁸⁸, A. Kazantsev ¹⁴, U. Kebschull ⁵⁹, R. Keidel ¹⁰², M.M. Khan ¹¹, P. Khan ⁵⁸, A. Khanzadeev ⁴⁷, Y. Kharlov ⁵⁵, B. Kileng ¹⁸, D.J. Kim ³³, D.S. Kim ¹², D.W. Kim ¹², J.H. Kim ⁹⁷, J.S. Kim ¹², M. Kim ¹⁰¹, S. Kim ⁹⁷, S.H. Kim ¹², S. Kirsch ^{6,x}, I. Kisel ¹⁹, S. Kiselev ¹³, A. Kisiel ⁶, J.L. Klay ¹⁰³, J. Klein ⁶⁴, C. Klein-Bösing ⁴¹, M. Kliemant ²⁹, A. Kluge ⁶, M.L. Knichel ²³, K. Koch ⁶⁴, M.K. Köhler ²³, A. Kolojvari ²¹, V. Kondratiev ²¹, N. Kondratyeva ⁵⁴, A. Konevskih ⁸⁸, E. Kornaś ⁴⁰, C. Kottachchi Kankanamge Don ⁵⁷, R. Kour ³⁹, M. Kowalski ⁴⁰, S. Kox ³⁰, G. Koyithatta Meethaleveedu ⁹⁹, K. Kozlov ¹⁴, J. Kral ³³, I. Králik ³⁷, F. Kramer ²⁹, I. Kraus ²³, T. Krawutschke ^{64,xi}, M. Kretz ¹⁹, M. Krivda ^{39,xii}, F. Krizek ³³, M. Krus ⁵¹, E. Kryshen ⁴⁷, M. Krzewicki ⁵², Y. Kucheriaev ¹⁴, C. Kuhn ⁴⁴, P.G. Kuijer ⁵², P. Kurashvili ⁸⁶, A. Kurepin ⁸⁸, A. Kuryakin ⁶³, S. Kushpil ³, V. Kushpil ³, M.J. Kweon ⁶⁴, Y. Kwon ¹⁰¹, P. La Rocca ³⁸, P. Ladrón de Guevara ⁷⁹, V. Lafage ⁶⁰, I. Lakomov ²¹, C. Lara ⁵⁹, D.T. Larsen ¹, C. Lazzeroni ³⁹, P. Ladrón de Guevara ⁷⁹, V. Lafage ⁶⁰, I. Lakomov ²¹, C. Lara ⁵⁹, D.T. Larsen ¹, C. Lazzeroni ³⁹, Y. Le Bornec ⁶⁰, R. Lea ⁶⁸, M. Lechman ⁶, K.S. Lee ¹², S.C. Lee ¹², F. Lefèvre ²⁸, J. Lehnert ²⁹, L. Leistam ⁶, M. Lenhardt ²⁸, V. Lenti ⁸⁴, I. León Monzón ⁹³, H. León Vargas ²⁹, P. Lévai ⁷, X. Li ¹⁰⁴, R. Lietava ³⁹, C. Lietava ³⁹, P. Lévai ⁷, X. Li ¹⁰⁴, R. Lietava ³⁹, P. Lévai ⁷, X. Li ¹⁰⁴, R. Lietava ³⁹, P. Lévai ⁷, X. Li ¹⁰⁴, R. Lietava ³⁹, P. Lévai ⁷, X. Li ¹⁰⁴, R. Lietava ³⁹, P. Lévai ⁷, X. Li ¹⁰⁴, R. Lietava ³⁹, P. Lévai ⁷, X. Li ¹⁰⁴, R. Lietava ³⁹, P. Lévai ⁷, X. Li ¹⁰⁴, R. Lietava ³⁹, P. Lévai ⁷, X. Li ¹⁰⁴, R. Lietava ³⁹, P. Lévai ⁷, X. Li ¹⁰⁴, R. Lietava ³⁹, P. Lévai ⁷, X. Li ¹⁰⁴, R. Lietava ³⁹, P. Lévai ⁷, X. Li ¹⁰⁴, R. Lietava ³⁹, P. Lévai ⁷, X. Li ¹⁰⁴, R. Lietava ³⁹, P. Lévai ⁷, X. Li ¹⁰⁴, R. Lietava ³⁹, P. Lévai ⁷, X. Li ¹⁰⁴, R. Lietava ³⁹, P. Lévai ⁷, Y. Lenti ⁸⁴, R. Lietava ⁸⁵, P. Lévai ⁷, X. Li ⁸⁴, R. Lietava ⁸⁵, P. Lévai ⁷, X. Li ⁸⁴, R. Lietava ⁸⁵, P. Lévai ⁷, X. Li ⁸⁵, R. Lietava ⁸⁶, R. Lietava ⁸⁷, P. Lévai ⁷, X. Li ⁸⁵, R. Lietava ⁸⁷, P. Lévai ⁷, Y. Lietava ⁸⁸, P. Lietava ⁸⁸, P. Lietava ⁸⁹, P. Lévai ⁷⁸, R. Lietava ⁸⁹, P. Lévai ⁷⁸, R. Lietava ⁸⁹, P. Lévai ⁷⁸, P. S. Lindal ⁶¹, V. Lindenstruth ¹⁹, C. Lippmann ²³, M.A. Lisa ²⁴, L. Liu ¹, V.R. Loggins ⁵⁷, V. Loginov ⁵⁴, S. Lohn ⁶, D. Lohner ⁶⁴, C. Loizides ⁹⁸, K.K. Loo ³³, X. Lopez ³⁶, M. López Noriega ⁶⁰, E. López Torres ², G. Løvhøiden ⁶¹, X.-G. Lu ⁶⁴, P. Luettig ²⁹, M. Lunardon ⁵⁰, G. Luparello ⁴⁶, L. Luquin ²⁸, C. Luzzi ⁶, K. Ma ⁶⁶, R. Ma ⁴, D.M. Madagodahettige-Don ⁴⁵, A. Maevskaya ⁸⁸, M. Mager ⁶, D.P. Mahapatra ⁸¹, A. Maire ⁴⁴, R. Ma ⁴, D.M. Madagodahettige-Don ⁴⁵, A. Maevskaya ⁸⁸, M. Mager ⁶, D.P. Mahapatra ⁸¹, A. Maire ⁴⁴, M. Malaev ⁴⁷, I. Maldonado Cervantes ⁷⁹, D. Mal'Kevich ¹³, P. Malzacher ²³, A. Mamonov ⁶³, L. Manceau ³⁶, V. Manko ¹⁴, F. Manso ³⁶, V. Manzari ⁸⁴, Y. Mao ^{66,xiii}, M. Marchisone ⁴⁶, J. Mareš ¹⁰⁵, G.V. Margagliotti ⁶⁸, A. Margotti ²⁷, A. Marín ²³, C. Markert ¹⁰⁶, I. Martashvili ¹⁰⁷, P. Martinengo ⁶, M.I. Martínez ⁷⁷, A. Martínez Davalos ⁹, G. Martínez García ²⁸, Y. Martynov ¹⁷, A. Mas ²⁸, S. Masciocchi ²³, M. Masera ⁴⁶, A. Masoni ⁷⁶, L. Massacrier ⁷¹, M. Mastromarco ⁸⁴, A. Mastroserio ⁶, Z.L. Matthews ³⁹, A. Matyja ⁴⁰, D. Mayani ⁷⁹, M.A. Mazzoni ⁸⁷, F. Meddi ¹⁰⁸, A. Menchaca-Rocha ⁹, P. Mendez Lorenzo ⁶, J. Mercado Pérez ⁶⁴, M. Meres ⁶², Y. Miake ⁷⁵, J. Midori ¹⁰⁹, L. Milano ⁴⁶, J. Milosevic ^{61,xiv}, A. Mischke ⁷³, D. Miśkowiec ^{6,xv}, C. Mitu ⁸⁰, J. Mlynarz ⁵⁷, B. Mohanty ¹⁰, L. Molnar ⁶, L. Montaño Zetina ⁶⁹, M. Monteno ¹⁵, E. Montes ⁵³, M. Morando ⁵⁰, D.A. Moreira De Godoy ⁸², S. Moretto ⁵⁰, A. Morsch ⁶, V. Muccifora ⁴⁹, F. Mudnic ⁹⁴, H. Müller ⁶, S. Muhuri ¹⁰, M.C. Munhoz ⁸², I. Musa ⁶, A. Musso ¹⁵ M. Monteno ¹⁵, E. Montes ⁵³, M. Morando ⁵⁰, D.A. Moreira De Godoy ⁸², S. Moretto ⁵⁰, A. Morsch ⁶, V. Muccifora ⁴⁹, E. Mudnic ⁹⁴, H. Müller ⁶, S. Muhuri ¹⁰, M.G. Munhoz ⁸², L. Musa ⁶, A. Musso ¹⁵, B.K. Nandi ⁹⁹, R. Nania ²⁷, E. Nappi ⁸⁴, C. Nattrass ¹⁰⁷, F. Navach ²⁰, S. Navin ³⁹, T.K. Nayak ¹⁰, S. Nazarenko ⁶³, G. Nazarov ⁶³, A. Nedosekin ¹³, F. Nendaz ⁷¹, M. Nicassio ²⁰, B.S. Nielsen ⁴³, S. Nikolaev ¹⁴, V. Nikolic ²⁵, S. Nikulin ¹⁴, V. Nikulin ⁴⁷, B.S. Nilsen ⁷⁰, M.S. Nilsson ⁶¹, F. Noferini ²⁷, G. Nooren ⁷³, N. Novitzky ³³, A. Nyanin ¹⁴, A. Nyatha ⁹⁹, C. Nygaard ⁴³, J. Nystrand ¹, H. Obayashi ¹⁰⁹, A. Ochirov ²¹, H. Oeschler ¹⁰⁰, S.K. Oh ¹², J. Oleniacz ⁹², C. Oppedisano ¹⁵, A. Ortiz Velasquez ⁷⁹, G. Ortona ^{6,v}, A. Oskarsson ⁷⁴, P. Ostrowski ⁹², I. Otterlund ⁷⁴, J. Otwinowski ²³, G. Øvrebekk ¹, K. Oyama ⁶⁴, K. Ozawa ⁹⁶, Y. Pachmayer ⁶⁴, M. Pachr ⁵¹, F. Padilla ⁴⁶, P. Pagano ⁸³, G. Paić ⁷⁹, F. Painke ¹⁹, C. Pajares ³¹, S. Pal ³⁵, S.K. Pal ¹⁰, A. Palaha ³⁹, A. Palmeri ³⁴, G.S. Pappalardo ³⁴, W.J. Park ²³, V. Paticchio ⁸⁴, A. Pavlinov ⁵⁷, T. Pawlak ⁹², T. Peitzmann ⁷³, D. Peresunko ¹⁴, C.E. Pérez Lara ⁵², D. Perini ⁶, D. Perrino ²⁰, W. Peryt ⁹², A. Pesci ²⁷, V. Peskoy ^{6,xvi}, Y. Pestoy ¹¹⁰, A.I. Peters ⁶, V. Petráček ⁵¹, M. Petra ⁵¹, M. Petris ²², P. Petroy ³⁹, A. Pesci ²⁷, V. Peskov ^{6,xvi}, Y. Pestov ¹¹⁰, A.J. Peters ⁶, V. Petráček ⁵¹, M. Petran ⁵¹, M. Petris ²², P. Petrov ³⁹, M. Petrovici ²², C. Petta ³⁸, S. Piano ⁸⁹, A. Piccotti ¹⁵, M. Pikna ⁶², P. Pillot ²⁸, O. Pinazza ⁶, L. Pinsky ⁴⁵, N. Pitz ²⁹, F. Piuz ⁶, D.B. Piyarathna ^{57,xvii}, R. Platt ³⁹, M. Płoskoń ⁹⁸, J. Pluta ⁹², T. Pocheptsov ^{42,xviii}, S. Pochybova ⁷, P.L.M. Podesta-Lerma ⁹³, M.G. Poghosyan ⁴⁶, B. Polichtchouk ⁵⁵, A. Pop ²², V. Pospíšil ⁵¹, B. Potukuchi ⁴⁸, S.K. Prasad ⁵⁷, R. Preghenella ¹⁶, F. Prino ¹⁵, C.A. Pruneau ⁵⁷, I. Pshenichnov ⁸⁸, G. Puddu ⁸⁵, A. Pulvirenti ^{38,ii}, V. Punin ⁶³, M. Putiš ⁵⁶, J. Putschke ⁴, E. Quercigh ⁶, H. Qvigstad ⁶¹, A. Rachevski ⁸⁹, A. Rademakers ⁶, S. Radomski ⁶⁴, T.S. Räihä ³³, J. Rak ³³, A. Rakotozafindrabe ³⁵, L. Ramello ⁷⁸, A. Ramírez Reyes ⁶⁹, M. Rammler ⁴¹, R. Raniwala ¹¹¹, S. Raniwala ¹¹¹, S.S. Räsänen ³³, D. Rathee ⁵, K.F. Read ¹⁰⁷, J.S. Real ³⁰, K. Redlich ⁸⁶, xix, R. Renfordt ²⁹, A.R. Reolon ⁴⁹, A. Reshetin ⁸⁸, F. Rettig ¹⁹, J.-P. Revol ⁶, K. Reygers ⁶⁴, H. Ricaud ¹⁰⁰, L. Riccati ¹⁵, R.A. Ricci ¹¹², M. Richter ¹, xx, P. Riedler ⁶, W. Riegler ⁶, F. Riggi ³⁸, M. Rodríguez Cahuantzi ⁷⁷, D. Rohr ¹⁹, D. Röhrich ¹, R. Romita ²³, F. Ronchetti ⁴⁹, P. Rosinský ⁶, P. Rosnet ³⁶, S. Rossegger ⁶, A. Rossi ⁵⁰, F. Roukoutakis ⁹⁰, S. Rousseau ⁶⁰, C. Roy ⁴⁴, P. Roy ⁵⁸, A.J. Rubio Montero ⁵³, R. Rui ⁶⁸, E. Ryabinkin ¹⁴, A. Rybicki ⁴⁰, S. Sadovsky ⁵⁵, K. Šafařík ⁶, R. Sahoo ⁵⁰, P.K. Sahu ⁸¹, P. Saiz ⁶, S. Sakai ⁹⁸, D. Sakata ⁷⁵, C.A. Salgado ³¹, S. Sambyal ⁴⁸, V. Samsonov ⁴⁷, L. Šándor ³⁷,

```
A. Sandoval <sup>9</sup>, M. Sano <sup>75</sup>, S. Sano <sup>96</sup>, R. Santo <sup>41</sup>, R. Santoro <sup>84</sup>, J. Sarkamo <sup>33</sup>, P. Saturnini <sup>36</sup>, E. Scapparone <sup>27</sup>, F. Scarlassara <sup>50</sup>, R.P. Scharenberg <sup>113</sup>, C. Schiaua <sup>22</sup>, R. Schicker <sup>64</sup>, C. Schmidt <sup>23</sup>, H.R. Schmidt <sup>23,xxi</sup>, S. Schreiner <sup>6</sup>, S. Schuchmann <sup>29</sup>, J. Schukraft <sup>6</sup>, Y. Schutz <sup>28,ii</sup>, K. Schwarz <sup>23</sup>, K. Schweda <sup>64</sup>, G. Scioli <sup>8</sup>, E. Scomparin <sup>15</sup>, P.A. Scott <sup>39</sup>, R. Scott <sup>107</sup>, G. Segato <sup>50</sup>, S. Senyukov <sup>78</sup>,
 K. Schweda <sup>64</sup>, G. Scioli <sup>8</sup>, E. Scomparin <sup>15</sup>, P.A. Scott <sup>39</sup>, R. Scott <sup>107</sup>, G. Segato <sup>50</sup>, S. Senyukov <sup>78</sup>, J. Seo <sup>12</sup>, S. Serci <sup>85</sup>, E. Serradilla <sup>53</sup>, A. Sevcenco <sup>80</sup>, I. Sgura <sup>84</sup>, G. Shabratova <sup>42</sup>, R. Shahoyan <sup>6</sup>, N. Sharma <sup>5</sup>, S. Sharma <sup>48</sup>, K. Shigaki <sup>109</sup>, M. Shimomura <sup>75</sup>, K. Shtejer <sup>2</sup>, Y. Sibiriak <sup>14</sup>, M. Siciliano <sup>46</sup>, E. Sicking <sup>6</sup>, T. Siemiarczuk <sup>86</sup>, D. Silvermyr <sup>32</sup>, G. Simonetti <sup>6</sup>, R. Singaraju <sup>10</sup>, R. Singh <sup>48</sup>, S. Singha <sup>10</sup>, B.C. Sinha <sup>10</sup>, T. Sinha <sup>58</sup>, B. Sitar <sup>62</sup>, M. Sitta <sup>78</sup>, T.B. Skaali <sup>61</sup>, K. Skjerdal <sup>1</sup>, R. Smakal <sup>51</sup>, N. Smirnov <sup>4</sup>, R. Snellings <sup>73</sup>, C. Søgaard <sup>43</sup>, R. Soltz <sup>114</sup>, H. Son <sup>97</sup>, J. Song <sup>115</sup>, M. Song <sup>101</sup>, C. Soos <sup>6</sup>, F. Soramel <sup>50</sup>, M. Spyropoulou-Stassinaki <sup>90</sup>, B.K. Srivastava <sup>113</sup>, J. Stachel <sup>64</sup>, I. Stan <sup>80</sup>, G. Stefanek <sup>86</sup>, G. Stefanini <sup>6</sup>, T. Steinbeck <sup>19</sup>, M. Steinpreis <sup>24</sup>, E. Stenlund <sup>74</sup>, G. Steyn <sup>65</sup>, D. Stocco <sup>28</sup>, R. Stock <sup>29</sup>, C.H. Stokkevag <sup>1</sup>, M. Stolpovskiy <sup>55</sup>, P. Strmen <sup>62</sup>, A.A.P. Suaide <sup>82</sup>, M.A. Subieta Vásquez <sup>46</sup>, T. Sugitate <sup>109</sup>, C. Suire <sup>60</sup>, M. Sukhorukov <sup>63</sup>, M. Šumbera <sup>3</sup>, T. Susa <sup>25</sup>, D. Swoboda <sup>6</sup>, T.J.M. Symons <sup>98</sup>, A. Szanto de Toledo <sup>82</sup>, L. Szarka <sup>62</sup>, A. Szostak <sup>1</sup>, C. Tagridis <sup>90</sup>, L. Takahashi <sup>72</sup>, LD. Tanja Takaki <sup>60</sup>, A. Tauro <sup>6</sup>, C. Tajeda Muñoc
  I. Szarka 62, A. Szostak 1, C. Tagridis 90, J. Takahashi 72, J.D. Tapia Takaki 60, A. Tauro 6, G. Tejeda Muñoz 77, A. Telesca 6, C. Terrevoli 20, J. Thäder 23, D. Thomas 73, J.H. Thomas 23, R. Tieulent 71, A.R. Timmins 45, D. Tlusty 51, A. Toia 6, H. Torii 109, F. Tosello 15, T. Traczyk 92, D. Truesdale 24, W.H. Trzaska 33, A. Tumkin 63, R. Turrisi 26, A.J. Turvey 70, T.S. Tveter 61, J. Ulery 29, K. Ullaland 1, A. Uras 85, J. Urbán 56, G.M. Urciuoli 87, G.L. Usai 85, M. Vajzer 51, M. Vala 42,xii, L. Valencia Palomo 60, S. Vallero 64, N. van der Kolk 52,
G.L. Usai <sup>85</sup>, M. Vajzer <sup>51</sup>, M. Vala <sup>42,xii</sup>, L. Valencia Palomo <sup>60</sup>, S. Vallero <sup>64</sup>, N. van der Kolk <sup>52</sup>, M. van Leeuwen <sup>73</sup>, P. Vande Vyvre <sup>6</sup>, L. Vannucci <sup>112</sup>, A. Vargas <sup>77</sup>, R. Varma <sup>99</sup>, M. Vasileiou <sup>90</sup>, A. Vasiliev <sup>14</sup>, V. Vechernin <sup>21</sup>, M. Veldhoen <sup>73</sup>, M. Venaruzzo <sup>68</sup>, E. Vercellin <sup>46</sup>, S. Vergara <sup>77</sup>, D.C. Vernekohl <sup>41</sup>, R. Vernet <sup>116</sup>, M. Verweij <sup>73</sup>, L. Vickovic <sup>94</sup>, G. Viesti <sup>50</sup>, O. Vikhlyantsev <sup>63</sup>, Z. Vilakazi <sup>65</sup>, O. Villalobos Baillie <sup>39</sup>, A. Vinogradov <sup>14</sup>, L. Vinogradov <sup>21</sup>, Y. Vinogradov <sup>63</sup>, T. Virgili <sup>83</sup>, Y.P. Viyogi <sup>10</sup>, A. Vodopyanov <sup>42</sup>, K. Voloshin <sup>53</sup>, S. Voloshin <sup>57</sup>, G. Volpe <sup>20</sup>, B. von Haller <sup>6</sup>, D. Vranic <sup>23</sup>, J. Vrláková <sup>56</sup>, B. Vulpescu <sup>36</sup>, A. Vyushin <sup>63</sup>, B. Wagner <sup>1</sup>, V. Wagner <sup>51</sup>, R. Wan <sup>44,xxii</sup>, D. Wang <sup>66</sup>, M. Wang <sup>66</sup>, Y. Wang <sup>64</sup>, Y. Wang <sup>66</sup>, K. Watanabe <sup>75</sup>, J.P. Wessels <sup>41,viii</sup>, U. Westerhoff <sup>41</sup>, J. Wiechula <sup>64,xxiii</sup>, J. Wikne <sup>61</sup>, M. Wilde <sup>41</sup>, A. Wilk <sup>41</sup>, G. Wilk <sup>86</sup>, M.C.S. Williams <sup>27</sup>, B. Windelband <sup>64</sup>, H. Yang <sup>35</sup>, S. Yasnopolskiy <sup>14</sup>, J. Yi <sup>115</sup>, Z. Yin <sup>66</sup>, H. Yokoyama <sup>75</sup>, I.-K. Yoo <sup>115</sup>, X. Yuan <sup>66</sup>, I. Yushmanov <sup>14</sup>, E. Zabrodin <sup>61</sup>, C. Zach <sup>51</sup>, C. Zampolli <sup>6</sup>, S. Zaporozhets <sup>42</sup>, A. Zarochentsev <sup>21</sup>, P. Závada <sup>105</sup>, N. Zaviyalov <sup>63</sup>, H. Zbroszczyk <sup>92</sup>, P. Zelnicek <sup>59,ii</sup>, A. Zenin <sup>55</sup>, I. Zgura <sup>80</sup>, M. Zhalov <sup>47</sup>, X. Zhang <sup>66,i</sup>, D. Zhou <sup>66</sup>, F. Zhou <sup>66</sup>, Y. Zhou <sup>73</sup>, X. Zhu <sup>66</sup>, A. Zichichi <sup>8,xxiv</sup>, G. Zinovjev <sup>17</sup>, Y. Zoccarato <sup>71</sup>, M. Zynovyev <sup>17</sup>
     ^{\rm 1} Department of Physics and Technology, University of Bergen, Bergen, Norway
     <sup>2</sup> Centro de Aplicaciones Tecnológicas y Desarrollo Nuclear (CEADEN), Havana, Cuba
     <sup>3</sup> Nuclear Physics Institute, Academy of Sciences of the Czech Republic, Řež u Prahy, Czech Republic
     <sup>4</sup> Yale University, New Haven, CT, United States
      <sup>5</sup> Physics Department, Panjab University, Chandigarh, India
     <sup>6</sup> European Organization for Nuclear Research (CERN), Geneva, Switzerland
     <sup>7</sup> KFKI Research Institute for Particle and Nuclear Physics, Hungarian Academy of Sciences, Budapest, Hungary
     <sup>8</sup> Dipartimento di Fisica dell'Università and Sezione INFN, Bologna, Italy
     <sup>9</sup> Instituto de Física, Universidad Nacional Autónoma de México, Mexico City, Mexico
     <sup>10</sup> Variable Energy Cyclotron Centre, Kolkata, India
```

¹¹ Department of Physics, Aligarh Muslim University, Aligarh, India

¹² Gangneung-Wonju National University, Gangneung, South Korea

¹³ Institute for Theoretical and Experimental Physics, Moscow, Russia

¹⁴ Russian Research Centre Kurchatov Institute, Moscow, Russia 15 Sezione INFN, Turin, Italy

¹⁶ Centro Fermi – Centro Studi e Ricerche e Museo Storico della Fisica "Enrico Fermi", Rome, Italy

¹⁷ Bogolyubov Institute for Theoretical Physics, Kiev, Ukraine

¹⁸ Faculty of Engineering, Bergen University College, Bergen, Norway

¹⁹ Frankfurt Institute for Advanced Studies, Johann Wolfgang Goethe-Universität Frankfurt, Frankfurt, Germany

²⁰ Dipartimento Interateneo di Fisica 'M. Merlin' and Sezione INFN, Bari, Italy

²¹ V. Fock Institute for Physics, St. Petersburg State University, St. Petersburg, Russia

²² National Institute for Physics and Nuclear Engineering, Bucharest, Romania

²³ Research Division and ExtreMe Matter Institute EMMI, GSI Helmholtzzentrum für Schwerionenforschung, Darmstadt, Germany

²⁴ Department of Physics, Ohio State University, Columbus, OH, United States

²⁵ Rudjer Bošković Institute, Zagreb, Croatia

²⁶ Sezione INFN, Padova, Italy

²⁷ Sezione INFN, Bologna, Italy

²⁸ SUBATECH, Ecole des Mines de Nantes, Université de Nantes, CNRS-IN2P3, Nantes, France

²⁹ Institut für Kernphysik, Johann Wolfgang Goethe-Universität Frankfurt, Frankfurt, Germany

³⁰ Laboratoire de Physique Subatomique et de Cosmologie (LPSC), Université Joseph Fourier, CNRS-IN2P3, Institut Polytechnique de Grenoble, Grenoble, France

³¹ Departamento de Física de Partículas and IGFAE, Universidad de Santiago de Compostela, Santiago de Compostela, Spain

³² Oak Ridge National Laboratory, Oak Ridge, TN, United States

³³ Helsinki Institute of Physics (HIP) and University of Jyväskylä, Jyväskylä, Finland

- ³⁴ Sezione INFN, Catania, Italy
- 35 Commissariat à l'Energie Atomique, IRFU, Saclay, France
- ³⁶ Laboratoire de Physique Corpusculaire (LPC), Clermont Université, Université Blaise Pascal, CNRS-IN2P3, Clermont-Ferrand, France
- ³⁷ Institute of Experimental Physics, Slovak Academy of Sciences, Košice, Slovakia
- ³⁸ Dipartimento di Fisica e Astronomia dell'Università and Sezione INFN, Catania, Italy
- ³⁹ School of Physics and Astronomy, University of Birmingham, Birmingham, United Kingdom
- ⁴⁰ The Henryk Niewodniczanski Institute of Nuclear Physics, Polish Academy of Sciences, Cracow, Poland
- ⁴¹ Institut für Kernphysik, Westfälische Wilhelms-Universität Münster, Münster, Germany
- ⁴² Joint Institute for Nuclear Research (JINR), Dubna, Russia
- ⁴³ Niels Bohr Institute, University of Copenhagen, Copenhagen, Denmark
- 44 Institut Pluridisciplinaire Hubert Curien (IPHC), Université de Strasbourg, CNRS-IN2P3, Strasbourg, France
- ⁴⁵ University of Houston, Houston, TX, United States
- ⁴⁶ Dipartimento di Fisica Sperimentale dell'Università and Sezione INFN, Turin, Italy
- ⁴⁷ Petersburg Nuclear Physics Institute, Gatchina, Russia
- ⁴⁸ Physics Department, University of Jammu, Jammu, India
- ⁴⁹ Laboratori Nazionali di Frascati, INFN, Frascati, Italy
- ⁵⁰ Dipartimento di Fisica dell'Università and Sezione INFN, Padova, Italy
- ⁵¹ Faculty of Nuclear Sciences and Physical Engineering, Czech Technical University in Prague, Prague, Czech Republic
- ⁵² Nikhef, National Institute for Subatomic Physics, Amsterdam, Netherlands
- ⁵³ Centro de Investigaciones Energéticas Medioambientales y Tecnológicas (CIEMAT), Madrid, Spain
- ⁵⁴ Moscow Engineering Physics Institute, Moscow, Russia
- ⁵⁵ Institute for High Energy Physics, Protvino, Russia
- ⁵⁶ Faculty of Science, P.J. Šafárik University, Košice, Slovakia
- ⁵⁷ Wayne State University, Detroit, MI, United States
- ⁵⁸ Saha Institute of Nuclear Physics, Kolkata, India
- ⁵⁹ Kirchhoff-Institut für Physik, Ruprecht-Karls-Universität Heidelberg, Heidelberg, Germany
- 60 Institut de Physique Nucléaire d'Orsay (IPNO), Université Paris-Sud, CNRS-IN2P3, Orsay, France
- ⁶¹ Department of Physics, University of Oslo, Oslo, Norway
- ⁶² Faculty of Mathematics, Physics and Informatics, Comenius University, Bratislava, Slovakia
- ⁶³ Russian Federal Nuclear Center (VNIIEF), Sarov, Russia
- ⁶⁴ Physikalisches Institut, Ruprecht-Karls-Universität Heidelberg, Heidelberg, Germany
- 65 Physics Department, University of Cape Town, iThemba LABS, Cape Town, South Africa
- ⁶⁶ Hua-Zhong Normal University, Wuhan, China
- ⁶⁷ Sección Física, Departamento de Ciencias, Pontificia Universidad Católica del Perú, Lima, Peru
- ⁶⁸ Dipartimento di Fisica dell'Università and Sezione INFN, Trieste, Italy
- ⁶⁹ Centro de Investigación y de Estudios Avanzados (CINVESTAV), Mexico City and Mérida, Mexico
- ⁷⁰ Physics Department, Creighton University, Omaha, NE, United States
- 71 Université de Lyon, Université Lyon 1, CNRS/IN2P3, IPN-Lyon, Villeurbanne, France
- ⁷² Universidade Estadual de Campinas (UNICAMP), Campinas, Brazil
- 73 Nikhef, National Institute for Subatomic Physics and Institute for Subatomic Physics of Utrecht University, Utrecht, Netherlands
- ⁷⁴ Division of Experimental High Energy Physics, University of Lund, Lund, Sweden
- ⁷⁵ University of Tsukuba, Tsukuba, Japan
- ⁷⁶ Sezione INFN, Cagliari, Italy
- ⁷⁷ Benemérita Universidad Autónoma de Puebla, Puebla, Mexico
- ⁷⁸ Dipartimento di Scienze e Tecnologie Avanzate dell'Università del Piemonte Orientale and Gruppo Collegato INFN, Alessandria, Italy
- ⁷⁹ Instituto de Ciencias Nucleares, Universidad Nacional Autónoma de México, Mexico City, Mexico
- ⁸⁰ Institute of Space Sciences (ISS), Bucharest, Romania
- ⁸¹ Institute of Physics, Bhubaneswar, India
- 82 Universidade de São Paulo (USP), São Paulo, Brazil
- ⁸³ Dipartimento di Fisica 'E.R. Caianiello' dell'Università and Gruppo Collegato INFN, Salerno, Italy
- 84 Sezione INFN, Bari, Italy
- ⁸⁵ Dipartimento di Fisica dell'Università and Sezione INFN, Cagliari, Italy
- 86 Soltan Institute for Nuclear Studies, Warsaw, Poland
- 87 Sezione INFN, Rome, Italy
- ⁸⁸ Institute for Nuclear Research, Academy of Sciences, Moscow, Russia
- ⁸⁹ Sezione INFN, Trieste, Italy
- ⁹⁰ Physics Department, University of Athens, Athens, Greece
- ⁹¹ Chicago State University, Chicago, IL, United States
- 92 Warsaw University of Technology, Warsaw, Poland
- ⁹³ Universidad Autónoma de Sinaloa, Culiacán, Mexico 94 Technical University of Split FESB, Split, Croatia
- ⁹⁵ Yerevan Physics Institute, Yerevan, Armenia
- ⁹⁶ University of Tokyo, Tokyo, Japan
- ⁹⁷ Department of Physics, Sejong University, Seoul, South Korea
- ⁹⁸ Lawrence Berkeley National Laboratory, Berkeley, CA, United States
- ⁹⁹ Indian Institute of Technology, Mumbai, India
- ¹⁰⁰ Institut für Kernphysik, Technische Universität Darmstadt, Darmstadt, Germany
- ¹⁰¹ Yonsei University, Seoul, South Korea
- ¹⁰² Zentrum für Technologietransfer und Telekommunikation (ZTT), Fachhochschule Worms, Worms, Germany
- ¹⁰³ California Polytechnic State University, San Luis Obispo, CA, United States
- ¹⁰⁴ China Institute of Atomic Energy, Beijing, China
- ¹⁰⁵ Institute of Physics, Academy of Sciences of the Czech Republic, Prague, Czech Republic
- ¹⁰⁶ The University of Texas at Austin, Physics Department, Austin, TX, United States
- 107 University of Tennessee, Knoxville, TN, United States
- ¹⁰⁸ Dipartimento di Fisica dell'Università 'La Sapienza' and Sezione INFN, Rome, Italy
- ¹⁰⁹ Hiroshima University, Hiroshima, Japan
- ¹¹⁰ Budker Institute for Nuclear Physics, Novosibirsk, Russia
- ¹¹¹ Physics Department, University of Rajasthan, Jaipur, India
- 112 Laboratori Nazionali di Legnaro, INFN, Legnaro, Italy

- ¹¹³ Purdue University, West Lafayette, IN, United States
- 114 Lawrence Livermore National Laboratory, Livermore, CA, United States
- 115 Pusan National University, Pusan, South Korea
- ¹¹⁶ Centre de Calcul de l'IN2P3, Villeurbanne, France
- * Corresponding author.

E-mail address: giuseppe.bruno@ba.infn.it (G.E. Bruno).

- i Also at Laboratoire de Physique Corpusculaire (LPC), Clermont Université, Université Blaise Pascal, CNRS-IN2P3, Clermont-Ferrand, France.
- $^{\rm ii}\,$ Also at European Organization for Nuclear Research (CERN), Geneva, Switzerland.
- iii Now at Physikalisches Institut, Ruprecht-Karls-Universität Heidelberg, Heidelberg, Germany.
- iv Also at Dipartimento di Fisica dell'Università, Udine, Italy.
- ^v Also at Dipartimento di Fisica Sperimentale dell'Università and Sezione INFN, Turin, Italy.
- vi Also at Benemérita Universidad Autónoma de Puebla, Puebla, Mexico.
- vii Deceased.
- viii Now at European Organization for Nuclear Research (CERN), Geneva, Switzerland.
- ix Also at Wayne State University, Detroit, Michigan, United States.
- x Also at Frankfurt Institute for Advanced Studies, Johann Wolfgang Goethe-Universität Frankfurt, Frankfurt, Germany.
- xi Also at Fachhochschule Köln, Köln, Germany.
- xii Also at Institute of Experimental Physics, Slovak Academy of Sciences, Košice, Slovakia.
- xiii Also at Laboratoire de Physique Subatomique et de Cosmologie (LPSC), Université Joseph Fourier, CNRS-IN2P3, Institut Polytechnique de Grenoble, Grenoble, France.
- xiv Also at "Vinča" Institute of Nuclear Sciences, Belgrade, Serbia.
- xv Also at Research Division and ExtreMe Matter Institute EMMI, GSI Helmholtzzentrum für Schwerionenforschung, Darmstadt, Germany.
- ^{xvi} Also at Instituto de Ciencias Nucleares, Universidad Nacional Autónoma de México, Mexico City, Mexico.
- xvii Also at University of Houston, Houston, Texas, United States.
- xviii Also at Department of Physics, University of Oslo, Oslo, Norway.
- xix Also at Institute of Theoretical Physics, University of Wroclaw, Wroclaw, Poland.
- xx Now at Department of Physics, University of Oslo, Oslo, Norway.
- $^{\rm xxi}\,$ Also at Eberhard Karls Universität Tübingen, Tübingen, Germany.
- xxii Also at Hua-Zhong Normal University, Wuhan, China.
- xxiii Now at Eberhard Karls Universität Tübingen, Tübingen, Germany.
- xxiv Also at Centro Fermi Centro Studi e Ricerche e Museo Storico della Fisica "Enrico Fermi", Rome, Italy.