

Online Resource II: References per method

The time of death in Dutch court; using the Daubert criteria to evaluate methods to estimate the PMI used in court

Legal Medicine

Tamara Gelderman^{1*}, Erik Stigter^{1*}, Tristan Krap^{1,2,3}, Jens Amendt⁴, Wilma Duijst^{1,3}

* Shared first author

¹ Faculty of Law and Criminology, Maastricht University, Maastricht, the Netherlands.
tgelderman@hotmail.com. Corresponding author.

² Amsterdam UMC, Location AMC, department of Medical Biology, Section Anatomy, Amsterdam, The Netherlands

³ Ars Cognoscendi Foundation for Legal and Forensic Medicine, Wezep, The Netherlands

⁴ Institute of Legal Medicine, University Hospital Frankfurt am Main, Goethe-University, Frankfurt am Main, Germany

Livor mortis

- Belenki L, Sterzik V, Bohnert M (2014) Similarity Analysis of Spectra Obtained via Reflectance Spectrometry in Legal Medicine. *J Lab Autom* 19:110–118.
<https://doi.org/10.1177/2211068213496089>
- Henßge C, Madea B (2004) Estimation of the time since death in the early post-mortem period. *Forensic Sci Int* 144:167–175. <https://doi.org/10.1016/j.forsciint.2004.04.051>
- v. Hunnius P, Mallach HJ, Mittmeyer H-J (1973) Über quantitative Druckmessungen an Totenflecken zur Bestimmung der Todeszeit. *Zeitschrift für Rechtsmedizin* 73:235–244.
<https://doi.org/10.1007/BF02116827>
- Inoue M, Suyama A, Matuoka T, et al (1994) Development of an instrument to measure postmortem lividity and its preliminary application to estimate the time since death. *Forensic Sci Int* 65:185–193. [https://doi.org/10.1016/0379-0738\(94\)90274-7](https://doi.org/10.1016/0379-0738(94)90274-7)
- Kaatsch HJ, Stadler M, Nietert M (1993) Photometric measurement of color changes in livor mortis as a function of pressure and time - Development of a computer-aided system for measuring pressure-induced blanching of livor mortis to estimate time of death. *Int J Legal Med* 106:91–97. <https://doi.org/10.1007/BF01225047>
- Kaatsch HJ, Schmidtke E, Nietsch W (1994) Photometric measurement of pressure-induced blanching of livor mortis as an aid to estimating time of death - Application of a new system for quantifying pressure-induced blanching in lividity. *Int J Legal Med* 106:209–214.
<https://doi.org/10.1007/BF01371339>
- Madea B, Henssge C, Reibe S, et al (2014) Postmortem Changes and Time Since Death. In: Madea B (ed) *Handbook of Forensic Medicine*, First. Wiley Blackwell, Bonn, pp 75–133.
- Madea B, Knight B (2016) Postmortem Lividity. In: Madea B (ed) *Estimation of the Time Since Death*, 3rd ed. CRC Press, Boca Raton, pp 59–62.
- Romanelli MC, Marrone M, Veneziani A, et al (2015) Hypostasis and time since death. *Am J Forensic Med Pathol* 36:99–103. <https://doi.org/10.1097/PAF.000000000000145>
- Saukko P, Knight B (2016) The Pathophysiology of Death. In: Knight's *Forensic Pathology*, 4th ed. CRC Press, Boca Raton, pp 55–94.
- Usumoto Y, Hikiji W, Sameshima N, et al (2010) Estimation of postmortem interval based on the spectrophotometric analysis of postmortem lividity. *Leg Med* 12:19–22.
<https://doi.org/10.1016/j.legalmed.2009.09.008>
- Vanezis P (1991) Assessing hypostasis by colorimetry. *Forensic Sci Int* 52:1–3.
[https://doi.org/10.1016/0379-0738\(91\)90089-2](https://doi.org/10.1016/0379-0738(91)90089-2)

- Vanezis P, Trujillo O (1996) Evaluation of hypostasis using a colorimeter measuring system and its application to assessment of the post-mortem interval (time of death). *Forensic Sci Int* 78:19–28. [https://doi.org/10.1016/0379-0738\(95\)01845-X](https://doi.org/10.1016/0379-0738(95)01845-X)

Rigor mortis

- Anders S, Kunz M, Gehl A, et al (2013) Estimation of the time since death-reconsidering the re-establishment of rigor mortis. *Int J Legal Med* 127:127–130. <https://doi.org/10.1007/s00414-011-0632-z>
- Crostack C, Sehner S, Raupach T, Anders S (2017) Re-establishment of rigor mortis: evidence for a considerably longer post-mortem time span. *Int J Legal Med* 131:1039–1042. <https://doi.org/10.1007/s00414-017-1558-x>
- Gorea R (2014) Study of Postmortem Interval from Rigor Mortis. *Journal of Punjab Academy of Forensic Medicine & toxicology*.
- Henßge C, Madea B (2004) Estimation of the time since death in the early post-mortem period. *Forensic Sci Int* 144:167–175. <https://doi.org/10.1016/j.forsciint.2004.04.051>
- Kobayashi M, Ikegaya H, Takase I, et al (2001) Development of rigor mortis is not affected by muscle volume. *Forensic Sci Int* 117:213–219. [https://doi.org/10.1016/S0379-0738\(00\)00406-0](https://doi.org/10.1016/S0379-0738(00)00406-0)
- Kobayashi M, Takemori S, Yamaguchi M (2004) Differential rigor development in red and white muscle revealed by simultaneous measurement of tension and stiffness. *Forensic Sci Int* 140:79–84. <https://doi.org/10.1016/j.forsciint.2003.11.026>
- Krompecher T (1981) Experimental evaluation of rigor mortis V. Effect of various temperatures on the evolution of rigor mortis. *Forensic Sci Int* 17:19–26. [https://doi.org/10.1016/0379-0738\(81\)90184-5](https://doi.org/10.1016/0379-0738(81)90184-5)
- Krompecher T (1994) Experimental evaluation of rigor mortis. VIII. Estimation of time since death by repeated measurements of the intensity of rigor mortis on rats. *Forensic Sci Int* 68:149–159. [https://doi.org/10.1016/0379-0738\(94\)90354-9](https://doi.org/10.1016/0379-0738(94)90354-9)
- Krompecher T (2016) Rigor Mortis. In: Madea B (ed) *Estimation of the Time Since Death*, 3rd ed. CRC Press, Boca Raton, pp 41–58.
- Krompecher T, Bergerioux C (1988) Experimental evaluation of rigor mortis. VII. Effect of ante- and post-mortem electrocution on the evolution of rigor mortis. *Forensic Sci Int* 38:27–35. [https://doi.org/10.1016/0379-0738\(88\)90006-0](https://doi.org/10.1016/0379-0738(88)90006-0)
- Krompecher T, Fryc O (1978) Experimental evaluation of rigor mortis III. Comparative study of the evolution of rigor mortis in different sized muscle groups in rats. *Forensic Sci Int* 12:97–102. [https://doi.org/10.1016/0379-0738\(78\)90016-6](https://doi.org/10.1016/0379-0738(78)90016-6)
- Krompecher T, Fryc O (1978) Experimental evaluation of rigor mortis IV. Change in strength and evolution of rigor mortis in the case of physical exercise preceding death. *Forensic Sci Int* 12:103–107. [https://doi.org/10.1016/0379-0738\(78\)90017-8](https://doi.org/10.1016/0379-0738(78)90017-8)
- Krompecher T, Gilles A, Brandt-Casadevall C, Mangin P (2008) Experimental evaluation of rigor mortis. IX. The influence of the breaking (mechanical solution) on the development of rigor mortis. *Forensic Sci Int* 176:157–162. <https://doi.org/10.1016/j.forsciint.2007.08.004>
- Madea B, Henssge C, Reibe S, et al (2014) Postmortem Changes and Time Since Death. In: Madea B (ed) *Handbook of Forensic Medicine*, First. Wiley Blackwell, Bonn, pp 75–133.
- Ozawa M, Iwadate K, Matsumoto S, et al (2013) The effect of temperature on the mechanical aspects of rigor mortis in a liquid paraffin model. *Leg Med* 15:293–297. <https://doi.org/10.1016/j.legalmed.2013.08.001>
- Saukko P, Knight B (2016) The Pathophysiology of Death. In: Knight's *Forensic Pathology*, 4th ed. CRC Press, Boca Raton, pp 55–94.
- Schuck M, Beier G, Liebhardt E, Spann W (1979) On the estimation of lay-time by measurements of rigor mortis. *Forensic Sci Int* 14:171–176. [https://doi.org/10.1016/0379-0738\(79\)90136-1](https://doi.org/10.1016/0379-0738(79)90136-1)

- Vain A, Kauppila R, Humal LH VE (1992) Grading Rigor Mortis With Myotonometry. *Forensic Sci Int* 56:147–150.
- Vain A, Kauppila R, Vuori E (1996) Estimation of the breaking of rigor mortis by myotonometry. *Forensic Sci Int* 79:155–161. [https://https://doi.org/10.1016/0379-0738\(96\)01902-0](https://doi.org/10.1016/0379-0738(96)01902-0)
- Varetto L, Curto O (2005) Long persistence of rigor mortis at constant low temperature. *Forensic Sci Int* 147:31–34. <https://https://doi.org/10.1016/j.forsciint.2004.03.033>

Algor Mortis

- Althaus L, Stückradt S, Henßge C, Bajanowski T (2007) Cooling experiments using dummies covered by leaves. *Int J Legal Med* 121:112–114. <https://https://doi.org/10.1007/s00414-006-0108-8>
- Baccino E, De Saint Martin L, Schuliar Y, et al (1996) Outer ear temperature and time of death. *Forensic Sci Int* 83:133–146. [https://https://doi.org/10.1016/S0379-0738\(96\)02027-0](https://https://doi.org/10.1016/S0379-0738(96)02027-0)
- Bakar ZA, Bin M, Abdullah C, et al (2018) Post-mortem Interval Estimation System (e-PMI) in Estimating the Time of Death. *Int J Inf Educ Technol* 8:46–50. <https://https://doi.org/10.18178/ijiet.2018.8.1.1010>
- Eagle MJ, Rooney P, Kearney JN (2007) Investigating the warming and cooling rates of human cadavers by development of a gel-filled model to validate core temperature. *Cell Tissue Bank* 8:297–302. <https://https://doi.org/10.1007/s10561-007-9038-z>
- Franchi A, Clerc-Urmès I, Martrille L (2019) Estimating the time of death with infrared tympanic thermometer: a new prospective study in France. *Int J Legal Med* 591–595. <https://https://doi.org/10.1007/s00414-019-02078-1>
- Henssge C (1988) Death time estimation in case work. I. The rectal temperature time of death nomogram. *Forensic Sci Int* 38:209–236. [https://https://doi.org/10.1016/0379-0738\(88\)90168-5](https://https://doi.org/10.1016/0379-0738(88)90168-5)
- Henssge C (1992) Rectal temperature time of death nomogram: dependence of corrective factors on the body weight under stronger thermic insulation conditions. *Forensic Sci Int* 54:51–66.
- Henssge C, Althaus L, Bolt J, et al (2000) Experiences with a compound method for estimating the time since death. I. Rectal temperature nomogram for time since death. *Int J Legal Med* 113:303–19.
- Henssge C, Madea B, Gallenkemper E (1988) Death time estimation in case work. II. Integration of different methods. *Forensic Sci Int* 39:77–87.
- Honjyo K, Yonemitsu K, Tsunenari S (2005) Estimation of early postmortem intervals by a multiple regression analysis using rectal temperature and non-temperature based postmortem changes. *J Clin Forensic Med* 12:249–253. <https://https://doi.org/10.1016/j.jcfm.2005.02.003>
- Hubig M, Muggenthaler H, Mall G (2011) Influence of measurement errors on temperature-based death time determination. *Int J Legal Med* 125:503–517. <https://https://doi.org/10.1007/s00414-010-0453-5>
- Hubig M, Muggenthaler H, Sinicina I, Mall G (2011) Body mass and corrective factor: Impact on temperature-based death time estimation. *Int J Legal Med* 125:437–444. <https://https://doi.org/10.1007/s00414-011-0551-z>
- Hubig M, Muggenthaler H, Sinicina I, Mall G (2015) Temperature based forensic death time estimation: The standard model in experimental test. *Leg Med* 17:381–387. <https://https://doi.org/10.1016/j.legalmed.2015.05.005>
- Kaliszán M (2011) Does a Draft Really Influence Postmortem Body Cooling? *J Forensic Sci* 56:1310–1314. <https://https://doi.org/10.1111/j.1556-4029.2011.01801.x>
- Kaliszán M, Hauser R, Kaliszán R, et al (2005) Verification of the exponential model of body temperature decrease after death in pigs. *Exp Physiol* 90:727–738. <https://https://doi.org/10.1113/expphysiol.2005.030551>
- Karhunen PJ, Goebeler S, Winberg O, Tuominen M (2008) Time of death of victims found in cold water environment. *Forensic Sci Int* 176:17–22. <https://https://doi.org/10.1016/j.forsciint.2007.06.014>

- Krap T, Meurs J, Duijst W (2015) Technical note: unsafe rectal temperature measurements due to delayed warming of the thermocouple by using a condom. An issue concerning the estimation of the postmortem interval by using Henßge's nomogram. *Int J Leg Med* 130:447-56. <https://doi.org/10.1007/s00414-015-1186-2>
- Nelson EL (2000) Estimation of short-term postmortem interval utilizing core body temperature: A new algorithm. *Forensic Sci Int* 109:31–38. [https://doi.org/10.1016/S0379-0738\(99\)00216-9](https://doi.org/10.1016/S0379-0738(99)00216-9)

Postmortem body cooling simulation

- Bartgis C, LeBrun AM, Ma R, Zhu L (2016) Determination of time of death in forensic science via a 3-D whole body heat transfer model. *J Therm Biol* 62:109–115. <https://doi.org/10.1016/j.jtherbio.2016.07.004>
- Ferreira MS, Yanagihara JI (2009) A transient three-dimensional heat transfer model of the human body. *Int Commun Heat Mass Transf* 36:718–724. <https://doi.org/10.1016/j.icheatmasstransfer.2009.03.010>
- den Hartog EA, Lotens WA (2004) Postmortem time estimation using body temperature and finite-element computer model. *Eur J Appl Physiol* 92:734–737. <https://doi.org/10.1007/s00421-004-1128-z>
- Hubig M, Muggenthaler H, Mall G (2016) Finite element method in temperature-based death time estimation. In: Madea B (ed) *Estimation of the Time Since Death*, 3rd ed. CRC Press, Boca Raton, pp 114–133.
- Kanawaku Y, Kanetake J, Komiya A, et al (2007) Computer simulation for postmortem cooling processes in the outer ear. *Leg Med* 9:55–62. <https://doi.org/10.1016/j.legalmed.2006.09.006>
- Mall G, Eisenmenger W (2005) Estimation of time since death by heat-flow Finite-Element model. Part I: Method, model, calibration and validation. *Leg Med* 7:1–14. <https://doi.org/10.1016/j.legalmed.2004.06.006>
- Mall G, Eisenmenger W (2005) Estimation of time since death by heat-flow Finite-Element model part II: Application to non-standard cooling conditions and preliminary results in practical casework. *Leg Med* 7:69–80. <https://doi.org/10.1016/j.legalmed.2004.06.007>
- Muggenthaler H, Hubig M, Schenkl S, et al (2017) Calibration and parameter variation using a finite element model for death time estimation: The influence of the substrate. *Leg Med* 25:23–28. <https://doi.org/10.1016/j.legalmed.2016.12.007>
- Muggenthaler H, Hubig M, Schenkl S, Mall G (2017) Influence of hypo- and hyperthermia on death time estimation – A simulation study. *Leg Med* 28:10–14. <https://doi.org/10.1016/j.legalmed.2017.06.005>
- Psikuta A, Fiala D, Laschewski G, et al (2012) Validation of the Fiala multi-node thermophysiological model for UTCI application. *Int J Biometeorol* 56:443–460. <https://doi.org/10.1007/s00484-011-0450-5>
- Schenkl S, Muggenthaler H, Hubig M, et al (2017) Automatic CT-based finite element model generation for temperature-based death time estimation: feasibility study and sensitivity analysis. *Int J Legal Med* 131:699–712. <https://doi.org/10.1007/s00414-016-1523-0>
- Smart JL, Kaliszan M (2013) Use of a Finite Element Model of Heat Transport in the Human Eye to Predict Time of Death. *J Forensic Sci* 58:S69-77. <https://doi.org/10.1111/1556-4029.12022>
- Weiser M, Erdmann B, Schenkl S, et al (2018) Uncertainty in temperature-based determination of time of death. *Heat Mass Transf und Stoffuebertragung* 54:2815–2826. <https://doi.org/10.1007/s00231-018-2324-4>
- Wilk LS, Hoveling RJM, Edelman GJ, et al (2020) Reconstructing the time since death using non-invasive thermometry and numerical analysis. *Sci Adv* 6:eaba4243.

Electrical stimulation of skeletal and facial muscle

- Cantürk I, Karabiber F, Çelik S, et al (2016) An experimental evaluation of electrical skin conductivity changes in postmortem interval and its assessment for time of death estimation. *Comput Biol Med* 69:92–96. <https://doi.org/10.1016/j.compbiomed.2015.12.010>
- Henssge C, Althaus L, Bolt J, et al (2000) Experiences with a compound method for estimating the time since death. II. Integration of non-temperature-based methods. *Int J Legal Med* 113:320–331. <https://doi.org/10.1007/s004149900090>
- Henssge C, Madea B, Gallenkemper E (1985) [Determination of the time of death--integration of various partial methods]. *Z Rechtsmed* 95:185–196. <https://doi.org/10.1007/BF00201077>
- Henssge C, Madea B, Gallenkemper E (1988) Death time estimation in case work. II. Integration of different methods. *Forensic Sci Int* 39:77–87.
- Madea B (1992) Estimating time of death from measurement of the electrical excitability of skeletal muscle. *J Forensic Sci Soc* 32:117–129. [https://doi.org/10.1016/S0015-7368\(92\)73061-8](https://doi.org/10.1016/S0015-7368(92)73061-8)
- Madea B, Henssge C (1990) Electrical excitability of skeletal muscle postmortem in casework. *Forensic Sci Int* 47:207–227. [https://doi.org/10.1016/0379-0738\(90\)90291-6](https://doi.org/10.1016/0379-0738(90)90291-6)
- Madea B, Rödiger A (2006) Precision of estimating the time since death using different criteria of supravital muscular excitability. *Forensic Sci Med Pathol* 2:127–133.
- Poposka V, Janeska B, Gutevska A, Duma A (2011) Estimation of time since death through electric and chemical excitability of muscles. *Prilozi* 32:211–218.

Mechanical stimulation of skeletal muscle

- Biermann FM, Potente S (2011) The deployment of conditional probability distributions for death time estimation. *Forensic Sci Int* 210:82–86. <https://doi.org/10.1016/j.forsciint.2011.02.007>
- Henssge C, Althaus L, Bolt J, et al (2000) Experiences with a compound method for estimating the time since death. II. Integration of non-temperature-based methods. *Int J Legal Med* 113:320–331. <https://doi.org/10.1007/s004149900090>
- Henßge C, Madea B (2004) Estimation of the time since death in the early post-mortem period. *Forensic Sci Int* 144:167–175. <https://doi.org/10.1016/j.forsciint.2004.04.051>
- Henssge C, Madea B, Gallenkemper E (1988) Death time estimation in case work. II. Integration of different methods. *Forensic Sci Int* 39:77–87.
- Warther S, Sehner S, Raupach T, et al (2012) Estimation of the time since death: Post-mortem contractions of human skeletal muscles following mechanical stimulation (idiomuscular contraction). *Int J Legal Med* 126:399–405. <https://doi.org/10.1007/s00414-011-0665-3>

Biomarkers

- Adjutantis G, Coutselinis A (1972) Estimation of the time of death by potassium levels in the vitreous humour. *Forensic Sci* 1:55–60. [https://doi.org/10.1016/0300-9432\(72\)90147-1](https://doi.org/10.1016/0300-9432(72)90147-1)
- Agrawal R, Gupta P, Bhasin S, Nagar C (1983) Determination of the time of death by estimating potassium level in the cadaver vitreous humour. *Indian J Ophthalmol* 31:528–531.
- Akhtar N, Ali R, Malik AR, et al (2015) Role of vitreous potassium level in postmortem interval estimation. *Pakistan J Med Heal Sci* 9:88–92.
- Arroyo A, Rosel P, Marron T (2005) Cerebrospinal fluid: postmortem biochemistry study. *J Clin Forensic Med* 12:153-156.
- Balasooriya BAW, St Hill CA, Williams AR (1984) The biochemical changes in pericardial fluid after death. An investigation of the relationship between the time since death and the rise or fall in electrolyte and enzyme concentrations and their possible usefulness in determining the time of death. *Forensic Sci Int* 26(2):93-102.

- Balasooriya BAW, St Hill CA, Williams AR (1984) The biochemistry of vitreous humour. A comparative study of the potassium, sodium and urate concentrations in the eyes at identical time intervals after death. *Forensic Sci Int* 26(2):85-91.
- Devgun MS, Dunbar JA (1986) Biochemical investigation of vitreous: applications in forensic medicine, especially in relation to alcohol. *Forensic Sci Int* 31(1):27-34.
- Barmate N, Shrigiriwar M, Batra A, Puppalwar P (2013) A Study of Correlation of Time Since Death with Vitreous Potassium and Sodium-Potassium Ratio. *Medico-Legal Updat* 13:66. <https://doi.org/10.5958/j.0974-1283.13.2.017>
- Bindu S, Amit K, Srivastava A. (2012) Potassium Ion & Alkaline Phosphatase Level in Vitreous Humour and their Role in Postmortem Interval. *J Indian Acad Forensic Med* 34:220–222
- Bocaz-Beneventi G, Tagliaro F, Bortolotti F, et al (2002) Capillary zone electrophoresis and artificial neural networks for estimation of the post-mortem interval (PMI) using electrolytes measurements in human vitreous humour. *Int J Legal Med* 116:5–11. <https://doi.org/10.1007/s004140100239>
- Bortolotti F, Pascali JP, Davis GG, et al (2011) Study of vitreous potassium correlation with time since death in the postmortem range from 2 to 110 hours using capillary ion analysis. *Med Sci Law* 51:20–23. <https://doi.org/10.1258/msl.2010.010063>
- Bray M (1984) The effect of chilling, freezing, and rewarming on the postmortem chemistry of vitreous humor. *J Forensic Sci* 29(2):404-411.
- Chandrakanth HV, Kanchan T, Balaraj BM et al (2013) Postmortem vitreous chemistry-An evaluation of sodium, potassium and chloride levels in estimation of time since death (during the first 36 h after death). *J Forensic Leg Med* 20:211–216. <https://doi.org/10.1016/j.jflm.2012.09.001>
- Coe JI (1973) Further thoughts and observations on postmortem chemistry. *Forensic Sci Gazette* 5:2-9.
- Coe JI (1969) Postmortem chemistries on human vitreous humor. *Am J Clin Pathol* 51(6):741-750.
- Coe JI (1989) Vitreous potassium as a measure of the postmortem interval: An historical review and critical evaluation. *Forensic Sci Int* 42:201–213. [https://doi.org/10.1016/0379-0738\(89\)90087-X](https://doi.org/10.1016/0379-0738(89)90087-X)
- Cordeiro C, Seoane R, Camba A et al (2015) The application of flow cytometry as a rapid and sensitive screening method to detect contamination of vitreous humor samples and avoid miscalculation of the postmortem interval. *J Forensic Sci* 60(5):1346-1349.
- Dhanak K, Johry A (2016) A Study on the Vitreous Potassium Ion Concentration in Relation with Time Since Death. *Indian J Forensic Med Toxicol* 10:250. <https://doi.org/10.5958/0973-9130.2016.00105.5>
- Ding Y, Li X, Guo Y, et al (2017) Estimation of postmortem interval by vitreous potassium evaluation with a novel fluorescence aptasensor. *Sci Rep* 7:. <https://doi.org/10.1038/s41598-017-02027-1>
- Farmer JG, Benomran F, Watson AA, Harland WA (1985) Magnesium, potassium, sodium and calcium in post-mortem vitreous humour from humans. *Forensic Sci Int* 27:1–13. [https://doi.org/10.1016/0379-0738\(85\)90099-4](https://doi.org/10.1016/0379-0738(85)90099-4)
- Foster SN, Smith PR, Biggs M, et al (2016) Estimation of postmortem interval using vitreous potassium levels in cases of fatal road traffic collision. *Arch Med Sadovej Kryminol* 66:71–82. <https://doi.org/10.5114/amsik.2016.63866>
- Gadzuric SB, Podunavac Kuzmanovic SO, Jokic AI, et al (2014) Chemometric estimation of post-mortem interval based on Na⁺ and K⁺ concentrations from human vitreous humour by linear least squares and artificial neural networks modelling. *Aust J Forensic Sci* 46:166–179. <https://doi.org/10.1080/00450618.2013.825812>
- Gamero Lucas JJ, Romero JL, Ramos HM, et al (1992) Precision of estimating time of death by vitreous potassium - Comparison of various equations. *Forensic Sci Int* 56:137–145. [https://doi.org/10.1016/0379-0738\(92\)90171-R](https://doi.org/10.1016/0379-0738(92)90171-R)

- Garg V, Oberoi S, Gorea R, Kaur K (2004) Changes in the levels of vitreous potassium with increasing time since death. *J Indian Acad Forensic Med* 26:136–139
- George GO, Bosede Ajayi O (2016) Investigation of Electrolyte Changes in Bovine Vitreous at Different Postmortem Interval. *J Clin Exp Ophthalmol* 07: <https://doi.org/10.4172/2155-9570.1000609>
- Hansson L, Uotila U, Lindfors R et al (1966) Potassium content of the vitreous body as an aid in determining the time of death. *J Forensic Sci* 11(3):390-394.
- Hughes WM (1965) Levels of potassium in the vitreous humour after death. *Med Sci Law* 5:150–156. <https://doi.org/10.1177/002580246500500306>
- James RA, Hoadley PA, Sampson BG (1997) Determination of Postmortem Interval by Sampling Vitreous Humour. *Am J Forensic Med Pathol* 18(2):158-162
- Jashnani KD, Kale SA, Rupani AB (2010) Vitreous humor: Biochemical constituents in estimation of postmortem interval. *J Forensic Sci* 55:1523–1527. <https://doi.org/10.1111/j.1556-4029.2010.01501.x>
- Kawamoto O, Michiue T, Ishikawa T et al (2013) Comprehensive evaluation of pericardial biochemical markers in death investigation. *Forensic Sci Int* 224(1):73-79.
- Komura S, Oshiro S. Potassium levels in the aqueous and vitreous humor after death. *Tohoku J Exp Med* 122(1):65-68.
- Jashnani KD, Kale SA, Rupani AB (2010) Vitreous humor: Biochemical constituents in estimation of postmortem interval. *J Forensic Sci* 55(6):1523-1527.
- Kulkarni C, Chavan G, Jadhao A (2016) Medicolegal study of determination of time since death from potassium level in vitreous humour. *Int J Adv Med* 3:1059–1064. <https://doi.org/10.18203/2349-3933.ijam20163749>
- Lange N, Swearer S, Sturner WQ (1994) Human postmortem interval estimation from vitreous potassium: an analysis of original data from six different studies. *Forensic Sci Int* 66:159–174. [https://doi.org/10.1016/0379-0738\(94\)90341-7](https://doi.org/10.1016/0379-0738(94)90341-7)
- Lie JT (1967) Changes of potassium concentration in the vitreous humor after death. *Am J Med Sci* 254:136–143
- Madea B (2005) Is there recent progress in the estimation of the postmortem interval by means of thanatochemistry? *Forensic Sci Int* 151:139–149. <https://doi.org/10.1016/j.forsciint.2005.01.013>
- Madea B, Henssge C, Hönig W, Gerbracht A (1989) References for determining the time of death by potassium in vitreous humor. *Forensic Sci Int* 40:231–243. [https://doi.org/10.1016/0379-0738\(89\)90181-3](https://doi.org/10.1016/0379-0738(89)90181-3)
- Madea B, Herrmann N, Henssge C (1990) Precision of estimating the time since death by vitreous potassium — comparison of two different equations. *Forensic Sci Int* 46:277–284
- Madea B, Käferstein H, Hermann N, Sticht G (1994) Hypoxanthine in vitreous humor and cerebrospinal fluid - a marker of postmortem interval and prolonged (vital) hypoxia? Remarks also on hypoxanthine in SIDS. *Forensic Sci Int* 65:19–31. [https://doi.org/10.1016/0379-0738\(94\)90296-8](https://doi.org/10.1016/0379-0738(94)90296-8)
- Madea B, Kreuser C, Banaschak S (2001) Postmortem biochemical examination of synovial fluid: a preliminary study. *Forensic Sci Int* 118(1):29-35.
- Madea B, Rödiger A (2006) Time of death dependent criteria in vitreous humor-Accuracy of estimating the time since death. *Forensic Sci Int* 164:87–92. <https://doi.org/10.1016/j.forsciint.2005.12.002>
- Meurs J, Krap T, Duijst W (2019) Evaluation of postmortem biochemical markers: Completeness of data and assessment of implication in the field. *Sci Justice* 59:177–180.
- Mihailovic Z, Atanasijevic T, Popovic V, et al (2012) Estimation of the postmortem interval by analyzing potassium in the vitreous humor: Could repetitive sampling enhance accuracy? *Am J Forensic Med Pathol* 33:400–403. <https://doi.org/10.1097/PAF.0b013e31826627d0>

- Mulla A, Massey KL, Kalra J (2005) Vitreous humor biochemical constituents: Evaluation of between-eye differences. *Am J Forensic Med Pathol* 26:146–149. <https://doi.org/10.1097/01.paf.0000164001.50097.4c>
- Muñoz JI, Suárez-Peñaranda JM, Otero XL, et al (2001) A new perspective in the estimation of postmortem interval (PMI) based on vitreous. *J Forensic Sci* 46:209–214
- Muñoz Barús JI, Suárez-Peñaranda JM, Otero XL, et al (2002) Improved estimation of postmortem interval based on differential behaviour of vitreous potassium and hypoxanthine in death by hanging. *Forensic Sci Int* 125:67–74. [https://doi.org/10.1016/S0379-0738\(01\)00616-8](https://doi.org/10.1016/S0379-0738(01)00616-8)
- Myo-Thaik-Oo, Tanaka E, Oikawa H, et al (2002) No significant differences in the postmortem interval in Myanmar and Japanese using vitreous potassium levels. *J Clin Forensic Med* 9:70–73. <https://doi.org/10.1054/jcfm.2002.0552>
- Ortmann J, Markwerth P, Madea B (2016) Precision of estimating the time since death by vitreous potassium—Comparison of 5 different equations. *Forensic Sci Int* 269:1–7. <https://doi.org/10.1016/j.forsciint.2016.10.005>
- Passos MLC, Santos AM, Pereira AI, Santos JR et al. (2009) Estimation of postmortem interval by hypoxanthine and potassium evaluation in vitreous humor with a sequential injection system. *Talanta*, 79(4):1094-1099.
- Patel U, Patel J, Prajapati P, Govekar G (2016) A study to evaluate of time since death from potassium level of vitreous humour. *Natl J Med Res* 6:255–258
- Prasad BK, Choudhary A, Sinha JN (2003) A study of correlation between vitreous potassium level and post mortem interval. *Kathmandu Univ Med J (KUMJ)* 1:132–134
- Prashad R, Parkash C, Bansal YS, Sharma SK et al. (2002) Linearization of the relationship between serum sodium, potassium concentration, their ratio and time since death in Chandigarh zone of north-west India. *Forensic Sci Int* 130(1):1-7.
- Ravindra D, Ajay S (2013) Estimation of post mortem interval by means of changes in the eye-vitreous humour potassium levels. *Indian J Forensic Med Toxicol* 7:232–236. <https://doi.org/10.5958/j.0973-9130.7.2.054>
- Rognum TO, Hauge S, Øyasaeter S, Saugstad OD (1991) A new biochemical method for estimation of postmortem time. *Forensic Sci Int* 51:139–146. [https://doi.org/10.1016/0379-0738\(91\)90214-4](https://doi.org/10.1016/0379-0738(91)90214-4)
- Rognum TO, Holmen S, Musse MA, et al (2016) Estimation of time since death by vitreous humor hypoxanthine, potassium, and ambient temperature. *Forensic Sci Int* 262:160–165. <https://doi.org/10.1016/j.forsciint.2016.03.001>
- Salam HA, Shaat EA, Aziz MHA, et al (2012) Estimation of postmortem interval using thanatochemistry and postmortem changes. *Alexandria J Med* 48:335–344. <https://doi.org/10.1016/j.ajme.2012.05.004>
- Sheikh NA (2007) Estimation of postmortem interval according to time course of potassium ion activity in cadaveric synovial fluid. *Indian J Med Forensic Med Toxicol* 1(2):45-49.
- Siddamsetty AK, Verma SK, Kohli A et al. (2014) Exploring time of death from potassium, sodium, chloride, glucose & calcium analysis of postmortem synovial fluid in semi-arid climate. *J Clin Forensic Med* 28:11-14.
- Siddamsetty AK, Verma SK, Kohli A, et al (2014) Estimation of time since death from electrolyte, glucose and calcium analysis of postmortem vitreous humour in semi-arid climate. *Med Sci Law* 54:158–166. <https://doi.org/10.1177/0025802413506424>
- Singh A, Goel S, Singh N, et al (2015) Vitreous potassium concentration as a predictor of postmortem interval in injury (trauma) cases: A cross-sectional study from a tertiary care center in rural Haryana. *Med J Dr DY Patil Univ* 8:315. <https://doi.org/10.4103/0975-2870.157072>
- Singh D, Prashad R, Sharma SK et al. (2006) Estimation of postmortem interval from human pericardial fluid electrolytes concentrations in Chandigarh zone of India: log transformed linear regression model. *Leg Med* 8(5):279-287.

- Sparks DL, Oeltgen PR, Kryscio RJ et al. (1989) Comparison of chemical methods for determining postmortem interval. *J Forensic Sci* 34(1):197-206.
- Stephens RJ, Richards RG (1987) Vitreous humor chemistry: the use of potassium concentration for the prediction of the postmortem interval. *J Forensic Sci* 32:503–509.
- Sturmer WQ (1963) the Vitreous Humour: Postmortem Potassium Changes. *Lancet* 281:807–808. [https://doi.org/10.1016/S0140-6736\(63\)91509-5](https://doi.org/10.1016/S0140-6736(63)91509-5)
- Sturmer WQ, Gantner GE Jr (1964) The postmortem interval. a study of potassium in the vitreous humor. *Am J Clin Pathol* 42:137-144.
- Swain R, Kumar A, Sahoo J, et al (2015) Estimation of post-mortem interval: A comparison between cerebrospinal fluid and vitreous humour chemistry. *J Forensic Leg Med* 36:144–148. <https://doi.org/10.1016/j.jflm.2015.09.017>
- Tagliaro F, Bortolotti F, Manetto G, et al (2001) Potassium concentration differences in the vitreous humour from the two eyes revisited by microanalysis with capillary electrophoresis. *J Chromatogr A* 924:493–498. [https://doi.org/10.1016/S0021-9673\(01\)00713-0](https://doi.org/10.1016/S0021-9673(01)00713-0)
- Tagliaro F, Manetto G, Cittadini F, et al (1999) Capillary zone electrophoresis of potassium in human vitreous humour: Validation of a new method. *J Chromatogr B Biomed Sci Appl* 733:273–279. [https://doi.org/10.1016/S0378-4347\(99\)00338-2](https://doi.org/10.1016/S0378-4347(99)00338-2)
- Tumram NK, Ambade VN, Dongre AP (2014) Thanatochemistry: Study of vitreous humor potassium. *Alexandria J Med* 50:365–368. <https://doi.org/10.1016/j.ajme.2013.12.002>
- Tumram NK, Bardale RV, Dongre AP (2011) Postmortem analysis of synovial fluid and vitreous humour for determination of death interval: A comparative study. *Forensic Sci Int* 204:186–190. <https://doi.org/10.1016/j.forsciint.2010.06.007>
- Yadav J, Deshpande A, Arora A et al. (2007) Estimation of time since death from CSF electrolyte concentration in Bhopal region of central India. *Leg Med* 9(6):309-313.
- Zhou B, Zhang L, Zhang G, et al (2007) The determination of potassium concentration in vitreous humor by low pressure ion chromatography and its application in the estimation of postmortem interval. *J Chromatogr B Anal Technol Biomed Life Sci* 852:278–281. <https://doi.org/10.1016/j.jchromb.2007.01.027>
- Zilg B, Bernard S, Alkass K, et al (2015) A new model for the estimation of time of death from vitreous potassium levels corrected for age and temperature. *Forensic Sci Int*. <https://doi.org/10.1016/j.forsciint.2015.07.020>

Decomposition

- Bachmann J, Simmons T (2010) The influence of preburial insect access on the decomposition rate. *J Forensic Sci* 55:893–900. <https://doi.org/10.1111/j.1556-4029.2010.01403.x>
- Bugelli V, Gherardi M, Focardi M, et al (2018) Decomposition pattern and insect colonization in two cases of suicide by hanging. *Forensic Sci Res* 3:94–102.
- Card A, Cross P, Moffatt C, Simmons T (2015) The Effect of Clothing on the Rate of Decomposition and Diptera Colonization on *Sus scrofa* Carcasses. *J Forensic Sci* 60:979–982.
- Ceciliason AS, Andersson MG, Lindström A, Sandler H (2018) Quantifying human decomposition in an indoor setting and implications for postmortem interval estimation. *Forensic Sci Int* 283:180–189. <https://doi.org/10.1016/j.forsciint.2017.12.026>
- Cockle DL, Bell LS (2015) Human decomposition and the reliability of a “Universal” model for post mortem interval estimations. *Forensic Sci Int* 253:136.e1-136.e9. <https://doi.org/10.1016/j.forsciint.2015.05.018>
- Cockle DL, Bell LS (2017) The environmental variables that impact human decomposition in terrestrially exposed contexts within Canada. *Sci Justice* 57:107–117. <https://doi.org/10.1016/j.scijus.2016.11.001>
- Connor M, Baigent C, Hansen ES (2018) Testing the Use of Pigs as Human Proxies in Decomposition Studies. *J Forensic Sci* 63:1350–1355. <https://doi.org/10.1111/1556-4029.13727>

- Connor M, Baigent C, Hansen ES (2019) Measuring Desiccation Using Qualitative Changes: A Step Toward Determining Regional Decomposition Sequences. *J Forensic Sci* 64:1004-1011. <https://doi.org/10.1111/1556-4029.14003>
- Cross P, Simmons T (2010) The influence of penetrative trauma on the rate of decomposition. *J Forensic Sci* 55:295–301. <https://doi.org/10.1111/j.1556-4029.2009.01277.x>
- van Daalen MA, de Kat DS, Oude Grotebevelsberg BFL, et al (2017) An Aquatic Decomposition Scoring Method to Potentially Predict the Postmortem Submersion Interval of Bodies Recovered from the North Sea. *J Forensic Sci* 62:369–373. <https://doi.org/10.1111/1556-4029.13258>
- Dabbs GR, Bytheway JA, Connor M (2017) Comparing the Scoring of Human Decomposition from Digital Images to Scoring Using On-site Observations. *J Forensic Sci* 62:1292–1296. <https://doi.org/10.1111/1556-4029.13409>
- Dabbs GR, Connor M, Bytheway JA (2016) Interobserver Reliability of the Total Body Score System for Quantifying Human Decomposition. *J Forensic Sci* 61:445-451. <https://doi.org/10.1111/1556-4029.12962>
- Dautartas A, Kenyhercz MW, Vidoli GM, et al (2018) Differential Decomposition Among Pig, Rabbit, and Human Remains. *J Forensic Sci* 63:1673-1683. <https://doi.org/10.1111/1556-4029.13784>
- Doberentz E, Madea B (2010) Estimating the time of immersion of bodies found in water - an evaluation of a common method to estimate the minimum time interval of immersion. *Rev Española Med Leg* 36:51–61. [https://doi.org/10.1016/S0377-4732\(10\)70045-6](https://doi.org/10.1016/S0377-4732(10)70045-6)
- De Donno A, Campobasso CP, Santoro V, et al (2014) Bodies in sequestered and non-sequestered aquatic environments: A comparative taphonomic study using decompositional scoring system. *Sci Justice* 54:439–446. <https://doi.org/10.1016/j.scijus.2014.10.003>
- Finaughty DA, Morris AG (2019) Precocious natural mummification in a temperate climate (Western Cape, South Africa). *Forensic Sci Int* 303:109948. <https://doi.org/10.1016/j.forsciint.2019.109948>
- Fitzgerald CM, Oxenham M (2009) Modelling time-since-death in Australian temperate conditions. *Aust J Forensic Sci* 41:27–41. <https://doi.org/10.1080/00450610902935981>
- Forbes MNS, Finaughty DA, Miles KL, Gibbon VE (2019) Inaccuracy of accumulated degree day models for estimating terrestrial post-mortem intervals in Cape Town, South Africa. *Forensic Sci Int* 296:67–73. <https://doi.org/10.1016/j.forsciint.2019.01.008>
- Galloway A, Birkby WH, Jones AM, et al (1989) Decay rates of human remains in an Arid Environment. *J Forensic Sci* 34:607–616.
- Gelderman HT, Boer L, Naujocks T, et al (2018) The development of a post-mortem interval estimation for human remains found on land in the Netherlands. *Int J Legal Med* 132:863-873. <https://doi.org/10.1007/s00414-017-1700-9>
- Gelderman HT, Kruiver CA, Oostra RJ, et al (2019) Estimation of the postmortem interval based on the human decomposition process. *J Forensic Leg Med* 61:122-127. <https://doi.org/10.1016/j.jflm.2018.12.004>
- Gruenthal A, Moffatt C, Simmons T (2012) Differential Decomposition Patterns in Charred Versus Un-Charred Remains. *J Forensic Sci* 57:12–18. <https://doi.org/10.1111/j.1556-4029.2011.01909.x>
- Hauteo C, Hamzah PA, Osman K, et al (2013) Post mortem changes in relation to different types of clothing. *Malaysian J Pathol* 35:77-85.
- Heaton V, Lagden A, Moffatt C, Simmons T (2010) Predicting the postmortem submersion interval for human remains recovered from U.K. waterways. *J Forensic Sci* 55:302–307. <https://doi.org/10.1111/j.1556-4029.2009.01291.x>
- Humphreys MK, Panacek E, Green W, Albers E (2013) Comparison of protocols for measuring and calculating postmortem submersion intervals for human analogs in fresh water. *J Forensic Sci* 58:513–517. <https://doi.org/10.1111/1556-4029.12033>

- Keough N, Myburgh J, Steyn M (2017) Scoring of Decomposition: A Proposed Amendment to the Method When Using a Pig Model for Human Studies. *J Forensic Sci* 62:986–993. <https://doi.org/10.1111/1556-4029.13390>
- Keyes CA (2019) How reliable is the charred body scale? An interobserver reliability study on scoring burned remains. *Burns* 45:1673–1679. <https://doi.org/10.1016/j.burns.2019.02.012>
- Lynch-Aird J, Moffatt C, Simmons T (2015) Decomposition Rate and Pattern in Hanging Pigs. *J Forensic Sci* 60:1155-63. <https://doi.org/10.1111/1556-4029.12796>
- Maile AE, Inoue CG, Barksdale LE, Carter DO (2017) Toward a universal equation to estimate postmortem interval. *Forensic Sci Int* 272:150–153. <https://doi.org/10.1016/j.forsciint.2017.01.013>
- Marais-Werner A, Myburgh J, Becker PJ, Steyn M (2018) A comparison between decomposition rates of buried and surface remains in a temperate region of South Africa. *Int J Legal Med* 132:301–309. <https://doi.org/10.1007/s00414-017-1618-2>
- Marais-Werner A, Myburgh J, Meyer A, et al (2017) Decomposition patterns of buried remains at different intervals in the Central Highveld region of South Africa. *Med Sci Law* 57:115–123. <https://doi.org/10.1177/0025802417705263>
- Marhoff SJ, Fahey P, Forbes SL, Green H (2016) Estimating post-mortem interval using accumulated degree-days and a degree of decomposition index in Australia: A validation study. *Aust J Forensic Sci* 48:24–36. <https://doi.org/10.1080/00450618.2015.1021378>
- Marhoff-Beard SJ, Forbes SL, Green H (2018) The validation of ‘universal’ PMI methods for the estimation of time since death in temperate Australian climates. *Forensic Sci Int* 291:158–166. <https://doi.org/10.1016/j.forsciint.2018.08.022>
- Matuszewski S, Konwerski S, Frątczak K, Szafałowicz M (2014) Effect of body mass and clothing on decomposition of pig carcasses. *Int J Legal Med* 128:1039-48. <https://doi.org/10.1007/s00414-014-0965-5>
- Megyesi MS, Nawrocki SP, Haskell NH (2005) Using Accumulated Degree-Days to Estimate the Postmortem Interval from Decomposed Human Remains. *J Forensic Sci* 50:618-26. <https://doi.org/10.1520/JFS2004017>
- Meyer J, Anderson B, Carter DO (2013) Seasonal variation of carcass decomposition and gravesoil chemistry in a cold (Dfa) climate. *J Forensic Sci* 58:1175–1182. <https://doi.org/10.1111/1556-4029.12169>
- Moffatt C, Simmons T, Lynch-Aird J (2016) An Improved Equation for TBS and ADD: Establishing a Reliable Postmortem Interval Framework for Casework and Experimental Studies. *J Forensic Sci* 61:201–207. <https://doi.org/10.1111/1556-4029.12931>
- Myburgh J, L’Abbé EN, Steyn M, Becker PJ (2013) Estimating the postmortem interval (PMI) using accumulated degree-days (ADD) in a temperate region of South Africa. *Forensic Sci Int* 229:165.e1-165.e6. <https://doi.org/10.1016/j.forsciint.2013.03.037>
- Nawrocka M, Frątczak K, Matuszewski S (2016) Inter-Rater Reliability of Total Body Score-A Scale for Quantification of Corpse Decomposition. *J Forensic Sci* 61:798-802. <https://doi.org/10.1111/1556-4029.13105>
- Palazzo C, Pelletti G, Fais P, et al (2020) Postmortem submersion interval in human bodies recovered from fresh water in an area of Mediterranean climate. Application and comparison of preexisting models. *Forensic Sci Int* 306:110051. <https://doi.org/10.1016/j.forsciint.2019.110051>
- Parks CL (2011) A Study of the Human Decomposition Sequence in Central Texas. *J Forensic Sci* 56:19–22. <https://doi.org/10.1111/j.1556-4029.2010.01544.x>
- Reh H (1967) Anhaltspunkte für die Bestimmung der Wasserzeit. *Dtsch Z Gesamte Gerichtl Med* 59:235–245.
- Reh H, Haarhoff K, Vogt C (1977) Die Schätzung der Todeszeit bei Wasserleichen (The estimation of the time of death of corpses recovered from water). *Zeitschrift für Rechtsmedizin* 79:261–266.

- Reijnen G, Gelderman HT, Oude Grotebevelsberg BFL, et al (2018) The correlation between the Aquatic Decomposition Score (ADS) and the post-mortem submersion interval measured in Accumulated Degree Days (ADD) in bodies recovered from fresh water. *Forensic Sci Med Pathol* 14:301-306. [https://https://doi.org/10.1007/s12024-018-9987-5](https://doi.org/10.1007/s12024-018-9987-5)
- Ribéreau-Gayon A, Rando C, Morgan RM, Carter DO (2018) The suitability of visual taphonomic methods for digital photographs: An experimental approach with pig carcasses in a tropical climate. *Sci Justice* 58:167–176. [https://https://doi.org/10.1016/j.scijus.2017.12.001](https://doi.org/10.1016/j.scijus.2017.12.001)
- Roberts LG, Dabbs GR (2015) A Taphonomic Study Exploring the Differences in Decomposition Rate and Manner between Frozen and Never Frozen Domestic Pigs (*Sus scrofa*). *J Forensic Sci* 60:588–594. [https://https://doi.org/10.1111/1556-4029.12717](https://doi.org/10.1111/1556-4029.12717)
- Roberts LG, Spencer JR, Dabbs GR (2017) The Effect of Body Mass on Outdoor Adult Human Decomposition. *J Forensic Sci* 62:1145-1150. [https://https://doi.org/10.1111/1556-4029.13398](https://doi.org/10.1111/1556-4029.13398)
- Ross, AH, Hale AR (2018) Decomposition of juvenile-sized remains: a macro- and microscopic perspective. *Forensic Sci Res* 3:294–303. [https://https://doi.org/10.1080/20961790.2018.1489362](https://doi.org/10.1080/20961790.2018.1489362)
- Simmons T, Adlam RE, Moffatt C (2010) Debugging decomposition data - Comparative taphonomic studies and the influence of insects and carcass size on decomposition rate. *J Forensic Sci* 55:8-13. [https://https://doi.org/10.1111/j.1556-4029.2009.01206.x](https://doi.org/10.1111/j.1556-4029.2009.01206.x)
- Simmons T, Cross PA, Adlam RE, Moffatt C (2010) The influence of insects on decomposition rate in buried and surface remains. *J Forensic Sci* 55:889–892. [https://https://doi.org/10.1111/j.1556-4029.2010.01402.x](https://doi.org/10.1111/j.1556-4029.2010.01402.x)
- Smith AC (2014) The Effects of Sharp-Force Thoracic Trauma on the Rate and Pattern of Decomposition. *J Forensic Sci* 59:319–326. [https://https://doi.org/10.1111/1556-4029.12338](https://doi.org/10.1111/1556-4029.12338)
- [https://https://doi.org/10.1111/1556-4029.12750](https://doi.org/10.1111/1556-4029.12750)
- Steadman DW, Dautartas A, Kenyhercz MW, et al (2018) Differential Scavenging Among Pig, Rabbit, and Human Subjects,. *J Forensic Sci* 63:1684–1691. [https://https://doi.org/10.1111/1556-4029.13786](https://doi.org/10.1111/1556-4029.13786)
- Suckling JK, Spradley MK, Godde K (2016) A Longitudinal Study on Human Outdoor Decomposition in Central Texas. *J Forensic Sci* 61:19-25. [https://https://doi.org/10.1111/1556-4029.12892](https://doi.org/10.1111/1556-4029.12892)
- Sutherland A, Myburgh J, Steyn M, Becker PJ (2013) The effect of body size on the rate of decomposition in a temperate region of South Africa. *Forensic Sci Int* 231:257–262. [https://https://doi.org/10.1016/j.forsciint.2013.05.035](https://doi.org/10.1016/j.forsciint.2013.05.035)
- Troutman L, Moffatt C, Simmons T (2014) A preliminary examination of differential decomposition patterns in mass graves. *J Forensic Sci* 59:621–626. [https://https://doi.org/10.1111/1556-4029.12388](https://doi.org/10.1111/1556-4029.12388)
- Vass AA (2011) The elusive universal post-mortem interval formula. *Forensic Sci Int* 204:34–40. [https://https://doi.org/10.1016/j.forsciint.2010.04.052](https://doi.org/10.1016/j.forsciint.2010.04.052)
- Weger M, Cross P, Cunliffe R (2020) The effect of plastic bag containment of the head on the rate and pattern of decomposition. *J Forensic Leg Med* 71:101922. [https://https://doi.org/10.1016/j.jflm.2020.101922](https://doi.org/10.1016/j.jflm.2020.101922)
- Wescott D, Steadman D, Miller N, et al (2018) Validation of the Total Body Score/Accumulated Degree-Day Model at Three Human Decomposition Facilities. *Forensic Anthropol* 1:143–149. [https://https://doi.org/10.5744/fa.2018.0015](https://doi.org/10.5744/fa.2018.0015)

Entomology

- Aballay F, Domínguez MC, Campón FF (2012) Adult Fanniidae associated to pig carcasses during the winter season in a semiarid environment: Initial examination of their potential as complementary PMI indicators. *Forensic Sci Int* 219:284.e1-284.e4. [https://https://doi.org/10.1016/j.forsciint.2011.11.019](https://doi.org/10.1016/j.forsciint.2011.11.019)

- Adams ZJ, Hall MJ. (2003) Methods used for the killing and preservation of blowfly larvae, and their effect on post-mortem larval length. *Forensic Sci Int* 138:50-61.
- Ahmad NW, Lim LH, Dhang CC, Chin HC, Abdullah AG, Norjuliana W, Mustaffa W, Kian CW, Jeffery J, Hashim R, Azirun SM (2011) Comparative insect fauna succession on indoor and outdoor monkey carrions in a semi-forested area in Malaysia. *Asian Pacific Journal of Tropical Biomedicine* 1:S232-S238. [https://doi.org/10.1016/S2221-1691\(11\)60161-5](https://doi.org/10.1016/S2221-1691(11)60161-5).
- Amendt J, Campobasso CP, Gaudry E, Reiter C, LeBlanc HN, Hall MJ (2007) Best practice in forensic entomology--standards and guidelines. *Int J Legal Med.* 121:90-104.
- Ames C, Turner BD, Daniel B (2006) Estimating the post-mortem interval (II) - The use of differential temporal gene expression to determine the age of blowfly pupae. *International Congress Series, Volume 1288*:861-863. <https://doi.org/10.1016/j.ics.2005.09.091>.
- Anderson GS (2004) Determining time of death using blow fly eggs in the early postmortem interval. *Int J Legal Med* 118:240-241.
- Andersson MG, Sundström A, Lindström A (2013) Bayesian networks for evaluation of evidence from forensic entomology. *Biosecur Bioterror.* 11 (Suppl 1):S64-77. <https://doi.org/10.1089/bsp.2012.0081>
- Archer MS, Jones SD, Wallman JF (2017) Delayed reception of live blowfly (*Calliphora vicina* and *Chrysomya rufifacies*) larval samples: implications for minimum postmortem interval estimates. *Forensic Sci Res.* 3:27-39. <https://doi.org/10.1080/20961790.2017.1408550>.
- Arnott S, Turner B (2008) Post-feeding larval behaviour in the blowfly, *Calliphora vicina*: effects on post-mortem interval estimates. *Forensic Sci Int* 177:162-167. <https://doi.org/10.1016/j.forsciint.2007.12.002>
- Aubernon C, Charabidzé D, Devigne C, Delannoy Y, Gosset D (2015) Experimental study of *Lucilia sericata* (Diptera Calliphoridae) larval development on rat cadavers: Effects of climate and chemical contamination. *Forensic Sci Int* 253:125-30. <https://doi.org/10.1016/j.forsciint.2015.05.032>.
- Avila FW, Goff ML (1998) Arthropod succession patterns onto burnt carrion in two contrasting habitats in the Hawaiian Islands. *J Forensic Sci* 43:581-586.
- Azwandi A, Nina Keterina H, Owen LC, Nurizzati MD, Omar B (2013) Adult carrion arthropod community in a tropical rainforest of Malaysia: analysis on three common forensic entomology animal models. *Trop Biomed.* 30:481-494.
- Baqué M, Amendt J (2013) Strengthen forensic entomology in court--the need for data exploration and the validation of a generalised additive mixed model. *Int J Legal Med* 127:213-223. <https://doi.org/10.1007/s00414-012-0675-9>.
- Baqué M, Filmann N, Verhoff MA, Amendt J (2015) Establishment of developmental charts for the larvae of the blow fly *Calliphora vicina* using quantile regression. *Forensic Sci Int* 248:1-9. <https://doi.org/10.1016/j.forsciint.2014.12.020>.
- Barrios M, Wolff M (2011) Initial study of arthropods succession and pig carrion decomposition in two freshwater ecosystems in the Colombian Andes. *Forensic Sci Int* 212:164-172. <https://doi.org/10.1016/j.forsciint.2011.06.008>.
- Barros LM, Ferreira-Keppler RL, Martins RT, Gutjahr ALN (2019) Bionomy of *Hermetia illucens* (Diptera: Stratiomyidae) on Decomposing Swine Carcass in an Urban Area of Central Amazon. *J Med Entomol* 56:681-689. <https://doi.org/10.1093/jme/tjz005>.
- Bauer A, Bauer AM, Tomberlin JK (2020) Impact of diet moisture on the development of the forensically important blow fly *Cochliomyia macellaria* (Fabricius) (Diptera: Calliphoridae). *Forensic Sci Int* 312:110333. <https://doi.org/10.1016/j.forsciint.2020.110333>
- Benbow ME, Lewis AJ, Tomberlin JK, Pechal JL (2013) Seasonal necrophagous insect community assembly during vertebrate carrion decomposition. *J Med Entomol* 50:440-450.
- Berg MC, Benbow ME (2013) Environmental factors associated with *Phormia regina* (Diptera: Calliphoridae) oviposition. *J Med Entomol* 50:451-457.

- Bernhardt V, Hannig L, Kinast R, Verhoff MA, Rothweiler F, Zehner R, Amendt J (2017) Quantitative pteridine fluorescence analysis: A possible age-grading technique for the adult stages of the blow fly *Calliphora vicina* (Diptera: Calliphoridae). *J Insect Physiol.* 98:356-359. <https://doi.org/10.1016/j.jinsphys.2017.03.002>.
- Bernhardt V, Pogoda W, Verhoff MA, Toennes SW, Amendt J (2017) Estimating the age of the adult stages of the blow flies *Lucilia sericata* and *Calliphora vicina* (Diptera: Calliphoridae) by means of the cuticular hydrocarbon n-pentacosane. *Science & Justice* 57:361-365. <https://doi.org/10.1016/j.scijus.2017.04.007>.
- Bernhardt V, Schomerus C, Verhoff MA, Amendt J (2017) Of pigs and men-comparing the development of *Calliphora vicina* (Diptera: Calliphoridae) on human and porcine tissue. *Int J Legal Med* 131:847-853. <https://doi.org/10.1007/s00414-016-1487-0>.
- Beuter L, Mendes J (2013) Development of *Chrysomya albiceps* (Wiedemann) (Diptera: Calliphoridae) in different pig tissues. *Neotrop Entomol* 42:426-430. <https://doi.org/10.1007/s13744-013-0134-4>.
- Boehme P, Spahn P, Amendt J, Zehner R. (2013) Differential gene expression during metamorphosis: a promising approach for age estimation of forensically important *Calliphora vicina* pupae (Diptera: Calliphoridae). *Int J Legal Med.* 127:243-249. <https://doi.org/10.1007/s00414-012-0699-1>.
- Boehme P, Spahn P, Amendt J, Zehner R (2014) The analysis of temporal gene expression to estimate the age of forensically important blow fly pupae: results from three blind studies. *Int J Legal Med* 128:565-573. <https://doi.org/10.1007/s00414-013-0922-8>.
- Bourel B, Callet B, Hédouin V, Gosset D (2003) Flies eggs: a new method for the estimation of short-term post-mortem interval? *Forensic Sci Int* 135:27-34. [https://doi.org/10.1016/S0379-0738\(03\)00157-9](https://doi.org/10.1016/S0379-0738(03)00157-9).
- Braga MV, Teixeira Pinto Z, de Carvalho Queiroz MM, Blomquist GJ (2016) Effect of age on cuticular hydrocarbon profiles in adult *Chrysomya putoria* (Diptera: Calliphoridae). *Forensic Sci Int* 259:e37-e47. <https://doi.org/10.1016/j.forsciint.2015.11.006>
- Brown K, Thorne A, Harvey M (2015) *Calliphora vicina* (Diptera: Calliphoridae) pupae: a timeline of external morphological development and a new age and PMI estimation tool. *Int J Legal Med.* 129:835-850. <https://doi.org/10.1007/s00414-014-1068-z>.
- Brown K, Thorne A, Harvey M (2012) Preservation of *Calliphora vicina* (Diptera: Calliphoridae) pupae for use in post-mortem interval estimation. *Forensic Sci Int.* 223:176-183. <https://doi.org/10.1016/j.forsciint.2012.08.029>.
- Brundage A, Bros S, Honda JY (2011) Seasonal and habitat abundance and distribution of some forensically important blow flies (Diptera: Calliphoridae) in Central California, *Forensic Sci Int* 212:115-120. <https://doi.org/10.1016/j.forsciint.2011.05.023>.
- Butcher JB, Moore HE, Day CR, Adam CD, Drijfhout FP (2013) Artificial neural network analysis of hydrocarbon profiles for the ageing of *Lucilia sericata* for post mortem interval estimation. *Forensic Sci Int* 232:25-31, <https://doi.org/10.1016/j.forsciint.2013.06.018>.
- Bygarski K, LeBlanc HN (2013) Decomposition and arthropod succession in Whitehorse, Yukon Territory, Canada. *J Forensic Sci* 58:413-418. <https://doi.org/10.1111/1556-4029.12032>.
- Byrd JH, Allen JC (2001) The development of the black blow fly, *Phormia regina* (Meigen). *Forensic Sci Int.* 120:79-88.
- Byrd JH, Butler JF (1996) Effects of temperature on *Cochliomyia macellaria* (Diptera:Calliphoridae) development. *J Med Entomol* 33:901-905.
- Byrd JH, Butler JF (1997) Effects of temperature on *Chrysomya rufifacies* (Diptera:Calliphoridae) development. *J Med Entomol* 34:353-358.
- Byrd JH, Butler JF (1998) Effects of temperature on *Sarcophaga haemorrhoidalis* (Diptera: Sarcophagidae) development. *J Med Entomol* 35:694-698.
- Cervantès L, Dourel L, Gaudry E, Pasquerault T, Vincent B (2017) Effect of low temperature in the development cycle of *Lucilia sericata* (Meigen) (Diptera, Calliphoridae): implications for the

minimum postmortem interval estimation. *Forensic Sci Res* 3:52-59.
<https://doi.10.1080/20961790.2017.1406839>.

- Charabidze D, Vincent B, Pasquerault T, Hedouin V (2015) The biology and ecology of *Necrodes littoralis*, a species of forensic interest in Europe. *Int J Legal Med* 130:273-280.
<https://doi.10.1007/s00414-015-1253-8>.
- Charabidze D, Hedouin V, Gosset D (2015) An experimental investigation into the colonization of concealed cadavers by necrophagous blowflies. *J Insect Sci.* 15:149. <https://doi.10.1093/jisesa/iev129>.
- Charabidze D, Bourel B, Gosset D (2011) Larval-mass effect: Characterisation of heat emission by necrophagous blowflies (Diptera: Calliphoridae) larval aggregates. *Forensic Sci Int.* 211:61-6.
<https://doi.10.1016/j.forsciint.2011.04.016>.
- Cherix D, Wyss C, Pape T (2012) Occurrences of flesh flies (Diptera: Sarcophagidae) on human cadavers in Switzerland, and their importance as forensic indicators. *Forensic Sci Int* 220:158-163
<https://doi.org/10.1016/j.forsciint.2012.02.016>.
- Costa Oliveira T, Vasconcelos SD (2010) Insects (Diptera) associated with cadavers at the Institute of Legal Medicine in Pernambuco, Brazil: Implications for forensic entomology. *Forensic Sci Int* 198:97-102. <https://doi.org/10.1016/j.forsciint.2010.01.011>.
- da Silva SM, Moura MO (2019) Intrapuparial Development of *Hemilucilia semidiaphana* (Diptera: Calliphoridae) and Its Use in Forensic Entomology. *J Med Entomol* 56:1623-1635.
<https://doi.10.1093/jme/tjz118>. PubMed PMID: 31290542.
- da Silva X, Barbosa RR, Guimarães Barbosa C, de Carvalho Queiroz MM (2015) Bionomy of two flies of sanitary and forensic importance: *Peckia* (Sarcodexia) *lambens* (Wiedemann) and *Oxysarcodexia amorosa* (Schiner) (Diptera, Sarcophagidae). *Revista Brasileira de Entomologia* 59:229-233. <https://doi.org/10.1016/j.rbe.2015.06.002>.
- Day DM, Wallman JF (2006) Influence of substrate tissue type on larval growth in *Calliphora augur* and *Lucilia cuprina* (Diptera: Calliphoridae). *J Forensic Sci* 51:657-663.
- Day DM, Wallman JF (2006) Width as an alternative measurement to length for post-mortem interval estimations using *Calliphora augur* (Diptera: Calliphoridae) larvae. *Forensic Sci Int* 159:158-167. <https://doi.org/10.1016/j.forsciint.2005.07.009>
- Day DM, Wallman JF (2008) Effect of preservative solutions on preservation of *Calliphora augur* and *Lucilia cuprina* larvae (Diptera: Calliphoridae) with implications for post-mortem interval estimates. *Forensic Sci Int* 179:1-10. <https://doi.org/10.1016/j.forsciint.2008.04.006>.
- Defilippo F, Bonilauri P, Dottori M (2013) Effect of temperature on six different developmental landmarks within the pupal stage of the forensically important blowfly *Calliphora vicina* (Robineau-Desvoidy) (Diptera: Calliphoridae). *J Forensic Sci* 58:1554-1557. <https://doi.10.1111/1556-4029.12199>.
- Díaz-Aranda LM, Martín-Vega D, Gómez-Gómez A, Cifrián B, Baz A (2018) Annual variation in decomposition and insect succession at a periurban area of central Iberian Peninsula. *J Forensic Legal Med* 56:21-31. <https://doi.org/10.1016/j.jflm.2018.03.005>
- Duarte JLP, Emmerich RF, Corrêa APF, Krüger RF (2015) Thermal requirements of *Ophyra albuquerquei* Lopes, 1985 (Diptera, Muscidae). *Forensic Sci Int* 254:227-230.
<https://doi.org/10.1016/j.forsciint.2015.07.014>.
- Faris AM, Wang HH, Tarone AM, Grant WE (2016) Forensic Entomology: Evaluating Uncertainty Associated With Postmortem Interval (PMI) Estimates With Ecological Models. *J Med Entomol* 53:1117-1130. <https://doi.10.1093/jme/tjw070>.
- Faris AM, West WR, Tomberlin JK, Tarone AM (2020) Field Validation of a Development Data Set for *Cochliomyia macellaria* (Diptera: Calliphoridae): Estimating Insect Age Based on Development Stage. *J Med Entomol* 57:39-49. <https://doi.10.1093/jme/tjz156>.
- Feddern N, Mitchell EAD, Amendt J, Szelecz I, Seppey CVW (2019) Decomposition and insect colonization patterns of pig cadavers lying on forest soil and suspended above ground. *Forensic Sci Med Pathol* 15:342-351. <https://doi.10.1007/s12024-019-00121-6>.

- Feng DX, Liu GC (2013) Pupal age estimation of forensically important *Megaselia spiracularis* Schmitz (Diptera: Phoridae). *Forensic Sci Int.* 231:199-203. <https://doi.org/10.1016/j.forsciint.2013.05.008>.
- Feng DX, Liu GC (2014) Pupal age estimation of forensically important *Megaselia scalaris* (Loew) (Diptera: Phoridae). *Forensic Sci Int* 236:133-137. <https://doi.org/10.1016/j.forsciint.2014.01.002>.
- Frątczak K, Matuszewski S (2014) Instar determination in forensically useful beetles *Necrodes littoralis* (Silphidae) and *Creophilus maxillosus* (Staphylinidae). *Forensic Sci Int.* 241:20-26. <https://doi.org/10.1016/j.forsciint.2014.04.026>.
- Frątczak-Łagiewska K, Matuszewski S (2018) Sex-specific developmental models for *Creophilus maxillosus* (L.) (Coleoptera: Staphylinidae): searching for larger accuracy of insect age estimates. *Int J Legal Med* 132:887-895. <https://doi.org/10.1007/s00414-017-1713-4>.
- García-Rojo AM, Martínez-Sánchez A, López R, García de la Vega JM, Rica M, González M, Disney RHL (2013) A mathematical model applied for assisting the estimation of PMI in a case of forensic importance. *Forensic Sci Int.* 231:e11-18. <https://doi.org/10.1016/j.forsciint.2013.05.038>.
- Gaudry E, Blais C, Maria A, Dauphin-Villemant C (2006) Study of steroidogenesis in pupae of the forensically important blow fly *Protophormia terraenovae* (Robineau-Desvoidy) (Diptera: Calliphoridae). *Forensic Sci Int* 160:27-34. <https://doi.org/10.1016/j.forsciint.2005.06.014>.
- Goff ML (1993) Estimation of Postmortem Interval Using Arthropod Development and Successional Patterns. *Forensic Sci Rev* 5:81-94.
- González Medina A, Soriano Hernando Ó, Jiménez Ríos G (2015) The Use of the Developmental Rate of the Aquatic Midge *Chironomus riparius* (Diptera, Chironomidae) in the Assessment of the Postsubmersion Interval. *J Forensic Sci.* 60:822-826. <https://doi.org/10.1111/1556-4029.12707>.
- Grassberger M, Reiter C (2001) Effect of temperature on *Lucilia sericata* (Diptera: Calliphoridae) development with special reference to the isomegalen- and isomorphen-diagram. *Forensic Sci Int* 120:32-36.
- Grassberger M, Reiter C (2002) Effect of temperature on development of the forensically important holarctic blow fly *Protophormia terraenovae* (Robineau-Desvoidy) (Diptera: Calliphoridae). *Forensic Sci Int* 128:177-182. [https://doi.org/10.1016/S0379-0738\(02\)00199-8](https://doi.org/10.1016/S0379-0738(02)00199-8).
- Greenberg B, Wells JD (1998) Forensic use of *Megaselia abdita* and *M. scalaris* (Phoridae: Diptera): case studies, development rates, and egg structure. *J Med Entomol* 35:205-209.
- Gruner SV, Slone DH, Capinera JL, Turco MP (2017) Development of the Oriental Latrine Fly, *Chrysomya megacephala* (Diptera: Calliphoridae), at Five Constant Temperatures. *J Med Entomol.* 54:290-298. <https://doi.org/10.1093/jme/tjw169>.
- Gruner SV, Slone DH, Capinera JL, Turco MP (2017) Volume of Larvae Is the Most Important Single Predictor of Mass Temperatures in the Forensically Important Calliphorid, *Chrysomya megacephala* (Diptera: Calliphoridae). *J Med Entomol* 54:30-34. <https://doi.org/10.1093/jme/tjw139>.
- Gruszka J, Matuszewski S (2020) Estimation of physiological age at emergence based on traits of the forensically useful adult carrion beetle *Necrodes littoralis* L. (Silphidae). *Forensic Sci Int* 314:110407. <https://doi.org/10.1016/j.forsciint.2020.110407>.
- Hu G, Wang M, Wang Y, Tang H, Chen R, Zhang Y, Zhao Y, Jin Y, Wang Y, Wu M, Wang J (2020) Development of *Necrobia rufipes* (De Geer, 1775) (Coleoptera: Cleridae) under constant temperatures and its implication in forensic entomology. *Forensic Sci Int* 311:110275. <https://doi.org/10.1016/j.forsciint.2020.110275>
- Harnden LM, Tomberlin JK (2016) Effects of temperature and diet on black soldier fly, *Hermetia illucens* (L.) (Diptera: Stratiomyidae), development. *Forensic Sci Int.* 266:109-116. <https://doi.org/10.1016/j.forsciint.2016.05.007>.
- Heo CC, Mohamad AM, Ahmad FM, Jeffery J, Kurahashi H, Omar B (2008) Study of insect succession and rate of decomposition on a partially burned pig carcass in an oil palm plantation in Malaysia. *Trop Biomed* 25:202-208.

- Hofer IMJ, Hart AJ, Martín-Vega D, Hall MJR (2020) Estimating crime scene temperatures from nearby meteorological station data. *Forensic Sci Int.* 306:110028. <https://doi.10.1016/j.forsciint.2019.110028>.
- Hu G, Wang Y, Sun Y, Zhang Y, Wang M, Wang J (2019) Development of *Chrysomya rufifacies* (Diptera: Calliphoridae) at Constant Temperatures Within its Colony Range in Yangtze River Delta Region of China. *J. Med. Entomol* 56:1215-1224. <https://doi.10.1093/jme/tjz052>.
- Huntington TE, Higley LG, Baxendale FP (2007) Maggot development during morgue storage and its effect on estimating the post-mortem interval. *J Forensic Sci* 52:453-458.
- Iancu L, Sahlean T, Purcarea C (2016) Dynamics of Necrophagous Insect and Tissue Bacteria for Postmortem Interval Estimation During the Warm Season in Romania. *J Med Entomol* 53:54-66. <https://doi.10.1093/jme/tjv156>.
- Jarmusz M, Grzywacz A, Bajerlein D(2020) A comparative study of the entomofauna (Coleoptera, Diptera) associated with hanging and ground pig carcasses in a forest habitat of Poland. *Forensic Sci Int.* 309:110212. <https://doi.10.1016/j.forsciint.2020.110212>.
- Johnson AP, Wallman JF (2014) Effect of massing on larval growth rate. *Forensic Sci Int* 241:141-149. <https://doi.10.1016/j.forsciint.2014.05.006>.
- Karabey T, Sert O (2018) The analysis of pupal development period in *Lucilia sericata* (Diptera: Calliphoridae) forensically important insect. *Int J Legal Med.* 132:1185-1196. <https://doi.10.1007/s00414-014-0968-2>.
- Kelly JA, van der Linde TC, Anderson GS (2009) The influence of clothing and wrapping on carcass decomposition and arthropod succession during the warmer seasons in central South Africa. *J Forensic Sci.* 54:1105-1112. <https://doi.10.1111/j.1556-4029.2009.01113.x>.
- Kotzé Z, Villet MH, Weldon CW (2016) Heat accumulation and development rate of massed maggots of the sheep blowfly, *Lucilia cuprina* (Diptera: Calliphoridae), *J Insect Physiol* 95:98-104. <https://doi.org/10.1016/j.jinsphys.2016.09.009>.
- Kulshrestha P, Satpathy DK (2001) Use of beetles in forensic entomology. *Forensic Sci Int* 120: 15-17.
- Lambiase S, Murgia G, Sacchi R, Ghitti M, di Lucia V (2018) Effects of Different Temperatures on the Development of *Dermestes Frischii* and *Dermestes undulatus* (Coleoptera, Dermestidae): Comparison Between Species. *J Forensic Sci* 63:469-473. <https://doi.10.1111/1556-4029.13580>.
- Lecheta MC, Moura MO (2019) Estimating the Age of Forensically Useful Blowfly, *Sarconesia chlorogaster* (Diptera: Calliphoridae), Using Larval Length and Weight. *J Med Entomol* 56:915-920. <https://doi.10.1093/jme/tjz028>. PubMed PMID: 30924860.
- Lecheta MC, Thyssen PJ, Moura MO (2015) The effect of temperature on development of *Sarconesia chlorogaster*, a blowfly of forensic importance. *Forensic Sci Med Pathol.* 11:538-543. <https://doi.10.1007/s12024-015-9727-z>.
- Lefebvre F, Pasquerault T (2004) Temperature-dependent development of *Ophyra aenescens* (Wiedemann, 1830) and *Ophyra capensis* (Wiedemann, 1818) (Diptera, Muscidae). *Forensic Sci Int* 139:75-79. <https://doi.org/10.1016/j.forsciint.2003.10.014>.
- Li L, Wang Y, Wang J, Ma M, Lai Y (2016) Temperature-dependent development and the significance for estimating postmortem interval of *Chrysomya nigripes* Aubertin, a new forensically important species in China. *Int J Legal Med* 130:1363-1370. <https://doi.10.1007/s00414-016-1315-6>.
- Li L, Wang Y, Wang J (2016) Intra-puparial development and age estimation of forensically important *Hermetia illucens* (L.). *J Asia-Pacific Entomol* 19:233-237. <https://doi.org/10.1016/j.aspen.2016.01.006>.
- Li X, Yang Y, Li G, Li H, Wang Q, Wan L (2014) The effect of dietary fat levels on the size and development of *Chrysomya megacephala* (Diptera: Calliphoridae). *J. Insect Sci* 14:174. <https://doi.10.1093/jisesa/ieu036>.

- Lin SW, Shiao SF (2013) Life history data on the fly parasitoids *Aleochara nigra* Kraatz and *A. asiatica* Kraatz (Coleoptera: Staphylinidae), and their potential application in forensic entomology. *Forensic Sci Int.* 232:46-55. <https://doi.org/10.1016/j.forsciint.2013.06.016>.
- López-Esclapez R, García MD, Arnaldos MI, Presa JJ, Ubero-Pascal N (2014) Are the evidences of forensic entomology preserved in ethanol suitable for SEM studies? *Micron* 62:43-51. <https://doi.org/10.1016/j.micron.2014.03.003>.
- Lutz L, Amendt J (2020) Precocious egg development in wild *Calliphora vicina* (Diptera: Calliphoridae) - An issue of relevance in forensic entomology? *Forensic Sci Int.* 306:110075. <https://doi.org/10.1016/j.forsciint.2019.110075>.
- Lutz L, Amendt J (2020) Stay cool or get hot? An applied primer for using temperature in forensic entomological case work. *Science & Justice* 60:415-422. <https://doi.org/10.1016/j.scijus.2020.05.003>.
- Ma T, Huang J, Wang JF (2015) Study on the pupal morphogenesis of *Chrysomya rufifacies* (Macquart) (Diptera: Calliphoridae) for postmortem interval estimation. *Forensic Sci Int.* 253:88-93. <https://doi.org/10.1016/j.forsciint.2015.06.005>.
- Mądra A, Konwerski S, Matuszewski S (2014) Necrophilous Staphylininae (Coleoptera: Staphylinidae) as indicators of season of death and corpse relocation. *Forensic Sci Int* 242:32-37. <https://doi.org/10.1016/j.forsciint.2014.06.011>
- Mądra A, Frątczak K, Grzywacz A, Matuszewski S (2015) Long-term study of pig carrion entomofauna. *Forensic Sci Int.* 252:1-10. <https://doi.org/10.1016/j.forsciint.2015.04.013>.
- Mądra-Bielewicz A, Frątczak-Łagiewska K, Matuszewski S (2017) Sex- and Size-Related Patterns of Carrion Visitation in *Necrodes littoralis* (Coleoptera: Silphidae) and *Creophilus maxillosus* (Coleoptera: Staphylinidae). *J Forensic Sci.* 62:1229-1233. <https://doi.org/10.1111/1556-4029.13376>.
- Mai M, Amendt J (2012) Effect of different post-feeding intervals on the total time of development of the blowfly *Lucilia sericata* (Diptera: Calliphoridae). *Forensic Sci Int* 221:65-69. <https://doi.org/10.1016/j.forsciint.2012.04.001>.
- Malainey SL, Anderson GS (2020) Effect of arson fires on survivability of entomological evidence on carcasses inside vehicle trunks. *Forensic Sci Int.* 306:110033. <https://doi.org/10.1016/j.forsciint.2019.110033>
- Mañas-Jordá S, León-Cortés JL, García-García MD, Caballero U, Infante F. (2018) Dipteran Diversity and Ecological Succession on Dead Pigs in Contrasting Mountain Habitats of Chiapas, Mexico. *J Med Entomol.* 55:59-68. <https://doi.org/10.1093/jme/tjx190>.
- Manlove JD, Disney RHL (2008) The use of *Megaselia abdita* (Diptera: Phoridae) in forensic entomology. *Forensic Sci Int* 175:83-84, <https://doi.org/10.1016/j.forsciint.2007.08.001>.
- Martínez E, Duque P, Wolff M (2007) Succession pattern of carrion-feeding insects in Paramo, Colombia. *Forensic Sci Int.* 166:182-189.
- Martín-Vega D, Baz A, Cifrián B, Gómez-Gómez A, Díaz-Aranda LM (2019) Long-term insect successional patterns on pig carcasses in central Spain. *Int J Legal Med* 133:1581-1592. <https://doi.org/10.1007/s00414-019-02088-z>.
- Martín-Vega D, Díaz-Aranda LM, Baz A, Cifrián B. (2017) Effect of Temperature on the Survival and Development of Three Forensically Relevant Dermestid Species (Coleoptera: Dermestidae). *J Med Entomol* 54:1140-1150. <https://doi.org/10.1093/jme/tjx110>.
- Martín-Vega D, Nieto CM, Cifrián B, Baz A, Díaz-Aranda LM (2017) Early colonisation of urban indoor carcasses by blow flies (Diptera: Calliphoridae): An experimental study from central Spain. *Forensic Sci Int* 278:87-94. <https://doi.org/10.1016/j.forsciint.2017.06.036>.
- Matuszewski S (2011) Estimating the pre-appearance interval from temperature in *Necrodes littoralis* L. (Coleoptera: Silphidae). *Forensic Sci Int* 212:180-188. <https://doi.org/10.1016/j.forsciint.2011.06.010>.
- Matuszewski S (2017) A general approach for postmortem interval based on uniformly distributed and interconnected qualitative indicators. *Int J Legal Med.* 131:877-884. <https://doi.org/10.1007/s00414-016-1520-3>.

- Matuszewski S, Frątczak-Łagiewska K (2018) Size at emergence improves accuracy of age estimates in forensically-useful beetle *Creophilus maxillosus* L. (Staphylinidae). *Sci Rep* 8:2390. [https://doi. 10.1038/s41598-018-20796-1](https://doi.org/10.1038/s41598-018-20796-1).
- Matuszewski S, Mądra A (2015) Factors affecting quality of temperature models for the pre-appearance interval of forensically useful insects. *Forensic Sci Int.* 247:28-35. [https://doi. 10.1016/j.forsciint.2014.11.026](https://doi.org/10.1016/j.forsciint.2014.11.026).
- Matuszewski S, Mądra-Bielewicz A (2016) Validation of temperature methods for the estimation of pre-appearance interval in carrion insects. *Forensic Sci Med Pathol.* 12:50-57. [https://doi. 10.1007/s12024-015-9735-z](https://doi.org/10.1007/s12024-015-9735-z).
- Matuszewski S, Szafałowicz M, Grzywacz A (2014) Temperature-dependent appearance of forensically useful flies on carcasses. *Int J Legal Med.* 128: 1013-1020. [https://doi. 10.1007/s00414-013-0921-9](https://doi.org/10.1007/s00414-013-0921-9).
- Matuszewski S, Bajerlein D, Konwerski S, Szpila K (2010) Insect succession and carrion decomposition in selected forests of Central Europe. Part 1: Pattern and rate of decomposition. *Forensic Sci Int.* 194:85-93. [https://doi. 10.1016/j.forsciint.2009.10.016](https://doi.org/10.1016/j.forsciint.2009.10.016).
- Matuszewski S, Bajerlein D, Konwerski S, Szpila K. (2010) Insect succession and carrion decomposition in selected forests of Central Europe. Part 2: Composition and residency patterns of carrion fauna. *Forensic Sci Int.* 195:42-51. [https://doi. 10.1016/j.forsciint.2009.11.007](https://doi.org/10.1016/j.forsciint.2009.11.007)
- Matuszewski S, Bajerlein D, Konwerski S, Szpila K (2011) Insect succession and carrion decomposition in selected forests of Central Europe. Part 3: Succession of carrion fauna. *Forensic Sci Int.* 207:150-163. [https://doi. .1016/j.forsciint.2010.09.022](https://doi.org/10.1016/j.forsciint.2010.09.022).
- Matuszewski S, Frątczak K, Konwerski S, Bajerlein D, Szpila K, Jarmusz M, Szafałowicz M, Grzywacz A, Mądra A (2016) Effect of body mass and clothing on carrion entomofauna. *Int J Legal Med* 130:221-232. [https://doi. 10.1007/s00414-015-1145-y](https://doi.org/10.1007/s00414-015-1145-y).
- Mello S, Aguiar-Coelho VM (2009) Durations of immature stage development period of *Nasonia vitripennis* (Walker) (Hymenoptera: Pteromalidae) under laboratory conditions: implications for forensic entomology. *Parasitol Res* 104:411-418. [https://doi. 10.1007/s00436-008-1213-z](https://doi.org/10.1007/s00436-008-1213-z).
- Michalski M, Nadolski J (2018) Thermal conditions in selected urban and semi-natural habitats, important for the forensic entomology. *Forensic Sci Int* 287:153-162. [https://doi.10.1016/j.forsciint.2018.03.042](https://doi.org/10.1016/j.forsciint.2018.03.042).
- Michaud JP, Moreau G (2009) Predicting the visitation of carcasses by carrion-related insects under different rates of degree-day accumulation. *Forensic Sci Int* 185:78-83. <https://doi.org/10.1016/j.forsciint.2008.12.015>.
- Michaud JP, Schoenly KG, Moreau G (2012) Sampling flies or sampling flaws? Experimental design and inference strength in forensic entomology. *J Med Entomol* 49:1-10.
- Midgley JM, Villet MH (2009) Development of *Thanatophilus micans* (Fabricius 1794) (Coleoptera: Silphidae) at constant temperatures. *Int J Legal Med.* 123:285-292. [https://doi. 10.1007/s00414-008-0280-0](https://doi.org/10.1007/s00414-008-0280-0).
- Moffatt C, Heaton V, De Haan D (2016) The distribution of blow fly (Diptera: Calliphoridae) larval lengths and its implications for estimating post mortem intervals. *Int J Legal Med.* 130:287-97. [https://doi. 10.1007/s00414-015-1289-9](https://doi.org/10.1007/s00414-015-1289-9).
- Moore HE, Adam CD, Drijfhout FP (2013) Potential use of hydrocarbons for aging *Lucilia sericata* blowfly larvae to establish the postmortem interval. *J Forensic Sci.* 58:404-412. [https://doi. 10.1111/1556-4029.12016](https://doi.org/10.1111/1556-4029.12016).
- Moore HE, Butcher JB, Adam CD, Day CR, Drijfhout FP (2016) Age estimation of *Calliphora* (Diptera: Calliphoridae) larvae using cuticular hydrocarbon analysis and Artificial Neural Networks. *Forensic Sci Int* 268:81-91. <https://doi.org/10.1016/j.forsciint.2016.09.012>.
- Moore HE, Pechal JL, Benbow ME, Drijfhout FP (2017) The potential use of cuticular hydrocarbons and multivariate analysis to age empty puparial cases of *Calliphora vicina* and *Lucilia sericata*. *Sci Rep.* 7:1933. [https://doi. .1038/s41598-017-01667-7](https://doi.org/10.1038/s41598-017-01667-7).

- Moore HE, Butcher JB, Day CR, Drijfhout FP (2017) Adult fly age estimations using cuticular hydrocarbons and Artificial Neural Networks in forensically important Calliphoridae species. *Forensic Sci Int* 280:233-244. <https://doi.org/10.1016/j.forsciint.2017.10.001>.
- Myskowiak J-B, Doums C (2002) Effects of refrigeration on the biometry and development of *Protophormia terraenovae* (Robineau-Desvoidy) (Diptera: Calliphoridae) and its consequences in estimating post-mortem interval in forensic investigations. *Forensic Sci Int* 125:254-261. [https://doi.org/10.1016/S0379-0738\(02\)00003-8](https://doi.org/10.1016/S0379-0738(02)00003-8).
- Nability PD, Higley LG, Heng-Moss TM (2006) Effects of temperature on development of *Phormia regina* (Diptera: Calliphoridae) and use of developmental data in determining time intervals in forensic entomology. *J Med Entomol* 43:1276-1286.
- Nability PD, Higley LG, Heng-Moss TM (2007) Light-induced variability in development of forensically important blow fly *Phormia regina* (Diptera: Calliphoridae). *J Med Entomol* 44:351-358.
- Nassu MP, Thyssen PJ, Linhares AX (2014) Developmental rate of immatures of two fly species of forensic importance: *Sarcophaga* (*Liopygia*) *ruficornis* and *Microcerella halli* (Diptera: Sarcophagidae). *Parasitol Res* 113:217-222. <https://doi.org/10.1007/s00436-013-3646-2>
- Neto-Silva A, Dinis-Oliveira RJ, Prado E Castro C (2017) Diversity of the Formicidae (Hymenoptera) carrion communities in Lisbon (Portugal): preliminary approach as seasonal and geographic indicators. *Forensic Sci Res*. 3:65-73. <https://doi.org/10.1080/20961790.2017.1372875>.
- Niederegger S, Pastuschek J, Mall G (2010) Preliminary studies of the influence of fluctuating temperatures on the development of various forensically relevant flies. *Forensic Sci Int* 199:72-78. <https://doi.org/10.1016/j.forsciint.2010.03.015>.
- Ortloff A, Peña P, Riquelme M (2012) Preliminary study of the succession pattern of necrobiont insects, colonising species and larvae on pig carcasses in Temuco (Chile) for forensic applications. *Forensic Sci Int*. 222:e36-41. <https://doi.org/10.1016/j.forsciint.2012.04.022>.
- Owings CG, Spiegelman C, Tarone AM, Tomberlin JK (2014) Developmental variation among *Cochliomyia macellaria* Fabricius (Diptera: Calliphoridae) populations from three ecoregions of Texas, USA. *Int J Legal Med*. 128:709-17. <https://doi.org/10.1007/s00414-014-1014-0>.
- Patrício Macedo M, Arantes LC, Tidon R (2018) Sexual size dimorphism in three species of forensically important blowflies (Diptera: Calliphoridae) and its implications for postmortem interval estimation. *Forensic Sci Int*. 293:86-90. <https://doi.org/10.1016/j.forsciint.2018.10.009>.
- Perez AE, Haskell NH, Wells JD (2014) Evaluating the utility of hexapod species for calculating a confidence interval about a succession based postmortem interval estimate. *Forensic Sci Int*. 241:91-95. <https://doi.org/10.1016/j.forsciint.2014.05.007>.
- Pérez SP, Duque P, Wolff M (2005) Successional behavior and occurrence matrix of carrion-associated arthropods in the urban area of Medellín, Colombia. *J Forensic Sci*. 50:448-454.
- Picard CJ, Deblois K, Tovar F, Bradley JL, Johnston JS, Tarone AM (2013) Increasing precision in development-based postmortem interval estimates: what's sex got to do with it? *J Med Entomol* 50:425-431.
- Pinilla YT, Patarroyo MA, Bello FJ (2013) *Sarconesiopsis magellanica* (Diptera: Calliphoridae) life-cycle, reproductive and population parameters using different diets under laboratory conditions. *Forensic Sci Int*. 233:380-386. <https://doi.org/10.1016/j.forsciint.2013.10.014>.
- Prado e Castro C, García MD, Martins da Silva P, Faria e Silva I, Serrano A (2013) Coleoptera of forensic interest: a study of seasonal community composition and succession in Lisbon, Portugal. *Forensic Sci Int*. 232:73-83. <https://doi.org/10.1016/j.forsciint.2013.06.014>.
- Rabêlo KCN, Thyssen PJ, Salgado RL, Araújo MSC, Vasconcelos SD (2011) Bionomics of two forensically important blowfly species *Chrysomya megacephala* and *Chrysomya putoria* (Diptera: Calliphoridae) reared on four types of diet. *Forensic Sci Int* 210:257-262. <https://doi.org/10.1016/j.forsciint.2011.03.022>

- Reibe S, Madea B (2010) How promptly do blowflies colonise fresh carcasses? A study comparing indoor with outdoor locations. *Forensic Sci Int* 195:52-57. <https://doi.10.1016/j.forsciint.2009.11.009>.
- Reibe S, Madea B (2010) Use of *Megaselia scalaris* (Diptera: Phoridae) for post-mortem interval estimation indoors. *Parasitol Res.* 106:637-640. <https://doi.10.1007/s00436-009-1713-5>.
- Reibe S, Doetinchem PV, Madea B (2010) A new simulation-based model for calculating post-mortem intervals using developmental data for *Lucilia sericata* (Dipt.: Calliphoridae). *Parasitol Res.* 107:9-16. <https://doi.1007/s00436-010-1879-x>.
- Reibe-Pal S, Madea B (2015) Calculating time since death in a mock crime case comparing a new computational method (ExLAC) with the ADH method. *Forensic Sci Int.* 248:78-81. <https://doi.10.1016/j.forsciint.2014.12.024>.
- Richards CS, Villet MH (2009) Data quality in thermal summation development models for forensically important blowflies. *Med Vet Entomol* 23:269-276. <https://doi.10.1111/j.1365-2915.2009.00819.x>.
- Richards CS, Villet MH (2008) Factors affecting accuracy and precision of thermal summation models of insect development used to estimate post-mortem intervals. *Int J Legal Med* 122:401-408. <https://doi.10.1007/s00414-008-0243-5>.
- Richards CS, Simonsen TJ, Abel RL, Hall MJ, Schwyn DA, Wicklein M (2012) Virtual forensic entomology: improving estimates of minimum post-mortem interval with 3D micro-computed tomography. *Forensic Sci Int.* 220:251-264. <https://doi.10.1016/j.forsciint.2012.03.012>
- Richards CS, Rowlinson CC, Cuttiford L, Grimsley R, Hall MJ (2013) Decomposed liver has a significantly adverse affect on the development rate of the blowfly *Calliphora vicina*. *Int J Legal Med.* 127:259-262. <https://doi.10.1007/s00414-012-0697-3>.
- Richards CS, Rowlinson CC, Hall MJ (2013) Effects of storage temperature on the change in size of *Calliphora vicina* larvae during preservation in 80% ethanol. *Int J Legal Med.* 127:231-241. <https://doi.10.1007/s00414-012-0683-9>.
- Richards CS, Paterson ID, Villet MH (2008) Estimating the age of immature *Chrysomya albiceps* (Diptera: Calliphoridae), correcting for temperature and geographical latitude. *Int J Legal Med* 122:271-279.
- Richards CS, Crous KL, Villet MH (2009) Models of development for blowfly sister species *Chrysomya chloropyga* and *Chrysomya putoria*. *Med Vet Entomol* 23:56-61. <https://doi.10.1111/j.1365-2915.2008.00767.x>.
- Roe A, Higley LG (2015) Development modeling of *Lucilia sericata* (Diptera: Calliphoridae). *PeerJ.* 3:e803. <https://doi.10.7717/peerj.803>.
- Rosilawati R, Baharudin O, Syamsa RA, Lee HL, Nazni WA (2014) Effects of preservatives and killing methods on morphological features of a forensic fly, *Chrysomya megacephala* (Fabricius, 1794) larva. *Trop Biomed* 31:785-791.
- Roux O, Gers C, Legal L (2008) Ontogenetic study of three Calliphoridae of forensic importance through cuticular hydrocarbon analysis. *Med Vet Entomol* 22:309-317. <https://doi.10.1111/j.1365-2915.2008.00752.x>.
- Salimi M, Chatrabgoun O, Akbarzadeh K, Oshaghi M, Falahati MH, Rafizadeh S, Yusuf MA, Rassi Y. (2018) Evaluation of Insect Succession Patterns and Carcass Weight Loss for the Estimation of Postmortem Interval. *J Med Entomol.* 55:1410-1422. <https://doi.10.1093/jme/tjy095>.
- Schoenly K (1992) A statistical analysis of successional patterns in carrion-arthropod assemblages: implications for forensic entomology and determination of the postmortem interval. *J Forensic Sci.* 37:1489-1513.
- Schoenly K, Goff ML, Early M (1992) A BASIC algorithm for calculating the postmortem interval from arthropod successional data. *J Forensic Sci.* 37:808-23.
- Schoenly K, Griest K, Rhine S (1991) An experimental field protocol for investigating the postmortem interval using multidisciplinary indicators. *J Forensic Sci* 36:1395-1415.

- Schroeder H, Klotzbach H, Püschel K (2003) Insects' colonization of human corpses in warm and cold season. *Leg Med* 5 Suppl 1:S372-374.
- Segura NA, Usaquén W, Sánchez MC, Chuaire L, Bello F (2009) Succession pattern of cadaverous entomofauna in a semi-rural area of Bogotá, Colombia. *Forensic Sci Int* 187:66-72. <https://doi.org/10.1016/j.forsciint.2009.02.018>.
- Sert O, Örsel GM, Şabanoğlu B, Özdemir S (2020) A Study of the pupal developments of *Sarcophaga argyrostoma* (Robineau-Desvoidy, 1830). *Forensic Sci Med Pathol.* 16:12-19. <https://doi.10.1007/s12024-019-00198-z>.
- Shang Y, Ren L, Yang L, Wang S, Chen W, Dong J, Ma H, Qi X, Guo Y (2020) Differential Gene Expression for Age Estimation of Forensically Important *Sarcophaga peregrina* (Diptera: Sarcophagidae) Intrapuparial. *J Med Entomol* 57:65-77. <https://doi.10.1093/jme/tjz137>.
- Shiravi A, Mostafavi R, Akbarzadeh K, Oshaghi M (2011) Temperature Requirements of Some Common Forensically Important Blow and Flesh Flies (Diptera) under Laboratory Conditions. *Iran J Arthropod Borne Dis* 5:54-62.
- Singh R, Sharma S, Sharma A (2016) Determination of post-burial interval using entomology: A review. *J Forensic Leg Med* 42:37-40. <https://doi.1016/j.jflm.2016.05.004>.
- Smith JL, Cook AK (2020) First Use of an Entire Age Cohort to Evaluate the Role of Sex in the Development of the Forensically Important *Chrysomya megacephala* (Diptera: Calliphoridae). *J Med Entomol* 57:641-644. <https://doi.10.1093/jme/tjz205>.
- Sukontason K, Piangjai S, Siri wattanarungsee S, Sukontason KL (2008) Morphology and developmental rate of blowflies *Chrysomya megacephala* and *Chrysomya rufifacies* in Thailand: application in forensic entomology. *Parasitol Res.* 102:1207-1216. <https://doi.10.1007/s00436-008-0895-6>.
- Tabor KL, Brewster CC, Fell RD (2004) Analysis of the successional patterns of insects on carrion in southwest Virginia. *J Med Entomol* 41:785-795.
- Tabor KL, Fell RD, Brewster CC (2005) Insect fauna visiting carrion in Southwest Virginia. *Forensic Sci Int* 150:73-80. <https://doi.org/10.1016/j.forsciint.2004.06.041>.
- Tarone AM, Foran DR (2011) Gene expression during blow fly development: improving the precision of age estimates in forensic entomology. *J Forensic Sci.* 56 Suppl 1:S112-122. <https://doi.10.1111/j.1556-4029.2010.01632.x>.
- Tarone AM, Foran DR (2008) Generalized additive models and *Lucilia sericata* growth: assessing confidence intervals and error rates in forensic entomology. *J Forensic Sci.* 53:942-948. <https://doi.10.1111/j.1556-4029.2008.00744.x>.
- Tarone AM, Picard CJ, Spiegelman C, Foran DR (2011) Population and temperature effects on *Lucilia sericata* (Diptera: Calliphoridae) body size and minimum development time. *J Med Entomol.* 48:1062-1068.
- Thevan K, Ahmad AH, Rawi CS, Singh B (2010) Growth of *Chrysomya megacephala* (Fabricius) maggots in a morgue cooler. *J Forensic Sci* 55:1656-1658. <https://doi.10.1111/j.1556-4029.2010.01485.x>.
- Thomas JK, Sanford MR, Longnecker M, Tomberlin JK (2016) Effects of Temperature and Tissue Type on the Development of *Megaselia scalaris* (Diptera: Phoridae). *J Med Entomol* 53:519-525.
- Turpin C, Kyle C, Beresford DV (2014) Postfeeding larval dispersal behavior of late season blow flies (calliphoridae) in Southern Ontario, Canada. *J Forensic Sci.* 59:1295-1302. <https://doi.10.1111/1556-4029.12469>.
- van der Ham JL (2016) Permutation Tests of Hierarchical Cluster Analyses of Carrion Communities and Their Potential Use in Forensic Entomology. *J Med Entomol.* 53:1238-1241. <https://doi.10.1093/jme/tjw069>.
- VanLaerhoven SL, Anderson GS (1999) Insect succession on buried carrion in two biogeoclimatic zones of British Columbia. *J Forensic Sci* 44:32-43.
- VanLaerhoven SL (2008) Blind validation of postmortem interval estimates using developmental rates of blow flies. *Forensic Sci Int* 180:76-80. <https://doi.10.1016/j.forsciint.2008.07.002>.

- Velásquez Y, Vilorio AL (2009) Effects of temperature on the development of the Neotropical carrion beetle *Oxelytrum discicolle* (Brullé, 1840) (Coleoptera: Silphidae). *Forensic Sci Int* 185:107-109. <https://doi.org/10.1016/j.forsciint.2008.12.020>.
- Velásquez Y, Vilorio A (2010) Instar determination of the neotropical beetle *Oxelytrum discicolle* (Coleoptera: Silphidae). *J Med Entomol* 47:723-726.
- Voss SC, Spafford H, Dadour IR (2009) Annual and seasonal patterns of insect succession on decomposing remains at two locations in Western Australia. *Forensic Sci Int* 193:26-36. <https://doi.org/10.1016/j.forsciint.2009.08.014>.
- Voss SC, Forbes SL, Dadour IR (2008) Decomposition and insect succession on cadavers inside a vehicle environment. *Forensic Sci Med Pathol* 4:22-32. <https://doi.org/10.1007/s12024-007-0028-z>.
- Wang J, Li Z, Chen Y, Chen Q, Yin X (2008) The succession and development of insects on pig carcasses and their significances in estimating PMI in south China. *Forensic Sci Int* 179:11-18. <https://doi.org/10.1016/j.forsciint.2008.04.014>.
- Wang Y, Gu ZY, Xia SX, Wang JF, Zhang YN, Tao LY (2018) Estimating the age of *Lucilia illustris* during the intrapuparial period using two approaches: Morphological changes and differential gene expression. *Forensic Sci Int* 287:1-11. <https://doi.org/10.1016/j.forsciint.2018.02.025>.
- Wang Y, Hu G, Zhang Y, Wang M, Amendt J, Wang J. (2019) Development of *Muscina stabulans* at constant temperatures with implications for minimum postmortem interval estimation. *Forensic Sci Int* 298:71-79. <https://doi.org/10.1016/j.forsciint.2019.02.051>.
- Wang Y, Li LL, Wang JF, Wang M, Yang LJ, Tao LY, Zhang YN, Hou YD, Chu J, Hou ZL (2016) Development of the green bottle fly *Lucilia illustris* at constant temperatures. *Forensic Sci Int* 267:136-144. <https://doi.org/10.1016/j.forsciint.2016.07.019>. Epub 2016 Aug 28. PubMed PMID: 27611954.
- Wang Y, Ma MY, Jiang XY, Wang JF, Li LL, Yin XJ, Wang M, Lai Y, Tao LY (2017) Insect succession on remains of human and animals in Shenzhen, China. *Forensic Sci Int* 271:75-86. <https://doi.org/10.1016/j.forsciint.2016.12.032>.
- Wang Y, Wang J, Wang Z, Tao L (2017) Insect succession on pig carcasses using different exposure time - A preliminary study in Guangzhou, China. *J Forensic Leg Med* 52:24-29. <https://doi.org/10.1016/j.jflm.2017.08.002>.
- Wang Y, Wang JF, Zhang YN, Tao LY, Wang M (2017) Forensically Important *Boettcherisca peregrina* (Diptera: Sarcophagidae) in China: Development Pattern and Significance for Estimating Postmortem Interval. *J Med Entomol* 54:1491-1497. <https://doi.org/10.1093/jme/tjx139>.
- Wang Y, Yang JB, Wang JF, Li LL, Wang M, Yang LJ, Tao LY, Chu J, Hou YD (2017) Development of the Forensically Important Beetle *Creophilus maxillosus* (Coleoptera: Staphylinidae) at Constant Temperatures. *J Med Entomol* 54:281-289. <https://doi.org/10.1093/jme/tjw193>.
- Wang Y, Yang L, Zhang Y, Tao L, Wang J (2018) Development of *Musca domestica* at constant temperatures and the first case report of its application for estimating the minimum postmortem interval. *Forensic Sci Int* 285:172-180. <https://doi.org/10.1016/j.forsciint.2018.02.004>.
- Wang Y, Zhang Y, Hu G, Wang M, Zhu R, Zhai Y, Sun J, Li X, Wang L, Wu M, Wang J (2020) Development of *Megaselia spiracularis* (Diptera: Phoridae) at different constant temperatures. *Journal of Thermal Biology* 93:102722. <https://doi.org/10.1016/j.jtherbio.2020.102722>
- Wang Y, Wang M, Hu G, Xu W, Wang Y, Wang J (2020) Temperature-dependent development of *Omosita colon* at constant temperature and its implication for PMImin estimation. *J Forensic and Legal Med* 72:101946. <https://doi.org/10.1016/j.jflm.2020.101946>.
- Wang Y, Gu Z, Li L, Wang J (2019) Gene expression during the intra-puparial stage of *Chrysomya megacephala*: Implications for postmortem interval estimation. *J Asia-Pacific Entomol* 22:841-846. <https://doi.org/10.1016/j.aspen.2019.06.012>.
- Warren GA, Anderson GS (2013) The development of *Protophormia terraenovae* (Robineau-Desvoidy) at constant temperatures and its minimum temperature threshold. *Forensic Sci Int* 233:374-379. <https://doi.org/10.1016/j.forsciint.2013.10.012>.

- Well J, Lamotte L (2017) The Role of a PMI-Prediction Model in Evaluating Forensic Entomology Experimental Design, the Importance of Covariates, and the Utility of Response Variables for Estimating Time Since Death. *Insects* 8:47. <https://doi.org/10.3390/insects8020047>
- Wells JD, Lecheta MC, Moura MO, LaMotte LR (2015) An evaluation of sampling methods used to produce insect growth models for postmortem interval estimation. *Int J Legal Med* 129:405-410. <https://doi.org/10.1007/s00414-014-1029-6>.
- Williams H (1984) A model for the aging of fly larvae in forensic entomology. *Forensic Sci Int* 25:191-199.
- Xu H, Ye GY, Xu Y, Hu C, Zhu GH (2014) Age-dependent changes in cuticular hydrocarbons of larvae in *Aldrichina grahami* (Aldrich) (Diptera: Calliphoridae). *Forensic Sci Int* 242:236-241. <https://doi.org/10.1016/j.forsciint.2014.07.003>.
- Yang L, Wang Y, Li L, Wang J, Wang M, Zhang Y, Chu J, Liu K, Hou Y, Tao L (2017) Temperature-dependent Development of *Parasarcophaga similis* (Meade 1876) and its significance in Estimating Postmortem Interval. *J Forensic Sci* 62:1234-1243. <https://doi.org/10.1111/1556-4029.13389>.
- Yang ST, Shiao SF (2012) Oviposition preferences of two forensically important blow fly species, *Chrysomya megacephala* and *C. rufifacies* (Diptera: Calliphoridae), and implications for postmortem interval estimation. *J Med Entomol* 49:424-435.
- Yang YQ, Li XB, Shao RY, Lyu Z, Li HW, Li GP, Xu LZ, Wan LH (2016) Developmental Times of *Chrysomya megacephala* (Fabricius) (Diptera: Calliphoridae) at Constant Temperatures and Applications in Forensic Entomology. *J Forensic Sci* 61:1278-1284. <https://doi.org/10.1111/1556-4029.13159>.
- Yang YQ, Lyu Z, Li XB, Li K, Yao L, Wan LH (2015) Technical note: Development of *Hemipyrellia ligurriens* (Wiedemann) (Diptera: Calliphoridae) at constant temperatures: Applications in estimating postmortem interval. *Forensic Sci Int* 253:48-54. <https://doi.org/10.1016/j.forsciint.2015.05.006>.
- Yanmanee S, Husemann M, Benbow ME, Suwannapong G (2016) Larval development rates of *Chrysomya rufifacies* Macquart, 1842 (Diptera: Calliphoridae) within its native range in South-East Asia. *Forensic Sci Int* 266:63-67. <https://doi.org/10.1016/j.forsciint.2016.04.033>.
- Zanetti NI, Visciarelli EC, Centeno ND (2016) The Effect of Temperature and Laboratory Rearing Conditions on the Development of *Dermestes maculatus* (Coleoptera: Dermestidae). *J Forensic Sci* 61:375-381. <https://doi.org/10.1111/1556-4029.12965>.
- Zehner R, Amendt J, Boehme P (2009) Gene expression analysis as a tool for age estimation of blowfly pupae. *Forensic Science International: Genetics Supplement Series* 2:292-293, <https://doi.org/10.1016/j.fsigss.2009.08.008>.
- Zehner R, Mösch S, Amendt J (2006) Estimating the postmortem interval by determining the age of fly pupae: Are there any molecular tools? *International Congress Series* 1288:619-621. <https://doi.org/10.1016/j.ics.2005.12.007>.
- Zhang X, Li Y, Shang Y, Ren L, Chen W, Wang S, Guo Y (2020) Development of *Sarcophaga dux* (Diptera: Sarcophagidae) at constant temperatures and differential gene expression for age estimation of the pupae. *Journal of Thermal Biology* 102735. <https://doi.org/10.1016/j.jtherbio.2020.102735>.
- Zhang Y, Wang Y, Liu C, Wang J, Hu G, Wang M, Yang L, Chu J (2019) Development of *Nasonia vitripennis* (Hymenoptera: Pteromalidae) at Constant Temperatures in China. *J Med Entomol* 56:368-377. <https://doi.org/10.1093/jme/tjy191>.
- Zhang Y, Wang Y, Sun J, Hu G, Wang M, Amendt J, Wang J (2020) Temperature-dependent development of the blow fly *Chrysomya pinguis* and its significance in estimating postmortem interval. *R Soc Open Sci* 6:190003. <https://doi.org/10.1098/rsos.190003>.
- Zhang Y, Wang Y, Yang L, Tao L, Wang J. (2017) Development of *Chrysomya megacephala* at constant temperatures within its colony range in Yangtze River Delta region of China. *Forensic Sci Res* 3:74-82. <https://doi.org/10.1080/20961790.2017.1403007>.

- Zhu GH, Jia ZJ, Yu XJ, Wu KS, Chen LS, Lv JY, Eric Benbow M (2017) Predictable weathering of puparial hydrocarbons of necrophagous flies for determining the postmortem interval: a field experiment using *Chrysomya rufifacies*. *Int J Legal Med* 131:885-894. [https://doi. 10.1007/s00414-016-1507-0](https://doi.org/10.1007/s00414-016-1507-0).
- Zhu GH, Xu XH, Yu XJ, Zhang Y, Wang JF (2007) Puparial case hydrocarbons of *Chrysomya megacephala* as an indicator of the postmortem interval. *Forensic Sci Int.* 169:1-5.
- Zhu GH, Ye GY, Li K, Hu C, Xu XH (2013) Determining the age of adult flesh flies, *Boettcherisca peregrina*, using pteridine fluorescence. *Med Vet Entomol* 27:59-63. [https://doi. 10.1111/j.1365-2915.2012.01021.x](https://doi.org/10.1111/j.1365-2915.2012.01021.x).
- Zuha RM, Omar B (2014) Developmental rate, size, and sexual dimorphism of *Megaselia scalaris* (Loew) (Diptera: Phoridae): its possible implications in forensic entomology. *Parasitol Res* 113:2285-94. [https://doi. 10.1007/s00436-014-3883-z](https://doi.org/10.1007/s00436-014-3883-z).
- Zuha RM, Razak TA, Ahmad NW, Omar B (2012) Interaction effects of temperature and food on the development of forensically important fly, *Megaselia scalaris* (Loew) (Diptera: Phoridae). *Parasitol Res.* 111:2179-87. [https://doi. 10.1007/s00436-012-3070-z](https://doi.org/10.1007/s00436-012-3070-z).