

Precise Measurements of Decay Parameters and CP Asymmetry with Entangled Λ - $\bar{\Lambda}$ Pairs

- M. Ablikim,¹ M. N. Achasov,^{11,b} P. Adlarson,⁷⁰ M. Albrecht,⁴ R. Aliberti,³¹ A. Amoroso,^{69a,69c} M. R. An,³⁵ Q. An,^{66,53} X. H. Bai,⁶¹ Y. Bai,⁵² O. Bakina,³² R. Baldini Ferroli,^{26a} I. Balossino,^{1,27a} Y. Ban,^{42,g} V. Batozskaya,^{1,40} D. Becker,³¹ K. Begzsuren,²⁹ N. Berger,³¹ M. Bertani,^{26a} D. Betttoni,^{27a} F. Bianchi,^{69a,69c} J. Bloms,⁶³ A. Bortone,^{69a,69c} I. Boyko,³² R. A. Briere,⁵ A. Brueggemann,⁶³ H. Cai,⁷¹ X. Cai,^{1,53} A. Calcaterra,^{26a} G. F. Cao,^{1,58} N. Cao,^{1,58} S. A. Cetin,^{57a} J. F. Chang,^{1,53} W. L. Chang,^{1,58} G. Chelkov,^{32,a} C. Chen,³⁹ Chao Chen,⁵⁰ G. Chen,¹ H. S. Chen,^{1,58} M. L. Chen,^{1,53} S. J. Chen,³⁸ S. M. Chen,⁵⁶ T. Chen,¹ X. R. Chen,^{28,58} X. T. Chen,¹ Y. B. Chen,^{1,53} Z. J. Chen,^{23,h} W. S. Cheng,^{69c} S. K. Choi,⁵⁰ X. Chu,³⁹ G. Cibinetto,^{27a} F. Cossio,^{69c} J. J. Cui,⁴⁵ H. L. Dai,^{1,53} J. P. Dai,⁷³ A. Dbeysi,¹⁷ R. E. de Boer,⁴ D. Dedovich,³² Z. Y. Deng,¹ A. Denig,³¹ I. Denysenko,³² M. Destefanis,^{69a,69c} F. De Mori,^{69a,69c} Y. Ding,³⁶ J. Dong,^{1,53} L. Y. Dong,^{1,58} M. Y. Dong,^{1,53,58} X. Dong,⁷¹ S. X. Du,⁷⁵ P. Egorov,^{32,a} Y. L. Fan,⁷¹ J. Fang,^{1,53} S. S. Fang,^{1,58} W. X. Fang,¹ Y. Fang,¹ R. Farinelli,^{27a} L. Fava,^{69b,69c} F. Feldbauer,⁴ G. Felici,^{26a} C. Q. Feng,^{66,53} J. H. Feng,⁵⁴ K. Fischer,⁶⁴ M. Fritsch,⁴ C. Fritzsch,⁶³ C. D. Fu,¹ H. Gao,⁵⁸ Y. N. Gao,^{42,g} Yang Gao,^{66,53} S. Garbolino,^{69c} I. Garzia,^{27a,27b} P. T. Ge,⁷¹ Z. W. Ge,³⁸ C. Geng,⁵⁴ E. M. Gersabeck,⁶² A. Gilman,⁶⁴ K. Goetzen,¹² L. Gong,³⁶ W. X. Gong,^{1,53} W. Gradl,³¹ M. Greco,^{69a,69c} L. M. Gu,³⁸ M. H. Gu,^{1,53} Y. T. Gu,¹⁴ C. Y. Guan,^{1,58} A. Q. Guo,^{28,58} L. B. Guo,³⁷ R. P. Guo,⁴⁴ Y. P. Guo,^{10,f} A. Guskov,^{32,a} T. T. Han,⁴⁵ W. Y. Han,³⁵ X. Q. Hao,¹⁸ F. A. Harris,⁶⁰ K. K. He,⁵⁰ K. L. He,^{1,58} F. H. Heinsius,⁴ C. H. Heinz,³¹ Y. K. Heng,^{1,53,58} C. Herold,⁵⁵ M. Himmelreich,^{31,d} G. Y. Hou,^{1,58} Y. R. Hou,⁵⁸ Z. L. Hou,¹ H. M. Hu,^{1,58} J. F. Hu,^{51,i} T. Hu,^{1,53,58} Y. Hu,¹ G. S. Huang,^{66,53} K. X. Huang,⁵⁴ L. Q. Huang,⁶⁷ L. Q. Huang,^{28,58} X. T. Huang,⁴⁵ Y. P. Huang,¹ Z. Huang,^{42,g} T. Hussain,⁶⁸ N. Hüskens,^{25,31} W. Imoehl,²⁵ M. Irshad,^{66,53} J. Jackson,²⁵ S. Jaeger,⁴ S. Janchiv,²⁹ E. Jang,⁵⁰ J. H. Jeong,⁵⁰ Q. Ji,¹ Q. P. Ji,¹⁸ X. B. Ji,^{1,58} X. L. Ji,^{1,53} Y. Y. Ji,⁴⁵ Z. K. Jia,^{66,53} H. B. Jiang,⁴⁵ S. S. Jiang,³⁵ X. S. Jiang,^{1,53,58} Y. Jiang,⁵⁸ J. B. Jiao,⁴⁵ Z. Jiao,²¹ S. Jin,³⁸ Y. Jin,⁶¹ M. Q. Jing,^{1,58} T. Johansson,⁷⁰ N. Kalantar-Nayestanaki,⁵⁹ X. S. Kang,³⁶ R. Kappert,⁵⁹ B. C. Ke,⁷⁵ I. K. Keshk,⁴ A. Khoukaz,⁶³ P. Kiese,³¹ R. Kiuchi,¹ R. Kliemt,¹² L. Koch,³³ O. B. Kolcu,^{57a} B. Kopf,⁴ M. Kuemmel,⁴ M. Kuessner,⁴ A. Kupsc,^{40,70} W. Kühn,³³ J. J. Lane,⁶² J. S. Lange,³³ P. Larin,¹⁷ A. Lavania,²⁴ L. Lavezzi,^{69a,69c} Z. H. Lei,^{66,53} H. Leithoff,³¹ M. Lellmann,³¹ T. Lenz,³¹ C. Li,⁴³ C. Li,³⁹ C. H. Li,³⁵ Cheng Li,^{66,53} D. M. Li,⁷⁵ F. Li,^{1,53} G. Li,¹ H. Li,⁴⁷ H. Li,^{66,53} H. B. Li,^{1,58} H. J. Li,¹⁸ H. N. Li,^{51,i} J. Q. Li,⁴ J. S. Li,⁵⁴ J. W. Li,⁴⁵ Ke Li,¹ L. J. Li,¹ L. K. Li,¹ Lei Li,³ M. H. Li,³⁹ P. R. Li,^{34,j,k} S. X. Li,¹⁰ S. Y. Li,⁵⁶ T. Li,⁴⁵ W. D. Li,^{1,58} W. G. Li,¹ X. H. Li,^{66,53} X. L. Li,⁴⁵ Xiaoyu Li,^{1,58} H. Liang,^{66,53} H. Liang,^{1,58} H. Liang,³⁰ Y. F. Liang,⁴⁹ Y. T. Liang,^{28,58} G. R. Liao,¹³ L. Z. Liao,⁴⁵ J. Libby,²⁴ A. Limphirat,⁵⁵ C. X. Lin,⁵⁴ D. X. Lin,^{28,58} T. Lin,¹ B. J. Liu,¹ C. X. Liu,¹ D. Liu,^{17,66} F. H. Liu,⁴⁸ Fang Liu,¹ Feng Liu,⁶ G. M. Liu,^{51,i} H. Liu,^{34,j,k} H. B. Liu,¹⁴ H. M. Liu,^{1,58} Huanhuan Liu,¹ Huihui Liu,¹⁹ J. B. Liu,^{66,53} J. L. Liu,⁶⁷ J. Y. Liu,^{1,58} K. Liu,¹ K. Y. Liu,³⁶ Ke Liu,²⁰ L. Liu,^{66,53} Lu Liu,³⁹ M. H. Liu,^{10,f} P. L. Liu,¹ Q. Liu,⁵⁸ S. B. Liu,^{66,53} T. Liu,^{10,f} W. K. Liu,³⁹ W. M. Liu,^{66,53} X. Liu,^{34,j,k} Y. Liu,^{34,j,k} Y. B. Liu,³⁹ Z. A. Liu,^{1,53,58} Z. Q. Liu,⁴⁵ X. C. Lou,^{1,53,58} F. X. Lu,⁵⁴ H. J. Lu,²¹ J. G. Lu,^{1,53} X. L. Lu,¹ Y. Lu,⁷ Y. P. Lu,^{1,53} Z. H. Lu,¹ C. L. Luo,³⁷ M. X. Luo,⁷⁴ T. Luo,^{10,f} X. L. Luo,^{1,53} X. R. Lyu,⁵⁸ Y. F. Lyu,³⁹ F. C. Ma,³⁶ H. L. Ma,¹ L. L. Ma,⁴⁵ M. M. Ma,^{1,58} Q. M. Ma,¹ R. Q. Ma,⁵⁸ R. T. Ma,^{1,53} Y. Ma,^{42,g} F. E. Maas,¹⁷ M. Maggiora,^{69a,69c} S. Maldaner,⁴ S. Malde,⁶⁴ Q. A. Malik,⁶⁸ A. Mangoni,^{26b} Y. J. Mao,^{42,g,g} Z. P. Mao,¹ S. Marcelli,^{69a,69c} Z. X. Meng,⁶¹ J. G. Messchendorp,^{59,12} G. Mezzadri,^{1,27a} H. Miao,¹ T. J. Min,³⁸ R. E. Mitchell,²⁵ X. H. Mo,^{1,53,58} N. Yu. Muchnoi,^{11,b} Y. Nefedov,³² F. Nerling,^{17,d} I. B. Nikolaev,¹¹ Z. Ning,^{1,53} S. Nisar,^{9,1} Y. Niu,⁴⁵ S. L. Olsen,⁵⁸ Q. Ouyang,^{1,53,58} S. Pacetti,^{26b,26c} X. Pan,^{10,f} Y. Pan,⁵² A. Pathak,¹ M. Pelizaeus,⁴ H. P. Peng,^{66,53} K. Peters,^{12,d} J. L. Ping,³⁷ R. G. Ping,^{1,58} S. Plura,³¹ S. Pogodin,³² V. Prasad,^{66,53} F. Z. Qi,¹ H. Qi,^{66,53} H. R. Qi,⁵⁶ M. Qi,³⁸ T. Y. Qi,^{10,f} S. Qian,^{1,53} W. B. Qian,⁵⁸ Z. Qian,⁵⁴ C. F. Qiao,⁵⁸ J. J. Qin,⁶⁷ L. Q. Qin,¹³ X. P. Qin,^{10,f} X. S. Qin,⁴⁵ Z. H. Qin,^{1,53} J. F. Qiu,¹ S. Q. Qu,⁵⁶ K. H. Rashid,⁶⁸ C. F. Redmer,³¹ K. J. Ren,³⁵ A. Rivetti,^{69c} V. Rodin,⁵⁹ M. Rolo,^{69c} G. Rong,^{1,58} Ch. Rosner,¹⁷ S. N. Ruan,³⁹ H. S. Sang,⁶⁶ A. Sarantsev,^{32,c} Y. Schelhaas,³¹ C. Schnier,⁴ K. Schönnung,⁷⁰ M. Scodeggio,^{27a,27b} K. Y. Shan,^{10,f} W. Shan,²² X. Y. Shan,^{66,53} J. F. Shangguan,⁵⁰ L. G. Shao,^{1,58} M. Shao,^{66,53} C. P. Shen,^{10,f} H. F. Shen,^{1,58} X. Y. Shen,^{1,58} B. A. Shi,⁵⁸ H. C. Shi,^{66,53} J. Y. Shi,¹ Q. Q. Shi,⁵⁰ R. S. Shi,^{1,58} X. Shi,^{1,53} X. D. Shi,^{66,53} J. J. Song,¹⁸ W. M. Song,^{1,30} Y. X. Song,^{42,g} S. Sosio,^{69a,69c} S. Spataro,^{69a,69c} F. Stieler,³¹ K. X. Su,⁷¹ P. P. Su,⁵⁰ Y. J. Su,⁵⁸ G. X. Sun,¹ H. Sun,⁵⁸ H. K. Sun,¹ J. F. Sun,¹⁸ L. Sun,⁷¹ S. S. Sun,^{1,58} T. Sun,^{1,58} W. Y. Sun,³⁰ X. Sun,^{23,h} Y. J. Sun,^{66,53} Y. Z. Sun,¹ Z. T. Sun,⁴⁵ Y. H. Tan,⁷¹ Y. X. Tan,^{66,53} C. J. Tang,⁴⁹ G. Y. Tang,¹ J. Tang,⁵⁴ L. Y. Tao,⁶⁷ Q. T. Tao,^{23,h} M. Tat,⁶⁴ J. X. Teng,^{66,53} V. Thoren,⁷⁰ W. H. Tian,⁴⁷ Y. Tian,^{28,58} I. Uman,^{57b} B. Wang,¹ B. L. Wang,⁵⁸ C. W. Wang,³⁸ D. Y. Wang,^{42,g} F. Wang,⁶⁷ H. J. Wang,^{34,j,k} H. P. Wang,^{1,58} K. Wang,^{1,53} L. L. Wang,¹ M. Wang,⁴⁵ M. Z. Wang,^{42,g} Meng Wang,^{1,58} S. Wang,¹³ S. Wang,^{10,f} T. Wang,^{10,f} T. J. Wang,³⁹ W. Wang,⁵⁴ W. H. Wang,⁷¹ W. P. Wang,^{66,53} X. Wang,^{42,g} X. F. Wang,^{34,j,k} X. L. Wang,^{10,f} Y. Wang,⁵⁶ Y. D. Wang,⁴¹ Y. F. Wang,^{1,53,58} Y. H. Wang,⁴³

Y. Q. Wang,¹ Yaqian Wang,^{1,16} Z. Wang,^{1,53} Z. Y. Wang,^{1,58} Ziyi Wang,⁵⁸ D. H. Wei,¹³ F. Weidner,⁶³ S. P. Wen,¹
 D. J. White,⁶² U. Wiedner,⁴ G. Wilkinson,⁶⁴ M. Wolke,⁷⁰ L. Wollenberg,⁴ J. F. Wu,^{1,58} L. H. Wu,¹ L. J. Wu,^{1,58} X. Wu,^{10,f}
 X. H. Wu,³⁰ Y. Wu,⁶⁶ Z. Wu,^{1,53} L. Xia,^{66,53} T. Xiang,^{42,g} D. Xiao,^{34,j,k} G. Y. Xiao,³⁸ H. Xiao,^{10,f} S. Y. Xiao,¹ Y. L. Xiao,^{10,f}
 Z. J. Xiao,³⁷ C. Xie,³⁸ X. H. Xie,^{42,g} Y. Xie,⁴⁵ Y. G. Xie,^{1,53} Y. H. Xie,⁶ Z. P. Xie,^{66,53} T. Y. Xing,^{1,58} C. F. Xu,¹ C. J. Xu,⁵⁴
 G. F. Xu,¹ H. Y. Xu,⁶¹ Q. J. Xu,¹⁵ X. P. Xu,⁵⁰ Y. C. Xu,⁵⁸ Z. P. Xu,³⁸ F. Yan,^{10,f} L. Yan,^{10,f} W. B. Yan,^{66,53} W. C. Yan,⁷⁵
 H. J. Yang,^{46,e} H. L. Yang,³⁰ H. X. Yang,¹ L. Yang,⁴⁷ S. L. Yang,⁵⁸ Tao Yang,¹ Y. F. Yang,³⁹ Y. X. Yang,^{1,58} Yifan Yang,¹
 M. Ye,^{1,53} M. H. Ye,⁸ J. H. Yin,¹ Z. Y. You,⁵⁴ B. X. Yu,^{1,53,58} C. X. Yu,³⁹ G. Yu,^{1,58} T. Yu,⁶⁷ C. Z. Yuan,^{1,58} L. Yuan,²
 S. C. Yuan,¹ X. Q. Yuan,¹ Y. Yuan,^{1,58} Z. Y. Yuan,⁵⁴ C. X. Yue,³⁵ A. A. Zafar,⁶⁸ F. R. Zeng,⁴⁵ X. Zeng,⁶ Y. Zeng,^{23,h}
 Y. H. Zhan,⁵⁴ A. Q. Zhang,¹ B. L. Zhang,¹ B. X. Zhang,¹ D. H. Zhang,³⁹ G. Y. Zhang,¹⁸ H. Zhang,⁶⁶ H. H. Zhang,⁵⁴
 H. H. Zhang,³⁰ H. Y. Zhang,^{1,53} J. L. Zhang,⁷² J. Q. Zhang,³⁷ J. W. Zhang,^{1,53,58} J. X. Zhang,^{34,j,k} J. Y. Zhang,¹
 J. Z. Zhang,^{1,58} Jianyu Zhang,^{1,58} Jiawei Zhang,^{1,58} L. M. Zhang,⁵⁶ L. Q. Zhang,⁵⁴ Lei Zhang,³⁸ P. Zhang,¹
 Q. Y. Zhang,^{35,75} Shuihan Zhang,^{1,58} Shulei Zhang,^{23,h} X. D. Zhang,⁴¹ X. M. Zhang,¹ X. Y. Zhang,⁴⁵ X. Y. Zhang,⁵⁰
 Y. Zhang,⁶⁴ Y. T. Zhang,⁷⁵ Y. H. Zhang,^{1,53} Yan Zhang,^{66,53} Yao Zhang,¹ Z. H. Zhang,¹ Z. Y. Zhang,⁷¹ Z. Y. Zhang,³⁹
 G. Zhao,¹ J. Zhao,³⁵ J. Y. Zhao,^{1,58} J. Z. Zhao,^{1,53} Lei Zhao,^{66,53} Ling Zhao,¹ M. G. Zhao,³⁹ Q. Zhao,¹ S. J. Zhao,⁷⁵
 Y. B. Zhao,^{1,53} Y. X. Zhao,^{28,58} Z. G. Zhao,^{66,53} A. Zhemchugov,^{32,a} B. Zheng,⁶⁷ J. P. Zheng,^{1,53} Y. H. Zheng,⁵⁸ B. Zhong,³⁷
 C. Zhong,⁶⁷ X. Zhong,⁵⁴ H. Zhou,⁴⁵ L. P. Zhou,^{1,58} X. Zhou,⁷¹ X. K. Zhou,⁵⁸ X. R. Zhou,^{66,53} X. Y. Zhou,³⁵ Y. Z. Zhou,^{10,f}
 J. Zhu,³⁹ K. Zhu,¹ K. J. Zhu,^{1,53,58} L. X. Zhu,⁵⁸ S. H. Zhu,⁶⁵ S. Q. Zhu,³⁸ T. J. Zhu,⁷² W. J. Zhu,^{10,f} Y. C. Zhu,^{66,53}
 Z. A. Zhu,^{1,58} B. S. Zou,¹ and J. H. Zou¹

(BESIII Collaboration)

¹*Institute of High Energy Physics, Beijing 100049, People's Republic of China*²*Beihang University, Beijing 100191, People's Republic of China*³*Beijing Institute of Petrochemical Technology, Beijing 102617, People's Republic of China*⁴*Bochum Ruhr-University, D-44780 Bochum, Germany*⁵*Carnegie Mellon University, Pittsburgh, Pennsylvania 15213, USA*⁶*Central China Normal University, Wuhan 430079, People's Republic of China*⁷*Central South University, Changsha 410083, People's Republic of China*⁸*China Center of Advanced Science and Technology, Beijing 100190, People's Republic of China*⁹*COMSATS University Islamabad, Lahore Campus, Defence Road, Off Raiwind Road, 54000 Lahore, Pakistan*¹⁰*Fudan University, Shanghai 200433, People's Republic of China*¹¹*G.I. Budker Institute of Nuclear Physics SB RAS (BINP), Novosibirsk 630090, Russia*¹²*GSI Helmholtzcentre for Heavy Ion Research GmbH, D-64291 Darmstadt, Germany*¹³*Guangxi Normal University, Guilin 541004, People's Republic of China*¹⁴*Guangxi University, Nanning 530004, People's Republic of China*¹⁵*Hangzhou Normal University, Hangzhou 310036, People's Republic of China*¹⁶*Hebei University, Baoding 071002, People's Republic of China*¹⁷*Helmholtz Institute Mainz, Staudinger Weg 18, D-55099 Mainz, Germany*¹⁸*Henan Normal University, Xinxiang 453007, People's Republic of China*¹⁹*Henan University of Science and Technology, Luoyang 471003, People's Republic of China*²⁰*Henan University of Technology, Zhengzhou 450001, People's Republic of China*²¹*Huangshan College, Huangshan 245000, People's Republic of China*²²*Hunan Normal University, Changsha 410081, People's Republic of China*²³*Hunan University, Changsha 410082, People's Republic of China*²⁴*Indian Institute of Technology Madras, Chennai 600036, India*²⁵*Indiana University, Bloomington, Indiana 47405, USA*^{26a}*INFN Laboratori Nazionali di Frascati, I-00044 Frascati, Italy*^{26b}*INFN Sezione di Perugia, I-06100 Perugia, Italy*^{26c}*University of Perugia, I-06100 Perugia, Italy*^{27a}*INFN Sezione di Ferrara, I-44122 Ferrara, Italy*^{27b}*University of Ferrara, I-44122 Ferrara, Italy*²⁸*Institute of Modern Physics, Lanzhou 730000, People's Republic of China*²⁹*Institute of Physics and Technology, Peace Avenue 54B, Ulaanbaatar 13330, Mongolia*³⁰*Jilin University, Changchun 130012, People's Republic of China*³¹*Johannes Gutenberg University of Mainz, Johann-Joachim-Becher-Weg 45, D-55099 Mainz, Germany*³²*Joint Institute for Nuclear Research, 141980 Dubna, Moscow region, Russia*³³*Justus-Liebig-Universitaet Giessen, II. Physikalisches Institut, Heinrich-Buff-Ring 16, D-35392 Giessen, Germany*

- ³⁴Lanzhou University, Lanzhou 730000, People's Republic of China
³⁵Liaoning Normal University, Dalian 116029, People's Republic of China
³⁶Liaoning University, Shenyang 110036, People's Republic of China
³⁷Nanjing Normal University, Nanjing 210023, People's Republic of China
³⁸Nanjing University, Nanjing 210093, People's Republic of China
³⁹Nankai University, Tianjin 300071, People's Republic of China
⁴⁰National Centre for Nuclear Research, Warsaw 02-093, Poland
⁴¹North China Electric Power University, Beijing 102206, People's Republic of China
⁴²Peking University, Beijing 100871, People's Republic of China
⁴³Qufu Normal University, Qufu 273165, People's Republic of China
⁴⁴Shandong Normal University, Jinan 250014, People's Republic of China
⁴⁵Shandong University, Jinan 250100, People's Republic of China
⁴⁶Shanghai Jiao Tong University, Shanghai 200240, People's Republic of China
⁴⁷Shanxi Normal University, Linfen 041004, People's Republic of China
⁴⁸Shanxi University, Taiyuan 030006, People's Republic of China
⁴⁹Sichuan University, Chengdu 610064, People's Republic of China
⁵⁰Soochow University, Suzhou 215006, People's Republic of China
⁵¹South China Normal University, Guangzhou 510006, People's Republic of China
⁵²Southeast University, Nanjing 211100, People's Republic of China
⁵³State Key Laboratory of Particle Detection and Electronics, Beijing 100049, Hefei 230026, People's Republic of China
⁵⁴Sun Yat-Sen University, Guangzhou 510275, People's Republic of China
⁵⁵Suranaree University of Technology, University Avenue 111, Nakhon Ratchasima 30000, Thailand
⁵⁶Tsinghua University, Beijing 100084, People's Republic of China
^{57a}Turkish Accelerator Center Particle Factory Group, Istinye University, 34010 Istanbul, Turkey
^{57b}Near East University, Nicosia, North Cyprus, Mersin 10, Turkey
⁵⁸University of Chinese Academy of Sciences, Beijing 100049, People's Republic of China
⁵⁹University of Groningen, NL-9747 AA Groningen, Netherlands
⁶⁰University of Hawaii, Honolulu, Hawaii 96822, USA
⁶¹University of Jinan, Jinan 250022, People's Republic of China
⁶²University of Manchester, Oxford Road, Manchester, M13 9PL, United Kingdom
⁶³University of Muenster, Wilhelm-Klemm-Strasse 9, 48149 Muenster, Germany
⁶⁴University of Oxford, Keble Road, Oxford OX13RH, United Kingdom
⁶⁵University of Science and Technology Liaoning, Anshan 114051, People's Republic of China
⁶⁶University of Science and Technology of China, Hefei 230026, People's Republic of China
⁶⁷University of South China, Hengyang 421001, People's Republic of China
⁶⁸University of the Punjab, Lahore-54590, Pakistan
^{69a}University of Turin and INFN, University of Turin, I-10125 Turin, Italy
^{69b}University of Eastern Piedmont, I-15121 Alessandria, Italy
^{69c}INFN, I-10125 Turin, Italy
⁷⁰Uppsala University, Box 516, SE-75120 Uppsala, Sweden
⁷¹Wuhan University, Wuhan 430072, People's Republic of China
⁷²Xinyang Normal University, Xinyang 464000, People's Republic of China
⁷³Yunnan University, Kunming 650500, People's Republic of China
⁷⁴Zhejiang University, Hangzhou 310027, People's Republic of China
⁷⁵Zhengzhou University, Zhengzhou 450001, People's Republic of China



(Received 26 April 2022; accepted 1 September 2022; published 22 September 2022)

Based on 10 billion J/ψ events collected at the BESIII experiment, a search for CP violation in Λ decay is performed in the difference between CP -odd decay parameters α_- for $\Lambda \rightarrow p\pi^-$ and α_+ for $\bar{\Lambda} \rightarrow \bar{p}\pi^+$ by using the process $e^+e^- \rightarrow J/\psi \rightarrow \Lambda\bar{\Lambda}$. With a five-dimensional fit to the full angular distributions of the daughter baryon, the most precise values for the decay parameters are determined to be $\alpha_- = 0.7519 \pm 0.0036 \pm 0.0024$ and $\alpha_+ = -0.7559 \pm 0.0036 \pm 0.0030$, respectively. The Λ and $\bar{\Lambda}$ averaged value of the decay parameter is extracted to be $\alpha_{\text{avg}} = 0.7542 \pm 0.0010 \pm 0.0024$ with unprecedented accuracy. The CP asymmetry $A_{CP} = (\alpha_- + \alpha_+)/(\alpha_- - \alpha_+)$ is determined to be $-0.0025 \pm 0.0046 \pm 0.0012$, which is

one of the most precise measurements in the baryon sector. The reported results for the decay parameter will play an important role in the studies of the polarizations and CP violations for the strange, charmed and beauty baryons.

DOI: 10.1103/PhysRevLett.129.131801

Charge-parity (CP) violation is a subject of continuing interest. To date, CP violation has been discovered in the K [1], B [2,3], and D meson [4] systems, but it has never been observed in the decays of any baryon. Hence, it is vital to search for additional sources of CP violation. Moreover, the standard model(SM) is difficult to explain the phenomenon of matter-antimatter asymmetry of the universe [5–7]. Thus, the CP test is an ideal and sensitive method to search for the physics beyond SM [8,9].

The promising CP -violating signature in spin- $\frac{1}{2}$ nonleptonic hyperon decays is the difference between hyperon and antihyperon decay distributions in their parity-violating two-body weak decays [10]. In such decays, the angular distribution of the daughter baryon is proportional to $(1 + \alpha_Y \mathbf{P}_Y \cdot \hat{\mathbf{p}}_d)$, where α_Y is the hyperon decay parameter, and \mathbf{P}_Y and $\hat{\mathbf{p}}_d$ are the hyperon polarization and the unit vector in the direction of the daughter baryon momentum, respectively, both in the hyperon rest frame. The CP asymmetry is defined as $A_{CP} = ([\alpha_Y + \alpha_{\bar{Y}}]/[\alpha_Y - \alpha_{\bar{Y}}])$. The parameters α_Y and $\alpha_{\bar{Y}}$ are CP odd so that a nonzero A_{CP} indicates CP violation. In the SM, the Cabibbo-Kobayashi-Maskawa mechanism predicts a tiny A_{CP} value of $\sim 10^{-4}$ [11]. Therefore, the hyperon decay is sensitive to the sources of CP asymmetry from physics beyond the SM [12,13]. A precise measurement of the $\Lambda(\bar{\Lambda})$ decay parameters is important for studies of spin polarization [14–17] and decay parameters [18–24] of many other baryons ($\Sigma^0, \Xi^0, \Xi^-, \Omega^-, \Lambda_c, \Lambda_b$, etc.) decays into final states involving Λ .

A total sample of 10 billion J/ψ events has been collected by the BESIII experiment, about 3.2 million quantum-entangled Λ - $\bar{\Lambda}$ pairs are expected to be fully reconstructed in the decay $J/\psi \rightarrow \Lambda\bar{\Lambda}$ with $\Lambda(\bar{\Lambda}) \rightarrow p\pi^-(\bar{p}\pi^+)$ [25]. Hence, in this Letter, we present the most precise measurements of Λ decay parameters and CP asymmetry with a five-dimensional fit to the full angular distributions of the daughter baryon.

Two real parameters, the $J/\psi \rightarrow \Lambda\bar{\Lambda}$ angular distribution parameter $\alpha_{J/\psi}$ and the helicity phase difference $\Delta\Phi$, describe the angular distribution and polarization of the produced Λ and $\bar{\Lambda}$ [26]. If the phase difference $\Delta\Phi$ is nonvanishing, the polarization of the Λ and $\bar{\Lambda}$ will be oriented perpendicular to the production plane. For the decay $\Lambda \rightarrow p\pi^-$, the angular distribution of the proton is $\frac{1}{4\pi}(1 + \alpha_- \mathbf{P}_\Lambda \cdot \hat{\mathbf{n}})$, where α_- is Λ decay parameter, \mathbf{P}_Λ is the polarization vector of the Λ , $\hat{\mathbf{n}}$ is the unit vector of the proton momentum in the Λ rest frame. The definition of the decay parameter α_+ for $\bar{\Lambda} \rightarrow \bar{p}\pi^+$ follows an analogous convention [27].

For the cascade decay $J/\psi \rightarrow \Lambda\bar{\Lambda}$ with $\Lambda(\bar{\Lambda}) \rightarrow p\pi^-(\bar{p}\pi^+)$, the angular distribution of each event is uniquely characterized by the kinematic variable $\xi = (\theta_\Lambda, \theta_p, \phi_p, \theta_{\bar{p}}, \phi_{\bar{p}})$, where θ_p, ϕ_p and $\theta_{\bar{p}}, \phi_{\bar{p}}$ are the polar and azimuth angles of the proton and antiproton in their mother particles' rest frames. The components of these vectors are expressed using a right-handed coordinate system $(\hat{x}, \hat{y}, \hat{z})$ shown in Fig. 1. The \hat{z} axis is taken along the Λ momentum $\mathbf{p}_\Lambda = -\mathbf{p}_{\bar{\Lambda}} \equiv \mathbf{p}$ in the e^+e^- center-of-mass system (CMS). The \hat{y} axis is perpendicular to the production plane and oriented along the vector $\mathbf{k} \times \mathbf{p}$, where $\mathbf{k}_{e^-} = -\mathbf{k}_{e^+} \equiv \mathbf{k}$ is the electron beam momentum. The scattering angle of the Λ is given by $\cos\theta_\Lambda = \hat{\mathbf{p}} \cdot \hat{\mathbf{k}}$.

The differential distribution function can be expressed as

$$\mathcal{W}(\xi)$$

$$\begin{aligned} &= \mathcal{F}_0(\xi) + \alpha_{J/\psi} \mathcal{F}_5(\xi) + \alpha_- \alpha_+ \\ &\quad \times \left[\mathcal{F}_1(\xi) + \sqrt{1 - \alpha_{J/\psi}^2} \cos(\Delta\Phi) \mathcal{F}_2(\xi) + \alpha_{J/\psi} \mathcal{F}_6(\xi) \right] \\ &\quad + \sqrt{1 - \alpha_{J/\psi}^2} \sin(\Delta\Phi) [\alpha_- \mathcal{F}_3(\xi) + \alpha_+ \mathcal{F}_4(\xi)], \end{aligned} \quad (1)$$

where the angular functions $\mathcal{F}_i(\xi)$ ($i = 0, 1, \dots, 6$) are described in detail in Ref. [26]. The terms proportional to $\alpha_- \alpha_+$ in Eq. (1) represent the contribution from Λ - $\bar{\Lambda}$ spin correlations, and the terms proportional to α_- and α_+ separately represent the contribution from the hyperon transverse polarization P_y , defined as

$$P_y(\cos\theta_\Lambda) = \frac{\sqrt{1 - \alpha_{J/\psi}^2} \sin(\Delta\Phi) \cos\theta_\Lambda \sin\theta_\Lambda}{1 + \alpha_{J/\psi} \cos^2\theta_\Lambda}. \quad (2)$$

The analysis presented here is based on the aforementioned sample of 10 billion J/ψ events [28] collected at the




FIG. 1. Definition of the right-hand coordinate system used to describe the $J/\psi \rightarrow \Lambda\bar{\Lambda}$ process. In the e^+e^- center-of-mass system, the Λ is emitted along the \hat{z} axis direction. \hat{y} axis is perpendicular to the plane of Λ and e^- . The hyperons are polarized along the \hat{y} direction.

BESIII detector [29,30]. A Monte Carlo (MC) simulation of J/ψ samples is used to determine the detector efficiency, optimize the event selection, and estimate the background. The simulation is performed by the GEANT4-based [31] BESIII Object Oriented Simulation Tool project [32], which includes the geometric description of the BESIII detector and the detector response. The MC event generators KKMC [33], BesEvtGen [34], and Lundcharm [35,36] are used to describe J/ψ production together with known and unknown decay modes. For the signal process, $J/\psi \rightarrow \Lambda\bar{\Lambda}$, the parameters of the angular distribution are obtained from previous measurements [12]. For the dominant background channel $J/\psi \rightarrow \gamma\eta_c(\eta_c \rightarrow \Lambda\bar{\Lambda})$, the decay $J/\psi \rightarrow \gamma\eta_c$ is generated with an angular distribution of $1 + \cos^2 \theta_\gamma$ [37], where θ_γ is the angle between the photon and positron beam direction in the CMS, and another background channel $J/\psi \rightarrow \gamma\Lambda\bar{\Lambda}$ is described by the phase space model.

The Λ and $\bar{\Lambda}$ baryons are reconstructed from their dominant hadronic decay mode, $\Lambda(\bar{\Lambda}) \rightarrow p\pi^-(\bar{p}\pi^+)$. Charged tracks detected in the main drift chamber must satisfy $|\cos(\theta)| < 0.93$, where θ is the angle between the charged track and the positron beam direction. Events with at least four charged tracks are retained. Tracks with momentum larger than $0.5 \text{ GeV}/c$ are considered as proton candidates, otherwise, as pion candidates. There are no further particle identification requirements. Vertex fits are performed by looping over all combinations with oppositely charged proton and pion candidates, constraining them to a common vertex. The pairs with vertex fit χ^2 lower than 200 and decay length larger than 0 are regarded as $\Lambda\bar{\Lambda}$ candidates. A four-momentum constrained kinematic fit is applied to the $p\bar{p}\pi^+\pi^-$ hypothesis, and events with a minimum χ^2 lower than 60 are selected as J/ψ candidates.

An inclusive MC sample of 10 billion J/ψ events is used for studying potential backgrounds. After applying the same selection criteria as for the data, the main backgrounds are divided into two types according to the shapes of $m_{p\pi^-}$ and $m_{\bar{p}\pi^+}$: (1)BKG I, nonpeaking backgrounds, including $J/\psi \rightarrow p\pi^-\bar{p}\pi^+$, $\Delta^{++}\bar{p}\pi^-$, $\bar{\Delta}^{++}p\pi^+$, and $\Delta^{++}\bar{\Delta}^{++}$; (2)BKG II, peaking backgrounds, including $J/\psi \rightarrow \gamma\Lambda\bar{\Lambda}$, $\gamma\eta_c(\eta_c \rightarrow \Lambda\bar{\Lambda})$. The number of nonpeaking backgrounds is estimated by the two-dimensional sideband regions of the $m_{p\pi^-}$ versus $m_{\bar{p}\pi^+}$ distribution from the data sample which is shown in Fig. 2. The signal region is defined as $m_{p\pi^-/\bar{p}\pi^+} \in [1.111, 1.121] \text{ GeV}/c^2$, and the lower and higher sideband regions are defined as $m_{p\pi^-/\bar{p}\pi^+} \in [1.098, 1.107] \text{ GeV}/c^2$ and $m_{p\pi^-/\bar{p}\pi^+} \in [1.125, 1.134] \text{ GeV}/c^2$, respectively. The yields of various peaking background sources are estimated by individual exclusive MC samples, then normalized to the data sample according to their branching fractions [27]. The final data sample contains 3231781 events including the estimated




FIG. 2. Distribution of the invariant mass spectra of $\bar{p}\pi^+$ versus the invariant mass spectra of $p\pi^-$ from the data. The signal and sideband regions are denoted by red and green boxes, respectively. The z axis is in logarithm style.

background yield of 3801 ± 63 events. The sample has a high purity of 99.9%.

Based on the joint angular distribution, a maximum likelihood fit with four free parameters ($\alpha_{J/\psi}$, $\Delta\Phi$, α_- , and α_+) is performed. The joint likelihood function is defined as

$$\begin{aligned} \mathcal{L} &= \prod_{i=1}^N \mathcal{P}(\xi^i; \alpha_{J/\psi}, \Delta\Phi, \alpha_-, \alpha_+) \\ &= \prod_{i=1}^N \mathcal{CW}(\xi^i; \alpha_{J/\psi}, \Delta\Phi, \alpha_-, \alpha_+) \epsilon(\xi^i), \end{aligned} \quad (3)$$

where $\mathcal{P}(\xi^i; \alpha_{J/\psi}, \Delta\Phi, \alpha_-, \alpha_+)$ is the probability density function of ξ^i , the kinematic variable of event i , and $\mathcal{W}(\xi^i; \alpha_{J/\psi}, \Delta\Phi, \alpha_-, \alpha_+)$ is given by Eq. (1). The detection efficiency is denoted by $\epsilon(\xi^i)$. The normalization factor $\mathcal{C}^{-1} = (1/N_{\text{MC}}) \sum_{j=1}^{N_{\text{MC}}} \mathcal{W}(\xi^j; \alpha_{J/\psi}, \Delta\Phi, \alpha_-, \alpha_+) \epsilon(\xi^j)$ is estimated with the N_{MC} events generated with the phase space model, applying the same event selection criteria as for the data. To improve the accuracy of the normalization factor, we generate a MC sample about 100 times larger than the selected experimental data. We use the ROOFIT package [38] to determine the fit parameters from the minimization of the function

$$\mathcal{S} = -\ln \mathcal{L}_{\text{data}} + \ln \mathcal{L}_{\text{BKG I}} + \ln \mathcal{L}_{\text{BKG II}}, \quad (4)$$

where $\mathcal{L}_{\text{data}}$ and $\mathcal{L}_{\text{BKG I}}$ are the likelihood function of events in the signal region and sideband regions, respectively. The $\mathcal{L}_{\text{BKG II}}$ is the likelihood function of background events obtained by exclusive MC samples. The likelihood function of background event is the same as the data. The results of

TABLE I. The angular distribution parameters, $\alpha_{J/\psi}$, $\Delta\Phi$ and the asymmetry parameters α_- for $\Lambda \rightarrow p\pi^-$, α_+ for $\bar{\Lambda} \rightarrow \bar{p}\pi^+$ obtained in this Letter and in previous BESIII measurements [12] for comparison. The first uncertainty is statistical, the second one is systematic.

Parameter	This Letter	Previous results [12]
$\alpha_{J/\psi}$	$0.4748 \pm 0.0022 \pm 0.0031$	$0.461 \pm 0.006 \pm 0.007$
$\Delta\Phi$	$0.7521 \pm 0.0042 \pm 0.0066$	$0.740 \pm 0.010 \pm 0.009$
α_-	$0.7519 \pm 0.0036 \pm 0.0024$	$0.750 \pm 0.009 \pm 0.004$
α_+	$-0.7559 \pm 0.0036 \pm 0.0030$	$-0.758 \pm 0.010 \pm 0.007$
A_{CP}	$-0.0025 \pm 0.0046 \pm 0.0012$	$0.006 \pm 0.012 \pm 0.007$
α_{avg}	$0.7542 \pm 0.0010 \pm 0.0024$...

the maximum likelihood fit of data are given in Table I, with the CP asymmetry given by $A_{CP} = (\alpha_- + \alpha_+)/(\alpha_- - \alpha_+)$, and the average value of the Λ and $\bar{\Lambda}$ decay parameters $\alpha_{avg} = (\alpha_- - \alpha_+)/2$. The correlation coefficient between α_- and α_+ is $\rho(\alpha_-, \alpha_+) = 0.850$.

The moment

$$\mu[\cos(\theta_\Lambda)] = (m/N) \sum_{i=1}^{N_k} (n_{1,y}^{(i)} - n_{2,y}^{(i)}), \quad (5)$$

which related to the polarization, is used to compare the consistency between the data and the fit results. Hereby, N is the total number of events in the data set, and $m = 100$ is the number of bins in $\cos(\theta_\Lambda)$ for calculating the moment. N_k denotes the number of events in the k th $\cos(\theta_\Lambda)$ bin. The expected angular dependence of the moment for the acceptance-corrected data reads

$$\mu(\cos \theta_\Lambda) = \frac{\alpha_- - \alpha_+}{2} \frac{1 + \alpha_{J/\psi} \cos^2 \theta_\Lambda}{3 + \alpha_{J/\psi}} P_y(\theta_\Lambda). \quad (6)$$




FIG. 3. Distribution of moment $\mu(\cos \theta_\Lambda)$ versus $\cos \theta_\Lambda$. The points with error bars are the data, and the red histogram is the signal MC sample with input parameters fixed to fit results. The blue histogram shows the result from phase space(PHSP) MC sample. The distribution of $\chi = (\mu_{data} - \mu_{MC})/\sigma(\mu_{data})$ is shown at the bottom, where μ_{data} and μ_{MC} are the moments of data and signal MC sample. The $\sigma(\mu_{data})$ is the statistical uncertainty of μ_{data} .

A significant transverse polarization of Λ and $\bar{\Lambda}$ can be seen in Fig. 3, in which the points with error bars are the data, and the solid line is obtained from signal MC sample generated by Eq. (1), where the input parameters are taken from fit results. The data are consistent with the fit results.

The systematic uncertainties in this analysis can be divided into two categories: (A) the uncertainties from event selection, including background estimation, tracking, the $\Lambda/\bar{\Lambda}$ vertex fit and kinematic fit; (B) the uncertainty associated with the fit procedure. The uncertainty from background is estimated by varying the input background numbers by 1 standard deviation. The differences on the fitted parameters are taken as the systematic uncertainty. For the tracking and $\Lambda/\bar{\Lambda}$ vertex fit and kinematic fit, a correction to the MC efficiency is made. We use control samples to get the efficiencies of the data and the MC simulation in tracking, $\Lambda/\bar{\Lambda}$ vertex fit, and kinematic fit, and use the data and MC difference to calibrate the MC sample. The uncertainty due to the charged particle tracking efficiency has been investigated with a $J/\psi \rightarrow p\pi^- \bar{p}\pi^+$ control sample. The systematic uncertainties due to the Λ and $\bar{\Lambda}$ vertex reconstruction and kinematic fits are estimated by a control sample $J/\psi \rightarrow \Lambda\bar{\Lambda} \rightarrow p\pi^- \bar{p}\pi^+$. In order to reduce the impact of statistical fluctuations, the fit with a corrected MC sample is performed 100 times by varying the correction factor randomly within 1 standard deviation. The differences between the mean value of the fit results with corrections and the nominal fit are taken as the systematic uncertainties. The MC simulation is used to estimate the uncertainty of the fit method. The sum of the differences between the input and output values and their uncertainty are regarded as systematic uncertainties. The absolute systematic uncertainties for various sources are summarized in Table II. The total systematic uncertainty of each parameter is obtained by summing the individual contributions in quadrature.

In summary, by analyzing 10 billion J/ψ events, we report the most precise measurements of the decay parameters of $\Lambda(\bar{\Lambda})$, with results given in Table I. The results are consistent with those in the previous analysis [12], however, with significantly improved accuracy. The measured

TABLE II. Absolute systematic uncertainties for the measured parameters $\alpha_{J/\psi}$, $\Delta\Phi$, the asymmetry parameters α_- for $\Lambda \rightarrow p\pi^-$, α_+ for $\bar{\Lambda} \rightarrow \bar{p}\pi^+$, the CP asymmetry value A_{CP} and the average value of the Λ asymmetry parameter α_{avg} .

Source (10^{-3})	$\alpha_{J/\psi}$	$\Delta\Phi$	α_-	α_+	A_{CP}	α_{avg}
Background	0.4	0.2	0.3	0.4	0.4	0.0
Tracking	1.7	0.6	1.7	2.1	0.2	1.9
$\Lambda/\bar{\Lambda}$ vertex fit	0.2	0.0	0.1	0.3	0.3	0.1
Kinematic fit	1.4	3.0	0.8	1.4	0.3	0.4
Fit method	2.1	5.8	1.5	1.6	1.0	1.4
Total	3.1	6.6	2.4	3.0	1.2	2.4

CP asymmetry provides a hunting ground for physics beyond the standard model [39]. A clear transverse polarization is observed for the Λ and $\bar{\Lambda}$ as shown in Fig. 3. The phase between helicity flip and helicity conserving transitions is determined to be $\Delta\Phi = 0.7521 \pm 0.0042 \pm 0.0066$, where the first uncertainty is statistical and the second one is systematic. The large value of the phase makes it possible to simultaneously determine the decay parameters of $\Lambda \rightarrow p\pi^-$ and $\bar{\Lambda} \rightarrow \bar{p}\pi^+$ to be $\alpha_- = 0.7519 \pm 0.0036 \pm 0.0024$ and $\alpha_+ = 0.7559 \pm 0.0036 \pm 0.0030$, which represents the most precise measurements to date. Owing to the large correlation coefficient of the two decay parameters $\rho(\alpha_-, \alpha_+) = 0.850$, the Λ and $\bar{\Lambda}$ averaged value is determined to be $\alpha_{\text{avg}} = 0.7542 \pm 0.0010 \pm 0.0024$ for the first time, which are the most precise measurements in the baryon sector. Being the lightest baryon with strangeness, the measurements of polarizations, decay parameters, and CP asymmetries of heavier baryons, therefore, implicitly depend on α_Λ [20–24].

Results of the Λ decay parameter from different experiments are shown in Fig. 4. The α_- value obtained in this Letter agrees with the previous BESIII measurements [12] and the BESIII result extracted from the $J/\psi \rightarrow \Xi^-\bar{\Xi}^+$




FIG. 4. Results of the Λ decay parameter from different experiments. The green band represents the PDG 2018 value, and the pink band represents the PDG 2022 value.

decay [40], but deviates from the CLAS result by 3.5σ . In addition, we obtain the value of CP violation for the Λ decay $A_{CP} = (\alpha_- + \alpha_+)/(\alpha_- - \alpha_+) = -0.0025 \pm 0.0046 \pm 0.0012$, which is compatible with zero, thereby, indicating a non- CP -violation scenario. The next generation of charm factories [41,42] will greatly improve the accuracy of the CP -violating measurements, and shed light on the mechanism of CP violation in the baryon sector.

The BESIII collaboration thanks the staff of BEPCII and the IHEP computing center for their strong support. This work is supported in part by the National Key R&D Program of China under Contracts No. 2020YFA0406300 and No. 2020YFA0406400; National Natural Science Foundation of China (NSFC) under Contracts No. 11635010, No. 11735014, No. 11835012, No. 11935015, No. 11935016, No. 11935018, No. 11961141012, No. 12022510, No. 12025502, No. 12035009, No. 12035013, No. 12192260, No. 12192261, No. 12192262, No. 12192263, No. 12192264, No. 12192265; the Chinese Academy of Sciences (CAS) Large-Scale Scientific Facility Program; Joint Large-Scale Scientific Facility Funds of the NSFC and CAS under Contract No. U1832207; CAS Key Research Program of Frontier Sciences under Contract No. QYZDJ-SSW-SLH040; 100 Talents Program of CAS; INPAC and Shanghai Key Laboratory for Particle Physics and Cosmology; Polish National Science Centre under Contract No. 2019/35/O/ST2/02907; ERC under Contract No. 758462; European Union’s Horizon 2020 research and innovation programme under Marie Skłodowska-Curie grant agreement under Contract No. 894790; German Research Foundation DFG under Contracts No. 443159800, Collaborative Research Center CRC 1044, GRK 2149; Istituto Nazionale di Fisica Nucleare, Italy; Ministry of Development of Turkey under Contract No. DPT2006K-120470; National Science and Technology fund; STFC (United Kingdom); The Royal Society, UK under Contracts No. DH140054 and No. DH160214; The Swedish Research Council; U.S. Department of Energy under Contract No. DE-FG02-05ER41374.

^aAlso at the Moscow Institute of Physics and Technology, Moscow 141700, Russia.

^bAlso at the Novosibirsk State University, Novosibirsk 630090, Russia.

^cAlso at the NRC “Kurchatov Institute,” PNPI, 188300 Gatchina, Russia.

^dAlso at Goethe University Frankfurt, 60323 Frankfurt am Main, Germany.

^eAlso at Key Laboratory for Particle Physics, Astrophysics and Cosmology, Ministry of Education; Shanghai Key Laboratory for Particle Physics and Cosmology; Institute of Nuclear and Particle Physics, Shanghai 200240, People’s Republic of China.

- ^fAlso at Key Laboratory of Nuclear Physics and Ion-beam Application (MOE) and Institute of Modern Physics, Fudan University, Shanghai 200443, People's Republic of China.
- ^gAlso at State Key Laboratory of Nuclear Physics and Technology, Peking University, Beijing 100871, People's Republic of China.
- ^hAlso at School of Physics and Electronics, Hunan University, Changsha 410082, China.
- ⁱAlso at Guangdong Provincial Key Laboratory of Nuclear Science, Institute of Quantum Matter, South China Normal University, Guangzhou 510006, China.
- ^jAlso at Frontiers Science Center for Rare Isotopes, Lanzhou University, Lanzhou 730000, People's Republic of China.
- ^kAlso at Lanzhou Center for Theoretical Physics, Lanzhou University, Lanzhou 730000, People's Republic of China.
- ^lAlso at the Department of Mathematical Sciences, IBA, Karachi, Pakistan.
- [1] J. H. Christenson, J. W. Cronin, V. L. Fitch, and R. Turlay, *Phys. Rev. Lett.* **13**, 138 (1964).
- [2] B. Aubert *et al.* (BABAR Collaboration), *Phys. Rev. Lett.* **87**, 091801 (2001).
- [3] K. Abe *et al.* (Belle Collaboration), *Phys. Rev. Lett.* **87**, 091802 (2001).
- [4] R. Aaij *et al.* (LHCb Collaboration), *Phys. Rev. Lett.* **122**, 211803 (2019).
- [5] A. D. Sakharov, Pis'ma Zh. Eksp. Teor. Fiz. **5**, 32 (1967) [Sov. Phys. Uspekhi **34**, 392 (1991)].
- [6] W. Bernreuther, *Lect. Notes Phys.* **591**, 237 (2002).
- [7] L. Canetti, M. Drewes, and M. Shaposhnikov, *New J. Phys.* **14**, 095012 (2012).
- [8] I. I. Bigi and A. I. Sanda, *CP Violation* (Cambridge University Press, Cambridge, England, 2009), Vol. 9.
- [9] I. Bediaga and C. Göbel, *Prog. Part. Nucl. Phys.* **114**, 103808 (2020).
- [10] T. D. Lee and C.-N. Yang, *Phys. Rev.* **108**, 1645 (1957).
- [11] J. F. Donoghue and S. Pakvasa, *Phys. Rev. Lett.* **55**, 162 (1985); J. F. Donoghue, X.-G. He, and S. Pakvasa, *Phys. Rev. D* **34**, 833 (1986).
- [12] M. Ablikim *et al.* (BESIII Collaboration), *Nat. Phys.* **15**, 631 (2019).
- [13] D. G. Ireland, M. Döring, D. I. Glazier, J. Haidenbauer, M. Mai, R. Murray-Smith, and D. Rönchen, *Phys. Rev. Lett.* **123**, 182301 (2019).
- [14] J. Adam *et al.* (STAR Collaboration), *Phys. Rev. Lett.* **123**, 132301 (2019).
- [15] F. Becattini and M. A. Lisa, *Annu. Rev. Nucl. Part. Sci.* **70**, 395 (2020).
- [16] M. Ablikim *et al.* (BESIII Collaboration), *Phys. Rev. Lett.* **123**, 122003 (2019).
- [17] M. S. Abdallah *et al.* (STAR Collaboration), *Phys. Rev. C* **104**, L061901 (2021).
- [18] T. Blake, S. Meinel, and D. van Dyk, *Phys. Rev. D* **101**, 035023 (2020).
- [19] T.-Z. Han, R.-G. Ping, T. Luo, and G.-Z. Xu, *Chin. Phys. C* **44**, 013002 (2020).
- [20] M. Ablikim *et al.* (BESIII Collaboration), *Phys. Rev. D* **100**, 072004 (2019).
- [21] J. M. Link *et al.* (FOCUS Collaboration), *Phys. Lett. B* **634**, 165 (2006).
- [22] R. Aaij *et al.* (LHCb Collaboration), *J. High Energy Phys.* **06** (2020) 110.
- [23] D. Wang, R.-G. Ping, L. Li, X.-R. Lyu, and Y.-H. Zheng, *Chin. Phys. C* **41**, 023106 (2017).
- [24] M. Ablikim *et al.* (BESIII Collaboration), *Phys. Rev. D* **103**, L091101 (2021).
- [25] H.-B. Li, *Front. Phys.* **12**, 121301 (2017); **14**, 64001(E) (2019).
- [26] A. Z. Dubnickova, S. Dubnicka, and M. P. Rekalo, *Nuovo Cimento A* **109**, 241 (1996); G. I. Gakh and E. Tomasi-Gustafsson, *Nucl. Phys.* **A771**, 169 (2006); H. Czyz, A. Grzelinska, and J. H. Kuhn, *Phys. Rev. D* **75**, 074026 (2007); G. Fäldt, *Eur. Phys. J. A* **51**, 74 (2015); **52**, 141 (2016); G. Fäldt and A. Kupsc, *Phys. Lett. B* **772**, 16 (2017).
- [27] R. L. Workman *et al.* (Particle Data Group), Review of Particle Physics, *Prog. Theor. Exp. Phys.* **2022**, 083C01 (2022).
- [28] M. Ablikim *et al.* (BESIII Collaboration), *Chin. Phys. C* **46**, 074001 (2022).
- [29] M. Ablikim *et al.* (BESIII Collaboration), *Nucl. Instrum. Methods Phys. Res., Sect. A* **614**, 345 (2010).
- [30] M. Ablikim *et al.* (BESIII Collaboration), *Chin. Phys. C* **44**, 040001 (2020).
- [31] S. Agostinelli *et al.*, *Nucl. Instrum. Methods Phys. Res., Sect. A* **506**, 250 (2003).
- [32] Z.-Y. Deng *et al.*, *Chin. Phys. C* **30**, 371 (2006).
- [33] S. Jadach, B. F. L. Ward, and Z. Was, *Phys. Rev. D* **63**, 113009 (2001).
- [34] R.-G. Ping, *Chin. Phys. C* **32**, 243 (2008).
- [35] J. C. Chen, G. S. Huang, X. R. Qi, D. H. Zhang, and Y. S. Zhu, *Phys. Rev. D* **62**, 034003 (2000).
- [36] R.-L. Yang, R.-G. Ping, and H. Chen, *Chin. Phys. Lett.* **31**, 061301 (2014).
- [37] G.-R. Liao, R.-G. Ping, and Y.-X. Yang, *Chin. Phys. Lett.* **26**, 051101 (2009).
- [38] W. Verkerke and D. P. Kirkby, eConf **C0303241**, MOLT007 (2003) [[arXiv:physics/0306116](https://arxiv.org/abs/physics/0306116)].
- [39] J. Tandean, *Phys. Rev. D* **69**, 076008 (2004).
- [40] M. Ablikim *et al.* (BESIII Collaboration), *Nature (London)* **606**, 64 (2022).
- [41] A. E. Bondar *et al.* (Charm-Tau Factory Collaboration), *Phys. At. Nucl.* **76**, 1072 (2013).
- [42] Q. Luo and D. R. Xu, in *Proc. 9th International Particle Accelerator Conference (IPAC'18)* (JACoW Publishing, Geneva, 2018), [10.18429/JACoW-IPAC2018-MOPML013](https://doi.org/10.18429/JACoW-IPAC2018-MOPML013).