

Paederus, sensu lato (Coleoptera: Staphylinidae): An Index and Review of the Taxa

J.H. Frank
Entomology and Nematology Department
3103 McCarty Hall
University of Florida
Gainesville, FL 32811 USA

Contents

Abstract	97
Introduction	98
Materials and methods	98
Results and discussion	99
Index to the names of <i>Paederus</i> (<i>sensu lato</i>)	99
Synonymy in genus-group names	137
<i>Geopaederus</i> and <i>Poederomorphus</i>	137
<i>Paederillus</i> and <i>Leucopaederus</i>	137
<i>Paederognathus</i> and <i>Neopaederus</i>	137
Standing names of genera distinguished	
from <i>Paederus</i>	138
<i>Paederidus</i>	138
<i>Allopaederus</i> , <i>Madecapaederus</i> , <i>Oreopaederus</i> , and-	
<i>Paederognathus</i>	138
Generic key to Afrotropical/Malagasy adult	
<i>Paederus sensu lato</i>	138
<i>Ctenopaederus</i>	138
<i>Diplopaederus</i> , <i>Eupaederus</i> , <i>Lobopaederus</i> , <i>Mega-</i>	
<i>lopaederus</i> , <i>Oncopaederus</i> , <i>Parameropaederus</i> , and	
<i>Uncopaederus</i>	139
Generic concepts	139
Relatives of <i>Paederus</i> (<i>sensu lato</i>)	140
Standing names of subgenera distinguished	
from <i>Paederus</i>	140
<i>Pseudopaederus</i> and <i>Gnathopaederus</i>	140
<i>Dioncopaederus</i> , <i>Eopaederus</i> , <i>Harpopaederus</i> , <i>Hetero-</i>	
<i>paederus</i> , and <i>Oedopaederus</i>	140
Subgeneric concepts	140
Homonymy in species-group names	142
Conclusion	144
Acknowledgments	144
Literature cited	145

Abstract

Generic, subgeneric, specific, subspecific, and infra-subspecific names in *Paederus* (*sensu lato*) are compiled alphabetically with literature references, showing that 622 specific names stand in the literature as valid. Five replacement names are required due to homonymy: *Paederus cumanus* Frank [*nom. nov.* for *Paederus bicolor* Wendeler *nec* Olivier], *Paederus sulawesi* Frank [*nom. nov.* for *Paederus melanocephalus* Heller *nec* Fabricius], *Paederus zairensis* Frank [*nom. nov.* for *Paederus orophilus* Fagel *nec* *Paederidus brunnescens orophilus* (Fagel)], *Paederus irianensis* Frank [*nom. nov.* for *Paederus litoreus* Last *nec* *Paederus littoreus* Austin], and *Oreopaederus manyemensis* Frank [*nom. nov.* for *Oreopaederus ater* (Bernhauer) *nec* *Paederidus rubrothoracicus ater* (Eichler)]. Four unjustified replacement names (*Paederus archeus* Blackwelder, *Paederus homonymus* Blackwelder, *Paederus erichsoni* Wollaston, and *Paederus samoensis* Fauvel) are here attributed to their respective senior synonyms (*Paederus elongatus* Wendeler, *Paederus tricolor* Erichson, *Paederus angolensis* Erichson, and *Paederus vitiensis* Fauvel). *Paederus limnophilus* Erichson is a junior synonym of *Paederus limophilus* Heer. Of the 622 names, one is that of a fossil species and 264 are of species which have not yet been attributed to taxa more precise than *Paederus* (*sensu lato*). Concepts of the Afrotropical/Malagasy species are due to Fagel (1958a-1973), and among the 281 species recognized, 157 are attributed to *Paederus* Fabricius (*sensu* Fagel) and the remainder to the related genera *Paederidus* Mulsant & Rey (36), *Allopaederus* Fagel

(3), *Madecapaederus* Fagel (11), *Pachypaederus* Fagel (19), and *Oreopaederus* Fagel (55). Concepts of classification of species from other parts of the world are due mainly to Scheerpeltz (1957); 14 are attributed to the Asian genera *Ctenopaederus* Fagel (1), *Eupaederus* Scheerpeltz (2), *Diplopaederus* Scheerpeltz (2), *Oncopaederus* Scheerpeltz (1) and *Megalopaederus* Scheerpeltz (8), while 46 species of the western Palearctic region are attributed to *Paederus* (*sensu* Scheerpeltz and Coiffait) (33) and related genera *Paederidus* (10), *Lobopaederus* Scheerpeltz (1), *Parameropaederus* Scheerpeltz (1), and *Uncopaederus* Korge (1). Eleven names stand as subgenera of *Paederus* (*Pseudopaederus* Bernhauer, *Gnathopaederus* Chapin, *Eopaederus* Scheerpeltz, *Harpopaederus* Scheerpeltz, *Heteropaederus* Scheerpeltz, *Dioncopaederus* Scheerpeltz, *Oedopaederus* Scheerpeltz, *Anomalopaederus* Scheerpeltz, *Oreinopaederus* Scheerpeltz, and *Nepalopaederus* Scheerpeltz), but none of these has many species attributed to it and only the western Palearctic species have been assigned systematically to subgenera. *Paederus* (*sensu lato*) is the subtribe Paederi (Casey 1905, Fagel 1958a) or Paederina (Herman 1981), with the subtribe Dolicaei (Fagel 1958a) or Dolicaonina (Herman 1981) as closest relative. The 2 major taxonomic needs are to review and redescribe the species of the Nearctic, Neotropical, Oriental, Oceanian and Australasian regions, and to extend Fagel's (1958a) methods to examine the higher classification of *Paederus* (*sensu lato*) worldwide.

Introduction

Frank & Kanamitsu (1987) reviewed the natural history and medical importance of *Paederus* (*sensu lato*), and listed 47 species about which some biological information had been made known. One of the reviewers of the manuscript (L.E. Munstermann) suggested expansion of the list to include all species of *Paederus* (*sensu lato*). I declined that suggestion on 3 grounds. First, I believed the group to have over 500 species, based in part on an estimate by Coiffait (1982c) of 450 species, and I thought a list of over 500 names was too long and inappropriate to include in the review. Second, I had not seen an estimated third of the taxonomic publications on the group and therefore I did not know all the names and was unsure of the status of several. Third, a review of the names of *Paederus* (*sensu lato*) was necessary as a project in its own right, as a background to taxonomic studies, and should be approached with that objective in mind.

I assembled the taxonomic literature on *Paederus* (*sensu lato*) and analyzed it as a background to future taxonomic studies. This paper is a result of the analysis. It is intended to provide answers to the following ques-

tions: 1) What names have been applied to *Paederus* (*sensu lato*) at the generic, subgeneric, specific, subspecific and infra-subspecific levels? 2) What are the taxonomic publications using these names? 3) Which of the names appear valid, and which are known to be synonyms, homonyms, nomina nuda, misspellings, and unpublished names? 4) How are the species bearing the valid names currently classified? 5) What is the basis of the classification? 6) What remains to be done with the classification of *Paederus* (*sensu lato*)? 7) What groups of Paederinae are closely related to *Paederus* (*sensu lato*)?

Materials and Methods

The taxonomic literature on *Paederus* (*sensu lato*) was examined and an alphabetical list of names was compiled, as follows.

All generic, subgeneric, specific, subspecific and infra-subspecific names are indexed. Disposition of these names as valid or not was determined by examination of the literature, with acceptance of the latest disposition over all earlier ones.

Generic and subgeneric names, indicated by a capitalized initial letter, are italicized when valid, but not italicized if invalid due to synonymy or homonymy. The author of each such name is given. Below and in brackets following the author's name are given the date and page of publication, original rank (genus or subgenus), and the name of the type species. This is followed (where applicable) by author, date and page of a later work assigning the name to synonymy or homonymy or changed status, followed (where applicable) by the senior synonym or replacement name.

Specific epithets are italicized when valid, but not italicized when they are (a) synonyms, homonyms, and nomina nuda or (b) unpublished or misidentifications, or (c) misspellings or *lapsi calami*. Those in category (a) are followed by the name of the author, those in (b) are followed by the abbreviation *auct.* and those in (c) by the word *sic*. For all valid names, there follow in chronological order the generic assignments with the latest (current) one italicized; subgeneric assignments follow, are treated in the same way as the generic assignments, but are placed in parentheses. Below and in brackets are given relevant citations in the taxonomic literature, beginning with the original description and including all citations which change the status, followed (in the case of junior synonyms or homonyms) by the senior synonym or replacement name.

Subspecific epithets are not italicized when indexed, but instead are referred to the correct specific epithet. However, subspecific epithets are also listed as binomina (with specific epithets) where applicable.

Infra-subspecific names are not italicized when indexed, but their current disposition is listed. Specific epithets that were at some time included in *Paederus* (*sensu lato*) (but were subsequently excluded) are indexed. The exclusion and current disposition (if different) are noted.

Results and Discussion

Index to the names of *Paederus* (*sensu lato*)

A

- abdominalis* Cameron, *Paederus*
[Cameron 1950c: 186, Fagel 1958a: 291, 1966: 261]
- abessynus* auct.
[Bernhauer *in litt.*, Fagel 1958a: 353, = *ignotus* Wendeler]
- abnormalis* Bernhauer, *Paederus*
[Bernhauer 1934a: 18]
- abyssinicus* Cameron
[Cameron 1950c: 186, Fagel 1958a: 311, = *fuscipes* Curtis, 1959: 62, 1960a: 121, 1961b: 268]
- adelaidae* Blackburn, *Paederus*
[Blackburn 1888: 10, 1900: 22, Bernhauer & Schubert 1912: 203, Lea 1923: 14]
- adumbratus* Wickham, *Paederus*
[Wickham 1913: 10, fossil]
- aequabilis* Last, *Paederus*
[Last 1979: 134]
- aequinoctialis* Dejean
[Dejean 1833: 65, *nomen nudum*]
- aequinoctialis* Erichson
[Erichson 1840: 657, = *columbinus* Laporte, Gemminger & Harold 1868: 626, Sharp 1876a: 285, 1886: 617, Bernhauer & Schubert 1912: 205, Blackwelder 1944: 122, Scheerpeltz 1965: 217]
- aereus* Fagel, *Allopaederus*
[Fagel 1968: 198]
- aestmans* sic
[Bernhauer 1915g: 362, misspelling of *aestuans*]
- aestuans* Erichson
[Erichson 1840: 655, Boheman 1848: 291 *partim* (*vide* Fagel 1958a: 275), Gemminger & Harold 1868: 626, Fauvel 1873: 334, = *fuscipes* Curtis, 1886a: 32, 1903: 154, 1904b: 290, 1907: 24, Czwalina 1889: 368, Ganglbauer 1895: 537, Bernhauer & Schubert 1912: 206, Cameron 1925b: 35, 1931a: 40, Fagel 1958a: 311, 1959: 62, 1960a: 121, 1961b: 268, Legosz-Owsianna 1963: 335, Scheerpeltz 1968: 47]
- aestuans* auct.
[Boheman 1848: 291 *partim*, misidentification, Fagel 1960a: 119, = *nakurensis* Fauvel]
- aestuans* auct.
[Boheman 1848: 291 *partim*, misidentification, Fagel 1958a: 275, = *bohemani* Fagel]
- aestuans* auct.
[Boheman 1848: 291, attribution to *sabaeus* Erichson (e.g., by Fauvel 1904b: 290, Bernhauer & Schubert 1912: 210, Coiffait 1982c: 61) is erroneous (*vide* Fagel 1958a: 275)]
- agnatus* Eppelsheim, *Paederus*
[Eppelsheim 1889b: 180, Bernhauer & Schubert 1912: 203, Wu 1937: 328, Scheerpeltz 1957a: 474]
- alberti* Fagel
[Fagel 1958a: 115, ssp. of *colonus* Wendeler, 1959: 38]
- albertisi* Fauvel
[Fauvel 1878a: 236, *nom. nov.* for *chilensis* Fauvel, 1879: 87, Bernhauer & Schubert 1912: 203, Bernhauer 1927c: 399, = *intermedius* Boheman]
- albipilis* (Solsky), *Paederus*, *Paederidus*
[Solsky 1871: 163, Fauvel 1873: 336, = *ruficollis* Fabricius, Eppelsheim 1892: 340, species, Bernhauer & Schubert 1912: 203, Kirschenblatt 1932: 216, Scheerpeltz 1933: 1220, 1963: 8, Tikhomirova 1973: 175, Tronquet 1981: 70, Coiffait 1982c: 27]
- albobubescens* (Bernhauer), *Paederus*, *Paederidus*
[Bernhauer 1927a: 117, Scheerpeltz 1933: 1220, Koch 1937: 25, Fagel 1958a: 120, 1959: 41]
- alfieri* sic
[Coiffait 1982c: 63, misspelling of *alfierii*]
- alfierii* Koch, *Paederus* (*Heteropaederus*)
[Koch 1934: 80, Scheerpeltz 1957a: 461, Fagel 1958a: 311, = *fuscipes* Curtis, 1959: 62, 1960a: 121, 1961b: 268, Coiffait 1982c: 63, species]
- algiricus* (Motschulsky), *Paederus*, *Paederidus*
[Motschulsky 1858: 635, Gemminger & Harold 1868: 626, Fauvel 1869b: 491, var. of *sanguinicollis* Stephens, 1886a: 32, = *ruficollis* Fabricius, Bernhauer & Schubert 1912: 210, = *rubrothoracicus* Goeze, Scheerpeltz 1957a: 451, ssp. of *ruficollis* Fabricius, Fagel 1958b: 234, Legosz-Owsianna 1963: 327, = *rubrothoracicus* Goeze, Coiffait 1982c: 24, species]
- algiricus antoinei* (Koch)
[Koch 1937: 24, species, Scheerpeltz 1957a: 450, ssp. of *ruficollis* Fabricius, Fagel 1958b: 234, = *algiricus* Motschulsky, Coiffait 1982c: 24, ssp. of *algiricus*]
- alliceptus* Cameron, *Paederus*
[Cameron 1914: 538, 1925b: 34, 1931a: 61]

- aliipennis* Cameron
[Cameron 1950c: 186, Fagel 1958a: 179, = *nakurensis* Fauvel, 1960a: 120]
- alinderi* (Wendeler), *Paederus*, *Oreopaederus*
[Wendeler 1933: 55, Fagel 1954a: 961, 1956: 188, 1958a: 102, 1961a: 265]
- allardianus* Fagel, *Oreopaederus*
[Fagel 1964: 225]
- Allopaederus* Fagel
[Fagel 1959: 104, *genus*, type *Allopaederus mirandus* Fagel]
- alluaudi* Fagel
[Fagel 1958a: 421, *ssp.* of *melanogaster* Fauvel]
- almorensis* Cameron, *Paederus*
[Cameron 1931a: 49, Scheerpeltz 1933: 1220]
- alternans* Walker, *Paederus*
[Walker 1858: 205, Gemminger & Harold 1868: 626, Fauvel 1904a: 54, Bernhauer & Schubert 1912: 203, Cameron 1925b: 34, 1930b: 332, 1931a: 41, Scheerpeltz 1933: 1220, 1935: 607, 1965b: 103, 1978: 196, Biswas & Sen Gupta 1982: 150]
- alticola* Sharp, *Paederus*
[Sharp 1886: 609, Fauvel 1908: 95, Bernhauer & Schubert 1912: 203, Blackwelder 1944: 122]
- alticola* Fauvel
[Fauvel 1907: 20, homonym, 1908: 95, = *altivagans* Fauvel, Bernhauer 1912c: 471, Bernhauer & Schubert 1912: 203, Fagel 1958a: 339, 1960a: 122]
- altivagans* Fauvel, *Paederus*
[Fauvel 1908: 95, replacement name for *alticola* Fauvel, Bernhauer & Schubert 1912: 203, Bernhauer 1912c: 471, 1940a: 135, Fagel 1958a: 339, 1960a: 122, 1966: 269]
- altivagans auct.*
[Fauvel 1908: 95, *partim*, Fagel 1960a: 122, = *kili-mandjarensis* Fagel]
- alutaceus* Fagel, *Paederus*
[Fagel 1958a: 167, 1959: 47, 1966: 272]
- alutipennis* Cameron, *Paederus*
[Cameron 1936: 43]
- alutithorax* (Bernhauer), *Paederus*, *Megalopaederus*
[Bernhauer 1943: 176, Shibata 1973: 44]
- alutiventris* Bernhauer
[Bernhauer 1931c: 572, Scheerpeltz 1933: 1220, Fagel 1958a: 351, = *duplex spectabilis* Bernhauer]
- amazonicus* Sharp
[Sharp 1876a: 287, Bernhauer & Schubert 1912: 203, Blackwelder 1944: 122]
- amicus* Bernhauer, *Paederus*
[Bernhauer 1915c: 292, 1942b: 357, Scheerpeltz 1933: 1220, Fagel 1958a: 225, 1959: 47, 1960b: 203, 1966: 260]
- amoenus sic*
[Cameron 1914: 540, *lapsus calami, nec* Erichson]
- amplicolis* Kraatz, *Paederus*
[Kraatz 1859a: 150, Gemminger & Harold 1868: 626, Bernhauer & Schubert 1912: 203, Cameron 1925b: 34, 1927: 115, 1931a: 58, Scheerpeltz 1933: 1220]
- amplicolis nigrescens* Cameron
[Cameron 1927: 115, *var.* of *amplicolis*, 1931a: 58, Scheerpeltz 1933: 1220, here treated as *ssp.* under Article 45(g) (ICZN 1985)]
- amplipennis* Fauvel
[Fauvel 1907: 23, Fagel 1958a: 208, = *eximius* Reiche]
- analis* Dahl
[Dahl 1823: 17, Dejean 1833: 65, = *Rugilus*]
- andinus auct.*
[Fauvel *in litt.*, Wendeler 1931: 44, = *andinus* Wendeler]
- andinus* Wendeler, *Paederus*
[Wendeler 1931: 44, Scheerpeltz 1933: 1220, Blackwelder 1944: 122]
- andrewesi* (Fauvel), *Paederus*, *Ctenopaederus*
[Fauvel 1904a: 53, Bernhauer & Schubert 1912: 203, Cameron 1925b: 35, 1931a: 59, Scheerpeltz 1933: 1220, Fagel 1958a: 69]
- angolanus* Fagel
[Fagel 1961a: 265, *nomen nudum*]
- angolanus* Fagel, *Oreopaederus*
[Fagel 1962b: 20]
- angolensis* Erichson
[Erichson 1843: 222, Wollaston 1867: 248, Gemminger & Harold 1868: 626, Fauvel 1903a: 154, = *fuscipes* Curtis, 1904b: 290, Bernhauer & Schubert 1912: 206, Cameron 1925b: 35, 1931a: 40, Scheerpeltz 1957a: 461, Fagel 1958a: 311, 1959: 62, 1960a: 121, 1961b: 268, Legosz-Owsianna 1963: 335]
- angolensis* Bernhauer
[Bernhauer 1936b: 262, homonym, Fagel 1958a: 279, = *plagiator* Kolbe]
- anguinus* Bernhauer, *Paederus*
[Bernhauer 1908a: 309, Bernhauer & Schubert 1912: 203, Blackwelder 1944: 122]
- angulatus sic*
[Lea 1904: 63, misspelling of *cingulatus*]
- angulicollis* MacLeay, *Paederus*
[MacLeay 1871: 146, Fauvel 1877a: 224, Blackburn 1899: 23, MacLeay 1904: 63, Bernhauer & Schubert 1912: 203, Lea 1923: 14]
- angustatus* (Paykull)

- [Paykull 1789: 36, 1800: 431, Fabricius 1792: 528, Panzer 1793b: 18, Fabricius 1801: 599, Gravenhorst 1802: 63, 1806: 141, Latreille 1806: 292, Gyllenhal 1810: 375, Olivier 1812: 630, Beck 1817: 45, Stephens 1829: 287, = *Sunius*, now in *Astenus*]
- angusticeps* Bernhauer, *Paederus*
[Bernhauer 1915g: 365, Scheerpeltz 1933: 1220, Fagel 1958a: 319, 1966: 267]
- angustipennis* Bernhauer, *Paederus* (*Paederus*)
[Bernhauer 1936a: 240, Cameron 1949b: 466, Scheerpeltz 1957a: 462, Shibata 1977: 26]
- annexus* Eppelsheim, *Paederus*
[Eppelsheim 1895a: 133, Bernhauer & Schubert 1912: 203, Fagel 1958a: 227, 1966: 259]
- Anomalopaederus* Scheerpeltz
[Scheerpeltz 1966: 400, *subgenus*, type *Paederus la-cordairei* Perroud]
- anthracinus* Bernhauer, *Paederus*
[Bernhauer 1927b: 157, Scheerpeltz 1933: 1220, Blackwelder 1944: 122]
- antipodum* Bernhauer & Schubert
[Bernhauer & Schubert 1912: 203, replacement name for *erichsoni* Bernhauer, Lea 1923: 15, = *mey-ricki* Blackburn, Scheerpeltz 1933: 1220]
- antiquus* Sharp, *Paederus*
[Sharp 1886: 615, Bernhauer & Schubert 1912: 203, Blackwelder 1944: 122]
- antoinei* Koch
[Koch 1937: 24, 1941: 56, Peyerimhoff 1949: 104, = *algericus* Motschulsky, Scheerpeltz 1957a: 450, ssp. of *ruficollis* Fabricius, Fagel 1958b: 234, = *algericus* Motschulsky, Coiffait 1982c: 24, ssp. of *algericus* Motschulsky]
- antoniensis* Sharp, *Paederus*
[Sharp 1886: 615, Bernhauer & Schubert 1912: 204, Blackwelder 1944: 122]
- apicalis* Sharp
[Sharp 1886: 617, Bernhauer & Schubert 1912: 204, Blackwelder 1939: 111, 1944: 123, = *ustus* LeConte]
- apteromelas* Lea, *Paederus*
[Lea 1923: 15, Scheerpeltz 1933: 1220]
- apterus* (Fauvel), *Paederus*, *Madecapaederus*
[Fauvel 1898: 180, Bernhauer & Schubert 1912: 204, Fagel 1958a: 457]
- aquaticus* Bernhauer, *Paederus*
[Bernhauer 1915a: 116, Scheerpeltz 1933: 1220, Fagel 1958a: 284, 1966: 258]
- aquatilis* Cameron, *Paederus*
[Cameron 1944: 710, Fagel 1958a: 281]
- aquatilis auct.*
[Bernhauer *in litt.*, Fagel 1958a: 281, = *aquatilis* Cameron]
- aquatilis marraensis* Fagel
[Fagel 1968: 197]
- arabicus* (Bernhauer), *Paederus*, *Paederidus*
[Bernhauer 1936a: 241, Fagel 1958a: 152, Coiffait 1982c: 28]
- archeus* Blackwelder
[Blackwelder 1944: 122, replacement name for *elongatus* Wendeler, unnecessary, junior synonym of *elongatus* Wendeler]
- arduus* Sharp, *Paederus*
[Sharp 1886: 615, Bernhauer & Schubert 1912: 204, Blackwelder 1944: 122]
- argentatus auct.*
[Fauvel *in litt.*, Cameron 1931a: 45, = *argentatus* Cameron]
- argentatus* Cameron, *Paederus*
[Cameron 1914: 540, 1925b: 35, 1931a: 45, Scheerpeltz 1933: 1220]
- argentinus* Bernhauer, *Paederus*
[Bernhauer 1911b: 407, Bernhauer & Schubert 1912: 204, Blackwelder 1944: 122]
- armstrongi* Steel, *Paederus*
[Steel 1955: 181]
- arrowi* (Bernhauer), *Paederus*, *Paederidus*
[Bernhauer 1927a: 116, Scheerpeltz 1933: 1220, Fagel 1958a: 142]
- arrowianus* Bernhauer, *Paederus*
[Bernhauer 1931c: 573, Fagel 1958a: 185, 1966: 270]
- asiaeminoris* Scheerpeltz
[Scheerpeltz 1957a: 456, ssp. of *rubrothoracicus* Goeze, Coiffait 1982c: 25, var. of *rubrothoracicus* Goeze]
- asperatus* Last
[Last 1950: 136, Fagel 1958a: 146, = *collaris* Boheman, 1961b: 267]
- ater* Eichler
[Eichler 1924: 65, var. of *ruficollis* Fabricius, Scheerpeltz 1933: 1227, var. of *rubrothoracicus* Goeze, Coiffait 1982c: 27, here treated as ssp. under Article 45(g) (ICZN 1985)]
- ater* Bernhauer
[Bernhauer 1927a: 107, homonym, Scheerpeltz 1933: 1220, Fagel 1954a: 913, Fagel 1958a: 101, here = *Oreopaederus manyemensis* Frank, *nom. nov.*]
- ater luvubuensis* Fagel
[Fagel 1954a: 914, 1958a: 101, here = *Oreopaederus manyemensis luvubuensis* (Fagel), new combination]

ater auct.

[Cameron 1950b: 25, misidentification, Fagel 1954a: 965, = *mikenensis* Fagel]

atratus Scheerpeltz

[Scheerpeltz 1957a: 453, Coiffait 1982c: 25, = *rubrothoracicus* var. *clermonti* Chapman]

atricolor (Fagel), *Paederus*, *Oreopaederus*

[Fagel 1954a: 909, 1958a: 101, 1961a: 265]

atriventris Cameron, *Paederus*

[Cameron 1930: 332, Scheerpeltz 1933: 1220, 1935: 609, 1957b: 265]

atrocyaneus Champion, *Paederus*

[Champion 1927: 50, Cameron 1931a: 36, Scheerpeltz 1933: 1220]

atropygus Fagel, *Oreopaederus*

[Fagel 1961a: 273]

aureipennis Bernhauer, *Paederus*

[Bernhauer 1942b: 356, Fagel 1958a: 194, 1966: 263]

australis Guerin, *Paederus*

[Guerin 1830: 63, Laporte 1840: 183, Erichson 1840: 666, Perroud 1865: 89, Gemminger & Harold 1868: 626, Fauvel 1877a: 223, 1878b: 516, Blackburn 1894: 92, 1900: 22, Lea 1904: 64, 1923: 14, Bernhauer & Schubert 1912: 204, Scheerpeltz 1933: 1220]

B*badius auct.*

[Knoch *in coll.*, Gravenhorst 1802: 53, = *Lathrobium*]

baeri Bernhauer, *Paederus*

[Bernhauer 1921: 72, Scheerpeltz 1933: 1220, Blackwelder 1944: 122]

bakeri Cameron, *Paederus*

[Cameron 1942a: 88]

bakerianus Bernhauer, *Paederus*

[Bernhauer 1927c: 401, Scheerpeltz 1933: 1220]

balachowskyi Jarrige, *Paederidus*, *Paederus* (*Eopaederus*)

[Jarrige 1971: 483, Coiffait 1982c: 49]

balcanicus Koch, *Paederus* (*Paederus*)

[Koch 1938: 103, ssp. of *riparius* Linnaeus, Szujecki 1965a: 83, species, 1965b: 28, Tikhomirova 1973: 175, Coiffait 1982c: 56, Bohac 1985: 370]

balcanicus Scheerpeltz

[Scheerpeltz 1957a: 455, ssp. of *rubrothoracicus* Goeze, Coiffait 1982c: 25, var. of *rubrothoracicus* Goeze]

balfourbrownei Fagel, *Paederidus*

[Fagel 1958a: 139, 1973: 154]

balfourbrownei Fagel, *Paederus*

[Fagel 1958a: 303, 1966: 260]

banghaasi Bernhauer

[Bernhauer 1927c: 400, Scheerpeltz 1933: 1221, Fagel 1958a: 179]

banghaasi Bernhauer

[Bernhauer 1927a: 111, homonym, Scheerpeltz 1933: 1221, = *banghaasianus* Scheerpeltz, replacement name]

banghaasianus Scheerpeltz

[Scheerpeltz 1933: 1221, replacement name for *banghaasi* Bernhauer 1927a, Fagel 1958a: 179, = *nakurensis* Fauvel, 1960a: 120]

basalis Bernhauer, *Paederus* (*Eopaederus*)

[Bernhauer 1914: 98, Cameron 1925b: 35, 1931a: 54, Scheerpeltz 1933: 1221, Blackwelder 1939: 111, Scheerpeltz 1957a: 474, 1963: 8]

basicornis auct.

[Fauvel *in litt.*, Wendeler 1931: 42, = *basicornis* Wendeler, Fagel 1958a: 121, = *schoutedeni* Bernhauer]

basicornis Wendeler

[Wendeler 1931: 42, Scheerpeltz 1933: 1221, Fagel 1958a: 121, = *schoutedeni* Bernhauer]

basilewskyi (Fagel), *Paederus*, *Oreopaederus*

[Fagel 1954a: 922, 1958a: 101, Gaedike 1981: 179]

basipes Fauvel, *Paederus*

[Fauvel 1907: 21, Bernhauer & Schubert 1912: 204, Fagel 1958a: 427, 1960a: 127, 1966: 268]

basipes basipes Fauvel

[Fauvel 1907: 21, Fagel 1960a: 127]

basipes hanangensis Fagel

[Fagel 1960a: 128, 1966: 269]

basipes ngorongoroensis Fagel

[Fagel 1960a: 129, 1966: 269]

basipes oldeaniensis Fagel

[Fagel 1960a: 128, 1966: 266]

basiventris Bernhauer, *Paederus*

[Bernhauer 1933a: 28, Scheerpeltz 1957a: 474, Tikhomirova 1973: 175]

baudii Fairmaire, *Paederus* (*Harpopaederus*)

[Fairmaire 1859: cxxxiv, replacement name for *lusitanicus auct.*, Gemminger & Harold 1868: 626, Fauvel 1866: 24, 1873: 329, Mulsant & Rey 1878: 250, Reitter 1884: 43, 1885: 154, 1889: 170, Czwalina 1889: 368, Ganglbauer 1895: 538, Bernhauer & Schubert 1912: 204, Cameron 1924: 180, Kirschenblatt 1932: 217, Scheerpeltz 1933: 1221, Aleksandrov 1934: 151, Koch 1938: 104, Blackwelder 1939: 112, Jarrige 1944: 9, Scheerpeltz 1957a: 460, Smetana 1962: 49, Szujecki 1965b: 30, Korge 1969: 60, Tikhomirova 1973: 175, Coiffait 1982c: 64]

- baudii* auct.
[Fauvel 1873: 329 *partim*, Smetana 1962: 49, = *schoenherri* Czwalina, Legosz-Owsianna 1963: 342, Scheerpeltz 1968: 47, Tikhomirova 1973: 175, Bohác 1985: 375]
- bayeri* Bernhauer, *Paederus*
[Bernhauer 1927a: 106, Scheerpeltz 1933: 1221, Fagel 1958a: 445, 1966: 265]
- beccarii* auct.
[Fauvel *in litt.*, Wendeler 1926b: 199, = *beccarii* Wendeler]
- beccarii* Wendeler, *Paederus*
[Wendeler 1926b: 199, Scheerpeltz 1933: 1221, 1935: 626, Cameron 1936: 42]
- benicki* Wendeler, *Paederus*
[Wendeler 1933: 56, Fagel 1958a: 447, 1966: 265]
- bernhaueri* (Fagel), *Paederus*, *Oreopaederus*
[Fagel 1954a: 964, 1958a: 102]
- bernhauerianus* Scheerpeltz, *Paederus*
[Scheerpeltz 1935: 624]
- bernieri* Laporte
[Laporte 1840: 183, Gemminger & Harold 1868: 626, Bernhauer & Schubert 1912: 204, Fagel 1958a: 457, = *coerulescens* Erichson]
- besucheti* Biswas & Sen Gupta, *Paederus*
[Biswas & Sen Gupta 1982: 146]
- bhutanicus* Coiffait, *Paederus*
[Coiffait 1978a: 116]
- bicolor* Olivier
[Olivier 1795: 7, Gravenhorst 1802: 59, Latreille 1804: 547, 1806: 290, = *Lathrobium*, now in *Sunius*]
- bicolor* Wendeler
[Wendeler 1942: 103, homonym, Gaedike 1981: 179, here = *Paederus cumanus* Frank, *nom. nov.*]
- bicoloriceps* Fagel, *Paederus*
[Fagel 1958a: 328, 1966: 268]
- bicoloripes* Cameron, *Paederus*
[Cameron 1937: 96]
- biguttatus* (Linnaeus)
[Linnaeus 1758: 422, Olivier 1795: 5, Latreille 1804: 352, = *Stenus*]
- binotatus* Say
[Say 1823: 154, Erichson 1840: 645, = *Sunius*, now in *Astenus*]
- birmanus* Fauvel, *Paederus*
[Fauvel 1895: 233, Bernhauer & Scheerpeltz 1912: 204, Cameron 1925b: 35, 1931a: 46, Scheerpeltz 1933: 1221, 1965b: 103, Coiffait 1976: 244, 1982b: 234, Gaedike 1981: 180]
- blukwensis* Fagel, *Paederus*
[Fagel 1958a: 402, 1966: 267]
- bohemani* Fagel, *Paederus* (*Heteropaederus*)
[Fagel 1958a: 275, 1959: 58, 1966: 258]
- bolivianus* Bernhauer, *Paederus*
[Bernhauer 1905b: 180, Bernhauer & Schubert 1912: 204, Blackwelder 1944: 122]
- bonariensis* Lynch, *Paederus*
[Lynch 1884: 294, Bernhauer & Schubert 1912: 204, Blackwelder 1944: 122]
- bonus* Last, *Paederus*
[Last 1980: 150]
- borneensis* Scheerpeltz, *Paederus*
[Scheerpeltz 1935: 605]
- brachycephalus* (Bernhauer), *Paederus*, *Oreopaederus*
[Bernhauer 1934d: 207, var. of *testaceus* Bernhauer, Fagel 1954a: 957, species, 1958a: 102]
- brachypterus* auct.
[Dahl *in litt.*, Gemminger & Harold 1868: 626, = *brevipennis* Lacordaire]
- brachypterus* Scheerpeltz, *Oncopaederus*
[Scheerpeltz 1957b: 265]
- brandti* Last, *Paederus*
[Last 1985: 334]
- brasiliensis* Dejean
[Dejean 1833: 65, *nomen nudum*]
- brasiliensis* Erichson, *Paederus*
[Erichson 1840: 658, Blanchard 1842: 85, Lucas 1857: 51, Burmeister 1865: 165, Gemminger & Harold 1868: 626, Fauvel 1877b: xxvi, 1901: 77, Lynch 1884: 290, Bernhauer & Schubert 1912: 204, Scheerpeltz 1933: 1221, Blackwelder 1939: 111, 1944: 122]
- braunsi* Bernhauer, *Paederus*
[Bernhauer 1915i: 316, Scheerpeltz 1933: 1221, Fagel 1958a: 298, 1966: 258]
- braunsi* auct.
[Cameron 1939: 4, misidentification, Fagel 1958a: 303, = *balfourbrowni* Fagel (*Paederus*)]
- brevelytratus* Coiffait
[Coiffait 1982c: 56, var. of *riparius* Linnaeus]
- breviceps* Bernhauer
[Bernhauer 1902: 37, Bernhauer & Schubert 1912: 206, = *fuscipes* Curtis, Wendeler 1926d: 328, Cameron 1925b: 36, 1931a: 40, Scheerpeltz 1935: 604, 1957a: 461]
- brevicollis* Fagel, *Paederus*
[Fagel 1958a: 337, 1966: 272]
- brevipennis* Dahl
[Dahl 1823: 17, *nomen nudum*, Lacordaire 1835: 430, = *brevipennis* Lacordaire, Heer 1839: 234]

brevipennis Lacordaire, *Paederus* (*Harpopaederus*)

[Lacordaire 1835: 430, Fairmaire & Laboulbène 1856: 570, Baudi 1857: 108, Kraatz 1857: 727, Gemminger & Harold 1868: 626, Fauvel 1873: 330, Mulsant & Rey 1878: 252, Reitter 1885: 154, 1889: 170, Ganglbauer 1895: 538, Bernhauer & Schubert 1912: 204, Scheerpeltz 1933: 1221, Jarrige 1944: 9, Scheerpeltz 1957a: 460, Smetana 1962: 49, Legosz-Owsianna 1963: 339, Lohse 1964: 134, Szujecki 1965b: 30, Scheerpeltz 1968: 47, Tikhomirova 1973: 175, Muona 1979: 16, Coiffait 1982c: 66, Bohác 1985: 373]

brittoni Fagel

[Fagel 1958a: 210, ssp. of *eximius* Reiche, 1966: 261]

bruchii Bernhauer, *Paederus*

[Bernhauer 1911b: 406, Bernhauer & Schubert 1912: 204, Blackwelder 1944: 122]

brunnescens (Fagel), *Paederus*, *Oreopaederus*

[Fagel 1954a: 917, 1958a: 101, 1960b: 196]

brunnescens orophilus (Fagel)

[Fagel 1954a: 919, 1958a: 101, 1960b: 196]

brunneus auct.

[Kirby *ms.*, Stephens 1829: 287, = *Sunius*]

brunneus (Bernhauer), *Paederus*, *Oreopaederus*

[Bernhauer 1927a: 108, Scheerpeltz 1933: 1221, Fagel 1954a: 925, 1958a: 101]

brunnipes Fabricius

[Fabricius 1792: 537 (*nec Staphylinus brunripes* Fabricius 1781: 336), 1801: 609, Paykull 1800: 429, Gravenhorst 1802: 57, = *Lathrobium*]

buehleri Scheerpeltz, *Eupaederus*

[Scheerpeltz 1957b: 269]

buluensis Last, *Paederus*

[Last 1979: 126]

burgeoni (Bernhauer), *Paederus*, *Paederidus*

[Bernhauer 1927a: 117, Scheerpeltz 1933: 1221, Fagel 1958a: 137, 1959: 39]

burgeoni auct.

[Cameron 1952b: 830, Fagel 1958a: 136, = *villiersi* Fagel]

burgeonianus Bernhauer

[Bernhauer 1934c: 232, Fagel 1954a: 961, = *alinderi* Wendeler, 1956: 188, 1958a: 102]

busalminensis Last, *Paederus*

[Last 1984: 109]

C*caeruleipennis* Boheman

[Boheman 1858: 33, Gemminger & Harold 1868: 626, = *cyanipennis* Guerin]

caffer Boheman, *Paederus*

[Boheman 1848: 290, Gemminger & Harold 1868: 626, Bernhauer & Schubert 1912: 204, Fagel 1958a: 244, 1961b: 268, 1965: 155]

caheni Fagel, *Pachypaederus*

[Fagel 1965: 137]

caligatus Erichson, *Paederus* (*Eopaederus*)

[Erichson 1840: 652, Lucas 1849: 121, Dietrich 1855: 203, Fairmaire & Laboulbène 1856: 570, Kraatz 1857: 729, Fauvel 1864: 14, 1873: 332, 1886a: 32, 1897: 281, Gemminger & Harold 1868: 626, Mulsant & Rey 1878: 264, Reitter 1889: 171, Czwalina 1889: 368, Ganglbauer 1895: 537, Bernhauer & Schubert 1912: 224, Kirschenblatt 1932: 219, Bernhauer 1931: 14, 1940b: 633, Scheerpeltz 1933: 1221, Jarrige 1944: 9, Scheerpeltz 1957a: 459, 1968: 47, Ferreira 1962: 14, Legosz-Owsianna 1963: 347, Lohse 1964: 135, Szujecki 1965b: 29, Tikhomirova 1973: 175, Hammond 1977: 27, Coiffait 1982c: 50, Bohác 1985: 372]

caligatus huetheri Bernhauer

[Bernhauer 1940b: 633, var. of *caligatus*, Scheerpeltz 1957a: 459, 1968: 47, Coiffait 1982c: 53, here treated as ssp. under Article 45(g) (ICZN 1985)]

caliginosus sic

[Shibata 1977: 23, misspelling of *caligatus*]

cameroni Last, *Paederus*

[Last 1979: 129]

canonicus (Casey), *Paederillus*, *Paederus*

[Casey 1905: 67, Bernhauer & Schubert 1912: 204, Leng 1920: 101, Blackwelder 1939: 111, Moore & Legner 1975: 133]

capensis Erichson, *Paederus*

[Erichson 1840: 651, Gemminger & Harold 1868: 626, Bernhauer & Schubert 1912: 204, Fagel 1958a: 242, 1966: 258]

capicola Fagel

[Fagel 1958a: 251, ssp. of *capitalis* Bernhauer, 1966: 258]

capillaris Fauvel, *Paederus*

[Fauvel 1895: 231, Bernhauer & Schubert 1912: 204, Cameron 1925b: 35, 1931a: 56, Scheerpeltz 1933: 1221, 1965b: 103, Gaedike 1981: 182]

capitalis Bernhauer, *Paederus*

[Bernhauer 1927a: 110, Scheerpeltz 1933: 1221, 1965b: 99, Cameron 1952b: 830, Fagel 1958a: 248, 1965: 155, 1966: 258]

capitalis capicola Fagel

[Fagel 1958a: 251, 1966: 258]

caprai Fagel, *Paederus*

[Fagel 1958a: 254, 1965: 155]

- captus* Wendeler, *Paederus*
[Wendeler 1931: 41, Scheerpeltz 1933: 1221, Fagel 1958a: 190, 1966: 272]
- carbonarius* Gautier
[Gautier 1861a: 394, Saulcy 1862: vi, Fauvel 1864: 13, Gautier 1866: 114, Marseul 1866: vi, = *longicornis* Aubé, Fauvel 1866: 25, Kraatz 1867: 415, Gemminger & Harold 1868: 628, = *sanguinicollis* Stephens, Mulsant & Rey 1878: 246, = *ruficollis* Fabricius, Bernhauer & Schubert 1912: 210, = *rubrothoracicus* Goeze, Legosz-Owsianna 1933: 327, Scheerpeltz 1968: 46, Coiffait 1982c: 25, var. of *rubrothoracicus* Goeze]
- cariniventris* (Cameron), *Paederus*, *Paederidus*
[Cameron 1948: 40, Fagel 1958a: 126]
- carolinae* (Casey), *Paederillus*, *Paederus*
[Casey 1905: 63, Bernhauer & Schubert 1912: 204, Leng 1920: 101, Blackwelder 1939: 111, Moore & Legner 1975: 133]
- carpathicola* Scheerpeltz
[Scheerpeltz 1957a: 455, ssp. of *rubrothoracicus* Goeze, Szujewski 1965b: 26, Coiffait 1982c: 25, var. of *rubrothoracicus* Goeze]
- carpathicus* Wendeler, *Paederus* (*Dioncopaederus*)
[Wendeler 1926c: 204, Scheerpeltz 1933: 1221, 1957a: 474, Coiffait 1982c: 60]
- carpaticus sic*
[Tikhomirova 1973: 175, misspelling of *carpathicus*]
- carpetanicus* Outerelo
[Outerelo 1980: 31, Coiffait 1982c: 68, = *lusitanicus* Aubé]
- cartaginis* Bernhauer, *Paederus*
[Bernhauer 1920a: 44, Scheerpeltz 1933: 1221, Blackwelder 1944: 122]
- castaneus auct.*
[Knoch in coll., Gravenhorst 1802: 60, = *castaneus* Gravenhorst]
- castaneus* Gravenhorst
[Gravenhorst 1802: 60, 1806: 139, Olivier 1812: 628, Mannerheim 1831: 454, = *Rugilus*, now in *Medon*]
- caucasicus* Scheerpeltz
[Scheerpeltz 1957a: 456, ssp. of *rubrothoracicus* Goeze, Coiffait 1982c: 25, var. of *rubrothoracicus* Goeze]
- celisianus* Fagel, *Oreopaederus*
[Fagel 1973: 152]
- celisianus peculiaris* Fagel
[Fagel 1973: 153]
- centralis* Last, *Paederus*
[Last 1985: 334]
- cephalicus auct.*
[Wendeler 1931: 46, *lapsus calami*, nec Fauvel]
- cephalotes* Motschulsky
[Motschulsky 1849: 86, Kraatz 1857: 725, Gemminger & Harold 1868: 626, Solsky 1871: 163, Fauvel 1873: 331, Ganglbauer 1895: 538, Bernhauer & Schubert 1912: 207, = *littoralis* Gravenhorst, Scheerpeltz 1933: 1225, 1957a: 465, Legosz-Owsianna 1963: 1222, Coiffait 1982c: 58]
- ceylonicus* Bernhauer, *Paederus*
[Bernhauer 1902: 37, Bernhauer & Schubert 1912: 205, Cameron 1925b: 35, 1931a: 38, Scheerpeltz 1933: 1222]
- chalybaeus sic*
[Bernhauer & Schubert 1912: 205, misspelling of *chalybeus*, Fagel 1958a: 457]
- chalybeus* (Erichson), *Paederus*, *Paederidus*
[Erichson 1840: 664, Gemminger & Harold 1868: 626, Fagel 1958a: 457]
- cheesmani* Cameron, *Paederus*
[Cameron 1937: 97]
- chilensis* Fauvel
[Fauvel 1868: 20, 1886b: 148, 1889a: 61, Gemminger & Harold 1868: 626, Bernhauer & Schubert 1912: 205, Bernhauer 1927c: 399, = *intermedius* Boheman, Scheerpeltz 1933: 1222]
- chimbuensis* Last, *Paederus*
[Last 1979: 134]
- chinensis* Bernhauer, *Paederus* (*Oedopaederus*)
[Bernhauer 1931b: 1, Wu 1937: 328, Scheerpeltz 1933: 1222, 1957a: 467]
- chinensis* Bernhauer
[Bernhauer 1934b: 6, homonym, ssp. of *fuscipes* Curtis, Scheerpeltz 1957a: 461, = *sinensis* Scheerpeltz replacement name]
- chopardi* Fagel, *Paederus*
[Fagel 1958a: 317, 1966: 259]
- cinctus* Say
[Say 1834: 457, Erichson 1840: 639, = *Sunius*, now in *Astenus*]
- cingulatus* MacLeay
[MacLeay 1871: 146, Fauvel 1877a: 223, = *cruenticollis* Germar, 1878a: 236, 1889b: 258, var. of *cruenticollis* Germar, 1903b: 265, Lea 1904: 63, species, Bernhauer & Schubert 1912: 205, var. of *cruenticollis* Germar, Bernhauer 1920c: 9, Lea 1923: 14, = *cruenticollis* Germar]
- citernii auct.*
[Gridelli in litt., Fagel 1958a: 185, = *arrowianus* Bernhauer]

- clavicornis* Lentz
[Lentz 1856: 57, here, possibly = *Cucujus clavicornis* Kugelann = *Nausibius clavicornis* (Kugelann) (Silvanidae)]
- clermonti* Chapman
[Chapman 1930: 102, ab. (infrasubspecific) of *ruficollis* Fabricius, Scheerpeltz 1933: 1227, Coiffait 1982c: 25, var. of *rubrothoracicus* Goeze]
- coarctatus* Erichson, *Paederus*
[Erichson 1840: 666, Gemminger & Harold 1868: 626, Fauvel 1877b: xxvi, Bernhauer & Schubert 1912: 205, Blackwelder 1944: 122]
- coelestinus* (Bernhauer), *Paederus*, *Paederidus*
[Bernhauer 1936a: 240, Fagel 1958a: 153, Coiffait 1982c: 19]
- coeruleipennis* sic
[Bernhauer & Schubert 1912: 205, misspelling of *caeruleipennis*, Blackwelder 1944: 122]
- coerulescens* (Erichson), *Paederus*, *Paederidus*
[Erichson 1840: 664, Gemminger & Harold 1868: 626, Bernhauer & Schubert 1912: 205, Fagel 1958a: 457]
- coerulipennis* sic
[Gemminger & Harold 1868: 626, misspelling of *caeruleipennis*]
- colasi* Fagel, *Paederus*
[Fagel 1958a: 237]
- collaris* (Boheman), *Paederus*, *Paederidus*
[Boheman 1848: 289, Gemminger & Harold 1868: 626, Fagel 1958a: 146, 1961b: 266]
- collaris auct.*
[Klug *in litt.*, Fagel 1958a: 457, = *perrieri* Fauvel]
- collarti* Cameron
[Cameron 1939: 3, Fagel 1958a: 269, = *xanthocerus* Eppelsheim, 1959: 50]
- colonus* (Wendeler), *Paederus*, *Paederidus*
[Wendeler 1931: 42, Scheerpeltz 1933: 1222, Fagel 1958a: 114, 1963: 343]
- colonus alberti* Fagel
[Fagel 1958a: 115, 1959: 38]
- columbinus* Laporte, *Paederus*
[Laporte 1835: 123, Gemminger & Harold 1868: 626, Sharp 1886: 617, Fauvel 1891: 100, 1901: 77, Bernhauer & Schubert 1912: 205, Blackwelder 1939: 112, 1944: 122, Scheerpeltz 1965a: 217]
- combustus* Fauvel, *Paederus*
[Fauvel 1879a: 88, Bernhauer & Schubert 1912: 205]
- comorensis* Bernhauer, *Paederus*
[Bernhauer 1917: 557, Fagel 1958a: 458, Jarrige 1970: 34]
- compactus* Wendeler
[Wendeler 1931: 39, Scheerpeltz 1933: 1222, Fagel 1958a: 73, = *crassus* Boheman, 1961b: 266]
- compotens* LeConte, *Paederus*, *Paederillus*, *Paederus*
[LeConte 1863: 48, 1878: 396, Gemminger & Harold 1868: 626, Austin 1876: 9, Casey 1905: 64, Bernhauer & Schubert 1912: 205, Leng 1920: 101, Scheerpeltz 1933: 1222, Blackwelder 1939: 112, Moore & Legner 1975: 133]
- condei* Wendeler, *Paederus*
[Wendeler 1933: 57, Blackwelder 1944: 122]
- confinis* Zetterstedt
[Zetterstedt 1838: 69, Erichson 1840: 654, = *riparius* Linnaeus, Kraatz 1857: 728, Thomson 1860: 196, = *littoralis* Gravenhorst, Gemminger & Harold 1868: 627, Fauvel 1869b: 491, Bernhauer & Schubert 1912: 207, Scheerpeltz 1957a: 465, Legosz-Owsianna 1963: 349, Coiffait 1982c: 58]
- confusus* Fagel, *Pachypaederus*
[Fagel 1958a: 75,]
- confusus* Fagel, *Paederus*
[Fagel 1958a: 433, 1959: 37, 1966: 269]
- congoensis* Bernhauer, *Paederus*
[Bernhauer 1915c: 291, Scheerpeltz 1933: 1222, Fagel 1958a: 265, 1959: 48, 1963: 344, 1966: 259]
- conicipennis auct.*
[Fauvel *in litt.*, Wendeler 1931: 47, = *conicipennis* Wendeler]
- conicipennis* Wendeler, *Paederus*
[Wendeler 1931: 47, Scheerpeltz 1933: 1222, Blackwelder 1944: 122, Scheerpeltz 1960: 81]
- conicollis* Motschulsky, *Paederus*
[Motschulsky 1860: 73, Gemminger & Harold 1868: 626, Fauvel 1903a: 153, 1904a: 54, Bernhauer & Schubert 1912: 205, Cameron 1925b: 35, 1931a: 51, Scheerpeltz 1933: 1222]
- conspicuous* Erichson, *Paederus*
[Erichson 1840: 660, Gemminger & Harold 1868: 626, Fauvel 1901: 77, Bernhauer & Schubert 1912: 205, Blackwelder 1944: 122, Scheerpeltz 1965a: 217]
- cooperi* Scheerpeltz, *Paederus*
[Scheerpeltz 1933: 1222, replacement name for *viridipennis* Bernhauer 1931, Fagel 1958a: 177, 1966: 261]
- copiosus* Last, *Paederus*
[Last 1979: 126, 1985: 329]
- corallifer auct.*
[Millet *in litt.*, Fauvel 1873: 335, = *sanguinicollis* Stephens]

- cordovens* Sharp, *Paederus*
[Sharp 1886: 616, Bernhauer & Schubert 1912: 205, Blackwelder 1944: 122]
- coriaceus* Fauvel, *Paederus*
[Fauvel 1869a: 57, 1869b: 491]
- corsicus* Gautier
[Gautier 1861a: 393, Saulcy 1862: vi, Fauvel 1864: 12, = *limnophilus* Erichson, Marseul 1866: vi, Gemminger & Harold 1868: 627, Fauvel 1873: 334, = *fuscipes* Curtis, Czwalina 1889: 368, Ganglbauer 1895: 537, Bernhauer & Schubert 1912: 206, Cameron 1925b: 35, 1931a: 40, Jarrige 1944: 8, Scheerpeltz 1957a: 461, Legosz-Owsianna 1963: 335, Coiffait 1982c: 60]
- corticinus auct.*
[Knoch *in coll.*, Gravenhorst 1802: 184, = *corticinus* Gravenhorst]
- corticinus* Gravenhorst
[Gravenhorst 1802: 184, 1806: 139, Olivier 1812: 629, Sharp 1886: 552, = *Aderocharis*, now in *Achenomorphus*]
- costaricensis* Sharp, *Paederus*
[Sharp 1886: 616, Bernhauer & Schubert 1912: 205, Blackwelder 1944: 122]
- coxalis* Fauvel, *Paederus* (*Paederus*)
[Fauvel 1895: 232, Bernhauer & Schubert 1912: 205, Cameron 1925b: 35, 1931a: 42, Scheerpeltz 1933: 1222, Shibata 1973: 44]
- crassicolis auct.*
[Fauvel *in litt.*, Wendeler 1931: 46, = *crassicolis* Wendeler]
- crassicolis* Wendeler, *Paederus*
[Wendeler 1931: 46, Scheerpeltz 1933: 1222, Blackwelder 1944: 122]
- crassus* (Boheman), *Paederus*, *Pachypaederus*
[Boheman 1848: 288, Gemminger & Harold 1868: 626, Bernhauer & Schubert 1912: 205, Blackwelder 1939: 112, Fagel 1958a: 73, 1961b: 266]
- crassus* Boheman *partim*
[Fagel 1958a: 75, = *confusus* Fagel]
- crassus* Boheman *partim*
[Fagel 1958a: 75, = *sparsicolis* Fagel]
- crassus* Boheman *partim*
[Fagel 1958a: 77, = *wendeleri* Fagel]
- crassus meersmanae* Fagel
[Fagel 1965: 144]
- crebrepunctatus* Eppelsheim
[Eppelsheim 1895b: 210, Fauvel 1904b: 290, Bernhauer 1908b: 105, 1931c: 575, Bernhauer & Schubert 1912: 205, Cameron 1950b: 25, Fagel 1958a: 208, = *eximius* Reiche]
- crebripunctatus sic*
[Chapin 1926: 369, misspelling of *crebrepunctatus*]
- cribratus auct.*
[Fauvel *in litt.*, Fagel 1958a: 89, = *puncticollis* Bernhauer]
- cribratus auct.*
[Bernhauer *in litt.*, Fagel 1958a: 141, = *rufofasciatus* Bernhauer]
- cribricollis auct.*
[Fauvel *in litt.*, Fagel 1958a: 124, = *cribricollis* Bernhauer]
- cribricollis* (Bernhauer), *Paederus*, *Paederidus*
[Bernhauer 1915c: 289, Scheerpeltz 1933: 1222, Fagel 1958a: 124]
- cribripennis* Fagel, *Paederus*
[Fagel 1959: 52, 1966: 258]
- crinitus* Fagel, *Paederidus*
[Fagel 1958a: 150]
- cruenticollis* Germar, *Paederus*
[Germar 1848: 174, Gemminger & Harold 1868: 626, MacLeay 1871: 146, Fauvel 1877a: 223, 1878b: 516, 1889b: 258, 1903b: 265, Blackburn 1891: 72, 1899: 23, Lea 1904: 64, 1923: 14, Bernhauer & Schubert 1912: 205, Blackwelder 1939: 112, Scheerpeltz 1978: 196]
- Ctenopaederus* Fagel
[Fagel 1958a: 69, *genus*, type *Paederus andrewesi* Fauvel]
- cumanus* Frank, *Paederus*
[here, *nom. nov.* for *bicolor* Wendeler, *nec* Olivier]
- currax* Sharp, *Paederus*
[Sharp 1886: 611, Bernhauer & Schubert 1912: 205, Blackwelder 1944: 122]
- curticeps* Bernhauer, *Paederus*
[Bernhauer 1910: 369, Bernhauer & Schubert 1912: 205, Blackwelder 1944: 122]
- curticollis* Bernhauer, *Paederus*
[Bernhauer 1905b: 182, Bernhauer & Schubert 1912: 205, Blackwelder 1944: 122]
- cyanelus* Fauvel, *Paederus*
[Fauvel 1879a: 88, Bernhauer & Schubert 1912: 205]
- cyaneipennis sic*
[Fauvel 1886b: 143, misspelling of *cyaneipennis*]
- cyanescens* Jarrige, *Madecapaederus*
[Jarrige 1978: 259]
- cyaneus* Cameron, *Paederus*
[Cameron 1937: 95, Puthz 1969: 529, Last 1985: 329]
- cyaneipennis* (Guerin), *Paederus*, *Eupaederus*
[Guerin 1830: 62, Erichson 1840: 933, Gemminger &

- Harold 1868: 626, Bernhauer & Schubert 1912: 205, Blackwelder 1944: 122, Scheerpeltz 1957b: 269]
- cyanocephalus sic*
[Motschulsky 1860a: 74, misspelling of *cyanocephalus*]
- cyanocephalus* Erichson, *Paederus*
[Erichson 1840: 662, Kraatz 1859a: 152, 1859b: xxxi, Gemminger & Harold 1868: 626, Fauvel 1878a: 236, Bernhauer & Schubert 1912: 205, Cameron 1925b: 35, 1931a: 44, Blackwelder 1939: 112, Scheerpeltz 1933: 1222, 1935: 608]
- cyanocephalus auct.*
[Baer 1886: 107, misidentification, Fauvel 1886b: 148, = *chilensis* Fauvel]
- cyanopterus* Cameron, *Paederus*
[Cameron 1942a: 87]
- cyanurus auct.*
[Kirby *ms*, Stephens 1829: 288, = *sanguinicollis* Stephens]
- D**
- dammermani* Cameron, *Paederus*
[Cameron 1925a: 180, Scheerpeltz 1933: 1222, 1935: 621]
- darfurensis* Bernhauer, *Paederus*
[Bernhauer 1937a: 294, Fagel 1958a: 338, 1966: 257]
- debiliior* Eppelsheim, *Paederus* (*Eopaederus*)
[Eppelsheim 1892: 339, Bernhauer & Schubert 1912: 205, Kirschenblatt 1932: 218, Scheerpeltz 1957a: 459, Tikhomirova 1973: 175, Coiffait 1982c: 54]
- decellei* Fagel
[Fagel 1958a: 134, ssp. of *montanellus* Bernhauer]
- decipiens* Wendeler
[Wendeler 1926b: 201, ssp. of *lutiventris* Wendeler]
- delagoanus* (Bernhauer), *Paederus*, *Pachypaederus*
[Bernhauer 1927a: 103, Scheerpeltz 1933: 1222, Fagel 1958a: 79]
- delegoanus sic*
[Fagel 1958a: 79, misspelling of *delagoanus*]
- delkeskampii* Fagel, *Paederus*
[Fagel 1958a: 290, 1966: 261]
- densipennis* Bernhauer, *Paederus*
[Bernhauer 1916: 30, Wu 1937: 328, Scheerpeltz 1957a: 474]
- densipennis auct.*
[Fauvel *in litt.*, Fagel 1958a: 159, = *minimus* Bernhauer]
- densiventris* Bernhauer, *Paederus*
[Bernhauer 1927a: 104, Scheerpeltz 1933: 1222, Fagel 1958a: 219, 1960a: 121, 1966: 262]
- densiventris katshensis* Fagel
[Fagel 1960a: 121]
- densiventris auct.*
[Bernhauer *in litt.*, Fagel 1958a: 447, = *benicki* Wendeler]
- descarpentriasi* Jarrige, *Madecapaederus*
[Jarrige 1978: 261]
- diamesus* Kirschenblatt, *Paederus* (*Paederus*)
[Kirschenblatt 1932: 221, Scheerpeltz 1957a: 462, Tikhomirova 1973: 175, Coiffait 1982c: 57]
- dilutipes* (Fauvel), *Paederus*, *Madecapaederus*
[Fauvel 1905b: 152, Bernhauer & Schubert 1912: 205, Fagel 1958a: 457]
- dilutus* Jarrige, *Madecapaederus*
[Jarrige 1978: 261]
- dimidiatus* Panzer
[Panzer 1796a: 24, Gravenhorst 1802: 40, = *Staphylinus*, now in *Philonthus*]
- dimidiatus auct.*
[Kirby *ms*, Stephens 1829: 288, = *Sunius*]
- Dioncopaederus* Scheerpeltz
[Scheerpeltz 1957a: 464, subgenus, type *Paederus littoralis* Gravenhorst, Szujewski 1965b: 29, Coiffait 1982c: 58]
- Diplopaederus* Scheerpeltz
[Scheerpeltz: 1957b: 272, genus, type *Paederus handschini* Scheerpeltz]
- discopunctatus* Say
[Say 1834: 457, Erichson 1840: 643, = *Sunius*, now in *Astenus*]
- distinctus* Cameron, *Paederus*
[Cameron 1931a: 57, Scheerpeltz 1933: 1222, 1957a: 474]
- diversiceps* Cameron, *Paederus*
[Cameron 1931a: 61, Scheerpeltz 1933: 1222]
- diversicollis auct.*
[Dahl *in litt.*, Gemminger & Harold 1868: 627, = *littoralis* Gravenhorst]
- diversipennis auct.*
[Bernhauer *in litt.*, Fagel 1958a: 220, = *rufocyanus* Bernhauer]
- dollmani* Bernhauer, *Paederus*
[Bernhauer 1927a: 112 *partim*, Scheerpeltz 1933: 1222, Fagel 1958a: 274, 1973: 157]
- dollmani* Bernhauer
[Bernhauer 1927a: 112 *partim*, Fagel 1958a: 272, = *simillimus* Bernhauer]
- dolosus* Jarrige, *Madecapaederus*
[Jarrige 1970: 33]

- doriae* Fauvel, *Paederus*
[Fauvel 1879a: 89, Bernhauer & Schubert 1912: 205]
- drakensbergensis* Fagel, *Paederus*
[Fagel 1965: 153, 1966: 258]
- drescheri* Cameron, *Paederus*
[Cameron 1936: 43]
- dubiosus* Fagel
[Fagel 1954a: 954, 1958a: 102, ssp. of *stricticornis* Fagel]
- dubius* Kraatz
[Kraatz 1859a: 151, Gemminger & Harold 1868: 628, Fauvel 1869b: 491, = *tamulus* Erichson, 1903a: 153, 1904a: 54, Bernhauer 1902: 38, Bernhauer & Schubert 1912: 211, Cameron 1925b: 37, 1931a: 43, Scheerpeltz 1935: 609, Wu 1937: 329, Shibata 1977: 23, Gaedike 1981: 187, Coiffait 1982c: 50]
- duplex* Eppelsheim, *Paederus*
[Eppelsheim 1895b: 211, Fauvel 1904b: 290, Bernhauer & Schubert 1912: 205, Bernhauer 1931c: 575, 1940a: 135, Scheerpeltz 1933: 1222, Cameron 1950b: 25, Fagel 1958a: 349, 1966: 269]
- duplex rothschildi* Fagel
[Fagel 1958a: 353, 1966: 266]
- duplex spectabilis* Bernhauer
[Bernhauer 1915a: 116, species, Fagel 1958a: 351, ssp.]
- duplex auct.*
[Fauvel *in litt.*, Fagel 1958a: 148, = *gracilior* Fagel]
- ## E
- ecuadorensis* Bernhauer, *Paederus*
[Bernhauer 1934e: 158, Blackwelder 1944: 122]
- edwardsi* Bernhauer, *Paederus*
[Bernhauer 1927a: 104, Scheerpeltz 1933: 1222, Fagel 1958a: 458]
- eidmanni auct.*
[Scheerpeltz *in litt.*, Fagel 1958a: 370, = *eidmanni* Fagel]
- eidmanni* Fagel, *Paederus*
[Fagel 1958a: 370, 1966: 264]
- elegans* Wendeler, *Paederus*
[Wendeler 1931: 47, Scheerpeltz 1933: 1222, Blackwelder 1944: 122]
- elegantulus* Wendeler, *Paederus*
[Wendeler 1931: 39, Scheerpeltz 1933: 1222, 1935: 633, Cameron 1936: 42]
- elongatus* (Linnaeus)
[Linnaeus 1767: 685, 1789: 415, Fabricius 1775: 268, 1781: 339, 1787: 223, 1792: 537, 1801: 609, Panzer 1793a: 12, Gravenhorst 1802: 55, = *Lathrobium* *elongatus auct.*
[Paykull 1789: 25, misidentification, 1800: 428, Gyllenhal 1810: 366, = *Lathrobium*]
- elongatus auct.*
[Ferrari *in litt.*, Kraatz 1857: 731, = *gemellus* Kraatz, Gemminger & Harold 1868: 627]
- elongatus* Wendeler, *Paederus*
[Wendeler 1931: 46, Scheerpeltz 1933: 1222, Blackwelder 1944: 122, = *archeus* Blackwelder, unnecessary replacement name]
- Eopaederus* Scheerpeltz
[Scheerpeltz 1957a: 458, subgenus, type *Paederus caligatus* Erichson, Szujewski 1965b: 29, Shibata 1973: 43, 1977: 23, Coiffait 1982c: 49]
- erichsoni* Wollaston
[Wollaston 1867: 247, unjustified replacement name for *angolensis* Erichson, Gemminger & Harold 1868: 626, Fauvel 1903a: 154, = *fuscipes* Curtis, 1904b: 290, Bernhauer & Schubert 1912: 206, Cameron 1925b: 35, 1931a: 40, Scheerpeltz 1957a: 461, Fagel 1958a: 311, 1959: 62, 1960a: 121, 1961b: 268, Legosz-Owsianna 1963: 335, Coiffait 1982c: 60]
- erichsoni* Bernhauer
[Bernhauer 1908c: 16, homonym, Bernhauer & Schubert 1912: 203, = *antipodum* Bernhauer & Schubert, replacement name, Lea 1923: 15, = *meyricki* Blackburn, Scheerpeltz 1933: 1220]
- ertli* Bernhauer, *Paederus*
[Bernhauer 1915c: 291, Scheerpeltz 1933: 1222, Fagel 1958a: 413]
- erythraeanus* Fagel, *Paederidus*
[Fagel 1958a: 143]
- erythraeanus kenyensis* Fagel
[Fagel 1958a: 145]
- erythraeanus piliventris* Fagel
[Fagel 1968: 197]
- erythroderus* Erichson, *Paederus*
[Erichson 1840: 663, Gemminger & Harold 1868: 627, Sharp 1886: 612, Bernhauer & Schubert 1912: 205, Blackwelder 1944: 122]
- Eupaederus* Scheerpeltz
[Scheerpeltz 1957b: 269, genus, type *Paederus cyanipennis* Guerin]
- Eupaederus auct.*
[Fagel *in litt.*, Fagel 1958a: 104, = *Allopaederus* Fagel]
- excellens* Bernhauer, *Paederus*
[Bernhauer 1942a: 12]
- eximius* Reiche, *Paederus*
[Reiche 1847: 281, Gemminger & Harold 1868: 627,

- Bernhauer & Schubert 1912: 205, Fagel 1958a: 208, 1966: 261]
- eximius brittoni* Fagel
[Fagel 1958a: 210, 1966: 261]
- extensus* Mannerheim
[Mannerheim 1831: 453, Dejean 1833: 66, = *Astenus*]
- extensus auct.*
[Gyllenhal unpublished, Mannerheim 1831: 453, = *extensus* Mannerheim]
- extraneus* Wiedemann, *Paederus*
[Wiedemann 1823: 133, Dejean 1833: 65, Erichson 1840: 661, Kraatz 1859a: 152, Gemminger & Harold 1868: 627, Bernhauer & Schubert 1912: 206, Cameron 1925b: 35, 1931a: 41, Scheerpeltz 1933: 1223]
- F**
- fageli* Jarrige, *Madecapaederus*
[Jarrige 1978: 260]
- fallaciosus* (Fagel), *Paederus*, *Oreopaederus*
[Fagel 1954a: 941, 1958a: 102]
- fallax* (Fagel), *Paederus*, *Oreopaederus*
[Fagel 1954a: 935, 1956: 187, 1958a: 101]
- falsus* Fagel, *Oreopaederus*
[Fagel 1961a: 268]
- fasciatus auct.*
[Bernhauer *in litt.*, Fagel 1958a: 141, = *rufofasciatus* Bernhauer]
- fastuosus* Klug, *Paederus*
[Klug 1834: 140, Erichson 1840: 665, Gemminger & Harold 1868: 627, Bernhauer & Schubert 1912: 206, Fagel 1958a: 458]
- fauveli* Quedenfeldt
[Quedenfeldt 1881: 292, Reitter 1889: 171, Czwalina 1889: 368, Fauvel 1886a: 32, 1897: 281, 1907: 23, Bernhauer & Schubert 1912: 205, Bernhauer 1931: 14, Peyerimhoff 1931: 29, Scheerpeltz 1933: 1223, Cameron 1952b: 830, Fagel 1958a: 129, = *luctuosus* Klug, 1960a: 117, 1961b: 266]
- faeae* Fauvel, *Paederus*
[Fauvel 1895: 232, Bernhauer & Schubert 1912: 206, Cameron 1925b: 35, 1931a: 50, Scheerpeltz 1933: 1223]
- femoralis* LeConte, *Paederus*
[LeConte 1858: 62, 1878: 395, Gemminger & Harold 1868: 627, Austin 1876: 8, Casey 1905: 61, Bernhauer & Schubert 1912: 206, Leng 1920: 101, Scheerpeltz 1933: 1223, Blackwelder 1939: 112, 1944: 122, Moore & Legner 1975: 133]
- femoratus* Cameron, *Paederus*
[Cameron 1931b: 357, Scheerpeltz 1933: 1223]
- fennicus* Sahlberg
[Sahlberg 1876: 38, Fauvel 1879a: 90, = *fuscipes* Curtis, Bernhauer & Schubert 1912: 206, Cameron 1925b: 35, 1931a: 40, Jarrige 1944: 9, Scheerpeltz 1957a: 461, Legosz-Owsianna 1963: 335, Coiffait 1982c: 60]
- ferreirai* Scheerpeltz, *Paederus*
[Scheerpeltz 1978: 197]
- ferus* Erichson, *Paederus*
[Erichson 1840: 659, Burmeister 1861: 484, Gemminger & Harold 1868: 627, Lynch 1884: 293, Bernhauer & Schubert 1912: 206, Blackwelder 1944: 122, Scheerpeltz 1965a: 216]
- fijiensis* Cameron, *Paederus*
[Cameron 1943b: 564]
- filicornis* Sharp, *Paederus*
[Sharp 1886: 612, Bernhauer & Schubert 1912: 206, Blackwelder 1944: 122]
- filicornis* (Fagel),
[Fagel 1954a: 949, homonym, = *minor* Bernhauer, 1958a: 102]
- filiformis* Fabricius
[Fabricius 1792: 538, 1801: 609, Gravenhorst 1802: 54, = *Lathrobium*]
- filiformis* Latreille
[Latreille 1806: 293, homonym, Olivier 1812: 629, Erichson 1839: 525, = *procerus* Gravenhorst, = *Sunius*, now in *Astenus*]
- filiformis auct.*
[Paykull 1800: 429, misidentification, Gravenhorst 1802: 55, = *Lathrobium*]
- filipes* (Fagel), *Paederus*, *Allopaederus*
[Fagel 1956: 190, 1958a: 106]
- filum* Walzl
[Walzl 1838: 267, Erichson 1840: 932, = *Sunius*, now in *Astenus*]
- finisterrae* Illiger
[Illiger 1807: 356, Gemminger & Harold 1868: 626, = *littoralis* Gravenhorst, Bernhauer & Schubert 1912: 207, Scheerpeltz 1957a: 465, Legosz-Owsianna 1963: 349]
- flaviceps* Bernhauer, *Paederus*
[Bernhauer 1927c: 401, Scheerpeltz 1933: 1223]
- flavivarsis* Cameron, *Paederus*
[Cameron 1949a: 316, Fagel 1958a: 322, 1963: 347, 1966: 259]
- flavocaudatus* Bernhauer, *Paederus*
[Bernhauer 1909: 241, Bernhauer & Schubert 1912: 206, Blackwelder 1944: 122]

- flavoides* (Fagel), *Paederus*, *Oreopaederus*
[Fagel 1954a: 947, 1958a: 102]
- flavoterminalis* (Cameron), *Paederus*, *Megalopaederus*
[Cameron 1949b: 466, Shibata 1973: 44]
- flavus* (Fagel), *Paederus*, *Oreopaederus*
[Fagel 1954a: 944, 1958a: 102]
- floridanus* Austin, *Paederus*, *Paederillus*, *Paederus*
[Austin 1876: 10, LeConte 1878: 396, Casey 1905: 62,
Bernhauer & Schubert 1912: 206, Leng 1920: 101,
Blackwelder 1939: 112, Moore & Legner 1975: 133]
- forficularius* (Fauvel), *Paederus*, *Madecapaederus*
[Fauvel 1905b: 152, Bernhauer & Schubert 1912: 206,
Fagel 1958a: 457]
- formosanus* (Adachi), *Paederus*, *Megalopaederus*
[Adachi 1939: 166, Cameron 1949b: 466, Shibata
1973: 45]
- fortior* Bernhauer
[Bernhauer *in litt.*, Fagel 1958a: 114, = *colonus*
Wendeler]
- fracticornis* (Paykull)
[Paykull 1790: 135, 1800: 430, Gravenhorst 1802: 54,
= *Lathrobium*, now in *Ochtheophilum*]
- fragilis* Latreille
[Latreille 1804: 347, 1806: 292, Olivier 1812: 629, =
fragilis Gravenhorst]
- fragilis* Gravenhorst
[Gravenhorst 1806: 140, homonym, Olivier 1812:
629, = *fragilis* Latreille, Stephens 1829: 288, =
Rugilus]
- fratellus* Bernhauer
[Bernhauer 1929: 123, replacement name for *socius*
Bernhauer 1927a, Scheerpeltz 1933: 1223, Fagel
1958a: 142, = *arrowi* Bernhauer]
- freudei* Fagel, *Paederus*
[Fagel 1958a: 253, 1966: 258]
- freyi* Fagel, *Paederus*
[Fagel 1958a: 203, 1966: 263]
- freyi* Fagel, *Paederidus*
[Fagel 1962a: 69]
- frontalis* auct.
[Fauvel *in litt.*, Fagel 1958a: 415, = *frontalis* Fagel]
- frontalis* Fagel, *Paederus*
[Fagel 1958a: 415, 1966: 265, Gaedike 1981: 191]
- fruhstorferi* auct.
[Fauvel *in litt.*, Wendeler 1931: 37, = *fruhstorferi*
Wendeler]
- fruhstorferi* Wendeler, *Paederus*
[Wendeler 1931: 37, Scheerpeltz 1933: 1223, 1935:
598]
- fulgidipennis* Fagel, *Paederus*
[Fagel 1958a: 398, 1966: 265]
- fulgidus* (Fabricius)
[Fabricius 1787: 220, 1792: 537, 1801: 609, Thomson
1860: 188, Smetana 1979: 51, = *Gauropterus*]
- fulgidus* auct.
[Paykull 1789: 22, misidentification, = *Othius punc-
tulatulus* (Goeze), Gravenhorst 1802: 48, misidentifi-
cation, = *Xantholinus glabratus* (Gravenhorst), *fide*
Smetana 1979: 44]
- fulvicornis* Erichson, *Paederus*
[Erichson 1840: 665, Kraatz 1859a: 150, Gemminger
& Harold 1868: 627, Bernhauer & Schubert 1912:
206, Scheerpeltz 1935: 627]
- fulvipennis* Fabricius
[Fabricius 1792: 537, 1801: 609, Gyllenhal 1810: 365,
= *Lathrobium*, now in *Othius*]
- fulvipennis* auct.
[Panzer 1796a: 23, misidentification, Gravenhorst
1802: 44, Gyllenhal 1810: 357, = *Staphylinus*]
- fulvipes* Stephens
[Stephens 1829: 288, misspelling of *fuscipes*]
- fulvocaudatus* Adachi
[Adachi 1939: 166, var. of *kosempoensis* Bernhauer,
here treated as ssp. under Article 45(g) (ICZN 1985)]
- funnebris* Fagel, *Paederidus*
[Fagel 1958a: 131]
- furcifer* (Fagel), *Paederus*, *Oreopaederus*
[Fagel 1954a: 959, 1958a: 102]
- furciferoides* (Fagel), *Paederus*, *Oreopaederus*
[Fagel 1956: 189, 1958a: 102]
- fuscipes* Curtis, *Paederus* (*Heteropaederus*)
[Curtis 1826: 108, Stephens 1833: 280, Fairmaire &
Laboulbène 1856: 570, Gemminger & Harold 1868:
627, Fauvel 1869b: 491, 1873: 334, 1879a: 90, 1886a:
32, 1886b: 148, 1897: 281, 1903a: 154, 1904a: 53, 1904b:
290, Ganglbauer 1895: 537, Bernhauer & Schubert
1912: 206, Bernhauer 1915g: 362, 1915i: 315, 1931c:
575, Lea 1923: 14, Cameron 1925b: 35, 1930b: 332,
1931a: 40, 1949b: 466, Wendeler 1926d: 328, Kir-
schenblatt 1932: 217, Roubal 1932: 60, Scheerpeltz
1933: 1223, Koch 1934: 82, Blackwelder 1939: 112,
1944: 123, Jarrige 1944: 8, 1971: 491, Hansen 1951: 224,
Fagel 1958a: 311, 1959: 62, 1960a: 121, 1961b: 268,
1963: 347, 1966: 259, Smetana 1959: 204, Scheerpeltz
1957a: 461, 1963: 8, 1965b: 103, 1968: 47, 1976: 82, Fer-
reira 1962: 14, Legosz-Owsianna 1963: 335, Szujewski
1965b: 29, Tikhomirova 1973: 175, Shibata 1973: 43,
1977: 24, Coiffait 1976: 243, 1982a: 34, 1982b: 60,
Hammond 1977: 27, Muona 1979: 16, Tronquet]

- 1981: 70, Biswas & Sen Gupta 1982: 153, Bohác 1985: 371]
- fuscipes chinensis* Bernhauer
[Bernhauer 1934b: 6, Scheerpeltz 1957a: 461, = *fuscipes sinensis* Scheerpeltz]
- fuscipes goudoti* Laporte
[Laporte 1840: 183, species, Fagel 1958a: 313, ssp.]
- fuscipes lindbergi* Fagel
[Fagel 1958a: 313]
- fuscipes sinensis* Scheerpeltz
[Scheerpeltz 1957a: 461, replacement name for *fuscipes chinensis* Bernhauer]
- fuscus* Dahl
[Dahl 1823: 17, *nomen nudum*; this may be *Rugilus fuscus* Mannerheim 1831: 454]
- fuscus* Last, *Paederus*
[Last 1979: 137]
- G**
- gabonicus* Fagel, *Paederus*
[Fagel 1958a: 200, 1966: 263]
- gallomeridionalis* Scheerpeltz
[Scheerpeltz 1957a: 450, ssp. of *ruficollis* Fabricius, Coiffait 1982c: 22, var. of *ruficollis* Fabricius]
- garambanus* Fagel, *Pachypaederus*
[Fagel 1959: 35]
- gebieni* Wendeler, *Paederus*
[Wendeler 1931: 40, Scheerpeltz 1933: 1223, Fagel 1958a: 189, 1966: 263]
- gedyei* Cameron, *Paederus*
[Cameron 1950c: 184, Fagel 1958a: 417, 1966: 270]
- gedyei auct.*
[Bernhauer *in litt.*, Fagel 1958a: 417, = *gedyei* Cameron]
- gemellus* Kraatz
[Kraatz 1857: 731, Baudi 1857: 109, Miller 1868: 16, Gemminger & Harold 1868: 627, Fauvel 1869b: 491, = *sanguinicollis* Stephens, 1873: 336, = *ruficollis* Fabricius, Mulsant & Rey 1878: 249, species, Reitter 1889: 171, Ganglbauer 1895: 536, Bernhauer & Schubert 1912: 210, = *ruficollis* Fabricius, Scheerpeltz 1933: 1227, Jarrige 1944: 8, Hansen 1951: 223, Scheerpeltz 1957a: 449, 1968: 46, Ferreira 1962: 14, = var. of *ruficollis* Fabricius, Legosz-Owsianna 1963: 323, species, Lohse 1964: 133, = *ruficollis* Fabricius, Tikhomirova 1973: 175, Bohác 1985: 357]
- geniculatus* Dietrich
[Dietrich 1855: 201, Kraatz 1857: 727, = *brevipennis* Lacordaire, Gemminger & Harold 1868: 626, Mulsant & Rey 1878: 253, Ganglbauer 1895: 538, Bernhauer & Scheerpeltz 1912: 204, Scheerpeltz 1933: 1221, 1957a: 460, 1968: 47, Smetana 1962: 49, Legosz-Owsianna 1963: 339, Coiffait 1982c: 66]
- geniculatus* Peyron
[Peyron 1858: 430, homonym, Gemminger & Harold 1868: 627, = *littoralis* Gravenhorst, Fauvel 1869b: 491, 1873: 331, Ganglbauer 1895: 538, Scheerpeltz 1957a: 465, 1968: 48, Legosz-Owsianna 1963: 349, Coiffait 1982c: 58]
- Geopaederus* Gistel
[Gistel 1848: x, unnecessary replacement name for *Paederus*, type = *Paederus riparius* (Linnaeus), Blackwelder 1952: 169]
- germainei sic*
[Scheerpeltz 1933: 1223, misspelling of *germairei*, Blackwelder 1944: 123]
- germairei auct.*
[Fauvel *in litt.*, Wendeler 1931: 44, = *germairei* Wendeler]
- germairei* Wendeler, *Paederus*
[Wendeler 1931: 44]
- germanus* Cameron, *Paederus*
[Cameron 1931a: 51, Scheerpeltz 1933: 1223]
- gestroi* Fauvel, *Paederus*
[Fauvel 1878a: 237, 1879: 89, Bernhauer & Schubert 1912: 206, Last 1979: 138, 1985: 328]
- ghesquierei* Bernhauer
[Bernhauer 1933b: 293, Fagel 1958a: 120, = *albopubescentis* Bernhauer, 1959: 41]
- gigus* Bernhauer, *Paederus*
[Bernhauer 1905a: 17, Bernhauer & Schubert 1912: 206, Blackwelder 1944: 122]
- glaberrimus* (Herbst)
[Herbst 1784: 151, Paykull 1790: 136, 1800: 430, = *fracticornis* Paykull, now in *Acylophorus*]
- globulicollis* Bernhauer, *Paederus*
[Bernhauer 1907: 283, Blackwelder 1944: 123]
- Gnathopaederus* Chapin
[Chapin 1927: 75, genus, type *Gnathopaederus szechuanus* Chapin, Scheerpeltz 1933: 1219, subgenus, Blackwelder 1939: 118, 1952: 172]
- Gnathopaederus* Wendeler
[Wendeler 1927: 1, homonym, subgenus, type *Paederus turrialbanus* Wendeler, 1928a: 37, = *Paederognathus* Wendeler, replacement name, Blackwelder 1939: 118, 1952: 172]
- godweyi sic*
[Fagel 1958a: 364, misspelling of *gowdeyi*]
- goeldii auct.*
[Göldi 1913: 13, *nomen nudum*, credited by Göldi to

- Wasmann, but no description by Wasmann appears to exist]
- goliath auct.*
[Fauvel *in litt.*, Fagel 1958a: 79, = *delagoanus* Bernhauer]
- gotschei sic*
[Scheerpeltz 1957a: 472, misspelling of *gotschei*]
- gotschei* Kolbe, *Paederus* (*Gnathopaederus*)
[Kolbe 1886: 179, Bernhauer & Schubert 1912: 206, Chapin 1927: 76, Scheerpeltz 1933: 1223, Wu 1937: 329, Tikhomirova 1973: 175]
- goudoti* Laporte
[Laporte 1840: 183, Gemminger & Harold 1868: 627, Bernhauer & Schubert 1912: 206, Fagel 1958a: 313, ssp. of *fuscipes* Curtis]
- gowdeyi* Bernhauer, *Paederus*
[Bernhauer 1927a: 105, Scheerpeltz 1933: 1223, Fagel 1956: 198, 1958a: 364, 1960b: 203, 1966: 266]
- gracilior* Fagel, *Paederidus*
[Fagel 1958a: 148]
- gracilis* (Paykull)
[Paykull 1789: 38, 1800: 432, Fabricius 1792: 528, 1801: 600, Gyllenhal 1810: 375, = *Astenus*]
- gracilis* Last, *Paederus*
[Last 1952: 91, Fagel 1958a: 309, 1966: 258]
- gragorius sic*
[Ferreira 1962: 14, misspelling of *gregarius*, as var. of *riparius*]
- grandicollis* Bernhauer, *Paederus*
[Bernhauer 1927c: 401, Scheerpeltz 1933: 1223]
- grandipennis* Koch
[Koch 1937: 25, 1941: 56, Peyerimhoff 1949: 104, = *meridionalis* Fauvel, Scheerpeltz 1957a: 475, Coiffait 1982c: 70, var. of *meridionalis* Fauvel]
- grandipennis megacephalus* Koch
[Koch 1941: 56, ssp., Peyerimhoff 1949: 104, = *meridionalis* Fauvel, Scheerpeltz 1957a: 475, species, Coiffait 1982c: 70, var. of *meridionalis* Fauvel]
- grandis* Austin, *Paederus*
[Austin 1876: 8, LeConte 1878: 395, Casey 1905: 60, Bernhauer & Schubert 1912: 206, Leng 1920: 101, Blackwelder 1939: 112, Moore & Legner 1975: 133]
- gratiosus* Fauvel, *Paederus*
[Fauvel 1904a: 54, Bernhauer & Schubert 1912: 207, Cameron 1925b: 36, 1931a: 57, Scheerpeltz 1933: 1223]
- graueri* Fagel, *Paederus*
[Fagel 1958a: 383, 1960b: 204, 1966: 267]
- greeni* Cameron, *Paederus*
[Cameron 1931a: 53, Scheerpeltz 1933: 1223]
- gregarius* (Scopoli)
[Scopoli 1763: 102, Fabricius 1775: 268, = *riparius* (Linnaeus), Olivier 1795: 4, Erichson 1840: 654, Kraatz 1857: 728, Gemminger & Harold 1868: 628, Ganglbauer 1895: 537, Bernhauer & Schubert 1912: 210, Jarrige 1944: 8, Scheerpeltz 1957a: 462, 1968: 47, Ferreira 1962: 14, var. of *riparius*, Legosz-Owsianna 1963: 344, = *riparius*, Coiffait 1982c: 56]
- gregarius auct.*
[Fauvel 1873: 331, Sahlberg 1876: 37, = *littoralis* Gravenhorst, Mulsant & Rey 1878: 256, Ganglbauer 1895: 538, Bernhauer & Schubert 1912: 207, Scheerpeltz 1933: 1225, Jarrige 1944: 8, Scheerpeltz 1957a: 465, 1968: 48, Legosz-Owsianna 1963: 349, Coiffait 1982c: 58]
- gressitti* Last, *Paederus*
[Last 1985: 336]
- gridellii* Fagel, *Paederus*
[Fagel 1958a: 289, 1966: 261]

H

- haematoderus* Gemminger & Harold
[Gemminger & Harold 1868: 627, replacement name for *sanguinicollis* Motschulsky, Fauvel 1873: 335, = *sanguinicollis* Stephens, Bernhauer & Schubert 1912: 210, = *rubrothoracicus* Goeze, Scheerpeltz 1968: 46, Coiffait 1982c: 24]
- hagenensis* Last, *Paederus*
[Last 1979: 135, 1985: 329]
- hanangensis* Fagel
[Fagel 1960: 128, ssp. of *basipes* Fauvel]
- handschini* (Scheerpeltz), *Paederus*, *Diplopaederus*
[Scheerpeltz 1935: 598, 1957b: 272]
- Harpopaederus* Scheerpeltz
[Scheerpeltz 1957a: 459, subgenus, type *Paederus schoenherri* Czwalińska, Szujewski 1965b: 30, Coiffait 1982c: 63]
- hebridensis* Cameron, *Paederus*
[Cameron 1934: 21]
- heteroderus auct.*
[Eppelsheim *in litt.*, Fagel 1958a: 145, = *parcepunctatus* Bernhauer]
- heterogaster auct.*
[Fauvel *in litt.*, Fagel 1958a: 458, = *insularis* Cameron]
- Heteropaederus* Scheerpeltz
[Scheerpeltz 1957a: 460, subgenus, type *Paederus fuscipes* Curtis, Szujewski 1965b: 29, Shibata 1973: 43, 1977: 24, Coiffait 1982c: 60]

- heynei* Wendeler, *Paederus*
[Wendeler 1926b: 202, Scheerpeltz 1933: 1223, Blackwelder 1944: 123, Gaedike 1981: 194]
- himalaycus sic*
[Coiffait 1982b: 234, misspelling of *himalayicus*]
- himalayicus* Bernhauer, *Paederus*
[Bernhauer 1914: 98, replacement name for *indicus* Bernhauer, Cameron 1925b: 36, 1931a: 39, Scheerpeltz 1933: 1223, Blackwelder 1939: 112, Scheerpeltz 1957a: 474, Coiffait 1982b: 234]
- himalayei* *sic*
[Bernhauer 1939d: 115, misspelling of *himalayicus*]
- hinalayicus sic*
[Bernhauer 1939d: 115, misspelling of *himalayicus*]
- hingstoni* Cameron, *Paederus*
[Cameron 1928: 561, 1931a: 62, Scheerpeltz 1933: 1224, 1957a: 474]
- hintzi* Bernhauer, *Paederus*
[Bernhauer 1915c: 290, Scheerpeltz 1933: 1224, Fagel 1958a: 211, 1966: 267]
- hispanolusitanicus* Scheerpeltz
[Scheerpeltz 1957a: 450, *ssp.* of *ruficollis* Fabricius, Coiffait 1982c: 23, *var.* of *ruficollis* Fabricius]
- homonymus* Blackwelder
[Blackwelder 1943: 323, replacement name for *tricolor* Erichson, 1944: 123, unnecessary, junior synonym of *tricolor* Erichson]
- hornabrooki* Last, *Paederus*
[Last 1979: 133, 1984: 109, 1985a: 327]
- horni* Bernhauer, *Paederus*
[Bernhauer 1902: 38, Bernhauer & Schubert 1912: 207, Cameron 1925b: 36, 1931a: 52, Scheerpeltz 1933: 1224, Gaedike 1981: 195]
- huetheri* Bernhauer
[Bernhauer 1940b: 633, *var.* of *caligatus* Erichson, Scheerpeltz 1957a: 459, 1968: 47, Coiffait 1982c: 53, here treated as *ssp.* under Article 45(g) (ICZN 1985)]
- huonensis* Last, *Paederus*
[Last 1979: 128, 1985: 329]
- hutheri sic*
[Coiffait 1982c: 53, misspelling of *huetheri*]
- I**
- iavanus sic*
[Kraatz 1959a: 150, variant spelling of *javanus*]
- ibericus* Scheerpeltz
[Scheerpeltz 1957a: 453, *ssp.* of *rubrothoracicus* Goeze, Coiffait 1982c: 25, *var.* of *rubrothoracicus* Goeze]
- idae auct.*
[Lewis *in litt.*, Sharp 1874: 75, = *idae* Sharp, Bernhauer & Schubert 1912: 206]
- idae* Sharp
[Sharp 1874: 75, Fauvel 1879: 90, 1903a: 154, = *fuscipes* Curtis, 1904b: 290, Cameron 1925b: 35, 1931a: 40, Blackwelder 1939: 112, Shibata 1977: 24]
- idea sic*
[Scheerpeltz 1957a: 461, misspelling of *idae*]
- ignotus* Wendeler, *Paederus*
[Wendeler 1931: 45, Scheerpeltz 1933: 1224, Fagel 1958a: 353, 1966: 266]
- itheringi* Bernhauer, *Paederus*
[Bernhauer 1911b: 407, Bernhauer & Schubert 1912: 207, Blackwelder 1944: 123]
- ikelaensis* Fagel, *Paederus*
[Fagel 1961a: 281, 1966: 260]
- iliensis* Coiffait, *Paederus* (*Heteropaederus*)
[Coiffait 1970: 102, 1982c: 62, Tikhomirova 1973: 175]
- illesei sic*
[Last 1985: 329, misspelling of *illiesi*]
- illiesi* Puthz, *Paederus*
[Puthz 1969: 529, Last 1979: 138, 1985: 329]
- ilsae* Bernhauer, *Paederus* (*Dioncopaederus*)
[Bernhauer 1931d: 233, Scheerpeltz 1933: 1224, 1957a: 465, 1963: 8, Tikhomirova 1973: 175, Coiffait 1982c: 59]
- imitator auct.*
[Fauvel *in litt.*, Wendeler 1931: 43, = *imitator* Wendeler]
- imitator* Wendeler, *Paederus*
[Wendeler 1931: 43, Scheerpeltz 1933: 1224]
- immaculatus auct.*
[Kirby *ms.*, Stephens 1829: 287, = *Sunius*]
- immunis auct.*
[Kirby *ms.*, Stephens 1829: 288, = *Rugilus*]
- impressipennis* Fauvel, *Paederus*
[Fauvel 1898: 180, Bernhauer & Schubert 1912: 207, Fagel 1958a: 457, Jarrige 1970: 34]
- impressipennis* Bernhauer
[Bernhauer 1942b: 357, Fagel 1958a: 106, = *mirandus* Fagel]
- impressus* Rossi
[Rossi 1790: 254, Gravenhorst 1802: 69, = *Aleochara*]
- incertus* Fagel, *Paederus*
[Fagel 1958a: 390, 1965: 265]
- incognitus* Bernhauer
[Bernhauer 1915i: 315, Scheerpeltz 1933: 1224, Fagel 1958a: 234, = *junodi* Bernhauer, 1959: 48, 1960a: 122]

inconspicuus (Cameron), *Paederus*, *Paederidus*
[Cameron 1944: 710, Fagel 1958a: 119, 1959: 45]

indicus Motschulsky
[Motschulsky 1858: 634, Kraatz 1859b: xxxi, = *cyanocephalus* Erichson, Motschulsky 1860a: 73, Gemminger & Harold 1868: 626, Fauvel 1878a: 236, Bernhauer & Schubert 1912: 205, Cameron 1925b: 35, 1931a: 44]

indicus Bernhauer
[Bernhauer 1911a: 61, homonym, Bernhauer & Schubert 1912: 207, Bernhauer 1914: 98, = *himalayicus* Bernhauer, replacement name, Cameron 1925b: 36, 1931a: 39, Scheerpeltz 1933: 1224]

inexpectatus Fagel, *Paederus*
[Fagel 1958a: 394, 1966: 266]

insidiosus Fagel, *Oreopaederus*
[Fagel 1973: 150]

insolitus Fagel
[Fagel 1960a: 120, ssp. of *nakurenensis* Fauvel]

insulanus auct.
[Bernhauer *in litt.*, Fagel 1958a: 319, = *angusticeps* Bernhauer]

insularis Cameron, *Paederus*
[Cameron 1938: 174, Fagel 1958a: 458]

intermedius Boheman, *Paederus*
[Boheman 1858: 32, Gemminger & Harold 1868: 627, Baer 1886: 107, Fauvel 1886b: 147, Bernhauer & Schubert 1912: 207, Bernhauer 1927c: 399, Scheerpeltz 1933: 1224, Blackwelder 1939: 112, Scheerpeltz 1935: 608, Cameron 1936: 43]

intolerans Fagel, *Oreopaederus*
[Fagel 1964: 228]

intricatus auct.
[Dohrn *in litt.*, Gemminger & Harold 1868: 628, = *peregrinus* Erichson]

iowaensis sic
[Moore & Legner 1975: 133, misspelling of *iowensis*]

iowensis (Casey), *Paederillus*, *Paederus*
[Casey 1905: 66, Leng 1920: 101, Blackwelder 1939: 112]

irebuanus Bernhauer
[Bernhauer 1927a: 118, Scheerpeltz 1933: 1224, Fagel 1958a: 120, = *albopubescens* Bernhauer, 1959: 41]

irianensis Frank, *Paederus*
[here, *nom. nov.* for *Paederus litoreus* Last, *nec Paederus littoreus* Austin]

irritans Chapin
[Chapin 1926: 370, Campos 1931: 24, = *ornaticornis* Sharp, Scheerpeltz 1933: 1224, Blackwelder 1939: 112, 1944: 123, Gaedike 1981: 198]

ismayi Last, *Paederus*
[Last 1985: 336]

itombwensis Fagel, *Paederus*
[Fagel 1961a: 284, 1966: 265]

J

jacobsoni Bernhauer, *Paederus*
[Bernhauer 1915e: 222, Scheerpeltz 1933: 1224, 1935: 629, Cameron 1936: 42, Last 1984: 109]

jaegeri Paulian, *Paederus*
[Paulian 1947: 28, Fagel 1958a: 205, 1966: 262]

jamaicensis Blackwelder, *Paederus*
[Blackwelder 1943: 324, 1944: 123, Sanderson 1967: 60]

jansei Fagel, *Paederus*
[Fagel 1959: 54, 1966: 258]

janssensi Fagel, *Paederus*
[Fagel 1958a: 166, 1966: 272]

japonicus Bernhauer, *Paederus* (*Paederus*)
[Bernhauer 1915h: 265, 1923: 180, Scheerpeltz 1933: 1224, 1957a: 462, Tikhomirova 1973: 175, Shibata 1977: 26]

jarrigeanus Fagel, *Paederus*
[Fagel 1958a: 199, 1966: 263]

jarrigei (Fagel), *Paederus*, *Oreopaederus*
[Fagel 1954a: 948, 1958a: 102]

javanus Laporte, *Paederus*
[Laporte 1835: 123, Gemminger & Harold 1868: 627, Bernhauer & Schubert 1912: 207, Scheerpeltz 1935: 633]

javanus Erichson
[Erichson 1840: 654, homonym, Kraatz 1859a: 150, Fauvel 1895a: 232, = *sondaicus* Fauvel, replacement name, 1904a: 53, Bernhauer & Schubert 1912: 211, Wendeler 1926d: 328, Cameron 1925b: 37, 1930b: 332, 1931a: 37, Scheerpeltz 1935: 609, 1957a: 462, Biswas & Sen Gupta 1982: 153]

jaegeri sic
[Paulian 1947: 28, misspelling of *jaegeri*]

jeanneli Fagel
[Fagel 1958a: 421, ssp. of *melanogaster* Fauvel, 1966: 268]

jessoensis Bernhauer, *Paederus*
[Bernhauer 1935a: 44, Scheerpeltz 1957a: 475, Tikhomirova 1973: 175, Shibata 1977: 28]

jowensis sic
[Bernhauer & Schubert 1912: 207, misspelling of *iowensis*]

junodi Bernhauer, *Paederus*
[Bernhauer 1912b: 181, Scheerpeltz 1933: 1224, Fagel 1958a: 234, 1959: 48, 1960a: 122, 1966: 258]

junodi auct.
[Fauvel *in litt.*, Fagel 1958a: 234, = *junodi* Bernhauer]

K

kabarensis Fagel, *Paederus*
[Fagel 1958a: 377, 1966: 266]

kaboboensis Fagel, *Pachypaederus*
[Fagel 1960b: 191]

kahuziensis (Fagel), *Paederus*, *Oreopaederus*
[Fagel 1954a: 915, 1958a: 101, 1960b: 195, Gaedike 1981: 198]

kaindiensis Last, *Paederus*
[Last 1979: 127]

kaiseri Bernhauer, *Paederus*
[Bernhauer 1934b: 6, Scheerpeltz 1957a: 474]

kalodanus Cameron, *Paederus*
[Cameron 1937: 96]

kalalovae Roubal
[Roubal 1932: 60, Scheerpeltz 1933: 1224, 1957a: 461, Coiffait 1982c: 62, Bohac 1985: 371, = *fuscipes* Curtis]

kambaitensis Scheerpeltz, *Paederus*
[Scheerpeltz 1965b: 174]

kamonoensis Last, *Paederus*
[Last 1979: 123, 1985: 328]

kampalanus Bernhauer
[Bernhauer 1927a: 114, Scheerpeltz 1933: 1224, Fagel 1958a: 326, 1959: 64, = *pretiosus* Bernhauer]

kaszabi Fagel, *Paederus*
[Fagel 1958a: 223, 1966: 261]

kateshensis Fagel
[Fagel 1960a: 121, ssp. of *densiventris* Bernhauer]

katubuensis Last, *Paederus*
[Last 1979: 128]

katumbuensis sic
[Last 1985: 327, misspelling of *katubuensis*]

kawaensis auct.
[Cameron *in litt.*, Fagel 1958a: 326, = *kampalanus* Bernhauer]

kenyanus Bernhauer
[Bernhauer 1940a: 135, Fagel 1958a: 179, = *nakurensis* Fauvel, 1960a: 120]

kenyensis Fagel
[Fagel 1958a: 145, ssp. of *erythraeanus* Fagel]

kilimandjarensis Fagel, *Paederus*
[Fagel 1958a: 343, 1960a: 122, 1966: 269]

kivuanus Fagel, *Paederus*
[Fagel 1956: 197, 1958a: 383, 1966: 265]

kivuensis Cameron
[Cameron 1950d: 93, Fagel 1958a: 364, = *gowdeyi* Bernhauer, 1960b: 204]

kiybiensis Fagel, *Oreopaederus*
[Fagel 1960b: 199]

klapperichi Bernhauer, *Paederus*
[Bernhauer 1941: 226, Scheerpeltz 1957a: 475]

klugi sic
[Gemminger & Harold 1868: 627, misspelling of *klugii*]

klugii Laporte
[Laporte 1840: 183, Gemminger & Harold 1868: 627, Bernhauer & Schubert 1912: 207, Fagel 1958a: 457, = *chalybeus* Erichson]

kochi Fagel, *Paederus*
[Fagel 1958a: 369, 1966: 268]

koebelei Blackburn, *Paederus*
[Blackburn 1899: 23, Bernhauer & Schubert 1912: 207, Lea 1923: 14]

koebeli sic
[Bernhauer 1920c: 9, misspelling of *koebelei*]

korbi Bernhauer
[Bernhauer 1935a: 44, ssp. of *riparius* Linnaeus, Scheerpeltz 1957a: 462, var. of *riparius* Linnaeus, Tikhomirova 1973: 175]

koreanus Scheerpeltz, *Paederidus*
[Scheerpeltz 1957a: 451]

korpi sic
[Scheerpeltz 1957a: 462, misspelling of *korbi*]

kosempoensis Bernhauer, *Paederus*
[Bernhauer 1922: 228, Scheerpeltz 1933: 1224, Adachi 1939: 166, Cameron 1949b: 466, Shibata 1973: 45]

kosempoensis fulvocaudatus Adachi
[Adachi 1939: 166, described as a variety but here treated as a subspecies under Article 45(g) (ICZN 1985)]

kozuensis Cameron, *Paederus*
[Cameron 1930a: 206, Scheerpeltz 1933: 1224, 1957a: 474, Tikhomirova 1973: 175, Shibata 1977: 28]

kraepelini Fauvel, *Paederus*
[Fauvel 1905a: 82, Bernhauer & Schubert 1912: 207, Scheerpeltz 1935: 622]

kuborensis Last, *Paederus*
[Last 1985: 330]

kuluensis Bernhauer, *Paederus* (*Eopaederus*)
[Bernhauer 1914: 99, Cameron 1925b: 36, 1931a: 55,

- Scheerpeltz 1933: 1224, 1957a: 474, Coiffait 1978b: 553, 1982c: 51, Tronquet 1981: 70]
kundelungensis Fagel, *Paederus*
 [Fagel 1958a: 163, 1966: 271]
kuntzeni Wendeler, *Paederus*
 [Wendeler 1926b: 201, Scheerpeltz 1933: 1224, 1935: 619]
kurosawai Watanabe, *Megalopaederus*
 [Watanabe 1986: 167]
- L**
- lacordairei* Perroud, *Paederus* (*Anomalopaederus*)
 [Perroud 1864: 88, Gemminger & Harold 1868: 627, Fauvel 1874: 433, 1889b: 258, 1903b: 265, Bernhauer & Schubert 1912: 207, Scheerpeltz 1966: 400]
laensis Last, *Paederus*
 [Last 1979: 136, 1985: 328]
laeticollis auct.
 [Fauvel *in litt.*, Fagel 1958a: 114, = *colonus* Wendeler]
laetipes Sharp, *Paederus*
 [Sharp 1886: 611, Bernhauer & Schubert 1912: 207, Blackwelder 1939: 112, 1944: 123]
laetus Erichson, *Paederus*
 [Erichson 1840: 660, Gemminger & Harold 1868: 627, Sharp 1886: 613, Bernhauer & Schubert 1912: 207, Blackwelder 1939: 112, 1944: 123, Scheerpeltz 1965: 215]
laevigatus Gyllenhal
 [Gyllenhal 1827: 483, Mannerheim 1831: 454, = *Rugilus*, now in *Scopaeus*]
lamottei Fagel, *Paederus*
 [Fagel 1963: 345, 1966: 257]
lasti Fagel, *Paederus*
 [Fagel 1958a: 292, 1966: 268]
lasti Fagel, *Pachypaederus*
 [Fagel 1961a: 276]
laticeps Fagel, *Paederus*
 [Fagel 1958a: 388, 1966: 267]
latipes auct.
 [Knoch *in coll.*, Gravenhorst 1802: 202, = *Pinophilus*]
lativentris Wendeler, *Paederus*
 [Wendeler 1926b: 200, Bernhauer 1927c: 400, Scheerpeltz 1933: 1224, Blackwelder 1939: 112, Gaedike 1981: 201]
lativentris decipiens Wendeler
 [Wendeler 1926b: 201]
latro Smetana, *Paederus* (*Harpopaederus*)
 [Smetana 1962: 46, Tikhomirova 1973: 175, = *us-suriensis* Kirschenblatt, Coiffait 1982c: 65, species]
latus auct.
 [Fagel *in litt.*, Fagel 1958a: 77, = *wendeleri* Fagel]
ledouxi Coiffait, *Paederus* (*Eopaederus*)
 [Coiffait 1978b: 561, 1982c: 52]
leleupi (Fagel), *Paederus*, *Oreopaederus*
 [Fagel 1954a: 927, 1956: 186, 1958a: 101]
leleupi Fagel, *Pachypaederus*
 [Fagel 1965: 146]
leleupi Fagel
 [Fagel 1958a: 135, ssp. of *montanellus* Bernhauer, 1960b: 202]
leleupianus Fagel, *Oreopaederus*
 [Fagel 1973: 148]
lemeranus Cameron
 [Cameron 1950b: 26, Fagel 1958a: 170, = *opucicollis* Bernhauer, 1960a: 119]
lenkoranus Scheerpeltz, *Paederus* (*Dioncopaederus*)
 [Scheerpeltz 1957a: 465, Tikhomirova 1973: 175, Coiffait 1982c: 59]
Leucopaederus Casey
 [Casey 1905: 59, *genus*, Bernhauer & Schubert 1912: 203, = *Paederus*, Blackwelder 1939: 119, type *Paederus ustus* LeConte]
leviter Wendeler
 [Wendeler 1933: 56, Fagel 1958a: 159, = *minimus* Bernhauer, 1959: 46]
lewisi (Cameron), *Paederus*, *Megalopaederus*
 [Cameron 1930a: 206, Scheerpeltz 1933: 1224, 1957a: 471, Tikhomirova 1973: 175, Shibata 1977: 29, Watanabe 1986: 162]
licenti Bernhauer, *Paederus*
 [Bernhauer 1938a: 51, Scheerpeltz 1957a: 475]
liepolti auct.
 [Bernhauer *in coll.*, Gaedike 1981: 230]
limnophilus Maerkel
 [limnophilus (or limophilus?) Maerkel unpubl., Erichson 1840: 653, = *limnophilus* Erichson]
limnophilus Erichson
 [Erichson 1840: 653, Heer 1841: 579, = *limophilus* Heer, Fairmaire & Laboulbène 1856: 571, Baudi 1857: 108, Kraatz 1857: 729, Gemminger & Harold 1868: 627, Fauvel 1864: 12, 1866: 25, 1873: 334, Mulsant & Rey 1878: 266, Reitter 1889: 170, Czwalina 1889: 368, Ganglbauer 1895: 537, Bernhauer & Schubert 1912: 207, Kirschenblatt 1932: 217, Jarrige 1944: 9, Scheerpeltz 1933: 1224, 1957a: 459, 1968: 47, Legosz-Owsianna 1963: 1224, Lohse 1964: 135, Szujecki 1965b: 29, Tikhomirova 1973: 175, Coiffait 1982c: 53, Bohác 1985: 372]

- limnophilus auct.*
[Chevrolat 1871: 136, misidentification, Fauvel 1873: 334, = *caligatus* Erichson]
- limophilus* Maerkel
[limophilus (or limnophilus?) Maerkel unpubl., Heer 1839: 235, = *limophilus* Heer]
- limophilus* Heer, *Paederus* (*Eopaederus*)
[Heer 1839: 235, 1841: 579, the original publication contains no admission of a misspelling, so under Article 32 (ICZN 1985) this name is valid and *limnophilus* Erichson is a synonym]
- lindbergi* Fagel
[Fagel 1958a: 313, ssp. of *fuscipes* Curtis, 1966: 262]
- linearis auct.*
[Hellwig in coll., Gravenhorst 1802: 55, = *Lathrobium*]
- lineiventris* Bernhauer, *Paederus*
[Bernhauer 1927a: 113, Scheerpeltz 1933: 1225, Fagel 1958a: 259, 1966: 259]
- lingualis* Sharp, *Paederus*
[Sharp 1876a: 286, Bernhauer & Schubert 1912: 207, Blackwelder 1944: 123]
- listeri* Gahan, *Paederus*
[Gahan 1900: 89, Bernhauer & Schubert 1912: 207]
- litorales sic*
[Ferreira 1962: 14, misspelling of *littoralis*]
- littoralis sic*
[Ganglbauer 1895: 538, misspelling of *littoralis*, Bernhauer & Schubert 1912: 207, Kirschenblatt 1932: 217, Scheerpeltz 1933: 1225, Hansen 1951: 224, Scheerpeltz 1957a: 465, 1968: 47, Smetana 1959: 204, Legosz-Owsianna 1963: 349, Lohse 1964: 134, Szujecki 1965: 30, Jarrige 1971: 491, Tikhomirova 1973: 175, Hammond 1977: 27, Bohác 1985: 373]
- litoreus* Last
[Last 1985: 331, homonym under Article 58(8) (ICZN 1985), here = *irianensis* Frank, *nom. nov.*]
- littoralis auct.*
[Knoch in coll., Gravenhorst 1802: 61, = *littoralis* Gravenhorst]
- littoralis* Gravenhorst, *Paederus* (*Dioncopaederus*)
[Gravenhorst 1802: 61, 1806: 143, Illiger 1807: 356, Dahl 1823: 17, Stephens 1829: 288, 1832: 279, Mannerheim 1831: 453, Dejean 1833: 65, Lacordaire 1835: 429, Faldermann 1839: 98, Heer 1839: 234, Laporte 1840: 183, Baudi 1857: 108, Kraatz 1857: 726, Thomson 1860: 196, Fauvel 1866: 25, Gemminger & Harold 1868: 627, Baudi 1869: 394, Mulsant & Rey 1878: 256, Reitter 1885: 154, 1889: 169, Czwalina 1889: 368, Eppelsheim 1892: 340, Bernhauer & Schubert 1912: 208, Scheerpeltz 1957a: 465, 1968: 47, Lohse 1964: 134, Hammond 1977: 27, Muona 1979: 16, Coiffait 1982c: 58]
- littoralis auct.*
[Gravenhorst 1802: 61 *partim*, Gemminger & Harold 1868: 626, = *cephalotes* Motschulsky]
- littoralis auct.*
[Olivier 1812: 627, misidentification, Erichson 1840: 656, = *littorarius* Gravenhorst, Bernhauer & Schubert 1912: 208]
- littoralis auct.*
[Lucas 1849: 121, misidentification, Fauvel 1886a: 32, = *meridionalis* Fauvel, 1897: 281]
- littorarius auct.*
[Knoch in coll., Gravenhorst 1806: 142, = *littorarius* Gravenhorst]
- littorarius* Gravenhorst, *Paederus*, *Paederillus*, *Paederus*
[Gravenhorst 1806: 142, Dejean 1833: 65, Erichson 1840: 656, Gemminger & Harold 1868: 627, Austin 1876: 11, LeConte 1878: 396, Casey 1905: 66, Bernhauer & Schubert 1912: 208, Leng 1920: 101, Cameron 1931a: 33, Scheerpeltz 1933: 1225, Blackwelder 1939: 112, Hatch 1957: 170, Moore & Legner 1975: 134]
- littoreus auct.*
[Zimmermann *ms.*, Austin 1876: 9, = *littoreus* Austin]
- littoreus* Austin, *Paederus*
[Austin 1876: 9, LeConte 1878: 396, Casey 1905: 61, Bernhauer & Schubert 1912: 208, Leng 1920: 101, Blackwelder 1939: 112, Moore & Legner 1975: 134]
- lividipennis* Bernhauer
[Bernhauer 1927c: 399, var. of *semiviridis* Bernhauer, here treated as ssp. under Article 45(g) (ICZN 1985)]
- lividus* Bernhauer, *Paederus*
[Bernhauer 1927c: 401, Scheerpeltz 1933: 1225]
- Lobopaederus* Scheerpeltz
[Scheerpeltz 1957a: 469, *genus*, type *Paederus meridionalis* Fauvel, Coiffait 1982c: 70]
- loebli* Biswas & Sen Gupta, *Paederus* (*Pseudopaederus*)
[Biswas & Sen Gupta 1982: 148]
- lombockianus* Cameron, *Paederus*
[Cameron 1925a: 181]
- lombokianus sic*
[Scheerpeltz 1933: 1225, misspelling of *lombockianus*, 1935: 611]
- longiceps* Bernhauer, *Paederus* (*Paederus*)
[Bernhauer 1901: 247, Bernhauer & Schubert 1912: 208, Roubal 1932: 61, Kirschenblatt 1932: 218, Scheerpeltz 1957: 459, Tikhomirova 1973: 175, Coiffait 1982c: 54]

longicollis Gautier

[Gautier 1861a: 393, Saulcy 1862: vi, Fauvel 1864: 13, = *riparius* Linnaeus, Marseul 1866: vi, Gautier 1866: 114, = *ruficeps* Baudi, Gemminger & Harold 1868: 627, = *melanurus* Aragona, Fauvel 1869b: 491, = *riparius* Linnaeus, Ganglbauer 1895: 537, Bernhauer & Schubert 1912: 210, Jarrige 1944: 8, Scheerpeltz 1957a: 462, 1968: 47, Legosz-Owsianna 1963: 335, Coiffait 1982c: 56]

longicornis Aube

[Aube 1850: 319, Fairmaire & Laboulbène 1856: 571, Baudi 1857: 109, Kraatz 1857: 730, Gemminger & Harold 1868: 626, = *sanguinicollis* Stephens, Fauvel 1866: 25, 1873: 335, Chevrolat 1871: 136, Mulsant & Rey 1878: 246, = *ruficollis* Fabricius, Reitter 1889: 171, Ganglbauer 1895: 536, = *sanguinicollis* Stephens, Bernhauer & Schubert 1912: 210, = *rubrothoracicus* Goeze, Scheerpeltz 1933: 1227, Jarrige 1944: 7, Scheerpeltz 1957a: 452, 1968: 46, Legosz-Owsianna 1963: 327, Coiffait 1982c: 25, var. of *rubrothoracicus* Goeze]

longicornis Motschulsky

[Motschulsky 1858: 634, homonym, Gemminger & Harold 1868: 627, Fauvel 1904b: 290, = *sabaeus* Erichson, 1907: 24, Bernhauer & Schubert 1912: 210, Scheerpeltz 1957a: 461, species, Coiffait 1982c: 61, = *sabaeus* Erichson]

longipennis Dahl

[Dahl 1823: 17, *nomen nudum*, Dejean 1833: 65, = *riparius* Linnaeus]

longipennis Erichson

[Erichson 1839: 517, 1840: 651, Heer 1839: 234, Fairmaire & Laboulbène 1856: 570, Baudi 1857: 108, Kraatz 1857: 728, = *fuscipes* Curtis, Thomson 1860: 196, Gemminger & Harold 1868: 627, Solsky 1871: 163, Fauvel 1873: 334, Mulsant & Rey 1878: 261, Reitter 1889: 170, Czwalina 1889: 368, Ganglbauer 1895: 537, Bernhauer & Schubert 1912: 206, Cameron 1925b: 36, 1931a: 40, Scheerpeltz 1933: 1223, Blackwelder 1939: 112, Jarrige 1944: 9, Scheerpeltz 1957a: 461, 1968: 47, Legosz-Owsianna 1963: 335, Coiffait 1982c: 60]

longiusculus auct.

[Harris unpublished, Mannerheim 1831: 453, = *longiusculus* Mannerheim]

longiusculus Mannerheim

[Mannerheim 1831: 453, Gemminger & Harold 1868: 625, = *Sunius*, now in *Astenus*]

longulus auct.

[Knoch *in coll.*, Gravenhorst 1802: 54, = *Lathrobium*]

lootensi Fagel, *Paederus*

[Fagel 1961a: 278, 1966: 260]

luberensis Fagel, *Paederus*

[Fagel 1954a: 921]

lubukensis (Fagel), *Paederus*, *Oreopaederus*

[Fagel 1954a: 931, 1958a: 101]

luctuosus (Klug), *Paederus*, *Paederidus*

[Klug 1855: 644, Gemminger & Harold 1868: 627, Bernhauer & Schubert 1912: 208, Bernhauer 1915: 315, Fagel 1956: 200, 1958a: 129, 1960a: 117, 1961b: 266, Coiffait 1982c: 21]

luffianus Fagel, *Paederus*

[Fagel 1967: 176]

lugubris auct.

[Erichson *in litt.*, Fagel 1958a: 129, = *luctuosus* Klug]

lugubris Motschulsky

[Motschulsky 1858: 635, Gemminger & Harold 1868: 628, = *rutilicornis* Erichson, Blackwelder 1944: 123]

luluanus auct.

[Bernhauer *in litt.*, Fagel 1958a: 269, = *xanthocerus* Eppelsheim]

luluensis (Bernhauer), *Paederus*, *Paederidus*

[Bernhauer 1937b: 299, Fagel 1958a: 123]

luluensis auct.

[Bernhauer *in litt. partim*, Fagel 1958a: 314, = *malaisei* Fagel]

luridiventris Sharp, *Paederus*

[Sharp 1886: 610, Bernhauer & Schubert 1912: 208, Blackwelder 1944: 123]

lusitanicus (Aube), *Paederus*, *Parameropaederus*

[Aubé 1842: 236, Kraatz 1857: 725, Fairmaire 1859: clxxxv, Gemminger & Harold 1868: 627, Czwalina 1889: 368, Bernhauer & Schubert 1912: 208, Scheerpeltz 1933: 1225, 1957: 468, Ferreira 1962: 12, Coiffait 1982c: 68]

lusitanicus Baudi

[Baudi 1857: 107, homonym, Fairmaire 1859: clxxxv, = *baudii* Fairmaire, replacement name, Gemminger & Harold 1868: 626, Fauvel 1873: 329, Mulsant & Rey 1878: 250, Reitter 1884: 44, 1889: 170, species, Ganglbauer 1895: 538, = *baudii* Fairmaire, Bernhauer & Schubert 1912: 204, Scheerpeltz 1957a: 460, Smetana 1962: 49]

luvubuensis Fagel

[Fagel 1954a: 914, 1958a: 101, ssp. of *ater* Bernhauer, here transferred to ssp. of *manyemensis* Frank]

M

- maai* Last, *Paederus*
[Last 1985: 332]
- macellus* Fauvel, *Paederus*
[Fauvel 1907: 22, Bernhauer & Schubert 1912: 208, Fagel 1958a: 355, 1960a: 123, 1966: 270]
- macellus macellus* Fauvel
[Fauvel 1907: 22, Fagel 1960a: 123]
- macellus rufobrunneus* Cameron
[Cameron 1950: 195, species, Fagel 1958a: 358, ssp.]
- macellus auct.*
[Fauvel 1907: 22, *partim*, Fagel 1958a: 362, = *meruensis* Fagel]
- machadoanus* Fagel, *Paederus*
[Fagel 1973: 154]
- machadoi* Scheerpeltz, *Paederus*
[Scheerpeltz 1965a: 223]
- machadoi* Fagel, *Oreopaederus*
[Fagel 1962b: 22]
- machadoi* Fagel
[Fagel 1961a: 265, *nomen nudum*]
- madagascariensis* Erichson, *Paederus*
[Erichson 1840: 652, Boheman 1848: 290, Gemminger & Harold 1868: 627, Bernhauer & Schubert 1912: 208, Fagel 1958a: 177]
- Madecapaederus* Fagel
[Fagel 1958a: 103, genus, type *Paederus metallicus* Fauvel]
- madudanus auct.*
[Cameron *in litt.*, Fagel 1958a: 204, = *madudanus* Fagel]
- madudanus* Fagel, *Paederus*
[Fagel 1958a: 204, 1966: 262]
- mafuluensis* Last, *Paederus*
[Last 1979: 137]
- magniceps* Bernhauer, *Paederus*
[Bernhauer 1905b: 181, Bernhauer & Schubert 1912: 208, Blackwelder 1944: 123, Scheerpeltz 1960: 81]
- magnificus* Bernhauer, *Paederus*
[Bernhauer 1912c: 474, Bernhauer & Schubert 1912: 208, Fagel 1958a: 405, 1960a: 125, 1966: 265]
- magnus* Fagel, *Paederus*
[Fagel 1958a: 400, 1966: 266]
- malaisei* Fagel, *Paederus*
[Fagel 1958a: 314, 1959: 62, 1966: 260]
- mandibularis* Erichson, *Paederus*
[Erichson 1840: 659, Lucas 1851: 51, Gemminger & Harold 1868: 627, Fauvel 1877b: xxvi, Lynch 1884: 292, Bernhauer & Schubert 1912: 208, Blackwelder 1944: 123]
- mangolae* Bernhauer, *Paederus*
[Bernhauer 1937a: 295, Fagel 1958a: 455, 1959: 65, 1966: 272]
- manyemensis* Frank, *Oreopaederus*
[here, = *nom. nov.* for *Paederus ater* Bernhauer 1927: 107 = *Oreopaederus ater* (Bernhauer), *nec Paederus ruficollis var. ater* Eichler = *Paederidus rubrothoracicus ater* (Eichler)]
- manyemensis luvubuensis* (Fagel), *Oreopaederus*
[here, new combination for *Oreopaederus ater luvubuensis* (Fagel)]
- maputoanus auct.*
[Scheerpeltz *in litt.*, Fagel 1958a: 234, = *junodi* Bernhauer]
- marcuzzii* Scheerpeltz, *Paederus*
[Scheerpeltz 1965a: 219, Marcuzzi 1977: 7]
- margelanicus* Bernhauer, *Paederus* (*Harpopaederus*)
[Bernhauer 1903: 187, Bernhauer & Schubert 1912: 208, Scheerpeltz 1957a: 460, Smetana 1962: 49, Tikhomirova 1973: 175, Coiffait 1982c: 66]
- margellanicus sic*
[Kirschenblatt 1932: 217, misspelling of *margelanicus*]
- mariepskopensis* Fagel, *Paederus*
[Fagel 1965: 151, 1966: 258]
- marquesanus auct.*
[Scheerpeltz *in litt.*, Fagel 1958a: 129, = *luctuosus* Klug]
- marraensis* Fagel
[Fagel 1968: 197, ssp. of *aquaticus* Cameron]
- marshalli* Bernhauer, *Paederus*
[Bernhauer 1937a: 293, Fagel 1958a: 437, 1966: 264]
- martensi* Coiffait, *Paederidus*
[Coiffait 1982a: 85, 1982b: 234]
- mateui* Fagel, *Paederus*
[Fagel 1958a: 193, 1966: 263]
- matumbiensis* Bernhauer
[Bernhauer 1927a: 110, Scheerpeltz 1933: 1225, Fagel 1958a: 248, = *capitalis* Bernhauer]
- mauritii auct.*
[Bernhauer *in litt.*, Bernhauer 1927a: 104, = *edwardsi* Bernhauer]
- mauritti sic*
[Fagel 1958a: 458, misspelling of *maurittii*]
- maxillosus* Lentz
[Lentz 1856: 57, here, possibly = *Cucujus maxillosus* Fabricius = *Leptochirus maxillosus* (Fabricius) (Staphylinidae)]

- maynei* Bernhauer, *Paederus*
[Bernhauer 1927a: 110, Scheerpeltz 1933: 1225, Fagel 1958a: 263, 1965: 156, 1966: 261]
- mayumbeanus* Cameron
[Cameron 1939: 4, Fagel 1958a: 311, = *fuscipes* Curtis, 1959: 62, 1960a: 121, 1961b: 268]
- meersmanae* Fagel, *Oreopaederus*
[Fagel 1961a: 263]
- meersmanae* Fagel
[Fagel 1965: 144, ssp. of *crassus* Boheman]
- megacephalus auct.*
[Knoch unpublished, Dahl 1823: 17, *nomen nudum*, Dejean 1833: 65, = *Rugilus*]
- megacephalus* Koch
[Koch 1941: 56, ssp. of *grandipennis* Koch, Scheerpeltz 1957a: 475, species, Coiffait 1982c: 70, var. of *meridionalis* Fauvel]
- megalocephalus sic*
[Coiffait 1982c: 70, misspelling of *megacephalus*]
- Megalopaederus* Scheerpeltz
[Scheerpeltz 1957a: 470, genus, type *Paederus poweri* Sharp, Shibata 1973: 44, 1977: 28, Watanabe 1986: 159]
- melampus* Erichson, *Paederus*
[Erichson 1840: 660, Kraatz 1859a: 153, Gemminger & Harold 1868: 627, Bernhauer & Schubert 1912: 208, Cameron 1925b: 36, 1931a: 43, Scheerpeltz 1933: 1225, Biswas & Sen Gupta 1982: 153]
- melanocephalus* Fabricius
[Fabricius 1792: 538, 1801: 610, Panzer 1796b: 22, Gravenhorst 1806: 138, Stephens 1829: 287, = *Sunius*]
- melanocephalus* Heller
[Heller 1889: 4, homonym, Bernhauer & Schubert 1912: 208, here, = *Paederus sulawesi* Frank, *nom. nov.*]
- melanogaster* Fauvel, *Paederus*
[Fauvel 1907: 21, Bernhauer & Schubert 1912: 208, Bernhauer 1940a: 135, Fagel 1958a: 419, 1960a: 127, 1966: 267]
- melanogaster alluaudi* Fagel
[Fagel 1958a: 421, 1966: 268]
- melanogaster janneli* Fagel
[Fagel 1958a: 421, 1966: 268]
- melanogaster porrectus* Wendeler
[Wendeler 1933: 56, species, Fagel 1958a: 422, ssp., 1966: 267]
- melanopus* Wendeler, *Paederus*
[Wendeler 1942: 102]
- melanurus* Aragona, *Paederus* (*Paederus*)
[Aragona 1830: 13, Dejean 1833: 65, Erichson 1840: 932, Kraatz 1857: 730, Gemminger & Harold 1868: 627, Reitter 1889: 171, Czwalina 1889: 368, Ganglbauer 1895: 537, Bernhauer & Schubert 1912: 208, Scheerpeltz 1933: 1225, Blackwelder 1939: 112, Scheerpeltz 1957a: 462, Coiffait 1982c: 57]
- memnomius sic*
[Coiffait 1982c: 21, misspelling of *memnonius*]
- memnonius* (Erichson), *Paederus*, *Paederidus*
[Erichson 1840: 664, Gemminger & Harold 1868: 628, Fauvel 1886a: 324, Bernhauer & Schubert 1912: 208, Scheerpeltz 1933: 1225, Koch 1934: 80, Blackwelder 1939: 112, Fagel 1958a: 127]
- mendax* (Fagel), *Paederus*, *Oreopaederus*
[Fagel 1954a: 960, 1956: 188, 1958a: 102]
- meridiogallicus* Scheerpeltz
[Scheerpeltz 1957a: 453, ssp. of *rubrothoracicus* Goeze, Coiffait 1982c: 25, var. of *rubrothoracicus* Goeze]
- meridioitalicus* Scheerpeltz
[Scheerpeltz 1957a: 454, ssp. of *rubrothoracicus* Goeze, Coiffait 1982c: 25, var. of *rubrothoracicus* Goeze]
- meridionalis auct.*
[Kraatz *in litt.*, Fauvel 1873: 331, = *meridionalis* Fauvel]
- meridionalis* (Fauvel), *Paederus*, *Lobopaederus*
[Fauvel 1873: 331, 1886a: 32, 1897: 281, Reitter 1885: 154, 1889: 169, Czwalina 1889: 368, Ganglbauer 1895: 538, Bernhauer & Schubert 1912: 208, Bernhauer 1931a: 14, Kirschenblatt 1932: 217, Koch 1937: 25, Jarige 1944: 8, Scheerpeltz 1957a: 470, Tikhomirova 1973: 175, Coiffait 1982c: 70]
- meruensis* Fagel
[Fagel 1958a: 362, species, 1960a: 125, ssp. of *superbus* Bernhauer, 1966: 270]
- mesopotamicus* Eppelsheim, *Paederus* (*Fopaederus*)
[Eppelsheim 1889a: 178, Bernhauer & Schubert 1912: 208, Roubal 1932: 61, Scheerpeltz 1957a: 459, Coiffait 1982c: 51]
- metallicus* (Fauvel), *Paederus*, *Madecapaederus*
[Fauvel 1905b: 151, Bernhauer & Schubert 1912: 208, Fagel 1958a: 103]
- methneri* Bernhauer, *Paederus*
[Bernhauer 1912c: 472, Bernhauer & Schubert 1912: 208, Fagel 1958a: 430, 1966: 270]
- mexicanus* Dejean
[Dejean 1833: 65, *nomen nudum*, Gemminger & Harold 1868: 628, = *tempestivus* Erichson]

- mexicanus* Erichson, *Paederus*
[Erichson 1840: 657, Gemminger & Harold 1868: 628, Sharp 1886: 617, Bernhauer & Schubert 1912: 208, Leng 1920: 101, Blackwelder 1939: 112, 1944: 123, Moore & Legner 1975: 134]
- meyricki* Blackburn, *Paederus*
[Blackburn 1891: 72, 1894: 92, 1899: 22, Bernhauer & Schubert 1912: 208, Lea 1904: 64, 1923: 15, Scheerpeltz 1933: 1225]
- michaelensis* Last, *Paederus*
[Last 1979: 127]
- micropterus* Lea, *Paederus*
[Lea 1931: 367, Scheerpeltz 1933: 1225]
- mikenensis* (Fagel), *Paederus*, *Oreopaederus*
[Fagel 1954a: 965, 1958a: 102]
- mimicus* Last, *Paederus*
[Last 1979: 133]
- minimus* Bernhauer, *Paederus*
[Bernhauer 1915g: 365, Fagel 1956: 200, 1958a: 159, 1959: 46, 1963: 343, 1966: 271]
- minor* (Bernhauer), *Paederus*, *Oreopaederus*
[Bernhauer 1942b: 357, var. of *testaceus* Bernhauer, Fagel 1954a: 949, species, senior synonym of *filicornis* Fagel]
- minutissimus* (Bernhauer), *Paederus*, *Oreopaederus*
[Bernhauer 1927a: 108, 1954a: 916, Scheerpeltz 1933: 1225, Fagel 1958a: 101, 1960a: 115]
- minutulus* Fagel, *Oreopaederus*
[Fagel 1960b: 193]
- minutus* Gautier
[Gautier 1861b: xxxvi, *nomen nudum*]
- minutus* Gautier
[Gautier 1862: 76, Fauvel 1864: 12, = *limnophilus* Erichson, 1866: 24, 1873: 334, Marseul 1866: vi, Kraatz 1867: 415, Gemminger & Harold 1868: 627, Mulsant & Rey 1878: 267, Ganglbauer 1895: 537, Bernhauer & Schubert 1912: 207, Jarrige 1944: 9, Scheerpeltz 1957a: 459, 1968: 47, Legosz-Owsianna 1963: 331, Coiffait 1982c: 53]
- minutus* Fagel, *Pachypaederus*
[Fagel 1965: 141]
- mirandus* Fagel, *Allopaederus*
[Fagel 1958a: 106]
- mixtus* Sharp
[Sharp 1874: 75, Fauvel 1903a: 153, = *tamulus* Erichson, 1904a: 54, Bernhauer & Schubert 1912: 211, Cameron 1925b: 37, 1931a: 43, 1949b: 466, Scheerpeltz 1935: 609, Wu 1937: 329, Blackwelder 1939: 112, Shibata 1977: 23, Coiffait 1982c: 50]
- monardi* Bernhauer
[Bernhauer 1936b: 263, Fagel 1958a: 453, 1959: 56, = *lineativentris* Bernhauer]
- monstratus* Wendeler, *Paederus*
[Wendeler 1930: 186, Scheerpeltz 1933: 1225, Blackwelder 1944: 123]
- montanellus* (Bernhauer), *Paederus*, *Paederidus*
[Bernhauer 1934c: 233, 1940a: 135, Fagel 1958a: 133]
- montanellus decellei* Fagel
[Fagel 1958a: 134]
- montanellus leleupi* Fagel
[Fagel 1958a: 135]
- montanellus montanellus* (Bernhauer)
[Bernhauer 1934: 233, Fagel 1958a: 134, 1960b: 202, 1973: 154]
- montanellus pelengensis* Fagel
[Fagel 1958a: 134, 1960b: 202]
- montanellus rubricollis* Fagel
[Fagel 1958a: 135]
- montanus* Wendeler, *Paederus*
[Wendeler 1931: 44, Scheerpeltz 1933: 1225, Fagel 1958a: 452, 1966: 272]
- monticola* Cameron, *Paederus*
[Cameron 1949a: 317, Fagel 1958a: 229, 1963: 344, 1966: 259]
- montislomaensis* Fagel, *Paederus*
[Fagel 1967: 173]
- montivagans* Cameron, *Paederus*
[Cameron 1950c: 185, Fagel 1958a: 423, 1966: 270]
- morio* Mannerheim, *Paederus*
[Mannerheim 1831: 453, Dejean 1833: 65, Erichson 1840: 667, Gemminger & Harold 1868: 628, Bernhauer & Schubert 1912: 208, Leng & Mutchler 1914: 404, Blackwelder 1939: 112, 1943: 324, 1944: 123, Sanderson 1967: 60]
- morosus* Cameron, *Paederus*
[Cameron 1952a: 247, Last 1985: 327]
- moses* Saulcy
[Saulcy 1864: 654, Gemminger & Harold 1868: 628, Fauvel 1873: 331, = *littoralis* Gravenhorst, Ganglbauer 1895: 538, Bernhauer & Schubert 1912: 207, Scheerpeltz 1957a: 465, Legosz-Owsianna 1963: 349, Coiffait 1982c: 58]
- mulengensis* Fagel, *Oreopaederus*
[Fagel 1960b: 197]
- multipunctus auct.*
[Knoch *in coll.*, Gravenhorst 1802: 53, = *Lathrobium*]
- mussardi* Biswas & Sen Gupta, *Paederus* (*Pseudopaederus*)
[Biswas & Sen Gupta 1982: 150]

mutans Sharp, *Paederus*

[Sharp 1876a: 286, Bernhauer & Schubert 1912: 209, Plavilstschikov 1929: 29, Blackwelder 1939: 112, 1943: 322, 1944: 123]

mucngensis Fagel, *Paederus*

[Fagel 1958a: 392, 1966: 266]

N*nakaoi* Last, *Paederus*

[Last 1966: 4]

nakurensis Fauvel, *Paederus*

[Fauvel 1907: 24, Bernhauer & Schubert 1912: 209, Fagel 1956: 199, 1958a: 179, 1960a: 119, 1966: 271]

nakurensis insolitus Fagel

[Fagel 1960a: 120]

natalensis (Last), *Paederus*, *Pachypaederus*

[Last 1950: 136, Fagel 1958a: 87]

natalensis auct.

[Bernhauer *in litt.*, Fagel 1958a: 248, = *capitalis* Bernhauer]

Negalopaederus sic

[Shibata 1977: 29, misspelling of *Megalopaederus*]

Neopaederus Blackwelder

[Blackwelder 1939: 120, *subgenus*, type *Paederus morio* Mannerheim, 1952: 260, Fagel 1958a: 155, = *Paederus*]

neotropicus Bernhauer & Schubert

[Bernhauer & Schubert 1912: 209, replacement name for *thoracicus* Erichson, Blackwelder 1944: 123]

nepalensis Bernhauer, *Paederus* (*Nepalopaederus*)

[Bernhauer 1911a: 60, Bernhauer & Schubert 1912: 209, Cameron 1925b: 36, 1931: 55, Scheerpeltz 1933: 1226, 1957a: 474, 1976: 111]

nepalicus Coiffait, *Paederus*

[Coiffait 1976: 249, 1982a: 34]

Nepalopaederus Scheerpeltz

[Scheerpeltz 1976: 111, *subgenus*, type *Paederus nepalensis* Bernhauer]

nevadensis Austin

[Austin 1876: 10, LeConte 1878: 396, = *computens* LeConte, Casey 1905: 64, Bernhauer & Schubert 1912: 209, Leng 1920: 101, Blackwelder 1939: 112, Moore & Legner 1975: 133]

newtoni (Last), *Paederus*, *Pachypaederus*

[Last 1950: 136, Fagel 1958a: 83]

ngorongoroensis Fagel

[Fagel 1960: 129, *ssp.* of *basipes* Fauvel]

ngoviensis Fagel, *Oreopaederus*

[Fagel 1958a: 97]

niger (Bernhauer), *Paederus*, *Paederidus*

[Bernhauer 1931c: 573, Scheerpeltz 1933: 1226, Fagel 1958a: 128]

niger Bernhauer

[Bernhauer 1931a: 14, *ab.* (*infrasubspecific*) of *ruficollis* Fabricius, Scheerpeltz 1933: 1228, Fagel 1958a: 128, Coiffait 1982c: 23]

nigerrimus Bernhauer, *Paederus* (*Pseudopaederus*)

[Bernhauer 1915b: 137, Cameron 1925b: 36, 1931a: 62, Scheerpeltz 1933: 1226, Blackwelder 1939: 121, Biswas & Sen Gupta 1982: 147]

nigrans Scheerpeltz

[Scheerpeltz 1957a: 455, *var.* of *rubrothoracicus balcanicus* Scheerpeltz, Smetana 1959: 204]

nigratus Scheerpeltz

[Scheerpeltz 1957a: 456, Coiffait 1982c: 25, *var.* of *rubrothoracicus* Goeze]

nigrescens Cameron

[Cameron 1927: 115, *var.* of *amplicollis* Kraatz, 1931a: 58, Scheerpeltz 1933: 1220, here treated as *ssp.* under Article 45(g) (ICZN 1985)]

nigricans Scheerpeltz

[Scheerpeltz 1957a: 454, Coiffait 1982c: 25, *var.* of *rubrothoracicus* Goeze]

nigricans Fagel, *Oreopaederus*

[Fagel 1961a: 265]

nigriceps auct.

[Kirby *ms.*, Stephens 1829: 287, = *Sunius*]

nigricollis Bernhauer

[Bernhauer 1938b: 328, *ssp.* of *riftensis* Fauvel, Fagel 1958a: 331, = *riftensis* Fauvel]

nigricornis Bernhauer, *Paederus* (*Eopaederus*)

[Bernhauer 1911a: 61, Bernhauer & Schubert 1912: 209, Cameron 1925b: 36, 1928: 560, 1931a: 47, Scheerpeltz 1933: 1226, Blackwelder 1939: 112, Scheerpeltz 1957a: 474, Coiffait 1978b: 553, 1982a: 34, 1982b: 234, 1982c: 49]

nigripennis Cameron, *Paederus*

[Cameron 1924a: 180, 1931a: 57, Scheerpeltz 1957a: 474, Coiffait 1976: 243, 1982b: 234]

nigripes Dejean

[Dejean 1833: 65, *nomen nudum*]

nigripes Bernhauer, *Paederus*

[Bernhauer 1932: 87, Scheerpeltz 1933: 1226, Cameron 1950b: 25, Fagel 1958a: 176, 1961b: 268, 1966: 258]

nigripes auct.

[Bernhauer *in litt.*, Fagel 1956: 198, = *gowdeyi* Bernhauer, Fagel 1958a: 364]

- nigrithorax auct.*
[Desbrochers *in litt.*, Fauvel 1873: 335, = *sanguinicolis* Stephens]
- nigriventris* Wendeler, *Paederus*
[Wendeler 1926a: 73, 1935: 612, Scheerpeltz 1933: 1226]
- nigriventris* Fagel, *Pachypaederus*
[Fagel 1965: 139]
- nigrolineatus* Bernhauer, *Paederus*
[Bernhauer 1915a: 118, var. of *fuscipes* Curtis, Scheerpeltz 1933: 1223, Fagel 1958a: 173, species]
- nigrothoracicus* Last
[Last 1952: 91, var. of *vulneratus* Bernhauer, Fagel 1958a: 117, 1961b: 266, here treated as ssp. under Article 45(g) (ICZN 1985)]
- nitidicollis* Fagel
[Fagel 1954a: 957, ssp. of *vulcanus* Fagel, 1954b: 237, ssp. of *vulcanicola* Fagel, 1958a: 102]
- nitidipennis* Dejean
[Dejean 1833: 65, *nomen nudum*]
- nitidus* (Fagel), *Paederus*, *Oreopaederus*
[Fagel 1954a: 945, 1958a: 102]
- nobilis* Bernhauer
[Bernhauer 1912c: 475, Bernhauer & Schubert 1912: 209, Fagel 1958a: 405, = *magnificus* Bernhauer, 1960a: 125]
- nyakagerensis* Fagel, *Oreopaederus*
[Fagel 1958a: 100]
- nyalengwensis* Fagel, *Oreopaederus*
[Fagel 1961a: 270]
- nyamukubiensis* Fagel, *Paederus*
[Fagel 1958a: 375, 1966: 266]
- O**
- obliteratus* LeConte, *Paederus*, *Paederillus*, *Paederus*
[LeConte 1878: 395, Casey 1905: 65, Bernhauer & Schubert 1912: 112, Leng 1920: 101, Blackwelder 1939: 112, Moore & Legner 1975: 134]
- obscura* Wendeler
[Wendeler 1926d: 329, *forma obscura* of *sondaicus* Fauvel, Scheerpeltz 1933: 1228 as *ab. obscurus*, 1933: 609 as *forma obscurus*]
- obscuricollis* (Bernhauer), *Paederus*, *Pachypaederus*
[Bernhauer 1927a: 103, Scheerpeltz 1933: 1226, Fagel 1958a: 81]
- obscuripennis* (Fagel), *Paederus*, *Oreopaederus*
[Fagel 1954a: 938, 1958a: 102]
- obscuripes* Bernhauer, *Paederus*
[Bernhauer 1927a: 106, Scheerpeltz 1933: 1226, Fagel 1954a: 938, 1958a: 218, 1966: 262]
- obscuripes auct.*
[Bernhauer *in litt.*, Fagel 1958a: 421, = *melanogaster jeanneli* Fagel]
- obscurus auct.*
[Kirby *ms.*, Stephens 1829: 287, = *Sunius*]
- obscurus* Wendeler
[Wendeler 1926d: 329, *forma obscura* of *sondaicus* Fauvel, Scheerpeltz 1933: 1228, *ab. obscurus*, 1935: 609, *forma obscurus*, here believed to indicate infra-specific rank]
- occidentalis* Scheerpeltz
[Scheerpeltz 1957a: 452, ssp. of *rubrothoracicus* Goeze]
- ochraceus auct.*
[Knoch *in coll.*, Gravenhorst 1802: 59, = *ochraceus* Gravenhorst]
- ochraceus* Gravenhorst
[Gravenhorst 1802: 59, 1806: 138, Latreille 1804: 347, Panzer 1809: 14, Olivier 1812: 630, Stephens 1829: 287, = *Sunius*, now in *Lithocharis*]
- Oedopaederus* Scheerpeltz
[Scheerpeltz 1957a: 466, *subgenus*, type *Paederus chinensis* Bernhauer]
- ohausi* Wendeler, *Paederus*
[Wendeler 1928c: 33, Scheerpeltz 1933: 1226, Blackwelder 1944: 123]
- okapaensis* Last, *Paederus*
[Last 1979: 125, 1985: 328]
- oldeaniensis* Fagel
[Fagel 1960a: 128, ssp. of *busipes* Fauvel]
- Oncopaederus* Scheerpeltz
[Scheerpeltz 1957b: 265, *genus*, type *Oncopaederus brachypterus* Scheerpeltz]
- opacicollis* Bernhauer, *Paederus*
[Bernhauer 1915d: 299, Scheerpeltz 1933: 1226, Fagel 1958a: 170, 1960a: 119, 1966: 271]
- opacinus auct.*
[Fauvel *in litt.*, Fagel 1958a: 147, = *opacus* Bernhauer]
- opacipennis* Bernhauer
[Bernhauer 1908b: 105, Bernhauer & Schubert 1912: 209, Fagel 1958a: 146, = *collaris* Boheman]
- opacus* (Bernhauer), *Paederus*, *Paederidus*
[Bernhauer 1908b: 105, Bernhauer & Schubert 1912: 209, Bernhauer 1915i: 314, Fagel 1958a: 147]
- orbiculatus* (Paykull)
[Paykull 1789: 35, 1800: 431, Fabricius 1792: 538, 1801: 609, Panzer 1797: 21, Latreille 1806: 291, Gyllenhal 1810: 374, Samouelle 1819: 173, Dahl 1823: 17, Stephens 1829: 288, = *Rugilus*]

orbiculatus auct.

[Olivier 1795: 7, misidentification, 1812: 628, Heer 1839: 232, Mulsant & Rey 1878: 238, = *Rugilus*]

orbiculatus auct.

[Gravenhorst 1802: 63, misidentification, 1806: 141, Illiger 1807: 356, Beck 1817: 45, Mulsant & Rey 1878: 227, = *Rugilus*]

Oreinopaederus Scheerpeltz

[Scheerpeltz 1976: 112, *subgenus*, type *Paederus vastus* Scheerpeltz]

Oreopaederus Fagel

[Fagel 1958a: 93, *genus*, type *Paederus testaceus* Bernhauer]

orientalis Dejean

[Dejean 1833: 65, *nomen nudum*, Gemminger & Harold 1868: 627, = *extraneus* Wiedemann]

ornaticornis Sharp, *Paederus*

[Sharp 1891: 41, Bernhauer & Schubert 1912: 209, Chapin 1926: 371, Campos 1931: 24, Blackwelder 1944: 123]

orophilus Fagel

[Fagel 1954a: 919, *ssp.* of *brunnescens* Fagel, 1958a: 101]

orophilus Fagel

[Fagel 1958a: 379, homonym, 1966: 265, here = *Paederus zairensis* Frank, *nom. nov.*]

ovaliceps Bernhauer, *Paederus*

[Bernhauer 1905b: 180, Bernhauer & Schubert 1912: 209, Blackwelder 1944: 123]

overlaeti Fagel, *Paederus*

[Fagel 1958a: 262, 1966: 259]

P*Pachypaederus* Fagel

[Fagel 1958a: 70, *genus*, type *Paederus crassus* Boheman]

Paderus sic

[Scheerpeltz 1965b: 174, misspelling of *Paederus*]

Paederidius sic

[Scheerpeltz 1957a: 458, misspelling of *Paederidus*]

Paederidus Mulsant & Rey

[Mulsant & Rey 1878: 245, *subgenus*, Ganglbauer 1895: 536, Casey 1905: 59, *genus*, Bernhauer & Schubert 1912: 203, *subgenus*, Chapin 1927: 76, *genus*, Scheerpeltz 1933: 1219, *subgenus*, Blackwelder 1939: 120, type *Paederus ruficollis* Fabricius, = *Paederus*, 1952: 285, Jarrige 1944: 7, *subgenus*, Focarile 1957: 66, Scheerpeltz 1957a: 448, *genus*, Fagel 1958a: 109, Scheerpeltz 1968: 46, Coiffait 1982c: 15]

Paederillus Casey

[Casey 1905: 59, *genus*, Bernhauer & Schubert 1912: 203, = *Paederus*, Blackwelder 1939: 120, type *Paederus littorarius* Gravenhorst, 1952: 285, Hatch 1957: 169]

Paederognathus Wendeler

[Wendeler 1928a: 37, *subgenus*, replacement name for *Gnathopaederus* Wendeler, type *Paederus turribanus* Wendeler, Blackwelder 1939: 120, 1952: 286, = *Paederus*]

Paederomorphus sic

[Marseul 1866: vi, misspelling of *Poederomorphus*, Blackwelder 1939: 120, Fagel 1958a: 155, Coiffait 1982c: 28]

Paederus Fabricius

[Fabricius 1775: 268, *genus*, Olivier 1795: 1, Latreille 1804: 343, 1810: 427, type *Staphylinus riparius* Linnaeus, Bernhauer & Schubert 1912: 203, Blackwelder 1939: 120]

Paederus Fabricius

[Fabricius 1775: 268, Ganglbauer 1895: 537, *subgenus*, Bernhauer & Schubert 1912: 203, Scheerpeltz 1957a: 461, Legosz-Owsianna 1963: 321, Shibata 1973: 44, 1977: 26, Coiffait 1982c: 54]

paeninsula Last, *Paederus*

[Last 1985: 332]

pallidicornis auct.

[Bernhauer *in litt.*, Fagel 1958a: 421, = *melanogaster jeanneli* Fagel]

pallidipes Bernhauer

[Bernhauer 1927a: 111, Scheerpeltz 1933: 1226, Fagel 1958a: 279, = *plagiator* Kolbe]

pallidulus Scheerpeltz, *Paederus* (*Pseudopaederus*)

[Scheerpeltz 1933: 1226, replacement name for *pallidus* Bernhauer]

pallidus Sharp, *Paederus*

[Sharp 1886: 618, Bernhauer & Schubert 1912: 209, Scheerpeltz 1933: 1226, Blackwelder 1944: 123]

pallidus Bernhauer

[Bernhauer 1915b: 138, homonym, Cameron 1925b: 36, 1931a: 63, Scheerpeltz 1933: 1226, = *pallidulus* Scheerpeltz, Biswas & Sen Gupta 1982: 148]

pallipes auct.

[Knoch *in coll.*, Gravenhorst 1802: 180, = *Lathrobium*]

paludosus Dietrich

[Dietrich 1855: 202, Kraatz 1857: 729, = *caligatus* Erichson, Gemminger & Harold 1868: 626, Mulsant & Rey 1878: 264, Ganglbauer 1895: 537, Bernhauer & Schubert 1912: 224, Jarrige 1944: 9, Scheerpeltz 1933: 1221, 1968: 47, Coiffait 1982c: 52]

- palustris* Austin, *Paederus*, *Paederillus*, *Paederus*
[Austin 1876: 9, LeConte 1878: 396, Casey 1905: 64,
Bernhauer & Schubert 1912: 112, Leng 1920: 120,
Blackwelder 1939: 209, Moore & Legner 1975: 134]
- papuanus* Cameron, *Paederus*
[Cameron 1937: 96]
- paraguayanus* Wendeler, *Paederus*
[Wendeler 1942: 102, Gaedike 1981: 209]
- parallelipennis auct.*
[Bernhauer *in litt. partim*, Fagel 1958a: 225, = *amicus* Bernhauer]
- parallelipennis auct.*
[Bernhauer *in litt. partim*, Fagel 1958a: 269, = *xanthocerus* Eppelsheim]
- parallelus* Weise, *Paederus* (*Paederus*)
[Weise 1877: 368, Bernhauer & Schubert 1912: 209,
Scheerpeltz 1957a: 462, Tikhomirova 1973: 175,
Shibata 1977: 27]
- Parameropaederus* Scheerpeltz
[Scheerpeltz 1957a: 467, *genus*, type *Paederus lusitanicus* Aube, Coiffait 1982c: 68]
- parcepunctatus* (Bernhauer), *Paederus*, *Paederidus*
[Bernhauer 1931c: 574, Scheerpeltz 1933: 1226, Fagel 1958a: 145]
- parvipennis auct.*
[Bernhauer *in litt.*, Fagel 1958a: 457, = *metallicus* Fauvel]
- parvipennis auct.*
[Bernhauer *in litt.*, Fagel 1958a: 262, = *overlaeti* Fagel]
- pauliani* Jarrige, *Madecapaederus*
[Paulian 1970: 32]
- pauloensis* Bernhauer, *Paederus*
[Bernhauer 1908a: 309, Bernhauer & Schubert 1912: 209, Scheerpeltz 1933: 1226, Blackwelder 1944: 123]
- peculiaris* Fagel
[Fagel 1973: 153, ssp. of *celisianus* Fagel]
- pedestris* Gerstaecker, *Paederus*
[Gerstaecker 1866: 29, 1873: 82, Gemminger & Harold 1868: 628, Fauvel 1904b: 290, 1907: 21, Bernhauer & Schubert 1912: 209, Fagel 1958a: 214, 1960a: 120, 1966: 262]
- pedestris auct.*
[Fauvel *in litt.*, Fagel 1958a: 220, = *rufocyanus* Bernhauer]
- pedoncularius* Gautier
[Gautier 1861b: xxxvi, *nomen nudum*]
- pedoncularius* Gautier
[Gautier 1862: 75, Fauvel 1864: 13, Marseul 1866: vi, = *littoralis* Gravenhorst, Fauvel 1866: 25, Kraatz 1867: 415, Gemminger & Harold 1868: 627, Scheerpeltz 1957a: 465, 1968: 48, Coiffait 1982c: 58]
- peduncularius sic*
[Marseul 1866: vi, misspelling of *pedoncularius*, Fauvel 1866: 25, Ganglbauer 1895: 538, Bernhauer & Schubert 1912: 207, Scheerpeltz 1957a: 465, 1968: 48, Legosz-Owsianna 1963: 349]
- pelengensis* Fagel
[Fagel 1958a: 134, ssp. of *montanellus* Bernhauer]
- pelikani* Reitter, *Paederus* (*Dioncopaederus*)
[Reitter 1884: 44, Fauvel 1884: 241, = *meridionalis* Fauvel, Reitter 1885: 153, species, 1889: 170, Czwalina 1889: 368, Bernhauer & Schubert 1912: 209, Scheerpeltz 1957a: 465, Coiffait 1982c: 59]
- pendleburyi* Cameron, *Paederus*
[Cameron 1950a: 11]
- pentagonalis* Wendeler, *Paederus*
[Wendeler 1931: 38, Scheerpeltz 1933: 1226, 1935: 628]
- peregrinus* Erichson, *Paederus* (*Heteropaederus*)
[Erichson 1840: 656, Kraatz 1959a: 151, Gemminger & Harold 1868: 628, Baer 1886: 107, Fauvel 1879: 89, = *fuscipes* Curtis, 1886b: 148, 1903a: 154, 1904b: 290, Bernhauer & Schubert 1912: 206, var. of *fuscipes* Curtis, Wendeler 1926d: 328, Cameron 1925b: 36, 1931a: 40, 1936: 43, Scheerpeltz 1933: 1223, Koch 1934 82, species, Blackwelder 1939: 112, Scheerpeltz 1935: 604, var. of *fuscipes* Curtis, 1957a: 461, Coiffait 1976: 244, species, 1982a: 34, 1982b: 234, 1982c: 61]
- perlongus* Last, *Paederus*
[Last 1979: 131, 1985: 327]
- permixtus* Fagel, *Paederidus*
[Fagel 1959: 43]
- perrieri* (Fauvel), *Paederus*, *Paederidus*
[Fauvel 1898: 180, Bernhauer & Schubert 1912: 209, Fagel 1958a: 457]
- peruanus* Wendeler, *Paederus*
[Wendeler 1931: 41, Scheerpeltz 1933: 1226, Blackwelder 1944: 123]
- philippinus* Bernhauer, *Paederus*
[Bernhauer 1912d: 250, 1927c: 399, Bernhauer & Schubert 1912: 209, Cameron 1930b: 332, Scheerpeltz 1933: 1226, Blackwelder 1939: 112]
- piceus* Fauvel, *Paederus*
[Fauvel 1907: 22, Bernhauer & Schubert 1912: 209, Fagel 1958a: 410, 1960a: 126]
- piceps* Bernhauer
[Bernhauer 1934c: 232, 1940a: 135, Fagel 1954a: 961, = *alinderi* Wendeler, 1956: 188, Fagel 1958a: 102]

- picipes sic*
[Fagel 1956: 188, misspelling of *piciceps*]
- pietschmanni* Bernhauer
[Bernhauer 1913b: 222, Roubal 1932: 61, Scheerpeltz 1933: 1226, 1957a: 459, Coiffait 1982c: 51, = *mesopotamicus* Eppelsheim]
- pilifer* Motschulsky, *Paederus* (*Paederus*)
[Motschulsky 1860: 74, epithet incorrectly formed as *piliferus*, Gemminger & Harold 1868: 628, spelled as *pilifer*, Bernhauer & Schubert 1912: 209, Cameron 1925b: 36, 1931a: 44, 1940: 88, Scheerpeltz 1933: 1226, Shibata 1973: 44]
- piliferus auct.*
[see *pilifer*]
- piliventris* Fagel
[Fagel 1968: 197, ssp. of *erythraeanus* Fagel]
- pilosus auct.*
[Knoch *in coll.*, Gravenhorst 1802: 56, = *Lathrobium*]
- plagiator* Kolbe, *Paederus*
[Kolbe 1883: 18, Bernhauer & Schubert 1912: 209, Bernhauer 1942b: 357, Fagel 1956: 199, 1958a: 279, 1966: 259]
- Poederomorphus* Gautier
[Gautier 1861b: xxxvi, *nomen nudum*]
- Poederomorphus* Gautier
[Gautier 1862: 75, *genus*, Bernhauer & Schubert 1912: 203, = *Paederus*, Blackwelder 1939: 120, type *Poederomorphus pedoncularius* Gautier, 1952: 316, Fagel 1958a: 155]
- Paederus sic*
[Baudi 1848: 138, misspelling of *Paederus*, Laporte 1840: 183, Peyron 1858: 430, Fairmaire 1859: clxxxiv, Thomson 1860: 195, Gautier 1861: 393, 1866: 114, Saulcy 1864: 654]
- politus* Fauvel
[Fauvel 1878a: 237, 1879: 88, Bernhauer & Schubert 1912: 209, Cameron 1931b: 357, 1937: 95, Puthz 1969: 529, Last 1979: 138, 1985: 328]
- porrectus* Wendeler
[Wendeler 1933: 56, *species*, Fagel 1958a: 422, = ssp. of *melanogaster* Fauvel]
- posticollis sic*
[Bernhauer 1942b: 357, misspelling of *usticollis*]
- posticus auct.*
[Fauvel *in litt.*, Fagel 1958a: 279, = *plagiator* Kolbe]
- poweri auct.*
[Lewis *in litt.*, Sharp 1874: 74, = *poweri* Sharp]
- poweri* (Sharp), *Paederus*, *Megalopaederus*
[Sharp 1874: 74, Bernhauer & Schubert 1912: 209, Scheerpeltz 1933: 1226, Wu 1937: 329, Blackwelder 1939: 113, Cameron 1949b: 466, Scheerpeltz 1957a: 471, Tikhomirova 1973: 175, Shibata 1977: 129, Watanabe 1986: 160]
- praecellens* Bernhauer, *Paederus*
[Bernhauer 1915e: 221, Scheerpeltz 1933: 1226, 1935: 622]
- praeses* Fagel, *Paederus*
[Fagel 1958a: 198, 1966: 263]
- praeses auct.*
[Fauvel *in litt.*, Fagel 1958a: 198, = *praeses* Fagel]
- preangeranus* Bernhauer, *Paederus*
[Bernhauer 1920b: 178, Scheerpeltz 1933: 1226, 1935: 607]
- pretiosus* Bernhauer, *Paederus*
[Bernhauer 1915c: 290, Scheerpeltz 1933: 1226, Fagel 1958a: 454, 1959: 64, 1966: 268]
- problematis* Last, *Paederus*
[Last 1979: 136, 1985: 327]
- proboscidens* Olivier
[Olivier 1795: 6, Gyllenhal 1810: 476, = *Stenus*]
- proboscideus sic*
[Gyllenhal 1810: 476, misspelling of *proboscidens*]
- proboscidus sic*
[Gyllenhal 1810: 476, misspelling of *proboscidens*]
- procerus auct.*
[Knoch *in coll.*, Gravenhorst 1806: 141, = *procerus* Gravenhorst]
- procerus* Gravenhorst
[Gravenhorst 1806: 141, Beck 1817: 45, Dahl 1823: 17, Dejean 1833: 66, = *Astenus*]
- procerus auct.*
[Lacordaire 1835: 436, synonymic homonym, = *Astenus*]
- protensus* Sharp
[Sharp 1876a: 287, Fauvel 1891: 100, = *columbinus* Laporte, Bernhauer & Schubert 1912: 205, Blackwelder 1939: 112, 1944: 122]
- pseudobaudii* Aleksandrov, *Paederus* (*Harpopaederus*)
[Aleksandrov 1934: 151, Scheerpeltz 1957a: 460, Smetana 1962: 49, Tikhomirova 1973: 175, Coiffait 1982c: 64]
- Pseudopaederus* Bernhauer
[Bernhauer 1915b: 137, *subgenus*, Scheerpeltz 1933: 1219, Blackwelder 1939: 121, type *Paederus nigerimus* Bernhauer, 1952: 328]
- pseustes* (Fagel), *Paederus*, *Oreopaederus*
[Fagel 1954a: 929, 1956: 186, 1958a: 101, 1960b: 196]
- puberulus* Motschulsky, *Paederus*
[Motschulsky 1860: 74, Gemminger & Harold 1868: 628, Bernhauer & Schubert 1912: 209, Cameron

- 1925b: 36, 1931a: 46, Scheerpeltz 1933: 1226]
pubescens Cameron, *Paederus* (*Eopaederus*)
 [Cameron 1914: 539, 1925b: 35, = *extraneus*
 Wiedemann, 1931a: 44, species, Scheerpeltz 1933:
 1226, Coiffait 1978b: 553, 1982c: 50]
- pugetanus sic*
 [Hatch 1957: 170, misspelling of *pugetensis*, Moore
 & Legner 1975: 134]
- pugetensis* (Casey)
 [Casey 1905: 66, Bernhauer & Schubert 1912: 209,
 Leng 1920: 101, Blackwelder 1939: 112, Hatch 1957:
 170, = *littorarius* Gravenhorst]
- pulchellus* Bernhauer, *Paederus*
 [Bernhauer 1927a: 116, Scheerpeltz 1933: 1226, Fagel
 1958a: 165, 1966: 272]
- pumilus* (Wendeler), *Paederus*, *Paederidus*
 [Wendeler 1928b: 263, Peyerimhoff 1931: 29,
 Scheerpeltz 1933: 1226, Coiffait 1982c: 22]
- punctatus* Last, *Paederus*
 [Last 1985: 337]
- puncticeps auct.*
 [Fauvel *in litt.*, Fagel 1958a: 269, = *xanthocerus* Ep-
 pelsheim]
- puncticollis* (Bernhauer), *Paederus*, *Pachypaederus*
 [Bernhauer 1912c: 471, Bernhauer & Schubert 1912:
 209, Bernhauer 1942b: 357, Fagel 1956: 199, 1958a:
 89, 1960a: 115, 1963: 343]
- punctiger* Sharp, *Paederus*
 [Sharp 1876a: 288, Blackwelder 1944: 123]
- punctipennis auct.*
 [Kirby *ms.*, Stephens 1829: 288, = *Rugilus*]
- punctipennis* (Fagel), *Paederus*, *Oreopaederus*
 [Fagel 1954a: 963, 1956: 189, 1958a: 102]
- punctiventris* Bernhauer, *Paederus*
 [Bernhauer 1908a: 310, Bernhauer & Schubert 1912:
 209, Blackwelder 1944: 123]
- punctiventris* Scheerpeltz, *Paederidus*
 [Scheerpeltz 1957a: 457, Coiffait 1982c: 27]
- punjabensis* Cameron, *Paederus*
 [Cameron 1945: 68]
- purpuripennis* Bernhauer, *Paederus*
 [Bernhauer 1915a: 118, Scheerpeltz 1933: 1227, Fagel
 1958a: 449, 1966: 270]

Q

- quadratus* (Paykull)
 [Paykull 1789: 29, *partim*, 1800: 429, = *filiformis* Fab-
 ricius]

- quadratus auct.*
 [Paykull 1789: 29, *partim*, 1800: 429, = *brunnipes* Fab-
 ricius]
- quadriiceps auct.*
 [Fagel 1958a: 329, *nomen nudum*]
- quadrifasciatus* Cameron
 [Cameron 1929: 57, Scheerpeltz 1933: 1227,
 Cameron 1950b: 26, var. of *riftensis* Fauvel, Fagel
 1958a: 331]
- quadrifasciatus auct.*
 [Bernhauer *in litt.*, Fagel 1958a: 331, = *riftensis* Fau-
 vel]

R

- raffrayi auct.*
 [Fauvel *in litt.*, Fagel 1958a: 295, = *raffrayi* Fagel]
- raffrayi* Fagel, *Paederus*
 [Fagel 1958a: 295, 1966: 257]
- raundanus sic*
 [Fagel 1956: 194, misspelling of *ruandanus*]
- reductus* Fagel, *Pachypaederus*
 [Fagel 1965: 143]
- republicanus* Bernhauer, *Paederus*
 [Bernhauer 1912a: 34, Scheerpeltz 1933: 1227, Black-
 welder 1944: 123]
- reticulatus* Last, *Paederus*
 [Last 1979: 125]
- reticuliventris* Fagel, *Paederus*
 [Fagel 1958a: 223, 1966: 262]
- rhodesianus* (Bernhauer), *Paederus*, *Pachypaederus*
 [Bernhauer 1927a: 102, Scheerpeltz 1933: 1227, Fagel
 1958a: 85]
- riftensis* Fauvel, *Paederus*
 [Fauvel 1907: 24, Bernhauer & Schubert 1912: 209,
 Cameron 1939: 4, 1950b: 26, Bernhauer 1926: 6,
 1940a: 135, Fagel 1956: 200, 1958a: 331, 1966: 261]
- riftensis nigricollis* Bernhauer
 [Bernhauer 1938b: 328, Fagel 1958a: 331, = *riftensis*
 Fauvel]
- riparius* (Linnaeus), *Staphylinus*, *Paederus* (*Paederus*)
 [Linnaeus 1758: 422, 1767: 684, DeGeer 1774: 28,
 Fabricius 1775: 268, Fabricius 1781: 339, 1787: 223,
 1792: 536, Harrer 1784: 253, Herbst 1784: 151, Lin-
 naeus 1789: 412, Panzer 1793a: 11, Paykull 1789: 27,
 1800: 427, Rossi 1790: 253, Fabricius 1801: 608, [? cor-
 rectly identified, Latreille 1806: 293, 1810: 427], Gyl-
 lenhal 1810: 372, Curtis 1826: 108, Stephens 1829:
 288, 1833: 282, Mannerheim 1831: 453, Dejean 1833:
 65, Lacordaire 1835: 428, Heer 1839: 234, Falder-
 mann 1839: 98, Erichson 1839: 518, 1840: 653, Fair-

- maire & Laboulbène 1856: 571, Kraatz 1857: 727, Thomson 1860: 196, Gemminger & Harold 1868: 628, Fauvel 1869b: 491, 1873: 332, Solsky 1871: 163, Austin 1876: 9, Sahlberg 1876: 37, Mulsant & Rey 1878: 259, LeConte 1878: 395, Fauvel 1882: 214, Eppelsheim 1891: 340, Reitter 1889: 170, Czwalina 1889: 368, Ganglbauer 1895: 537, Fauvel 1897: 281, Casey 1905: 61, Bernhauer & Schubert 1912: 209, Leng 1920: 101, Kirschenblatt 1932: 218, Scheerpeltz 1933: 1227, Aleksandrov 1934: 150, Blackwelder 1939: 112, Jarrige 1944: 8, Hansen 1951: 223, Scheerpeltz 1957a: 462, 1968: 47, Ferreira 1962: 14, Legosz-Owsianna 1963: 344, Lohse 1964: 134, Szujewski 1965b: 28, Jarrige 1971: 491, Tikhomirova 1973: 175, Moore & Legner 1975: 134, Coiffait 1978c: 165, 1982c: 55, Bohác 1985: 370]
- riparius auct.*
[Schrank 1781: 233, misidentification, Olivier 1795: 4, 1812: 627, Erichson 1839: 519, = *littoralis* Gravenhorst, Fairmaire & Laboulbène 1856: 570, Kraatz 1857: 726, Gemminger & Harold 1868: 627, Mulsant & Rey 1878: 256, Ganglbauer 1895: 538, Bernhauer & Schubert 1912: 207, Jarrige 1944: 8, Scheerpeltz 1957a: 465, Legosz-Owsianna 1963: 349]
- riparius auct.*
[Gravenhorst 1802: 62, misidentification, 1806: 143, Illiger 1807: 356, Dahl 1823: 17, Erichson 1839: 51, Kraatz 1857: 728, = *fuscipes* Curtis, Gemminger & Harold 1868: 627, Mulsant & Rey 1878: 262, Ganglbauer 1895: 537, Bernhauer & Schubert 1912: 206, Cameron 1925b: 36, 1931a: 40, Legosz-Owsianna 1963: 335, Scheerpeltz 1968: 47, Hammond 1977: 27, Muona 1979: 16, Coiffait 1982c: 61]
- riparius balcanicus* Koch
[Koch 1938: 103, ssp., Szujewski 1965: 83, species]
- riparius korbi* Bernhauer
[Bernhauer 1935a: 44, ssp., Scheerpeltz 1957a: 462, var. of *riparius* Linnaeus]
- roepkei* Bernhauer, *Paederus*
[Bernhauer 1915e: 220, Scheerpeltz 1933: 1227, 1935: 614]
- rothschildi* Fagel
[Fagel 1958a: 353, ssp. of *duplex* Eppelsheim]
- ruandanus* Fagel, *Paederus*
[Fagel 1956: 191, 1958a: 381, 1966: 265]
- rubidus* Cameron, *Paederus*
[Cameron 1947: 114]
- rubricollis auct.*
[Knoch *ms.*, Gravenhorst 1806: 138, = *rubricollis* Gravenhorst]
- rubricollis* Gravenhorst
[Gravenhorst 1806: 138, Gyllenhal 1810: 376, Olivier 1812: 630, Stephens 1829: 287, = *Sunius*, now in *Lithocharis*]
- rubricollis* Fagel
[Fagel 1958a: 135, ssp. of *montanellus* Bernhauer, 1960b: 202]
- rubrothoracicus* (Goeze), *Staphylinus*, *Paederus*, *Paederidus*
[Goeze 1777: 731, Bernhauer & Schubert 1912: 210, Kirschenblatt 1932: 216, Jarrige 1944: 7, Focarile 1957: 66, Scheerpeltz 1933: 1227, 1957a: 452, 1968: 46, Ferreira 1962: 14, Legosz-Owsianna 1963: 327, Lohse 1964: 133, Tikhomirova 1973: 175, Hammond 1977: 27, Bohác 1985: 367]
- rubrothoracicus asiaeminoris* Scheerpeltz
[Scheerpeltz 1957a: 456, Coiffait 1982c: 25, var.]
- rubrothoracicus ater* (Eichler)
[Eichler 1924: 65, var. of *ruficollis* Fabricius, Scheerpeltz 1933: 1227, var. of *rubrothoracicus* Goeze, Coiffait 1982c: 27, here treated as ssp. under Article 45(g) (ICZN 1985)]
- rubrothoracicus balcanicus* Scheerpeltz
[Scheerpeltz 1957a: 455, Coiffait 1982c: 25, var.]
- rubrothoracicus carpathicola* Scheerpeltz
[Scheerpeltz 1957a: 455, Coiffait 1982c: 25, var., Bohác 1985: 369, ssp.]
- rubrothoracicus caucasicus* Scheerpeltz
[Scheerpeltz 1957a: 456, Coiffait 1982c: 25, var.]
- rubrothoracicus ibericus* Scheerpeltz
[Scheerpeltz 1957a: 453, Coiffait 1982c: 25, var.]
- rubrothoracicus meridiogalicus* Scheerpeltz
[Scheerpeltz 1957a: 453, Coiffait 1982c: 25, var.]
- rubrothoracicus meridionalis* Scheerpeltz
[Scheerpeltz 1957a: 454, Coiffait 1982c: 25, var.]
- rubrothoracicus occidentalis* Scheerpeltz
[Scheerpeltz 1957a: 452]
- rubrothoracicus rubrothoracicus* (Goeze)
[Goeze 1777: 731, Szujewski 1965b: 26, Bohác 1985: 367]
- rubrothoracicus sardous* Scheerpeltz
[Scheerpeltz 1957a: 454, Coiffait 1982c: 25, var.]
- rubrothoracicus sic*
[Scheerpeltz 1957a: 455, misspelling of *rubrothoracicus*]
- rudebecki* Fagel, *Paederus*
[Fagel 1961b: 268, 1966: 264]
- ruficeps* Baudi
[Baudi 1848: 138, Kraatz 1857: 730, Fauvel 1873: 332, Reitter 1889: 171, Czwalina 1889: 368, Bernhauer & Schubert 1912: 20, = var. of *melanurus* Aragona,

- Scheerpeltz 1957a: 462, Smetana 1959: 204, species, Coiffait 1982c: 57, var. of *melanurus* Aragona]
- ruficollis sic*
[Coiffait 1982c: 26, misspelling of *ruficollis*]
- ruficollis* (Fabricius), *Paederus*, *Paederidus*
[Fabricius 1776: 243, 1781: 339, Herbst 1786: 180, Fabricius, 1787: 223, 1792: 537, 1801: 608, Olivier 1795: 4, Panzer 1796a: 22, Paykull 1789: 26, 1800: 427, Gravenhorst 1802: 185, 1806: 143, Gyllenhal 1810: 373, Dahl 1823: 17, Stephens 1829: 288, 1833: 281, Mannerheim 1831: 453, Dejean 1833: 65, Lacordaire 1835: 430, Faldermann 1839: 98, Heer 1839: 233, Laporte 1840: 183, Erichson 1840: 662 *partim*, Lucas 1849: 121, Fairmaire & Laboulbène 1856: 571, Kraatz 1857: 731, Thomson 1860: 196, Gemminger & Harold 1868: 628, Fauvel 1873: 336, 1882: 214, 1886b: 32, 1904b: 290, Mulsant & Rey 1878: 246, Reitter 1889: 171, Eppelsheim 1895b: 211, Ganglbauer 1895: 536, Bernhauer & Schubert 1912: 210, Chapman 1930: 102, Kirschenblatt 1932: 216, Koch 1937: 24, Blackwelder 1939: 112, Jarrige 1944: 8, Hansen 1951: 223, Focarile 1957: 69, Scheerpeltz 1957a: 449, 1968: 46, Ferreira 1962: 14, Legosz-Owsianna 1963: 323, Lohse 1964: 133, Szujecki 1965b: 25, Tikhomirova 1973: 175, Muona 1979: 16, Coiffait 1982c: 22, Bohác 1985: 367]
- ruficollis algiricus* Motschulsky
[Motschulsky 1858: 635, species, Scheerpeltz 1957a: 451, ssp., Coiffait 1982c: 24, species]
- ruficollis antoinei* Koch
[Koch 1937: 24, species, Scheerpeltz 1957a: 450, ssp., Coiffait 1982c: 24, ssp. of *algiricus*]
- ruficollis gallomeridionalis* Scheerpeltz
[Scheerpeltz 1957a: 450, ssp., Coiffait 1982c: 22, var.]
- ruficollis hispanolusitanus* Scheerpeltz
[Scheerpeltz 1957a: 450, ssp., Coiffait 1982c: 23, var.]
- ruficollis auct.*
[Erichson 1840: 662 *partim*, Baudi 1857: 109 *partim*, Gemminger & Harold 1868: 628, = *sanguinicollis* Stephens, Fauvel 1869b: 491, 1873: 335, Bernhauer & Schubert 1912: 210, = *rubrothoracicus* Goeze, Jarrige 1944: 7, Scheerpeltz 1957a: 452, 1968: 46, Legosz-Owsianna 1963: 327, Hammond 1977: 27]
- ruficollis auct.*
[Baudi 1857: 109, *partim*, Gemminger & Harold 1868: 627, = *gemellus* Kraatz]
- ruficoxis* Kraatz
[Kraatz 1859a: 151, Gemminger & Harold 1868: 628, Bernhauer 1902: 38, Fauvel 1904a: 54, = *alternans* Walker, Bernhauer & Schubert 1912: 203, Cameron 1925b: 34, 1931a: 41, Scheerpeltz 1935: 607, 1978: 196, Gaedike 1981: 216, Biswas & Sen Gupta 1982: 150]
- rufipes* (Fauvel), *Paederus*, *Madecapaederus*
[Fauvel 1898: 181, var. of *apterus* Fauvel, Bernhauer & Schubert 1912: 204, Fagel 1958a: 457, species]
- rufitarsis* Solsky, *Paederus*
[Solsky 1866: 106, Sharp 1886: 611, Bernhauer & Schubert 1912: 210, Blackwelder 1944: 123]
- rufitartis sic*
[Gemminger & Harold 1868: 628, misspelling of *rufitarsis*]
- rufiventris* Bernhauer, *Paederus*
[Bernhauer 1915a: 116, Scheerpeltz 1933: 1228, Fagel 1958a: 335, 1966: 263]
- rufobrunneus* Cameron
[Cameron 1950c: 185, species, Fagel 1958a: 358, ssp. of *macellus* Fauvel]
- rufocyaneus* Bernhauer, *Paederus*
[Bernhauer 1912c: 475, 1931c: 575, Bernhauer & Schubert 1912: 210, Fagel 1958a: 220, 1966: 262]
- rufocyaneus auct.*
[Fauvel *in litt.*, Fagel 1958a: 220, = *rufocyaneus* Bernhauer]
- rufofasciatus* (Bernhauer), *Paederus*, *Paederidus*
[Bernhauer 1915a: 115, Scheerpeltz 1933: 1228, Fagel 1958a: 141, 1960a: 119]
- rufotestaceus* Cameron, *Paederus*
[Cameron 1942b: 841]
- rugegensis* (Fagel), *Paederus*, *Oreopaederus*
[Fagel 1954a: 923, 1958a: 101]
- rugepunctatus auct.*
[Chevrolat *in litt.*, Sharp 1886: 614, = *yucateca* Sharp]
- rugepunctatus auct.*
[Chevrolat *in coll.*, Gemminger & Harold 1868: 628, = *tempestivus* Erichson]
- rugipennis* Motschulsky
[Motschulsky 1860: 74, Gemminger & Harold 1868: 628, Fauvel 1903a: 153, = *tamulus* Erichson, 1904a: 54, Bernhauer & Schubert 1912: 211, Cameron 1925b: 37, 1931a: 43, Wu 1937: 329, Scheerpeltz 1935: 609, Coiffait 1982c: 50]
- ruhembeanus* Bernhauer, *Paederus*
[Bernhauer 1927a: 115, Scheerpeltz 1933: 1228, Fagel 1959: 161, 1966: 272]
- rutilicornis* Erichson, *Paederus*
[Erichson 1840: 666, Gemminger & Harold 1868: 628, Bernhauer & Schubert 1912: 210, Blackwelder 1944: 123]
- ruwenzoricus* Fagel, *Paederus*
[Fagel 1956: 193, 1958a: 381, 1966: 266, 1973: 158]

S

sabaeus Erichson, *Paederus* (*Heteropaederus*)

[Erichson 1840: 655, Gerstaecker 1867: 29, 1873: 81, Gemminger & Harold 1868: 628, Eppelsheim 1895b: 210, Fauvel 1904b: 290, 1907: 24, Bernhauer & Schubert 1912: 210, Bernhauer 1912c: 471, 1915g: 362, 1915i: 315, 1926: 6, 1931c: 575, 1940a: 135, Eichelbaum 1913: 125, Roubal 1932: 61, Blackwelder 1939: 112, Cameron 1950b: 25, 1952b: 829, Fagel 1956: 199, 1958a: 307, 1959: 59, 1960b: 203, 1963: 346, 1966: 269, Coiffait 1982c: 61]

sachtlebeni Fagel, *Paederus*

[Fagel 1958a: 247, 1966: 260]

saegeri Fagel, *Paederidus*

[Fagel 1959: 44]

saengeri Coiffait, *Paederus* (*Paederus*)

[Coiffait 1979: 167, 1982c: 55]

saginatus (Casey), *Paederillus*, *Paederus*

[Casey 1905: 63, Bernhauer & Schubert 1912: 210, Leng 1920: 101, Blackwelder 1939: 112, Moore & Legner 1975: 134]

salvini Sharp, *Paederus*

[Sharp 1876b: 431, 1886: 610, Bernhauer & Schubert 1912: 210, Blackwelder 1944: 123, Scheerpeltz 1965: 215]

samarensis Bernhauer, *Paederus*

[Bernhauer 1927c: 399, Scheerpeltz 1933: 1228]

samoensis Fauvel, *Paederus*

[Fauvel 1877a: 224, 1878b: 517, = *vitiensis* Fauvel, Fairmaire 1881: 251, Bernhauer & Schubert 1912: 211]

samuelseni Last, *Paederus*

[Last 1985: 331]

sanguinicollis Dahl

[Dahl 1823: 17, *nomen nudum*, Dejean 1933: 65, = *Rugilus*]

sanguinicollis Stephens

[Stephens 1829: 288, *nomen nudum*]

sanguinicollis Stephens

[Stephens 1833: 281, Gemminger & Harold 1868: 628, Fauvel 1869b: 491, Ganglbauer 1895: 536, Bernhauer & Schubert 1912: 210, = *rubrothoracicus* Goeze, Scheerpeltz 1933: 1227, Blackwelder 1939: 112, Focarile 1957: 66, Scheerpeltz 1957a: 452, 1968: 46, Lohse 1964: 133, Tikhomirova 1973: 175, Hammond 1977: 27]

sanguinicollis Motschulsky

[Motschulsky 1860b: 559, homonym, Gemminger & Harold 1868: 627, = *haematoderus* Gemminger & Harold, replacement name]

sanguinicollis sic

[Legosz-Owsianna 1963: 327, misspelling of *sanguinicollis*]

santoensis Cameron, *Paederus*

[Cameron 1942a: 87]

sardous Scheerpeltz

[Scheerpeltz 1957a: 454, ssp. of *rubrothoracicus* Goeze, Coiffait 1982c: 25, var. of *rubrothoracicus* Goeze]

scabricollis Dahl

[Dahl 1823: 17, *nomen nudum*, Dejean 1833: 65, = *Rugilus*]

scabricollis sic

[Bernhauer 1942b: 357, misspelling of *scabripennis*]

scabripennis (Fauvel), *Paederus*, *Paederidus*

[Fauvel 1907: 22, Bernhauer & Schubert 1912: 211, Peyerimhoff 1931: 29, Bernhauer 1931c: 575, Fagel 1958a: 116, 1960a: 117]

scabripennis zanzibaricus Fagel

[Fagel 1958a: 117]

scheerpeltzi Fagel, *Paederus*

[Fagel 1958a: 276, 1959: 58, 1966: 260]

schmidti auct.

[Bernhauer *in coll.*, Gaedike 1981: 231]

schoenheri sic

[Tikhomirova 1973: 175, misspelling of *schoenherri*]

schoenherri auct.

[Faldermann unpublished, Czwalina 1889: 368, = *schoenherri* Czwalina]

schoenherri Czwalina, *Paederus* (*Harpopaederus*)

[Czwalina 1889: 368, Ganglbauer 1895: 538, = *baudii* Fairmaire, Bernhauer & Schubert 1912: 204, Koch 1938: 104, species, Jarrige 1944: 9, Scheerpeltz 1957a: 460, 1968: 47, Smetana 1962: 49, Legosz-Owsianna 1963: 342, Lohse 1964: 134, Szujewski 1965b: 30, Scheerpeltz 1968: 47, Tikhomirova 1973: 175, Coiffait 1982c: 63, Bohác 1985: 375]

schoenherri schulzei Korge

[Korge 1969: 61]

schonheri sic

[Coiffait 1978c: 165, misspelling of *schoenherri* and misidentification, 1978d: 387]

schoutedeni (Bernhauer), *Paederus*, *Paederidus*

[Bernhauer 1915d: 300, Scheerpeltz 1933: 1228, Fagel 1958a: 121]

schuberti auct.

[Fauvel *in litt.*, Fagel 1958a: 442, = *usambaricus* Schubert]

schultheissi Fauvel, *Paederus*

[Fauvel 1895: 233, Bernhauer & Schubert 1912: 211,

- Cameron 1930b: 332, 1936: 43, Scheerpeltz 1935: 606]
schulzei Korge
 [Korge 1969: 61, ssp. of *schoenherri* Czwalina]
- schusteri* Bernhauer, *Paederus*
 [Bernhauer 1908a: 310, Bernhauer & Schubert 1912: 211, Blackwelder 1944: 123]
- scottianus* Fagel, *Paederus*
 [Fagel 1958a: 184, 1966: 271]
- secretus* Bernhauer, *Paederus*
 [Bernhauer 1939c: 227, Fagel 1958a: 372, 1966: 266, 1973: 158]
- sedlaceki* Last, *Paederus*
 [Last 1985: 333]
- segmentarius* Cameron
 [Cameron 1933: 40, Fagel 1958a: 331, = *riftensis* Fauvel]
- semaranganus* Bernhauer, *Paederus*
 [Bernhauer 1915e: 220, Scheerpeltz 1933: 1228, 1935: 615]
- semicyaneus* Perty
 [Perty 1830: 33, Laporte 1835: 120, = *Taenodema*]
- semicyaneus* Cameron
 [Cameron 1950c: 185, homonym, Fagel 1958a: 129, = *luctuosus* Klug, 1960a: 117]
- semiviridis* Bernhauer, *Paederus*
 [Bernhauer 1927c: 400, Scheerpeltz 1933: 1228]
- semiviridis lividipennis* Bernhauer
 [Bernhauer 1927c: 399, var., here treated as ssp. under Article 45(g) (ICZN 1985)]
- senegalensis* Laporte, *Paederus*
 [Laporte 1835: 123, Gemminger & Harold 1868: 628, Bernhauer & Schubert 1912: 211, Fagel 1958a: 454, 1966: 272]
- senegalensis* Fagel, *Paederidus*
 [Fagel 1962a: 67]
- septimus* Schaeffer
 [Schaeffer 1779: lxxi, Harrer 1784: 253, = *riparius* Linnaeus, Erichson 1840: 654, Kraatz 1857: 728, Gemminger & Harold 1868: 628, Ganglbauer 1895: 537, Bernhauer & Schubert 1912: 210, Scheerpeltz 1957a: 462, Legosz-Owsianna 1963: 344]
- sericeicollis* Scheerpeltz, *Paederus*
 [Scheerpeltz 1935: 616]
- setifer* Cameron, *Paederus*
 [Cameron 1914: 538, 1925b: 36, 1931a: 60, Scheerpeltz 1933: 1228, Biswas & Sen Gupta 1982: 146]
- setosus* Scheerpeltz, *Paederus*
 [Scheerpeltz 1935: 630]
- seydeli* Fagel, *Paederus*
 [Fagel 1959: 56, 1965: 155]
- sharpi* Cameron, *Paederus*
 [Cameron 1914: 539, 1925b: 36, 1931a: 38, Scheerpeltz 1933: 1228]
- signaticornis* Sharp, *Paederus*
 [Sharp 1886: 613, Bernhauer & Schubert 1912: 211, Blackwelder 1939: 112, 1944: 123, Scheerpeltz 1965a: 215]
- signiventris* (Smetana), *Paederus*, *Uncopaederus*
 [Smetana 1962: 45, Korge 1969: 60, Tikhomirova 1973: 175, Coiffait 1978d: 387, 1982c: 67]
- sijthoffi* Bernhauer, *Paederus*
 [Bernhauer 1915e: 219, Scheerpeltz 1933: 1228, 1935: 617, Cameron 1936: 42]
- sikorae* auct.
 [Fauvel in litt., Fagel 1958a: 457, = *dilutipes* Fauvel]
- simillimus* Bernhauer, *Paederus*
 [Bernhauer 1927a: 109, Scheerpeltz 1933: 122, Fagel 1958a: 272, 1966: 258]
- simsoni* Blackburn, *Paederus*
 [Blackburn 1894: 92, 1899: 22, Lea 1904: 63, 1923: 14, Bernhauer & Schubert 1912: 211, Blackwelder 1939: 112]
- sinensis* Scheerpeltz
 [Scheerpeltz 1957a: 461, replacement name for *chinensis* Bernhauer 1934, ssp. of *fuscipes* Curtis]
- sjoestedti* Bernhauer, *Paederus* (*Pseudopaederus*)
 [Bernhauer 1920c: 9, Lea 1923: 14, Scheerpeltz 1933: 1228]
- smithi* Bernhauer
 [Bernhauer 1937b: 300, Fagel 1958a: 457, = *coerule-scens* Erichson]
- socius* Bernhauer, *Paederus* (*Paederus*)
 [Bernhauer 1916: 29, Wu 1937: 329, Scheerpeltz 1933: 1228, 1957a: 462]
- socius* Bernhauer
 [Bernhauer 1927a: 117, homonym, 1929: 123, = *fratellus* Bernhauer replacement name, Scheerpeltz 1933: 1223, Fagel 1958a: 142, = *arrowi* Bernhauer]
- socius* auct.
 [Bernhauer in coll., Fagel 1958a: 141, = *rufofasciatus* Bernhauer]
- solidus* Sharp, *Paederus*
 [Sharp 1876a: 284, Bernhauer & Schubert 1912: 211, Blackwelder 1944: 123, Scheerpeltz 1960: 81]
- sondaicus* Fauvel, *Paederus* (*Paederus*)
 [Fauvel 1895: 232, replacement name for *javanus* Erichson, 1904a: 53, Bernhauer & Schubert 1912: 211, Wendeler 1926d: 328, Cameron 1925b: 37, 1930b: 332, 1931a: 37, 1936: 42, 1949b: 467, Scheerpeltz 1933: 1228, Blackwelder 1939: 112, Scheerpeltz 1935: 609,

- 1957a: 462, 1965b: 103, Shibata 1973: 44, Biswas & Sen Gupta 1982: 153]
- sordidus* sic
[Sharp 1886: 616, misspelling of *solidus*]
- sparsicollis* Fagel, *Pachypaederus*
[Fagel 1958a: 78, 1959: 37]
- sparsus* Fauvel, *Paederus*
[Fauvel 1878b: 516, Blackburn 1899: 22, Lea 1904: 63, 1923: 14, Bernhauer & Schubert 1912: 211]
- spectabilis* Bernhauer
[Bernhauer 1915a: 116, Scheerpeltz 1933: 1228, Fagel 1958a: 351, ssp. of *duplex* Eppelsheim]
- speculator* Dahl
[Dahl 1823: 17, *nomen nudum*, Dejean 1833: 65, = *littoralis* Gravenhorst, Gemminger & Harold 1868: 627, Fauvel 1869b: 491]
- speculicollis* auct.
[Scheerpeltz *in litt.*, Fagel 1958a: 117, = *vulneratus* Bernhauer]
- speculifrons* Scheerpeltz, *Paederus*
[Scheerpeltz 1935: 613]
- steeleanus* Bernhauer
[Bernhauer 1937a: 294, Fagel 1958a: 300, = *sudanensis* Bernhauer]
- steelei* Bernhauer
[Bernhauer 1939b: 255, Bernhauer 1942b: 354, Fagel 1958a: 211, = *hintzi* Bernhauer; Fagel *loc. cit.* pointed out that the species was named for a Miss Steele and should therefore have been named *steeleae*: this emendation should be made, under Article 32c (ICZN), if the name is removed from synonymy]
- stenopterus* Lea, *Paederus*
[Lea 1923: 15, Scheerpeltz 1933: 1228]
- stigmula* auct.
[Fauvel *in litt.*, Quedenfeldt 1881: 292, = *fauveli* Quedenfeldt]
- straeleni* (Fagel), *Paederus*, *Oreopaederus*
[Fagel 1954a: 942, 1958a: 102, 1959: 58]
- straelenianus* Fagel, *Paederus*
[Fagel 1958a: 286, 1966: 261]
- stricticornis* (Fagel), *Paederus*, *Oreopaederus*
[Fagel 1954a: 953, 1958a: 102]
- stricticornis dubiosus* (Fagel)
[Fagel 1954a: 954, 1958a: 102]
- strictus* auct.
[Fauvel *in litt.*, Baudi 1869: 394, = *strictus* Baudi]
- strictus* Baudi
[Baudi 1869: 394, = *littoralis* Gravenhorst, Fauvel 1873: 331, Bernhauer & Schubert 1912: 207, Scheerpeltz 1957a: 465, 1968: 48, Legosz-Owsianna 1963: 349, Coiffait 1982c: 58]
- studti* Wendeler, *Paederus*
[Wendeler 1931: 38, Scheerpeltz 1933: 1228, 1935: 626]
- subcoriaceus* Cameron, *Paederus*
[Cameron 1942b: 840]
- subnitidicollis* (Bernhauer), *Paederus*, *Oreopaederus*
[Bernhauer 1934c: 232, var. of *testaceus* Bernhauer, Fagel 1954a: 937, species, 1958a: 101, 1960b: 197]
- subnitidus* Bernhauer, *Paederus*
[Bernhauer 1905b: 181, Bernhauer & Schubert 1912: 211, Blackwelder 1944: 123]
- subopacus* Bernhauer, *Paederus*
[Bernhauer 1915a: 117, Scheerpeltz 1933: 1228, Fagel 1958a: 435, 1966: 269]
- subtilis* Dahl
[Dahl 1823: 17, *nomen nudum*, Dejean 1833: 65, = *Rugilus*]
- sudanensis* Bernhauer, *Paederus*
[Bernhauer 1927a: 113, Scheerpeltz 1933: 1228, 1957a: 474, Fagel 1958a: 300, 1966: 259, Gaedike 1981: 223]
- sulawesi* Frank, *Paederus*
[here, *nom. nov.* for *melanocephalus* Heller, *nec* Fabricius]
- sulcicollis* auct.
[Kirby *ms.*, Stephens 1829: 288, = *Sunius*]
- sumbaensis* Scheerpeltz, *Paederidus*
[Scheerpeltz 1957b: 263]
- superbus* Bernhauer, *Paederus*
[Bernhauer 1912c: 474, Bernhauer & Schubert 1912: 211, Fagel 1958a: 360, 1960a: 124, 1966: 270]
- superbus meruensis* Fagel
[Fagel 1958a: 362, species, 1960a: 125, ssp., 1966: 270]
- sutteri* Scheerpeltz, *Diplopaederus*
[Scheerpeltz 1957b: 272]
- syriacus* Reitter
[Reitter 1889: 170, 1890: 14, = *mesopotamicus* Eppelsheim, Bernhauer & Schubert 1912: 208, Scheerpeltz 1957a: 459, Gaedike 1981: 223, Coiffait 1982c: 51]
- sythoffi* sic
[Last 1985: 335, misspelling of *sijthoffi*]
- szechuanus* sic
[Scheerpeltz 1957a: 475, misspelling of *szechuanus*]
- szechuanus* (Chapin), *Gnathopaederus*, *Paederus*
(*Gnathopaederus*)
[Chapin 1927: 76, Scheerpeltz 1933: 1228, Wu 1937: 329, Blackwelder 1939: 112, Scheerpeltz 1957a: 475]

T

- tambulensis* Last, *Paederus*
[Last 1985: 335]
- tamulus* Erichson, *Paederus* (*Eopaederus*)
[Erichson 1840: 661, Kraatz 1859a: 153, Gemminger & Harold 1868: 628, Fauvel 1869b: 491, 1903: 153, Bernhauer 1902: 38, Bernhauer & Schubert 1912: 211, Cameron 1925b: 37, 1930: 332, 1931a: 43, 1936: 43, 1949b: 467, Wu 1937: 329, Blackwelder 1939: 112, Scheerpeltz 1933: 1228, 1935: 609, 1957a: 459, 1957b: 229, 1976: 82, Shibata 1973: 43, 1977: 23, Coiffait 1982b: 234]
- tandalensis* Bernhauer, *Paederus*
[Bernhauer 1927a: 109, Scheerpeltz 1933: 1228, Fagel 1958c: 239, 1966: 260]
- taprobanus* Cameron, *Paederus*
[Cameron 1931a: 61, Scheerpeltz 1933: 1228]
- tempestivus* Erichson, *Paederus*
[Erichson 1840: 658, Gemminger & Harold 1868: 628, Bernhauer & Schubert 1912: 211, Blackwelder 1944: 123, Scheerpeltz 1965: 218]
- tempestivus auct.*
[Erichson 1840: 658 *partim*, Sharp 1886: 614, = *yucateca* Sharp, Bernhauer & Schubert 1912: 212, Blackwelder 1939: 112, Scheerpeltz 1965a: 216]
- tenuicornis auct.*
[Dahl *in litt.*, Gemminger & Harold 1868: 626, = *brevicornis* Lacordaire]
- tenuicornis* Fauvel
[Fauvel 1877a: 224, 1878b: 517, Blackburn 1899: 23, = *angulicollis* MacLeay, Bernhauer & Schubert 1912: 211, species, Lea 1923: 14, = *angulicollis* MacLeay]
- tenuis* Fagel, *Paederus*
[Fagel 1958a: 305, 1966: 259]
- terminatus auct.*
[Knoch *in coll.*, Gravenhorst 1802: 55, = *Lathrobium*]
- termitophilus* (Wasmann), *Paederus*, *Oreopaederus*
[Wasmann 1911: 101, Scheerpeltz 1933: 1229, Fagel 1954a: 952, 1958a: 102]
- testaceitarsis* Sharp, *Paederus*
[Sharp 1886: 610, Bernhauer & Schubert 1912: 211, Blackwelder 1944: 123]
- testaceopiceus* Bernhauer, *Paederus*
[Bernhauer 1915c: 292, Scheerpeltz 1933: 1229, Fagel 1958a: 324, 1959: 62, 1963: 348, 1966: 264]
- testaceus* (Paykull)
[Paykull 1789: 28, Olivier 1795: 5, Paykull 1800: 429, Gravenhorst 1802: 44, 1806: 138, Olivier 1812: 628, Stephens 1829: 286, = *Lathrobium*]
- testaceus* (Bernhauer), *Paederus*, *Oreopaederus*
[Bernhauer 1934c: 231, 1934d: 207, Fagel 1954a: 932, 1958a: 101]
- testaceus auct.*
[Cameron 1950b: 26, misidentification, Fagel 1954a: 956, = *vulcanus* Fagel]
- texanus* (Casey)
[Casey 1905: 65, 1910: 184, = *saginitus* Casey, Bernhauer & Schubert 1912: 210, Leng 1920: 101, Blackwelder 1939: 112, Moore & Legner 1975: 134]
- thoracicus* (Fourcroy)
[Fourcroy 1785: 170, Kraatz 1857: 731, = *ruficollis* Fabricius, Gemminger & Harold 1868: 628, Fauvel 1873: 336, = *rubrothoracicus* Goeze, Mulsant & Rey 1878: 246, = *ruficollis* Fabricius, Bernhauer & Schubert 1912: 210, = *rubrothoracicus* Goeze, Jarrige 1944: 7, Legosz-Owsianna 1963: 327]
- thoracicus* Erichson
[Erichson 1840: 663, homonym, Gemminger & Harold 1868: 628, Sharp 1886: 612, Bernhauer & Schubert 1912: 209, = *neotropicus* Bernhauer & Schubert, Blackwelder 1944: 123]
- thoracicus auct.*
[Marshall 1878: xxx, misidentification, Leng & Mutchler 1917: 198, Blackwelder 1943: 323, = *homonymus* Blackwelder, 1944: 123]
- tibestiensis* Fagel, *Paederidus*
[Fagel 1958a: 151]
- tibetanus* Cameron, *Paederus*
[Cameron 1928: 560, 1931a: 55, Wu 1937: 329, Scheerpeltz 1933: 1229, 1957a: 474, Coiffait 1982b: 234, 282]
- togoensis auct.*
[Bernhauer *in litt.*, Fagel 1958a: 227, = *annexus* Ep-pelsheim]
- togoensis* Fauvel
[Fauvel *in litt.*, Fagel 1958a: 227, = *annexus* Ep-pelsheim]
- tongyai* Bernhauer, *Paederus*
[Bernhauer 1939d: 115]
- toxopei* Wendeler, *Paederus*
[Wendeler 1926d: 329, Scheerpeltz 1933: 1229, 1935: 619, Gaedike 1981: 224]
- transvaalensis* Fagel, *Paederus*
[Fagel 1965: 148, 1966: 258]
- trapeziceps* Scheerpeltz
[Scheerpeltz 1957a: 463, 1968: 47, Smetana 1959: 204, 1964: 165, Lohse 1964: 134, Szujewski 1965a: 83, = *balcanicus* Koch, Tikhomirova 1973: 175, Coiffait 1982c:

- 56, var. of *balcanicus* Koch, Bohác 1985: 370, = *balcanicus* Koch]
- tricolor* (Fabricius)
[Fabricius 1787: 221, 1792: 537, 1801: 609, Paykull 1789: 23, 1800: 378, Gravenhorst 1802: 46, Erichson 1839: 428, = *Xantholinus*]
- tricolor* Erichson, *Paederus*
[Erichson 1840: 663, Reiche 1847: 281, Gemminger & Harold 1868: 628, Bernhauer & Schubert 1912: 211, Leng & Mutchler 1914: 404, Blackwelder 1939: 112, 1943: 323, = *homonymus* Blackwelder, unnecessary replacement name, 1944: 123]
- tricoloricornis* Fagel, *Oreopaederus*
[Fagel 1958a: 99]
- tridens* Sharp, *Paederus*
[Sharp 1876a: 285, Bernhauer & Schubert 1912: 211, Plavilstschikov 1929: 29, Blackwelder 1944: 123]
- trimerus* Fauvel, *Paederus*
[Fauvel 1904c: 294, Bernhauer & Schubert 1912: 211]
- tristis* Dejean
[Dejean 1833: 65, *nomen nudum*]
- tristis* Bernhauer, *Paederus*
[Bernhauer 1915f: 183, Scheerpeltz 1933: 1229]
- tropicus* Bernhauer, *Paederus*
[Bernhauer 1912c: 473, Bernhauer & Schubert 1912: 211, Fagel 1958a: 441, 1966: 269]
- tsaratananus* Jarrige, *Paederus*
[Jarrige 1970: 33]
- tshibindensis* (Fagel), *Paederus*, *Oreopaederus*
[Fagel 1954a: 966, 1958a: 102]
- tumbaensis* Fagel, *Oreopaederus*
[Fagel 1958a: 94, 1961a: 265]
- tumidicollis* Gerstaecker, *Paederus*
[Gerstaecker 1867: 28, 1873: 81, Gemminger & Harold 1868: 628, Fauvel 1904b: 290, Bernhauer & Schubert 1912: 211, Fagel 1958a: 347, 1966: 269]
- turneri* sic
[Fagel 1958a: 175, misspelling of *turneris*, 1966: 211]
- turneris* Cameron, *Paederus*
[Cameron 1950c: 186]
- turrialbanus* Wendeler, *Paederus*
[Wendeler 1927: 2, Scheerpeltz 1933: 1229, Blackwelder 1944: 123]
- tweedensis* Blackburn, *Paederus*
[Blackburn 1899: 22, Lea 1904: 63, Bernhauer & Schubert 1912: 211, Bernhauer 1920c: 9, Lea 1923: 14]
- U**
- ugandae* Bernhauer
[Bernhauer 1912c: 473, Fagel 1958a: 214, = *pedestris* Gerstaecker, 1960a: 120]
- ugandensis* Fagel, *Paederus*
[Fagel 1958a: 396, 1966: 267]
- uganensis* Fagel, *Paederus*
[Fagel 1958a: 367, 1966: 267]
- ulindiensis* Fagel, *Paederus*
[Fagel 1958a: 385, 1960b: 204, 1966: 267]
- uluguruensis* Fagel, *Paederus*
[Fagel 1960a: 129, 1966: 265]
- umbrosus* (Fagel), *Paederus*, *Oreopaederus*
[Fagel 1954a: 934, 1958a: 101]
- Uncopaederus* Korge
[Korge 1969: 60, *genus*, type *Paederus signiventris* Smetana, Coiffait 1982c: 67]
- ungandae* sic
[Cameron 1949a: 317, misspelling of *ugandae*]
- unicolor* Cameron, *Paederus*
[Cameron 1942a: 88]
- upembaensis* Fagel, *Oreopaederus*
[Fagel 1958a: 95]
- uruguayensis* Bernhauer, *Paederus*
[Bernhauer 1912a: 33, Scheerpeltz 1933: 1229, Blackwelder 1944: 123]
- usagarae* Bernhauer, *Paederus*
[Bernhauer 1908b: 105, Bernhauer & Schubert 1912: 211, Fagel 1958a: 334, 1966: 263]
- usambarae* (Fauvel), *Paederus*, *Paederidus*
[Fauvel 1904b: 290, 1907: 23, Bernhauer & Schubert 1912: 211, Fagel 1958a: 132, 1960a: 118]
- usambarae usambarae* (Fauvel)
[Fauvel 1904b: 290, 1907: 23, Fagel 1960a: 118]
- usambarae xeuxes* (Tottenham)
[Tottenham 1953: 490, Fagel 1958a: 123, = *usambarae* (Fauvel), 1960a: 118, ssp. of *usambarae*]
- usambaricus* Schubert, *Paederus*
[Schubert 1906: 375, Bernhauer & Schubert 1912: 211, Fagel 1958a: 442, 1966: 268]
- usambicus* sic
[Eichelbaum 1913: 125, misspelling of *usambaricus*]
- ussuriensis* Kirschenblatt, *Paederus* (*Harpopaederus*)
[Kirschenblatt 1932: 219, Scheerpeltz 1957a: 460, Smetana 1962: 50, Tikhomirova 1973: 175]
- usticollis* Fauvel
[Fauvel 1907: 23, var. of *scabripennis* Fauvel, Bernhauer & Schubert 1912: 211, Peyerimhoff 1931: 29, Bernhauer 1931c: 515, Blackwelder 1939: 112, Cameron 1950b: 25, species, Fagel 1956: 200, 1958a: 116, var. of *scabripennis* Fauvel, 1960a: 117]

usticollis auct.

[Fauvel *in litt.*, Fagel 1958a: 146, = *collaris* Boheman]

ustus LeConte, *Paederus*, *Leucopaederus*, *Paederus*

[LeConte 1858: 62, Gemminger & Harold 1868: 628, Casey 1905: 67, Bernhauer & Schubert 1912: 211, Leng 1920: 101, Cameron 1931: 33, Blackwelder 1939: 112, 1944: 123, Moore & Legner 1975: 134]

V*vagus* Wendeler

[Wendeler 1933: 57, Fagel 1958a: 307, 1959: 59, = *sabaicus* Erichson, 1960b: 203]

variceps Kraatz, *Paederus*

[Kraatz 1859a: 152, Gemminger & Harold 1868: 628, Bernhauer & Schubert 1912: 211, Cameron 1925b: 37, 1931a: 53, Scheerpeltz 1933: 1229, Gaedike 1981: 226]

variiceps sic

[Cameron 1931a: 53, misspelling of *variceps*]

variicornis Fauvel, *Paederus*

[Fauvel 1903a: 154, 1904a: 54, Bernhauer & Schubert 1912: 211, Cameron 1925b: 37, 1931a: 49, Scheerpeltz 1933: 1229, Biswas & Sen Gupta 1982: 152]

vastus Scheerpeltz, *Paederus* (*Oreinopaederus*)

[Scheerpeltz 1976: 112]

velox Sharp, *Paederus*

[Sharp 1886: 611, Bernhauer & Schubert 1912: 211, Blackwelder 1944: 123]

ventricosus Gautier

[Gautier 1861b: xxxvi, *nomen nudum*]

ventricosus Gautier

[Gautier 1862: 77, Fauvel 1864: 13, = '*brevipennis* Erichson', Gautier 1866: 114, = *baudii* Fairmaire, Marseille 1866: vi, = *lusitanicus* Baudi, Fauvel 1866: 24, = *baudii* Fairmaire, 1873: 329, Gemminger & Harold 1868: 626, Kraatz 1867: 415, Mulsant & Rey 1878: 250, Reitter 1884: 44, Ganglbauer 1895: 538, Bernhauer & Schubert 1912: 204, Scheerpeltz 1957a: 460, Smetana 1962: 49, Coiffait 1982c: 64]

ventricosus sic

[Coiffait 1982c: 64, misspelling of *ventricosus*]

versicolor Fagel, *Paederus*

[Fagel 1956: 195, 1958a: 382, 1966: 266]

vestitus Gravenhorst

[Gravenhorst 1806: 140, Olivier 1812: 629, *nomen oblitum*]

vianai Bernhauer, *Paederus*

[Bernhauer 1939a: 242]

villiersi (Cameron), *Paederus*, *Pachypaederus*

[Cameron 1949a: 317, Fagel 1958a: 92, 1963: 342]

villiersi Fagel, *Paederidus*

[Fagel 1958a: 136]

violaceipennis Fagel, *Paederus*

[Fagel 1958a: 181, 1966: 271]

viridipennis Bernhauer, *Paederus*

[Bernhauer 1915a: 117, Scheerpeltz 1933: 1229, Fagel 1958a: 408, 1966: 269]

viridipennis Bernhauer

[Bernhauer 1931c: 574, homonym, Scheerpeltz 1933: 1229, = *cooperi* Scheerpeltz, replacement name, Fagel 1958a: 177, 1960a: 126]

vitiensis Fauvel

[Fauvel 1878b: 517, unjustified replacement name for *samoensis* Fauvel, 1879b: 84, Fairmaire 1881: 251, Bernhauer & Schubert 1912: 211]

vulcanicola (Fagel), *Paederus*, *Oreopaederus*

[Fagel 1954b: 237, repl. name for *vulcanus* Fagel, 1958a: 102]

vulcanicola nitidicollis (Fagel)

[Fagel 1954b: 957, 1958a: 102]

vulcanus Wendeler, *Paederus*

[Wendeler 1926b: 203, Scheerpeltz 1933: 1229, Blackwelder 1944: 123, Fagel 1954b: 237, 1958a: 102]

vulcanus Fagel

[Fagel 1954a: 956, homonym, 1954b: 237, 1958a: 102, = *vulcanicola* Fagel, replacement name]

vulcanus nitidicollis Fagel

[Fagel 1954a: 957, 1954b: 237, = *vulcanicola nitidicollis* Fagel]

vulcanus sic

[Fagel 1954b: 237, misspelling of *vulcanus*]

vulgaris Miller

[Miller 1852: 27, Kraatz 1857: 726, = *littoralis* Gravenhorst, Gemminger & Harold 1868: 627, Ganglbauer 1895: 538, Bernhauer & Schubert 1912: 207, Scheerpeltz 1933: 1225, 1968: 48, Legosz-Owsianna 1963: 349, Coiffait 1982c: 58]

vulneratus (Bernhauer), *Paederus*, *Paederidus*

[Bernhauer 1927a: 114, Scheerpeltz 1933: 1229, Last 1952: 91, Fagel 1958a: 117, 1961b: 266]

vulneratus nigrothoracicus Last

[Last 1952: 91, var. of *vulneratus*, Fagel 1958a: 117, 1961b: 266, here treated as ssp. under Article 45(g) (ICZN 1985)]

W*wabasensis* Last

[Last 1984: 109, var. of *hornabrooki* Last]

wadai Scheerpeltz, *Megalopaederus*
[Scheerpeltz 1957a: 472, Shibata 1977: 30, Watanabe
1986: 165]

wauensis Last, *Paederus*
[Last 1979: 124, 1985: 329]

weisei Schubert, *Paederus*
[Schubert 1906: 375, Bernhauer & Schubert 1912:
212, Fagel 1958a: 443, 1966: 268]

weisei auct.
[Fauvel *in litt.*, Fagel 1958a: 443, = *weisei* Schubert]

wendeleri Fagel, *Pachypaederus*
[Fagel 1958a: 77, Gaedike 1981: 228]

wenzeli Fagel, *Paederus*
[Fagel 1958a: 294, 1966: 258]

wilhelmi Bernhauer
[Bernhauer 1926: 5, Scheerpeltz 1933: 1229, Fagel
1958a: 225, = *amicus* Bernhauer, 1959: 47, 1960b:
203]

wilsoni Lea, *Paederus*
[Lea 1923: 16, Scheerpeltz 1933: 1229]

witteanus Bernhauer
[Bernhauer 1935b: 99, Cameron 1950b: 25, Fagel
1958a: 257, 1965: 155, = *capitalis* Bernhauer]

wittei Bernhauer
[Bernhauer 1933b: 293, Fagel 1958a: 121, =
schoutedeni Bernhauer]

wittei auct.
[Cameron *in litt.*, Fagel 1958a: 170, = *opacicollis*
Bernhauer]

X

xanthocerus Eppelsheim, *Paederus*
[Eppelsheim 1895b: 210, Fauvel 1904b: 290, Bern-
hauer & Schubert 1912: 212, Fagel 1956: 199, 1958a:
269, 1959: 50, 1963: 344, 1966: 258]

xeuxes (Tottenham)
[Tottenham 1953: 490, Fagel 1958a: 132, = *usambarae*
Fauvel, 1960a: 118, ssp. of *usambarae*]

Y

yangasanus sic
[Scheerpeltz 1933: 1229, misspelling of *yuangasanus*,
Blackwelder 1944: 123]

yaninensis (Fagel), *Paederus*, *Oreopaederus*
[Fagel 1956: 187, 1958a: 102]

yaoundensis Fagel, *Paederus*
[Fagel 1958a: 191, 1966: 263]

yuangasanus Wendeler, *Paederus*
[Wendeler 1931: 48]

yucateca Sharp, *Paederus*
[Sharp 1886: 614, Fauvel 1891: 100, Bernhauer &
Schubert 1912: 212, Blackwelder 1939: 112, 1944: 123,
Scheerpeltz 1965a: 216]

Z

zairensis Frank, *Paederus*
[here, *nom. nov.* for *Paederus orophilus* Fagel 1958a:
379, *nec* Fagel 1954a: 919]

zanzibaricus Fagel
[Fagel 1958a: 117, ssp. of *scabripennis* Fauvel]

zayasi Sanderson, *Paederus*
[Sanderson 1967: 61]

zulu Fagel, *Paederus*
[Fagel 1965: 150, 1966: 259]

Synonymy

in genus-group names

Geopaederus and Poederomorphus. *Geopaederus* was proposed by Gistel (1848) as a replacement name for *Paederus* in the belief that *Paederus* was preoccupied by *Paederia* Linnaeus (plant kingdom). *Geopaederus* is an objective junior synonym of *Paederus*. *Poederomorphus* was proposed by Gautier (1862) as the name of a new genus with *Poederomorphus pedoncularius* as type species. *Poederomorphus pedoncularius* was relegated to synonymy with *Paederus littoralis* by Marseul (1866); thus *Poederomorphus* is a subjective junior synonym of *Paederus*.

Paederillus and Leucopaederus. Casey (1905) considered the species of America north of Mexico to belong to three distinct genera (*Paederus*, *Paederillus* and *Leucopaederus*) but he failed to contrast the distinguishing characters unambiguously. Adult *Paederillus* and *Leucopaederus* have a truncate, edentate labrum with median emargination, while adult *Leucopaederus* are entirely testaceous (Casey 1905). Blackwelder (1939, 1952) relegated *Paederillus* and *Leucopaederus* to synonymy with *Paederus* s.str. Although Fagel (1958a) thought they might prove to be subgenera of *Paederus*, he did not resurrect them.

Paederognathus and Neopaederus. Wendeler (1927) proposed the subgenus *Gnathopaederus* for *Paederus turrialbanus*. However, on finding the name preoccupied by *Gnathopaederus* Chapin, he replaced it with *Paederognathus* (Wendeler 1928a). Characters were stated to be presence of a carina between the metacoxae, labrum deeply emarginate, and mandible without a median tooth (Wendeler 1928a). Blackwelder (1939) proposed a new subgeneric classifica-

tion for *Paederus* in which macropterous adults with elytra of normal size were assigned to subgenus *Paederus*, whereas wingless adults with reduced elytra were assigned to subgenus *Gnathopaederus* (mandible with an additional tooth), or to the new subgenus *Neopaederus* (mandible without additional teeth); *Paederidus*, *Paederillus*, *Leucopaederus*, and *Pseudopaederus* were relegated to synonymy with *Paederus* s. str. Later, Blackwelder (1952) relegated *Paederognathus* to synonymy with *Paederus* s. str. However, Fagel (1958a) suggested that *Paederognathus* may prove to be a useful subdivision of *Paederus*. Furthermore, because he considered Blackwelder's (1939) arrangement highly artificial due to the probably polyphyletic origins of aptery and elytral reduction, he relegated *Neopaederus* to synonymy with *Paederus*. *Paederognathus* has not been studied by later authors and remains a synonym of *Paederus*.

Standing names of genera distinguished from *Paederus*

***Paederidus*.** Differentiated from *Paederus* as a subgenus by Mulsant & Rey (1878), *Paederidus* was raised to generic status by Casey (1905), reduced to a subgenus by Bernhauer & Schubert (1912), placed into synonymy with *Paederus* s. str. by Blackwelder (1939), considered again a subgenus by Jarrige (1944), and re-elevated to generic rank by Scheerpeltz (1957a), followed by later authors. It currently includes 44 species. Adults can be distinguished from those of *Paederus* and some related genera by the key below.

***Allopaederus*, *Madecapaederus*, *Oreopaederus* and *Pachypaederus*.** Described by Fagel (1958a), these four genera are based on Afrotropical (including Malagasy) species. They currently contain 3, 11, 19, and 55 species respectively. Their adults are differentiated from those of *Paederidus* and *Paederus* in the following key translated and modified from Fagel (1958a).

Generic key to Afrotropical/Malagasy adult *Paederus* (*sensu lato*)

1. Mentum with a convexity (tooth, tubercle or carina) 2
- 1'. Mentum without a convexity 3
- 2(1). Mentum with a median tooth or tubercle. Median lobe of aedeagus with ventral orifice partially obscured by internal structure hinged by a membrane and forming a valve; parameres styliform, symmetrical, fused only at base.

2. Large, apterous adults of the Afrotropical fauna proper *Allopaederus*
Mentum with a fine, median, longitudinal carina. Median lobe of aedeagus indented, with ventral orifice partially obscured by free internal structure; parameres either styliform and highly asymmetrical or finer and more symmetrical. Small, apterous adults of the Malagasy fauna *Madecapaederus*
- 3(1'). Prosteronum strongly carinate. Gular sutures narrowly separated, parallel for much of their length. Aedeagus like that of *Allopaederus*. Most adults large and apterous or brachypterous, a few winged and smaller *Pachypaederus*
- 3'. Prosteronum not or feebly carinate 4
- 4(3'). Gular sutures widely separated, more or less parallel. Aedeagus like that of *Pachypaederus* but internal structure not hinged by a membrane so not forming a valve. Adults small or of medium size, apterous or brachypterous *Oreopaederus*
- 4'. Gular sutures strongly convergent over part of their length, parallel or not. Median lobe of aedeagus with median orifice and with dorsal and ventral plate; parameres of several forms but not styliform and in some species asymmetrical, fused to the median lobe for part of their length 5
- 5(4'). Basal abdominal sternite (sternite III) with strong carina between posterior coxae. In most species mandible near midpoint of its length with bicuspid tooth (but in some species with a pair of contiguous teeth), in some species with an additional dorsal tooth. Most adults winged, but some apterous or brachypterous *Paederus*
- 5'. Basal abdominal sternite (sternite III) without carina between posterior coxae, at most with a small lobe. Mandible with two large, well-separated teeth near midpoint of its length. Adults slender, winged *Paederidus*

***Ctenopaederus*.** Fagel (1958a) based this monotypic genus on *Paederus andrewesi* from India. Adults differ from those of related genera in having serrate tarsal claws (Fagel 1958a).

***Diplopaederus*, *Eupaederus*, *Lobopaederus*, *Megalopaederus*, *Oncopaederus*, *Parameropaederus* and *Uncopaederus*.** Scheerpeltz (1957a) erected *Megalopaederus* for three species from Japan. Shibata (1973, 1977) added four more species and extended the range

to Taiwan. Scheerpeltz (1957b) based *Parameropaederus* on a species from southern Europe, *Lobopaederus* on a species from southern Europe/northern Africa, *Diplopaederus* on two species from South America, Oceania and Indonesia, *Eupaederus* on two species from Indonesia, and *Oncopaederus* on another Indonesian species. Korge (1969) erected *Uncopaederus* for a species from Europe. All of these genera were differentiated from *Paederus* on the basis of aedeagal structure (Scheerpeltz 1957a,b, Korge 1969).

Generic concepts.

The generic concepts of Fagel (1958a) contrast with those of Scheerpeltz (1957a,b) and Korge (1969) and later authors (Coiffait 1982c, Bohác 1985). Whereas Fagel (1958a) was unwilling even to name new species on the basis of aedeagal structure alone, other authors erected genera solely on the basis of aedeagal structure. There are thus three current concepts of *Paederus*: (1) a concept unchanged since the 19th century of *Paederus* (*sensu lato*), still employed by some authors (e.g., Last 1979-1985) in which there is no attempt to define segregates from or within *Paederus*, (2) the concept of Fagel (1958a-1973) in which genera are separated from *Paederus* only after consistent external distinguishing structures are recognized, and (3) the concept of Scheerpeltz-Korge-Coiffait in which larger numbers of genera are recognized, some of them based only on differences in aedeagal structure. Thirteen genera stand apart from *Paederus* in the literature; the species attributed to them are listed below.

Paederidus Mulsant & Rey: *albipilis* (Solsky), *albopubescentis* (Bernhauer), *algericus* (Motschulsky), *arabicus* (Bernhauer), *arrowi* (Bernhauer), *balfourbrowni* Fagel, *burgeoni* (Bernhauer), *cariniventris* (Cameron), *chalybeus* (Erichson), *coelestinus* (Bernhauer), *coerulescens* (Erichson), *collaris* (Boheman), *colonus* (Wendeler), *cribricollis* (Bernhauer), *crinitus* (Fagel), *erythraeanus* Fagel, *freyi* Fagel, *funebri* Fagel, *gracilior* Fagel, *inconspicuus* (Cameron), *koreanus* Scheerpeltz, *luctuosus* (Klug), *luluensis* (Bernhauer), *martensi* Coiffait, *memnonius* (Erichson), *montanellus* (Bernhauer), *niger* (Bernhauer), *opacus* (Bernhauer), *parcepunctatus* (Bernhauer), *permixtus* Fagel, *perrieri* (Fauvel), *pumilus* (Wendeler), *punctiventris* Scheerpeltz, *ruficollis* (Fabricius), *rufofasciatus* (Bernhauer), *saegeri* Fagel, *scabripennis* (Fauvel), *schoutedeni* (Bernhauer), *senegalensis* Fagel, *sumbuensis* Scheerpeltz, *tibestiensis* Fagel, *usambarae* (Fauvel), *villiersi* Fagel, *vulneratus* (Bernhauer).

Parameropaederus Scheerpeltz: *lusitanicus* (Aube).

Lobopaederus Scheerpeltz: *meridionalis* (Fauvel).

Megalopaederus Scheerpeltz: *alutithorax* (Bernhauer), *flavoterminalis* (Cameron), *formosanus* (Adachi), *kosempoensis* (Bernhauer), *kurosawai* Watanabe, *lewisi* (Cameron), *poweri* (Sharp), *wadai* Scheerpeltz.

Eupaederus Scheerpeltz: *buehleri* Scheerpeltz, *cyanipennis* (Guerin).

Diplopaederus Scheerpeltz: *handschini* (Scheerpeltz), *sutteri* Scheerpeltz.

Oncopaederus Scheerpeltz: *brachypterus* Scheerpeltz.

Allopaederus Fagel: *aereus* Fagel, *filipes* (Fagel), *mirandus* Fagel.

Ctenopaederus Fagel: *andrewesi* (Fauvel).

Madecapaederus Fagel: *apterus* (Fauvel), *cyanescens* Jarrige, *descarpentriensis* Jarrige, *dilutipes* (Fauvel), *dilutus* Jarrige, *dolosus* Jarrige, *fugeli* Jarrige, *forficularius* (Fauvel), *metallicus* (Fauvel), *pauliani* Jarrige, *rufipes* (Fauvel).

Oreopaederus Fagel: *alinderi* (Wendeler), *allardianus* Fagel, *angolanus* Fagel, *atricolor* (Fagel), *atropygus* Fagel, *basilewskyi* (Fagel), *bernhaueri* (Fagel), *brachycephalus* (Bernhauer), *brunnescens* (Fagel), *brunneus* (Bernhauer), *celisianus* Fagel, *fallax* (Fagel), *falsus* Fagel, *flavoides* (Fagel), *flavus* (Fagel), *furcifer* (Fagel), *furciferoides* (Fagel), *insidiosus* Fagel, *intolerans* Fagel, *jarrigei* (Fagel), *kahuziensis* (Fagel), *kiyambiensis* Fagel, *leleupi* (Fagel), *leleupianus* Fagel, *luberensis* Fagel, *lubukensis* (Fagel), *machadoi* Fagel, *manyemensis* Frank, *meersmanae* Fagel, *mendax* (Fagel), *mikenensis* (Fagel), *minor* (Bernhauer), *minutissimus* (Bernhauer), *minutus* Fagel, *mulengensis* Fagel, *ngoviensis* Fagel, *nigricans* Fagel, *nitidus* (Fagel), *nyakagerensis* Fagel, *nyalengwensis* Fagel, *obscuripennis* (Fagel), *pseustes* (Fagel), *punctipennis* (Fagel), *rugegensis* (Fagel), *straeleni* (Fagel), *stricticornis* (Fagel), *subnitidicollis* (Bernhauer), *termitophilus* (Wasmann), *testaceus* (Bernhauer), *tricoloricornis* Fagel, *tshibindensis* (Fagel), *umbrosus* (Fagel), *upembaensis* Fagel, *vulcanicola* (Fagel), *yaninensis* (Fagel).

Pachypaederus Fagel: *caheni* Fagel, *confusus* Fagel, *crassus* (Boheman), *delagoanus* (Bernhauer), *gurambanus* Fagel, *kaboboensis* Fagel, *lasti* Fagel, *leleupi* Fagel, *minutus* Fagel, *natalensis* (Last), *newtoni* (Last), *nigriventris* Fagel, *obscuricollis* (Bernhauer), *puncticollis* (Bernhauer), *reductus* Fagel, *rhodesianus* (Bernhauer), *sparisicollis* Fagel, *villiersi* (Cameron), *wendeleri* Fagel.

Uncopaederus Korge: *signiventris* (Smetana).

Relatives of *Paederus* (*sensu lato*)

Concepts of the tribe Paederini have been restricted after Blackwelder (1952) excluded from it only the tribe Pinophilini of the subfamily Paederinae. Jeanne & Jarrige (1949) grouped within Paederini only those genera in which adults have the last maxillary palpomere short and squamiform, and the aedeagal parameres well developed and piliferous; Fagel (1958a) accepted this diagnosis and added that the internal margin of the anterior tibia bears a ctenidium, and the posterior border of abdominal sternite VIII of the male is very deeply and narrowly emarginate, almost dividing the sternite. Fagel (1958a) recognized two subtribes within Paederini: Dolicaei (*Dolicaon* Laporte and allies) and Paederi (containing only *Paederus* (*sensu lato*)). Fagel's (1958a) concept of the subtribe Paederi is exactly that of Casey (1905) who even used the same subtribal name.

Coiffait (1982c) reverted to a broader definition of Paederini [the same as that by Blackwelder (1952)] which included subtribe Paederi (both Paederi and Dolicaei of Fagel) and others. After studying the New World species of genera related to *Dolicaon*, Herman (1981) showed a close relationship (sister group) of two subtribes: Dolicaonina (*Dolicaon* and allies, i.e., the Dolicaei of Fagel) and Paederina (*Paederus* (*sensu lato*)), thus supporting Fagel's (1958a) concept. In addition, he found that these two subtribes together have a sister group relationship to the subtribe Cryptobiina (*Cryptobium* Mannerheim and allies) because all adults of these taxa have a gland on the anterior margin of abdominal sternite IV. If the Fagel-Herman concept is accepted, then *Paederus* (*sensu lato*) may conveniently be called subtribe Paederi (or Paederina); its closest allies are *Dolicaon* and relatives in which the the 4th anterior tarsomere of the adult is entire (cf. apically emarginate or bilobed).

Standing names of subgenera distinguished from *Paederus*

***Pseudopaederus* and *Gnathopaederus*.** Bernhauer (1915b) founded *Pseudopaederus* on *Paederus nigerrimus* and *P. pallidulus* from India, with *P. nigerrimus* as typus subgeneris (Blackwelder 1939). Bernhauer (1915b) stated that distinguishing characters are a short, square head, a short, round, strongly developed prothorax, and a generally compact body. Although an Australian species was added by Bernhauer (1920c) and two more species from India by Biswas & Sen Gupta (1982), there has been no further discussion of subgeneric characters of *Pseudopaederus*. Chapin (1927) described *Gnathopaederus* as a genus, with *G. sze-*

chuanus as its type species; characters were stated to be gular sutures confluent or nearly so at base of head, mandible with prominent dorsal tooth, 4th tarsomere bilobed, elytral punctation coarse and sparse. Scheerpeltz (1933) placed *Gnathopaederus* as a subgenus of *Paederus*, where it has been retained by subsequent authors without further study and without incorporation of additional species.

***Dioncopaederus*, *Eopaederus*, *Harpopaederus*, *Heteropaederus*, *Oedopaederus*, *Anomalopaederus*, *Oreinopaederus* and *Nepalopaederus*.** *Eopaederus*, *Harpopaederus*, *Heteropaederus*, *Dioncopaederus* and *Oedopaederus* were named and distinguished from *Paederus* s.str. by Scheerpeltz (1957a) on the basis of aedeagal structure, as were *Anomalopaederus* (Scheerpeltz 1966), *Oreinopaederus* and *Nepalopaederus* (Scheerpeltz 1976).

Subgeneric concepts.

Subgenera have been attributed to the genus *Paederus* only, and not to the other genera which have been segregated. It is questionable whether subgeneric separation should be pursued until generic concepts have become more stable. Nevertheless, ten subgenera stand apart from *Paederus* s.str. in the literature, and species attributed to them are listed below. There follows a listing of species attributed to *Paederus* s.str. (concept of Scheerpeltz and Coiffait), then of species (from the Afrotropical region) attributed to the genus *Paederus* (concept of Fagel), then of species (from various parts of the world) attributed to *Paederus* (*sensu lato*) and not assigned to any finer division, and finally the single fossil species is listed.

Pseudopaederus Bernhauer: *loebli* Biswas & Sen Gupta, *mussardi* Biswas & Sen Gupta, *nigerrimus* Bernhauer, *pallidulus* Scheerpeltz, *sjoestedti* Bernhauer.

Gnathopaederus Chapin: *gottschei* Kolbe, *szechuanus* Chapin.

Eopaederus Scheerpeltz: *balachowskyi* Jarrige, *basalis* Bernhauer, *caligatus* Erichson, *debilior* Eppelsheim, *kuluensis* Bernhauer, *ledouxi* Coiffait, *limnophilus* Erichson, *mesopotamicus* Eppelsheim, *nigricornis* Bernhauer, *tamulus* Erichson.

Harpopaederus Scheerpeltz: *baudii* Fairmaire, *brevipennis* Lacordaire, *latro* Smetana, *margelanicus* Bernhauer, *pseudobaudii* Aleksandrov, *schoenherri* Czwalińska, *ussuriensis* Kirschenblatt.

Heteropaederus Scheerpeltz: *alfierii* Koch, *bohemani* Fagel, *fuscipes* Curtis, *iliensis* Coiffait, *peregrinus* Erichson, *sabaesus* Erichson.

Dioncopaederus Scheerpeltz: *carpathicus* Wendeler, *ilsae* Bernhauer, *lenkoranus* Scheerpeltz, *littoralis* Gravenhorst, *pelikani* Reitter.

Oedopaederus Scheerpeltz: *chinensis* Bernhauer.

Anomalopaederus Scheerpeltz: *lacordairei* Perroud.

Oreinopaederus Scheerpeltz: *vastus* Scheerpeltz.

Nepalopaederus Scheerpeltz: *nepalensis* Bernhauer.

Paederus s.str., sensu Scheerpeltz and Coiffait: *balkanicus* Koch, *coxalis* Fauvel, *diamesus* Kirschenblatt, *japonicus* Bernhauer, *longiceps* Bernhauer, *melanurus* Aragona, *parallelus* Weise, *pilifer* Motschulsky, *riparius* (Linnaeus), *saengeri* Coiffait, *socius* Bernhauer, *sondaicus* Fauvel.

Paederus sensu Fagel: *abdominalis* Cameron, *altivagans* Fauvel, *alutaceus* Fagel, *amicus* Bernhauer, *angusticeps* Bernhauer, *Annexus* Eppelsheim, *aquaticus* Bernhauer, *aqualitis* Cameron, *arrowianus* Bernhauer, *aureipennis* Bernhauer, *balfourbrownei* Fagel, *banghaasi* Bernhauer, *basipes* Fauvel, *bayeri* Bernhauer, *benicki* Wendeler, *bicoloriceps* Fagel, *blukwensis* Fagel, *braunsi* Bernhauer, *brevicollis* Fagel, *caffer* Boheman, *capensis* Erichson, *capitalis* Bernhauer, *caprai* Fagel, *chopardi* Fagel, *colasi* Fagel, *confusus* Fagel, *congoensis* Bernhauer, *cooperi* Scheerpeltz, *cribripennis* Fagel, *darfurensis* Bernhauer, *delkeskampii* Fagel, *densiventris* Bernhauer, *dollmani* Bernhauer, *drakensbergensis* Fagel, *duplex* Eppelsheim, *edwardsi* Bernhauer, *eidmanni* Fagel, *ertli* Bernhauer, *eximius* Reiche, *fastuosus* Klug, *flavitaris* Cameron, *freudei* Fagel, *freyi* Fagel, *frontalis* Fagel, *fulgidipennis* Fagel, *gabonicus* Fagel, *gebieni* Wendeler, *gedyei* Cameron, *gowdeyi* Bernhauer, *gracilis* Last, *graueri* Fagel, *gridellii* Fagel, *hintzi* Bernhauer, *ignotus* Wendeler, *ikelaensis* Fagel, *impressipennis* Fauvel, *incertus* Fagel, *inexpectatus* Fagel, *insidiosus* Fagel, *insularis* Cameron, *itombwensis* Fagel, *jaegeri* Paulian, *jansei* Fagel, *janssensi* Fagel, *jarrigeanus* Fagel, *jarrigei* Fagel, *junodi* Bernhauer, *kabarensis* Fagel, *kaszabi* Fagel, *kilimandjarensis* Fagel, *kiouanus* Fagel, *kochi* Fagel, *kundelungensis* Fagel, *lamottei* Fagel, *lusti* Fagel, *laticeps* Fagel, *lineativentris* Bernhauer, *lootensi* Fagel, *luffianus* Fagel, *macellus* Fagel, *machadoanus* Fagel, *madagascariensis* Erichson, *madudanus* Fagel, *magnificus* Bernhauer, *magnus* Fagel, *malaisei* Fagel, *mangolae* Bernhauer, *mariepskopensis* Fagel, *marshalli* Bernhauer, *mateui* Fagel, *maynei* Bernhauer, *melano-*

gaster Fauvel, *methneri* Fagel, *minimus* Bernhauer, *montanus* Wendeler, *monticola* Cameron, *montislomaensis* Fagel, *montivagans* Cameron, *mwengensis* Fagel, *nakurensis* Fagel, *ngoviensis* Fagel, *nigripes* Bernhauer, *nigrolincatus* Bernhauer, *nyamukubiensis* Fagel, *obscuripes* Bernhauer, *opacicollis* Bernhauer, *overlaeti* Fagel, *pedestris* Gerstaecker, *piceus* Fauvel, *plagiator* Kolbe, *praeses* Fagel, *pretiosus* Bernhauer, *pulchellus* Bernhauer, *purpuripennis* Bernhauer, *raffrayi* Fagel, *reticuliventris* Fagel, *riftensis* Fauvel, *ruandanus* Fagel, *rudebecki* Fagel, *rufiventris* Bernhauer, *rufocyaneus* Bernhauer, *ruhembanus* Bernhauer, *ruwenzoricus* Fagel, *sachtlebeni* Fagel, *scheerpeltzi* Fagel, *scottianus* Fagel, *secretus* Bernhauer, *seydeli* Fagel, *simillimus* Bernhauer, *straelenianus* Fagel, *subopacus* Bernhauer, *sudanensis* Bernhauer, *superbus* Bernhauer, *tandalensis* Bernhauer, *tenuis* Fagel, *testaceopiceus* Bernhauer, *tibestiensis* Fagel, *transvaalensis* Fagel, *tropicus* Bernhauer, *tumidicollis* Gerstaecker, *turneris* Cameron, *ugandensis* Fagel, *uganensis* Fagel, *ulindiensis* Fagel, *uluguruensis* Fagel, *usagarae* Bernhauer, *usambaricus* Schubert, *versicolor* Fagel, *violaceipennis* Fagel, *viridipennis* Bernhauer, *vulcanus* Wendeler, *weisei* Schubert, *wenzeli* Fagel, *xanthocerus* Eppelsheim, *y-oundensis* Fagel, *zairensis* Frank, *zulu* Fagel.

Unassigned Paederus (sensu lato): *abdominalis* Cameron, *abnormalis* Bernhauer, *adelaidae* Blackburn, *aequabilis* Last, *agnatus* Eppelsheim, *aliiceps* Cameron, *almorensis* Cameron, *alternans* Walker, *alticola* Sharp, *alutipennis* Cameron, *amazonicus* Sharp, *amplicollis* Kraatz, *andinus* Wendeler, *anguinus* Bernhauer, *angulicollis* MacLeay, *anthracinus* Bernhauer, *antiquus* Sharp, *antoniensis* Sharp, *apteromelas* Lea, *arduus* Sharp, *argentatus* Cameron, *argentinus* Bernhauer, *armstrongi* Steel, *atriventris* Cameron, *atrocyaneus* Champion, *australis* Guerin, *baeri* Bernhauer, *bakeri* Cameron, *bakerianus* Bernhauer, *basiventris* Bernhauer, *beccarii* Wendeler, *bernhauerianus* Scheerpeltz, *besucheti* Biswas & Sen Gupta, *bhutanicus* Coiffait, *bicoloripes* Cameron, *birmanus* Fauvel, *bolivianus* Bernhauer, *bonariensis* Lynch, *bonus* Last, *borneensis* Scheerpeltz, *brandti* Last, *brasiliensis* Erichson, *bruchii* Bernhauer, *buluensis* Last, *busalminensis* Last, *cameroni* Last, *canonicus* (Casey), *capillaris* Fauvel, *captus* Wendeler, *carolinae* (Casey), *cartaginis* Bernhauer, *centralis* Last, *ceylonicus* Bernhauer, *cheesmani* Cameron, *chimbuenensis* Last, *coarctatus* Erichson, *columbinus* Laporte, *combustus* Fauvel, *comorensis* Bernhauer, *compotens* LeConte, *condei* Wendeler, *conicipennis* Wendeler, *conicollis* Motschulsky, *conspicuus* Erichson, *cooperi* Scheerpeltz, *copiosus* Last, *cordovens* Sharp, *coriaceus* Fauvel, *costaricensis* Sharp, *crassicollis* Wendeler,

cruenticollis Germar, *cumanus* Frank, *currax* Sharp, *curticeps* Bernhauer, *curticollis* Bernhauer, *cyanellus* Fauvel, *cyaneus* Cameron, *cyanopterus* Cameron, *dammermani* Cameron, *densipennis* Bernhauer, *distinctus* Cameron, *diversiceps* Cameron, *doriae* Fauvel, *drescheri* Cameron, *ecuadorensis* Bernhauer, *elegans* Wendeler, *elegantulus* Wendeler, *elongatus* Wendeler, *erythroderus* Erichson, *excellens* Bernhauer, *extraneus* Wiedemann, *faeae* Fauvel, *femoralis* LeConte, *femoratus* Cameron, *ferreirai* Scheerpeltz, *ferus* Erichson, *fijiensis* Cameron, *filicornis* Sharp, *flaviceps* Bernhauer, *flavo-caudatus* Bernhauer, *floridanus* Austin, *fruhstorferi* Wendeler, *fulvicornis* Erichson, *fuscus* Last, *germairi* Wendeler, *germanus* Cameron, *gestroi* Fauvel, *gigas* Bernhauer, *globulicollis* Bernhauer, *grandicollis* Bernhauer, *grandis* Austin, *gratiosus* Fauvel, *greeni* Cameron, *gressitti* Last, *hagenensis* Last, *hebridensis* Cameron, *heynei* Wendeler, *himalayicus* Bernhauer, *hingstoni* Cameron, *hornabrooki* Last, *horni* Bernhauer, *huonensis* Last, *iheringi* Bernhauer, *illiesi* Puthz, *imitator* Wendeler, *intermedius* Boheman, *iowensis* (Casey), *irianensis* Frank, *ismayi* Last, *jacobsoni* Bernhauer, *jamaicensis* Blackwelder, *javanus* Laporte, *jessoensis* Bernhauer, *kaindiensis* Last, *kaiseri* Bernhauer, *kakodanus* Cameron, *kambaitensis* Scheerpeltz, *kamonoensis* Last, *katubuensis* Last, *klapperichi* Bernhauer, *koebelci* Blackburn, *kozuensis* Cameron, *kraepelini* Fauvel, *kuborensis* Last, *kuntzeni* Wendeler, *luensis* Last, *laetipes* Sharp, *laetus* Erichson, *lativentris* Wendeler, *licenti* Bernhauer, *lingualis* Sharp, *listeri* Gahan, *littorarius* Gravenhorst, *littoreus* Austin, *lividus* Bernhauer, *lombocianus* Cameron, *luridiventris* Sharp, *maai* Last, *machadoi* Scheerpeltz, *mafuluensis* Last, *magniceps* Bernhauer, *mandibularis* Erichson, *marcuzzii* Scheerpeltz, *melampus* Erichson, *melanopus* Wendeler, *mexicanus* Erichson, *meyricki* Blackburn, *michaelensis* Last, *micropterus* Lea, *mimicus* Last, *montanus* Wendeler, *morio* Mannerheim, *mosus* Cameron, *mutans* Sharp, *nakaoi* Last, *neotropicus* Bernhauer & Schubert, *nepalicus* Coiffait, *nigripennis* Cameron, *nigriventris* Wendeler, *obliteratus* LeConte, *ohausi* Wendeler, *okapaensis* Last, *ornaticornis* Sharp, *ovaliceps* Bernhauer, *paeninsula* Last, *pallidus* Sharp, *palustris* Austin, *papuanus* Cameron, *paraguayanus* Wendeler, *pauloensis* Bernhauer, *pendleburyi* Cameron, *pentagonalis* Wendeler, *perlongus* Last, *peruanus* Wendeler, *philippinus* Bernhauer, *plagiator* Kolbe, *politus* Fauvel, *praecellens* Bernhauer, *preangeranus* Bernhauer, *problematis* Last, *puberulus* Motschulsky, *punctatus* Last, *punctiger* Sharp, *punctiventris* Bernhauer, *punjabensis* Cameron, *republicanus* Bernhauer, *reticulatus* Last, *roepkei* Bernhauer, *rubidus* Cameron, *rufitarsis*

Solsky, *rufotestaceus* Cameron, *rutilicornis* Erichson, *saginatulus* (Casey), *salvini* Sharp, *samarensis* Bernhauer, *samoensis* Fauvel, *samuelseni* Last, *santoensis* Cameron, *schultheissi* Fauvel, *schusteri* Bernhauer, *sedlaceki* Last, *semaranganus* Bernhauer, *semiviridis* Bernhauer, *senegalensis* Laporte, *sericeicollis* Scheerpeltz, *setifer* Cameron, *sharpi* Cameron, *signaticornis* Sharp, *sijthoffi* Bernhauer, *simsoni* Blackburn, *solidus* Sharp, *sparsus* Fauvel, *speculifrons* Scheerpeltz, *stenopterus* Lea, *studti* Wendeler, *subcoriaceus* Cameron, *subnitidus* Bernhauer, *sulawesi* Frank, *tambulensis* Last, *taprobanus* Bernhauer, *tempestivus* Erichson, *testaceitarsis* Sharp, *tibetanus* Cameron, *tongyai* Bernhauer, *toxopei* Wendeler, *tricolor* Erichson, *tridens* Sharp, *trimerus* Fauvel, *tristis* Bernhauer, *tsaratananus* Jarrige, *turrialbanus* Wendeler, *tweedensis* Blackburn, *unicolor* Cameron, *uruguayensis* Bernhauer, *ustus* LeConte, *variceps* Kraatz, *variicornis* Fauvel, *velox* Sharp, *vianai* Bernhauer, *wauensis* Last, *wilsoni* Lea, *yuangasanus* Wendeler, *yucateca* Sharp, *zayusi* Sanderson.

Fossils: *adumbratus* Wickham.

Homonymy in species-group names

The International Code of Zoological Nomenclature (ICZN 1964) recognizes two forms of homonymy, primary and secondary, and requires replacement names for junior primary homonyms and some junior secondary homonyms. The forms of homonymy are illustrated by diagrams by Blackwelder (1967: 551).

1: Primary homonyms. Some junior primary homonyms (e.g., *Paederus erichsoni* Bernhauer) already have been replaced deliberately (Table 1). Others (e.g., *Paederus angolensis* Bernhauer) have available senior synonyms, so do not require replacement names (Table 1). The senior synonym of *Oreopaederus filicornis* (Fagel) (originally *Paederus filicornis* Fagel) is *Oreopaederus minor* (Bernhauer) (originally *Paederus testaceus minor* Bernhauer); this synonymy was admitted by Fagel (1954a, 1958a), though he created the unnecessary new epithet *filicornis*. Four replacement names are required: *Paederus cumanus* Frank [*nom. nov.* for *Paederus bicolor* Wendeler, *nec* Olivier], *Paederus sulawesi* Frank [*nom. nov.* for *Paederus melanocephalus* Heller, *nec* Fabricius], *Paederus zairensis* Frank [*nom. nov.* for *Paederus orophilus* Fagel, *nec* *Paederidus brunnescens orophilus* (Fagel), originally *Paederus brunnescens orophilus* Fagel], and *Paederus irianensis* Frank [*nom. nov.* for *Paederus litoreus* Last, *nec* *Paederus littoreus* Austin, replacement required under Article 58(b) (ICZN 1964)] and are of geographical origin. Another geographical name, *Oreopaederus manyemensis* Frank, is used here for *Oreopaederus ater* (Bernhauer),

Table 1. Primary homonyms (1), secondary homonyms (2) and non-homonyms (3) of species-group names in *Paederus* sensu lato. Replacement epithets in square brackets are senior synonyms of the junior homonyms, making new replacement epithets unnecessary.

Epithet	First Author	Second Author	Form	Replacement Epithet	Epithet	First Author	Second Author	Form	Replacement Epithet
<i>angolensis</i>	Erichson 1843: 222	Bernhauer 1936b: 262	(1A)	[plagiator Kolbe]	<i>longicornis</i>	Aube 1850: 319	Motschulsky 1858: 634	(1b)	[sabaeus Erichson]
<i>ater</i>	Eichler 1924: 65	Bernhauer 1927a: 107	(1B)	manyemensis Frank	<i>lusitanicus</i>	Aube 1842: 236	Baudi 1857: 107	(1A)	baudii Fairmaire
<i>balcanicus</i>	Koch 1938: 103	Scheerpeltz 1957a: 455	(3)	-----	<i>machadoi</i>	Fagel 1962b: 22	Scheerpeltz 1965a: 223	(3)	-----
<i>ballfourbrownei</i>	Fagel 1958a: 139	Fagel 1958a: 303	(3)	-----	<i>melanocephalus</i>	Fabricius 1792: 538	Heller 1899: 4	(1C)	sulawesi Frank
<i>banghaasi</i>	Bernhauer 1927c: 400	Bernhauer 1927a: 111	(1A)	banghaasianus Scheerpeltz	<i>meersmanae</i>	Fagel 1961a: 263	Fagel 1965: 141	(3)	-----
<i>bicolor</i>	Olivier 1795: 7	Wendeler 1942: 103	(1C)	cumanus Frank	<i>minutus</i>	Gautier 1862: 76	Fagel 1965: 141	(3)	-----
<i>chinensis</i>	Bernhauer 1931b: 1	Bernhauer 1934b: 6	(1A)	sinensis Scheerpeltz	<i>nigricans</i>	Scheerpeltz 1957a: 454	Fagel 1961a: 265	(3)	-----
<i>confusus</i>	Fagel 1958a: 75	Fagel 1958a: 433	(3)	-----	<i>nigriventris</i>	Wendeler 1926a: 73	Fagel 1965: 139	(3)	-----
<i>elongatus</i>	(Linnaeus) 1767: 685	Wendeler 1931: 46	(2F)	archeus * Blackwelder	<i>orophilus</i>	Fagel 1954a: 919	Fagel 1958a: 379	(1C)	zairensis Frank
<i>erichsoni</i>	Wollaston 1867: 247	Bernhauer 1908c: 16	(1A)	antipodum Bernh. & Schubert	<i>pallidus</i>	Sharp 1886: 618	Bernhauer 1915b: 457	(1A)	pallidulus Scheerpeltz
<i>filicornis</i>	Sharp 1886: 612	Fagel 1954a: 949	(1b)	[minor Bernhauer]	<i>punctiventris</i>	Bernhauer 1908a: 310	Scheerpeltz 1957a: 457	(3)	-----
<i>filiformis</i>	Fabricius 1792: 538	Latreille 1806: 293	(1C)	[procerus Gravenhorst]	<i>rubricollis</i>	Gravenhorst 1806: 138	Fagel 1958a: 135	(3)	-----
<i>freyi</i>	Fagel 1958a: 203	Fagel 1962a: 69	(3)	-----	<i>sanguinicollis</i>	Stephens 1833: 281	Motschulsky 1860b: 559	(1A)	haematoderus Gemm. & Harold
<i>geniculatus</i>	Dietrich 1855: 201	Peyron 1858: 430	(1A)	[littoralis Gravenhorst]	<i>semicyaneus</i>	Perty 1830: 33	Cameron 1950c: 185	(1C)	[luctuosus Klug]
<i>gracilis</i>	(Paykull) 1789: 38	Last 1952: 91	(2F)	unnecessary	<i>senegalensis</i>	Laporte 1835: 123	Fagel 1962a: 67	(3)	-----
<i>indicus</i>	Motschulsky 1858: 634	Bernhauer 1911a: 61	(1A)	himalayicus Bernhauer	<i>socius</i>	Bernhauer 1916: 29	Bernhauer 1927a: 117	(1A)	fratellus Bernhauer
<i>javanus</i>	Laporte 1835: 123	Erichson 1840: 654	(1A)	sondaicus Fauvel	<i>testaceus</i>	(Paykull) 1789: 28	Bernhauer 1934c: 231	(2F)	unnecessary
<i>lasti</i>	Fagel 1958a: 292	Fagel 1961a: 276	(3)	-----	<i>thoracicus</i>	Fourcroy 1785: 170	Erichson 1840: 663	(2D)	neotropicus Bernh. & Schubert
<i>leleupi</i>	Fagel 1954a: 927	Fagel 1958a: 135	(3)	-----	<i>tricolor</i>	(Fabricius) 1787: 221	Erichson 1840: 663	(2F)	homonymus * Blackwelder
<i>leleupi</i>	Fagel 1954a: 927	Fagel 1965: 146	(3)	-----	<i>villersi</i>	Cameron 1949a: 317	Fagel 1958a: 136	(3)	-----
<i>leleupi</i>	Fagel 1958a: 135	Fagel 1965: 146	(3)	-----	<i>viridipennis</i>	Bernhauer 1915a: 117	Bernhauer 1931c: 574	(1A)	cooperi Scheerpeltz
<i>littoreus</i>	Austin 1876: 9	Last 1985: 331	(§)	irianensis Frank	<i>vulcanus</i>	Wendeler 1926b: 203	Fagel 1954a: 956	(1A)	vulcanicola Fagel

Form of homonymy is specified according to diagrams lettered in capitals A-H (here in parentheses) by Blackwelder (1967: 551); form (1b) is a derivative of Fig. B of Blackwelder (1967) and has the junior homonym (not the senior) transferred to another genus. Replacement epithets with an asterisk were unnecessary at the time they were named, so they are junior objective synonyms. § To be treated as homonyms under Article 58(8) (ICZN 1985).

originally *Paederus ater* Bernhauer, which is preoccupied by *Paederidus rubrothoracicus ater* (Eichler), originally *Paederus ruficollis var. ater* Eichler; this need for replacement arose because of Article 45(g) (ICZN 1985).

2: Secondary homonyms. The name *Paederus thoracicus* Erichson required replacement [form D in Blackwelder (1967)] and this was done by Bernhauer & Schubert (1912). The names *Paederus testaceus* Bernhauer and *Paederus gracilis* Last do not require replacement [form F in Blackwelder (1967)]. The names *Paederus elongatus* Wendeler and *Paederus tricolor* Erichson did not require replacement [form F in Blackwelder (1967)] but were replaced by the names *Paederus archeus* (Blackwelder 1944) and *Paederus homonymus* (Blackwelder 1943) respectively. *Paederus archeus* Blackwelder thus becomes a junior objective synonym of *Paederus elongatus* Wendeler, and *Paederus homonymus* Blackwelder becomes a junior objective synonym of *Paederus tricolor* Erichson.

3: Non-homonyms. The 16 pairs of names marked by a number (3) in Table 1 are not considered homonyms because they were originally attributed to, and remain in, different genera. However, all of the genera belong to *Paederus sensu lato*. Three of these non-homonyms were published by Scheerpeltz and the remainder by Fagel. The International Code [ICZN 1964: Appendix D, recommendation 5(a)] urges against the use of such identical epithets in closely related genera. Other pairs of names (*ugandensis* Fagel, *uganensis* Fagel; *saegeri* Fagel, *saengeri* Coiffait) contravene recommendation 5(b) of Appendix D in having small differences in spelling, while yet other pairs of names (*angolensis* Erichson, *angolanus* Fagel; *chinensis* Bernhauer, *sinensis* Scheerpeltz) contravene recommendation 6 of Appendix D in being derived from the same geographical names (ICZN 1964).

Conclusion

There is a wide range of aedeagal structure in *Paederus sensu lato*, providing important taxonomic indicators at the species level. Classification at generic and subgeneric levels has proven difficult because of the large number of species and because of limited understanding of external characters.

After studies by Fagel (1958a-1973) using diagnostic external characters, augmented by Jarrige (1970, 1978) and Coiffait (1982c), the species of the Afrotropical (including Malagasy) region are grouped as *Paederidus* (36 spp.), *Allopaederus* (3 spp.), *Madecapaederus* (11 spp.), *Pachypaederus* (19 spp.), *Oreopaederus* (55 spp.), and

Paederus sensu Fagel (157 spp.). Fagel (1958a) also described an Oriental genus, *Ctenopaederus* (1 sp.). However, this includes only 45% of the world's known fauna. Other generic groupings are questionable due to lack of defined external characters, and 264 spp. have been assigned only to *Paederus sensu lato*. The 47 species of the western Palearctic region are grouped as *Paederidus* (10 spp.), *Parameropaederus* (1 sp.), *Lobopaederus* (1 sp.), *Uncopaederus* (1 sp.), and *Paederus sensu Scheerpeltz-Coiffait* (33 spp.) (Coiffait 1982c).

Fagel (1958a) did not separate the 157 Afrotropical (including Malagasy) *Paederus sensu Fagel* into subgenera, though other authors have attempted to define subgenera (mainly Palearctic) largely on aedeagal characters. There seems now to be a reasonably good understanding of the Palearctic fauna at the species level. The world fauna, as currently understood, contains 621 extant species classified as follows, plus one fossil.

Various genera with numbers of species included:

Paederidus (44), *Parameropaederus* (1), *Lobopaederus* (1), *Megalopaederus* (8), *Eupaederus* (2), *Diplopaederus* (2), *Oncopaederus* (1), *Allopaederus* (3), *Ctenopaederus* (1), *Madecapaederus* (11), *Oreopaederus* (55), *Pachypaederus* (19), *Uncopaederus* (1), Total = 149 spp.

Various subgenera of *Paederus* with numbers of species included: *Paederus s.str.* (12), *Pseudopaederus* (5), *Gnathopaederus* (2), *Eopaederus* (10), *Harpopaederus* (7), *Heteropaederus* (6), *Dioncopaederus* (5), *Oedopaederus* (1), *Anomalopaederus* (1), *Oreinopaederus* (1), *Nepalopaederus* (1), Total = 51 spp.

Species assigned to *Paederus* by Fagel: Total = 157 spp.

Unassigned species of *Paederus (sensu lato)*:

Total = 264 spp.

There are now two major taxonomic needs. These are to review and redescribe the species of the Nearctic, Neotropical, Oriental, Oceanian and Australasian regions, and to extend Fagel's (1958a) methods to examine the higher classification of *Paederus sensu lato* worldwide.

Acknowledgments

Gratitude is owed to Lee Herman (American Museum of Natural History) and Randall Lundgren (Gainesville) for reviewing manuscript drafts. They and Ruthie Brumbaugh and Gail Wilson (Central Science Library, University of Florida) provided originals or photocopies of about 150 of the references cited. This is University of Florida, Institute of Food & Agricultural Sciences, Journal Series no. 8316.

Addendum

The following paper was noticed while this paper was in press: Lecoq, J.L. 1986. Les Paederinae des Îles Mascareignes (Coleoptera, Staphylinidae). *Nouv. Rev. Ent. (n.s.)* 3:417-437. It contains descriptions of 2 new species: *Paederus vinsoni* Lecoq from Mauritius (p. 419) and *Paederus colettae* Lecoq from Reunion (p. 419). It also gives records of *P. insularis* Cameron from Reunion (p. 418), *P. edwardsi* Bernhauer from Mauritius (p. 419), *P. rubidus* Cameron, and *P. fuscipes* Curtis from Reunion (p. 419).

Literature Cited

- Adachi, T.
1939. New Staphylinidae from Formosa. *Kontyû* 13: 165-166.
- Aleksandrov, A.I.
1934. Towards a list of beetles of the family Staphylinidae (Coleoptera) of Ghirinsk Province, with descriptions of new forms [in Russian]. *Ann. Club Sci. Geogr. YMCA* 1: 150-155.
- Aragona, A.
1830. De quibusdam Coleopteris Italiae novis aut rarioribus. 31 p. Bizzoni; Ticino.
- Aubé, C.
1842. Notes sur quelques coléoptères nouveaux. *Ann. Soc. Ent. France* 11: 225-237, pl. 9.
1850. Description de quelques insectes coléoptères appartenant à l'Europe et à l'Algérie. *Ann. Soc. Ent. France (ser. 2)* 8: 299-346.
- Austin, E.P.
1876. On the species of *Sunius* and *Paederus* found in the United States. *Proc. Boston Soc. Nat. Hist.* 19: 4-11.
- Baer, G.-A.
1886. Catalogue des coléoptères des Îles Philippines. *Ann. Soc. Ent. France (ser. 6)* 6: 97-108.
- Baudi Di Selve, F.
1848. Alcune specie nuove di stafilini. *Studi Entomologici* 1: 113-148.
1857. Coleoptera quaedam e Staphylinorum familia nova vel minus cognita cum observationibus. *Berliner Ent. Ztschr.* 1: 97-115.
1869. Coleopterorum messis in insula Cypro et Asia minore ab Eugenio Truqui congregatae recensio: de Europaeis notis quibusdam additis (Pars altera). *Berliner Ent. Ztschr.* 13: 369-418.
- Beck, L. von
1817. Beiträge zur bayerischen Insectenfaune, oder Beschreibung und Abbildung neuentdeckter Käfer, mit angehangtem Namensverzeichnisse der Eleuteraten des Landgerichtbezirks Zusmeshausen. 47 p. Wolff; Augsburg.
- Bernhauer, M.
1901. Neue Staphyliniden der paläarktischen Fauna, nebst Bemerkungen. *Deutsche Ent. Ztschr.* (1901): 241-251.
1902. Zur Staphyliniden-Fauna von Ceylon. *Deutsche Ent. Ztschr.* (1902): 17-45.
1903. Neue Staphyliniden der paläarktischen Fauna. *Munch. Koleopt. Ztschr.* 1: 186-192.
1905a. Neue exotische Staphyliniden. *Deutsche Ent. Ztschr.* (1905): 9-21.
1905b. Neue Staphyliniden aus Südamerika. *Deutsche Ent. Ztschr.* (1905): 177-187.
1907. Neue Staphyliniden aus Südamerika. 4. Stück. *Wiener Ent. Ztg.* 26: 281-289.
1908a. Beitrag zur Staphylinidenfauna von Südamerika. *Arch. Naturges.* 74: 283-372.
1908b. Neue Staphylinidae aus dem subtropischen und tropischen Afrika. *Denkschr. Med.-Naturwiss. Ges. Jena* 13: 101-118.
1908c. Staphylinidae [p. 15-23, in] J.W. Michaelsen & R. Hartmeyer (eds.) *Die Fauna Südwest-Australiens. Vol. 2, Pt. 2.* Fischer; Jena.
1909. Zur Staphylinidenfauna von Südamerika. 6. Beitrag. *Boll. Soc. Ent. Ital.* 40: 225-251.
1910. Beitrag zur Kenntnis der Staphyliniden-Fauna von Zentralamerika. *Verh. Zool. Bot. Ges. Wien* 60: 350-393.
1911a. Zur Staphylinidenfauna Ostindiens und der Sundainseln (3. Beitrag). *Ent. Bl. Biol. Syst. Käfer* 7: 55-62.
1911b. Zur Staphylinidenfauna von Süd-Amerika (Col.). (7. Beitrag). *Deutsche Ent. Ztschr.* (1911): 403-422.
1912a. Zur Staphylinidenfauna von Südamerika (10. Beitrag). *Verh. Zool.-Bot. Ges. Wien* 62: 26-48.
1912b. Beitrag zur Staphylinidenfauna von Afrika. *Ent. Mitt.* 1: 177-183, 203-209.
1912c. Neue Staphyliniden aus Zentral- und Deutsch-Ostafrika [p. 469-486, in] *Wissenschaftliche Ergebnisse der Deutschen Zentral-Afrika-Expedition 1907-1908 unter Führung Adolf Friedrich, Herzogs zu Mecklenburg. Vol. 3 (Zool. 1).* Klinkhardt & Biermann; Leipzig.
1912d. Neue Staphyliniden der Philippinen. *Philippine J. Sci. (ser. D)* 7: 245-254.

Bernhauer, M. (continued)

- 1913a. Neue Staphyliniden der paläarktischen Fauna. Münchener Kol. Ztschr. 1: 186-192.
- 1913b. Beitrag zur Staphylinidenfauna der paläarktischen Region. Ent. Bl. 9: 219-224.
1914. Neue Staphylinen der indo-malaiischen Fauna. Verh. Zool.-Bot. Ges. Wien 64: 76-109.
- 1915a. Zur Staphyliniden-Fauna des tropischen Afrika. Ann. Hist. Nat. Mus. Natn. Hungarici 13: 95-189.
- 1915b. Neue Staphyliniden der indo-malaiischen Fauna, insbesondere der Sunda-Insel Borneo. Verh. Zool.-Bot. Ges. Wien 65: 134-158.
- 1915c. Neue Staphyliniden des tropischen Afrika. (10. Beitrag). Verh. Zool. Bot. Ges. Wien 65: 287-321.
- 1915d. Neue Staphyliniden aus dem Kongogebiet (11. Beitrag zur afrikanischen Staphylinidenfauna). Wiener Ent. Ztg. 34: 298-300.
- 1915e. Neue Staphyliniden aus Java und Sumatra. (7. Beitrag zur indo-malayischen Staphylinidenfauna). Tijdschr. Ent. 58: 213-243.
- 1915f. Beitrag zur Staphylinidenfauna von Neu-Guinea. Deutsche Ent. Ztschr. (1915): 179-202.
- 1915g. Staphyliniden [p. 361-368, *in*] Ergebnisse der II Deutsche Zentral Afrika Expedition 1910-1911 unter Führung Adolf Friedrichs, Herzogs zu Mecklenburg.
- 1915h. Beiträge zur Kenntnis der paläarktischen Staphyliniden-Fauna. Münchener Kol. Ztschr. 4 (1914): 262-270.
- 1915i. Coleoptera vii: Staphylinidae. Beitr. Kenntnis Land-Süßwasserf. Deutsch-Südwestafr. 1: 313-321.
1916. Kurzflügler aus dem deutschen Schutzgebiete Kiautschau und China. Arch. Naturges. 81(A): 27-34.
1917. Staphyliniden von Madagascar, den Comoren und Zanzibar [Vol. 3, p. 552-560, *in*] A. Voeltzkow (ed.) Reise in Ostafrika in den Jahren 1903-1905, Wissenschaftliche Ergebnisse.
- 1920a. Neue Staphyliniden aus Mittelamerika. Wiener Ent. Ztg. 38: 44-47.
- 1920b. Neue Staphyliniden des indo-malayischen Gebietes. 16. Beitrag. Arch. Naturges. 84(A): 177-188.
- 1920c. Results of Dr. E. Mjöberg's Swedish scientific expeditions to Australia 1910-1913. 22. Staphylinidae. Ark. Zool. 13(8): 1-27.
1921. Zur Staphylinidenfauna von Südamerika (24. Beitrag). Deutsche Ent. Ztschr. (1921): 65-77.
1922. Sauter's Formosa-Ausbeute: Staphylinidae. I. Teil. Arch. Naturges. 88(A): 220-237.
1923. Zur palaearktischen Staphylinidenfauna. Ent. Bl. 19: 176-181.
1926. Staphylinidae [p. 4-6, *in*] Y. Sjöstedt (ed.) Zoological results of the Swedish expedition to Central Africa 1921. Insecta. 23. Abschluss und Zusammenfassung. Ark. Zool. 18a(24): 1-13.
- 1927a. Neue afrikanische *Paederus* (sechzehnter Beitrag zur Staphylinidenfauna Afrikas). Verh. Zool. Bot. Ges. Wien 77: 102-118.
- 1927b. 32. Beitrag zur Staphylinidenfauna Südamerikas insbesondere Brasiliens. Mem. Soc. Ent. Ital. 5: 152-169.
- 1927c. Die Staphyliniden der Philippinen. 24. Beitrag zur indo-malayischen Staphylinidenfauna. Philippine J. Sci. 34: 399-401.
1929. Richtigestellung. Ent. Nachrichtenbl. 3: 123.
- 1931a. Staphylinidae [*in*] H. Lindberg (ed.) Inventaria entomologica itineris Hispanici et Marocani, quod a. 1926 fecerunt Harald et Hakan Lindberg. VII Soc. Sci. Fenn. Comm. Biol. 3: 1-29.
- 1931b. Neue Staphyliniden aus China von der Stotner'schen Expedition. Ent. Nachrichtenbl. 5: 1-3.
- 1931c. Entomological Expedition to Abyssinia, 1926-1927: Coleoptera, Staphylinidae. Systematic Report by Dr. Max Bernhauer (Horn, Niederösterreich, Austria). J. Linn. Soc. London, Zool. 37: 559-605.
- 1931d. Neuheiten der paläarktischen Staphylinidenfauna. Kol. Rdsch. 17: 232-245.
1932. Neue Kurzflügler aus dem belgischen Kongostaate. (24. Beitrag zur afrikanischen Fauna). Rev. Zool. Bot. Afr. 22: 72-90.
- 1933a. Neuheiten der chinesischen Staphylinidenfauna. Wiener Ent. Ztg. 50: 25-48.
- 1933b. Zur Staphylinidenfauna des belgischen Kongostaates. (30. Beitrag zur afrikanischen Fauna). Rev. Zool. Bot. Afr. 23: 291-301.
- 1934a. Neue Staphyliniden von den Neuen Hebriden im British Museum. Stylops 3: 18-19.
- 1934b. Siebenter Beitrag zur Staphylinidenfauna Chinas. Ent. Nachrichtenbl. 8: 1-20.
- 1934c. 31. Beitrag zur Staphylinidenfauna Afrika's. Rev. Zool. Bot. Afr. 24: 229-248.
- 1934d. Neue Kurzflügler vom Ruwenzori-Kivu Gebiet. 34. Beitrag zur afrikanischen Staphylinidenfauna. Rev. Zool. Bot. Afr. 25: 206-217.
- 1934e. Neue Staphyliniden (Col.) aus Südamerika. Stylops 3: 156-162.
- 1935a. Neuheiten der paläarktischen Staphylinidenfauna. I. Koleopt. Rundschau 21: 39-48.
- 1935b. Neue Staphyliniden vom belgischen Kongo.

- (37. Beitrag zur Staphylinidenfauna Afrikas). Rev. Zool. Bot. Afr. 27: 97-106.
- 1936a. Neuheiten der palaearktischen Staphylinidenfauna. I. Pubbl. Mus. Ent. Pietro Rossi Duino 1: 237-254.
- 1936b. Neue Staphylinidae aus Angola. Novit. Zool. 39: 261-267.
- 1937a. 44. Beitrag zur afrikanischen Staphylinidenfauna. Ann. Mag. Nat. Hist. (ser. 10) 20: 289-315.
- 1937b. Zur Staphylinidenfauna des belgischen Kongostaates. Rev. Zool. Bot. Afr. 30: 298-303.
- 1938a. Neue Staphyliniden aus China aus dem Ausbeuten von E. Licent S.J. vom Hoang-ho Pai-ho Museum in Tientsin und B. Becquart S.J. vom Philosophischen Institut der Jesuiten in Sienhsien-Hopeh. (11. Beitrag zur chinesischen fauna). Not. Ent. Chinoise, Mus. Heude 5: 49-57.
- 1938b. Neuheiten vom Belgischen Kongo. (49. Beitrag zur afrikanischen Staphylinidenfauna). Rev. Zool. Bot. Afr. 31: 326-333.
- 1939a. Zur Staphylinidenfauna Argentiniens und Brasiliens (Col.). Rev. Ent., Rio de J. 10: 231-249.
- 1939b. Zur Staphylinidenfauna von Kamerun. 42. Beitrag zur afrikanischen Fauna. Ent. Bl. 35: 252-262.
- 1939c. Zur Staphylinidenfauna des Belgischen Kongostaates. Rev. Zool. Bot. Afr. 32: 226-235.
- 1939d. Neue Staphyliniden aus Thailand. J. Thailand Res. Soc. Nat. Hist. 12 (suppl.): 115-116.
- 1940a. Staphylinidae. (51. Beitrag zur Staphylinidenfauna des tropischen Afrika). British Mus. Nat. Hist. Ruwenzori Exped. 1934/1935. 3: 129-144.
- 1940b. Neuheiten der paläarktischen Staphylinidenfauna (Col. Staph.). Mitt. Münchener Ent. Ges. 30: 622-642, 1025-1047
1941. Neue Staphyliniden aus China. (114. Beitrag). Ent. Bl. 37: 226-228.
- 1942a. Neue Staphyliniden aus Kostarika. Zool. Anz. 138: 1-27.
- 1942b. [Descriptions d'espèces nouvelles, in] M. Bernhauer & R. Paulian. Coléoptères staphylinides du Cameroun (Mission P. Lapesme, R. Paulian et A. Villiers, 1939). Rev. Zool. Bot. Afr. 35: 344-375.
1943. Neuheiten der palaearktischen Staphylinidenfauna (Zugleich 15. Beitrag zur japanisch-chinesischen Fauna). Mitt. Münchener Ent. Ges. 33: 169-188.
- Bernhauer, M. & K. Schubert.
1912. Staphylinidae I [in] Coleopterorum Catalogus (W. Junk, Berlin) 40: 191-288.
- Biswas, D.N. & T. Sen Gupta.
1982. New species and new records of Staphylinidae (Coleoptera) from India and Sri Lanka. Rev. Suisse Zool. 89: 135-154.
- Blackburn, T.
1888. Descriptions of twenty new species of South Australian Coleoptera. Trans. R. Soc. South Australia 10: 1-11.
1891. Further notes on Australian Coleoptera, with descriptions of new genera and species. Trans. R. Soc. South Australia 14: 65-153.
1894. Notes on Australian Coleoptera, with descriptions of new genera and species. Proc. Linn. Soc. New South Wales (ser. 2) 9: 85-108.
1899. Further notes on Australian Coleoptera, with descriptions of new genera and species. Trans. R. Soc. South Australia 23: 22-101.
- Blackwelder, R.E.
1939. A generic revision of the staphylinid beetles of the tribe Paederini. Proc. U.S. Natn. Mus. 87: 93-125.
1943. Monograph of the West Indian beetles of the family Staphylinidae. U.S. Natn. Mus. Bull. 182: i-vii, 1-658.
1944. Checklist of the coleopterous insects of Mexico, Central America, the West Indies, and South America. Pt. 1. U.S. Natn. Mus. Bull. 185: i-xii, 1-188.
1952. The generic names of the beetle family Staphylinidae with an essay on monotypy. U.S. Natn. Mus. Bull. 200: i-iv, 1-483.
1967. Taxonomy: A text and reference book. xiv + 698 p. Wiley; New York.
- Blanchard, C.E.
1842. Famille des brachélytres [p. 76-88, in] C. E. Blanchard & G.A. Brullé. Insectes de l'Amérique méridionale recueillis par Alcide d'Orbigny [in] Voyage dans l'Amérique méridionale (le Brésil, la République Orientale de l'Uruguay, la République Argentine, la Patagonie, la République du Chili, la République de Bolivia, la République du Pérou), exécuté pendant les années 1826, 1827, 1828, 1829, 1830, 1831, 1832 et 1833 par Alcide d'Orbigny. Vol. 6, pars 2, 222 p. Bertrand; Paris.
- Bohác, J.
1985. Review of the subfamily Paederinae (Coleoptera Staphylinidae) in Czechoslovakia. Acta Ent. Bohemoslov. 82: 369-385.

Boheman, C.

1848. *Insecta caffraria annis 1838-1845 a J.A. Wahlberg collecta. Pars 1, fasc. 1, viii + 297 p. Coleoptera (Carabici, Hydrocanthari, Gyrinii et Staphylinii).* Norstedt; Stockholm.
1858. *Coleoptera. Species novas descripsit [p. 1-112, in] Kongliga Svenska Fregatten Eugenie's resa omkring Jorden. Zoologi 1, Insekter. Kongl. Svenska Vetensk.- Akad.; Stockholm.*

Burmeister, H.

- 1861-1865. *Reise durch die La Plata- Staaten, mit besonderer Rücksicht auf die physische Beschaffenheit und den Culturzustand der Argentinischen Republik. Aufgeführt in den Jahren 1857, 1858, 1859 und 1860. Vol. 1, 504 p. + 1 pl. + 1 map (1861), Vol. 2, v + 540 p. + 1 map ['1861'] (1865). Schmidt; Halle.*

Cameron, M.

1914. *Descriptions of new species of Staphylinidae from India. Trans. Ent. Soc. London (1913): 525-544.*
1924. *New species of Staphylinidae from India. Trans. Ent. Soc. London (1924): 160-198.*
- 1925a. *New Staphylinidae from the Dutch East Indies. Treubia 6: 174-198.*
- 1925b. *Catalogue of Indian insects. Pt. 6 - Staphylinidae, 126 p. Govt. India Central Publ. Br.; Calcutta.*
1927. *New Staphylinidae from Ceylon. Ceylon J. Sci. Sect. B. Zool. Geol. (= Spolia Zeylan.) 14: 115.*
1928. *The Staphylinidae (Coleoptera) of the Third Mount Everest Expedition. Ann. Mag. Nat. Hist. (ser. 10) 2: 558-569.*
1929. *New species of Staphylinidae from Belgian Congo. Rev. Zool. Bot. Afr. 18: 56-65.*
- 1930a. *New species of Staphylinidae from Japan. Entomologist's Mon. Mag. 66: 181-185, 205-208.*
- 1930b. *Fauna Sumatrensis (Bijdrage Nr. 67). Staphylinidae (Col.). Tijdschr. Ent. 73: 325-348.*
- 1931a. *The fauna of British India, including Ceylon and Burma. Coleoptera. Staphylinidae, Vol. 2, viii + 257 p. + 2 pl. Taylor & Francis; London.*
- 1931b. *Staphylinidae (Coleoptera) from New Guinea, in the South Australian Museum. Rec. South Australian Mus. 4: 355-364.*
1933. *New species of Staphylinidae (Col.) from the Belgian Congo. Bull. Ann. Soc. Ent. Belg. 73: 35-53.*
1934. *The staphylinid Coleoptera of the New Hebrides. Stylops 3: 20-24.*
1936. *Fauna Javanica. The Staphylinidae (Col.) col-*

lected by Mr. F. C. Drescher. Tijdschr. Ent. 79: 25-54.

1937. *Staphylinidae (Col.) collected by Miss L. E. Cheesman in eastern New Guinea. Nova Guinea (n.s.) 1: 83-111.*
1938. *New species of Staphylinidae from Mauritius and Reunion, collected by Mr. J. Vinson. Entomologist's Mon. Mag. 74: 173-174.*
1939. *New species of Staphylinidae (Col.) from the Belgian Congo. Bull. Mus. Hist. Nat. Belgique 15: 1-7.*
1940. *Descriptions of some of Motschulsky's types of oriental Staphylinidae (Col.). Proc. R. Ent. Soc. London (B) 9: 85-89.*
- 1942a. *New species of Paederus (Col., Staphylinidae) from the New Hebrides. Entomologist's Mon. Mag. 78: 87-88.*
- 1942b. *New species of Staphylinidae (Col.) from the Moluccas and Celebes. Ann. Mag. Nat. Hist. (ser. 11) 9: 838-847.*
1943. *New species of Staphylinidae (Col.) from the Gilbert Islands and Fiji. Ann. Mag. Nat. Hist. (ser. 11) 10: 564-566.*
1944. *New species of South African Staphylinidae (Col.). Ann. Mag. Nat. Hist. (ser. 11) 11: 705-730.*
1945. *Descriptions of new Staphylinidae (Coleoptera). Proc. R. Ent. Soc. London (B) 14: 63-69.*
1947. *New species of Staphylinidae (Col.) from Mauritius. Proc. R. Ent. Soc. London (B) 16: 114-119.*
1948. *New species of Staphylinidae from Africa. Rev. Française Ent. 15: 39-43.*
- 1949a. *New species of African Staphylinidae. Bull. Inst. Français Afrique Noire, Dakar 11: 313-326.*
- 1949b. *New species and records of staphylinid beetles from Formosa, Japan, and south China. Proc. U.S. Natn. Mus. 99: 455-477.*
- 1950a. *New species of Staphylinidae (Col.) from the Malay peninsula. Ann. Mag. Nat. Hist. (ser. 12) 3: 1-40.*
- 1950b. *Staphylinidae (Coleoptera Polyphaga). in Exploration du Parc National Albert, Mission G.F. de Witte. Fasc. 59: 1-85. Institut des Parcs Nationaux du Congo Belge; Brussels.*
- 1950c. *New species of African Staphylinidae (Coleoptera). Part I. J. East Africa Uganda Nat. Hist. Soc. 19: 179-191.*
- 1950d. *New Staphylinidae from the Belgian Congo. Rev. Zool. Bot. Afr. 43: 92-98.*
- 1952a. *Results of the Archbold Expeditions. New species of Staphylinidae (Col.) from New Guinea and Misool. Treubia 21: 241-256.*

- 1952b. Mission A. Villiers au Togo et au Dahomey (1950) xvii. Coléoptères Staphylinidae. Bull. Inst. Français Afrique Noire 14: 827-836.
- Campos, F.
1931. Nuevas capturas del *Paederus ornaticornis* Sharp, en Guayaquil: Su identidad específica con la forma *Paederus irritans* Chap. Rev. Col. Nac. Vicente Rocafuerte 13: 23-25.
- Casey, T.L.
1905. A revision of the American Paederini. Trans. Acad. Sci. St. Louis 15: 17-248.
1910. Synonymic and descriptive notes on the Paederini and Pinophilini. Memoirs on the Coleoptera (New Era; Lancaster, Pennsylvania) 1: 184-201.
- Champion, G.C.
1927. Some Indian Coleoptera (22). Entomologist's Mon. Mag. 63: 49-57.
- Chapin, E.A.
1926. A new *Paederus* (Coleopt.: Staphylinidae) causing vesicular dermatitis in man. Arch. Schiffs-Tropenhyg. 30: 369-372.
1927. A new genus and species of Staphylinidae from Sze-Chuan, China. Proc. Biol. Soc. Washington 40: 75-77.
- Chapman, W.
1930. Description d'une aberration nouvelle de *Paederus ruficollis* (Col. Staphylinidae). Misc. Ent. 32: 101-102.
- Chevrolat, L.A.A.
1871. [Nouvelles acquisitions pour la faune française]. Petit. Nouv. Ent. (1871): 136.
- Coiffait, H.
1970. Staphylinides nouveaux ou mal connus de la région paléarctique occidentale. Bull. Soc. Hist. Nat. Toulouse 106: 99-111.
1976. Staphylinides récoltés au Népal par le professeur Franz. Bull. Soc. Hist. Nat. Toulouse 112: 243-275.
1978a. Ergebnisse der Bhutan-Expedition 1972 des Naturhistorischen Museums in Basel. Coleoptera: fam. Staphylinidae subfam. Paederinae, Euaesthetinae, Piestinae, Osoriinae et Omaliinae. Entomologica Basiliensia 3: 109-150.
1978b. Staphylinides du Nouristan (Afghanistan). Ann. Soc. Ent. France (n.s.) 14: 551-569.
1978c. Staphylinides récoltés par T. Deuve en Anatolie septentrionale (Col. Staph.). Nouv. Rev. Ent. 8: 163-175.
1978d. Staphylinides récoltés par T. Deuve en Anatolie septentrionale (erratum). Nouv. Rev. Ent. 8: 387.
1979. Insects of Saudi Arabia: Fam. Staphylinidae, Subfam. Xantholininae, Staphylininae, Paederinae, Oxytelinae, Aleocharinae [p. 162-180, in] W. Wittmer & W. Buttiker (eds.) Fauna of Saudi Arabia, Vol. 1, 372 + 17 p. Ciba-Geigy; Basle.
- 1982a. Contribution à la connaissance des staphylinides de l'Himalaya (Népal, Ladakh, Cachemire) (Insecta: Coleoptera: Staphylinidae). Senckenberg Biol. 62: 21-179.
1982b. Staphylinides (Col.) de la région himalayenne et de l'Inde (1. Xantholininae, Staphylininae et Paederinae). Ent. Basiliensia 7: 231-302.
1982c. Coléoptères Staphylinidae de la région paléarctique occidentale IV. Sous famille Paederinae tribu Paederinae 1 (Paederi, Lathrobii). Nouv. Rev. Ent. 12(4) suppl.: 1-440.
- Curtis, J.
1826. British entomology; being illustrations and descriptions of the genera of insects found in Great Britain and Ireland: containing coloured figures from nature of the most rare and beautiful species, and in many instances of the plants upon which they are found. Vol. 3, pl. 99-146. Curtis; London.
- Czwalina, G.
1889. [Forcipes einiger *Paederus* Arten]. Deutsche Ent. Ztschr. (1889): 368, pl. 2.
- Dahl, G.
1823. Coleoptera und Lepidoptera. Ein systematisches Verzeichniss, mit beygesetzten Preisen der Vorrathe. vi + 105 p. Akkermann; Vienna.
- Degeer, C.
1774. Mémoires pour servir à l'histoire des insectes. Vol. 4, xii + 456 p. + 19 pl. Hesselberg; Stockholm.
- Dejean, P.F.M.A.
1833. Catalogue des coléoptères de la collection de M. le comte Dejean [2d. ed.]. 443 p. Méquignon-Marvis; Paris.
- Dietrich, J.K.
1855. Beschreibung von zwei neuen Käferarten aus der Gattung *Paederus*. Stettin. Ent. Ztg. 16: 200-203.
- Eichelbaum, F.
1913. Verzeichnis der von mir in den Jahren 1903 und 1904 in Deutsch- und Britisch-Ostafrika eingesammelten Staphylinidae. Arch. Naturges. 79(A): 114-168.
- Eichler, W.
1924. Nouvelles espèces et variétés de coléoptères

- Eichler, W. (continued)
du Caucase et de l'Asie-mineure. Polsk. Pismo Ent. 3: 61-68.
- Eppelsheim, E.
1889a. Neue Staphylinen Europas und der angrenzenden Lander. Deutsche Ent. Ztschr. (1889): 161-183.
1889b. Insecta, a Cl. G. N. Potanin in China et in Mongolia novissime lecta. V. Neue Staphylinen. Hor. Soc. Ent. Rossicae 23: 169-184.
1892. Zur Staphylininenfauna Turkestan's. Deutsche Ent. Ztschr. (1892): 321-346.
1895a. Beitrag zur Staphylinenfauna West-Afrika's. Zweites Stuck. Deutsche Ent. Ztschr. (1895): 113-144.
1895b. Esplorazione del Giuba e dei suoi affluenti compiuta dal Cap. V. Bottego durante gli anni 1892-93 sotto gli auspicii della Società Geografica Italiana. Risultati Zoologici. XII: Staphylinidae. Ann. Mus. Civ. Stor. Nat. Genova 35: 197-213.
- Erichson, W.F.
1839. Die Käfer der Mark Brandenburg. Vol. 1, Pt. 2, p. 385-740. Morin; Berlin.
1840. Genera et species Staphylinorum Coleopterorum familiae. Pt. 2, p. 401-954. Morin; Berlin.
1843. Beitrag zur Insecten-Fauna von Angola, in besonderer Beziehung zur geographischen Verbreitung der Insecten in Afrika. Arch. Naturg. 9: 199-267.
- Fabricius, J.C.
1775. Systema entomologiae, sistens Insectorum classes, ordines, genera, species, adiectis synonymis, locis, descriptionibus, observationibus. 832 p. Kortius; Flensburg.
1776. Genera Insectorum eorumque characteres naturales secundum numerum, figuram, situm et proportionem omnium partium oris adiecta mantissa specierum nuper detectarum. 16 + 310 p. Bartsch; Kiel.
1781. Species Insectorum exhibentes eorum differentias specificas, synonyma auctorum, loca natalia, metamorphosin adiectis observationibus, descriptionibus. Vol. 1, viii + 552 p. Bohn; Hamburg.
1787. Mantissa Insectorum sistens eorum species nuper detectas adiectis characteribus genericis, differentiis specificis, emendationibus, observationibus. Vol. 1, xx + 348 p. Proft; Copenhagen.
1792. Entomologia systematica emendata et aucta. Secundum classes, ordines, genera, species adiectis synonymis, locis, observationibus, descriptionibus. Vol. 1, pt. 2, 538 p. Proft; Copenhagen.
1801. Systema eleutheratorum secundum ordines, genera, species: adiectis synonymis, locis, observationibus, descriptionibus. Vol. 2, 687 p. Bibliop. Acad. Nov.; Kiel.
- Fagel, G.
1954a. Contribution à la connaissance des Staphylinidae XXVII. Le complexe des *Paederus ater* Bernh. - *P. testaceus* Bernh. Volume Jubilaire Victor van Straelen 1989-1259.2: 899-967.
1954b. Homonymie. Bull. Ann. Soc. Ent. Belg. 90: 237.
1956. Contributions à l'étude de la faune entomologique de Ruanda Urundi (Mission P. Basilewsky 1953) LXXXV. Coleoptera Staphylinidae Paederini. Ann. Mus. Congo Belge Sci. Zool. 51: 184-200.
1958a. Paederini (Coleoptera Polyphaga) Fam. Staphylinidae. in Exploration du Parc National de l'Upemba, Mission G.F. de Witte. Fasc. 51: 1-470. Institut des Parcs Nationaux du Congo Belge; Brussels.
1958b. Contribution à la connaissance des Staphylinidae. LII. Sur quelques espèces du bassin méditerranéen. Bull. Ann. Soc. R. Ent. Belg. 94: 232-248.
1959. Paederini (Coleoptera Polyphaga) Fam. Staphylinidae. in Exploration du Parc National de la Garamba, Mission H. de Saeger. Fasc. 16: 1-70. Institut des Parcs Nationaux du Congo Belge; Brussels.
1960a. Mission zoologique de l'I.R.S.A.C. en Afrique orientale (R. Basilewsky et N. Leleup 1957) X. Coleoptera Staphylinidae Paederinae. Ann. Mus. Congo Belge, Zool. 81: 114-131.
1960b. Contribution à la connaissance des Staphylinidae LX. Sur quelques Paederini de la dorsale congolaise. Rev. Zool. Bot. Afr. 61: 189-204.
1961a. Contribution à la connaissance des Staphylinidae LXXV. Paederini éthiopiens nouveaux ou mal connus. Bull. Ann. Soc. R. Ent. Belg. 97: 257-286.
1961b. Coleoptera (Staphylinidae): Paederinae (Results of the Lund University Expedition in 1950-1951). South African Animal Life 8: 258-295.
1962a. Contribution à la connaissance des Staphylinidae LXXIV - *Paederidus* nouveaux d'Afrique occidentale. Ent. Arb. Mus. Frey 14: 67-70.
1962b. Contribution à la connaissance des Staphylinidae LXXVI. Paederini nouveaux de l'Angola. *Jeanneliusa* nouveaux de l'Angola. Publ.

- Cult. Cia. Diamant. Angola, Lisbon 58: 13-23, 25-36.
1963. La réserve naturelle intégrale du Mont Nimba XII. Coleoptera Staphylinidae (Piestinae, Osoriinae, Oxytelinae, Paederinae). Mem. Inst. Français Afrique Noire 66: 337-356.
1964. Contribution à la connaissance des Staphylinidae LXXXVII. Deux *Oreopaederus* nouveaux du Katanga. Bull. Ann. Soc. Ent. Belg. 100: 225-229.
1965. Contribution à la connaissance des Staphylinidae LXXXVIII. Sur quelques Paederini africains intéressants. Rev. Zool. Bot. Afr. 71: 129-156.
1966. Contribution à la connaissance des Staphylinidae XCV. Les *Paederus* éthiopiens. Rev. Zool. Bot. Afr. 73: 255-272.
1967. Contribution à la connaissance des Staphylinidae XCVIII. Deux remarquables *Paederus* d'Afrique Occidentale. Rev. Zool. Bot. Afr. 76: 173-177.
1968. Contribution à la connaissance des Staphylinidae CV. Paederini africains nouveaux. Rev. Zool. Bot. Afr. 78: 197-200.
1973. Contribution à la connaissance des Staphylinidae (Coleoptera) CXV. Sur quelques Paederini africains. Rev. Zool. Bot. Afr. 87: 146-158.
- Fairmaire, L.
1859. [Diagnoses de quelques coléoptères européens nouveaux]. Bull. Soc. Ent. France (1859): clxxxiv - clxxxv.
1881. Essai sur les coléoptères des Îles Viti (Fidji). Ann. Soc. Ent. France (sér. 6) 1: 243-318.
- Fairmaire, L. & A. Laboulbène.
1856. Faune entomologique française ou descriptions des insectes qui se trouvent en France. Coléoptères. Vol. 1, p. 371-665. Deyrolle; Paris.
- Faldermann, F.
1839. Fauna entomologica Trans-Caucasica. Pars III. Mem. Soc. Imp. Nat. Moscou 12 (n.s. 6): 1-310.
- Fauvel, A.
1864. Note synonymique sur quelques *Paederus* européens. Bull. Soc. Linn. Normandie 8: 12-14.
1866. Remarques synonymiques. Bull. Soc. Linn. Normandie 10: 22-26.
1868. Faune du Chili. Insectes coléoptères. Staphylinides. (Suite et fin). Bull. Soc. Linn. Normandie (ser. 2) 1: 6-67, pl. 1.
- 1869a. Description d'une espèce nouvelle de *Paederus*. Tijdschr. Ent. 12: 134-135, pl. 7.
- 1869b. Remarques synonymiques sur les staphylinides du catalogus Coleopterorum de MM. v. Harold et Gemminger. Abeille 5: 479-494.
1873. Faune Gallo-Rhénane ou species des insectes qui habitent la France, la Belgique, la Hollande, le Luxembourg, la Prusse Rhénane, le Nassau et le Valais. Coléoptères, Vol. 3, p. 215-390. Le Blanc - Hardel; Caen.
1874. Les staphylinides de la Nouvelle-Calédonie. Ann. Soc. Ent. France (sér. 5) 4: 432-438.
- 1877a. Les staphylinides de l'Australie et de la Polynésie. Ann. Mus. Civ. Stor. Nat. Genova 10: 168-298.
- 1877b. Staphylinides du Brésil et de la Plata recueillis par Camille van Volxem. Ann. Soc. Ent. Belg. (Comptes Rendus) 20: xxiv-xxviii.
- 1878a. Les staphylinides des Moluques et de la Nouvelle Guinée. Ann. Mus. Civ. Stor. Nat. Genova 12: 171-315, pl. 1-2.
- 1878b. Les staphylinides de l'Australie et de la Polynésie (2e mémoire). Ann. Mus. Civ. Stor. Nat. Genova 13: 465-598.
- 1879a. Les staphylinides des Moluques et de la Nouvelle Guinée (2e mémoire). Ann. Mus. Civ. Stor. Nat. Genova 15: 63-121.
- 1879b. Staphylinides nouveaux de la coll. Godeffroy [p. 83-84, in] L. Fairmaire. Descriptions de coléoptères nouveaux ou peu connus du Musée Godeffroy. J. Mus. Godeffroy 14: 80-114.
1882. Les staphylinides du Systema Eleutheratorum de Fabricius (1801). Rev. Ent., Caen 1: 211-214.
1884. Réponse aux auteurs du Catalogus Coleopterorum (1883). Rev. Ent., Caen 3: 240-245.
- 1886a. Les staphylinides du nord de l'Afrique. Rev. Ent., Caen 5: 9-100.
- 1886b. Staphylinides des Iles Philippines. Rev. Ent., Caen 5: 143-150.
- 1889a. Catalogo de los coleópteros de Chile por Federico Philippi. Remarques sur les staphylinides. Rev. Ent., Caen 8: 60-61.
- 1889b. Les coléoptères de la Nouvelle-Calédonie et dépendances avec descriptions, notes et synonymies nouvelles (suite). Staphylinides. Rev. Ent., Caen 8: 242-271, 277-285.
1891. Voyage de M. E. Simon au Venezuela (Décembre 1887 - Avril 1888). Rev. Ent., Caen 10: 87-127.
1895. Staphylinides nouveaux de l'Inde et de la Malaisie. Rev. Ent., Caen 14: 180-286.
1897. Catalogue des staphylinides de Barbarie et des Îles Açores, Madères, Salvage et Canaries (4e édition). Rev. Ent., Caen 16: 237-371.

Fauvel, A. (continued)

1901. Voyage de M. le Dr. Ed. Bugnion au Vénézuéla, en Colombie et aux Antilles. Staphylinides. Rev. Ent., Caen 20: 69-91, pl. 1-2.
- 1903a. Mission de M. Maurice Maindron dans l'Inde méridionale. Staphylinides. Rev. Ent., Caen 22: 149-163.
- 1903b. Faune analytique des coléoptères de la Nouvelle-Calédonie. Rev. Ent., Caen 22: 203-378.
- 1904a. Staphylinides de l'Hindoustan et de la Birmanie. Rev. Ent., Caen 23: 43-70.
- 1904b. Les staphylinides du Thierwelt Deutsch-Ost-Afrika: notes et descriptions. Rev. Ent., Caen 23: 284-294.
- 1904c. Staphylinides de Nouvelle Guinée recueillis par l'expédition hollandaise (1903). Rev. Ent., Caen 23: 294-296.
- 1905a. Staphylinides de Java recueillis par M. le Dr. K. Kraepelin et M. le Dr. Koenigsberger en 1904. Mitt. Naturhist. Mus. Hamburg 22: 75-86.
- 1905b. Staphylinides nouveaux de Madagascar, 2e partie. Rev. Ent., Caen 24: 149-184.
1907. Voyage de M. Ch. Alluaud dans l'Afrique orientale. Staphylinidae. Rev. Ent., Caen 26: 10-70.
1908. Rectification. Rev. Ent., Caen 27: 95.

Ferreira, R. do Nascimento.

1962. Contribuições para um catalogo da familia Staphylinidae da fauna Portuguesa I - Estado actual do conhecimento dos estafilinideos Portugueses. Mem. Estudos Mus. Zool. Univ. Coimbra 280: 1-21.

Focarile, A.

1957. Sistematica, ecologia e geonemia dei *Paederus* del. subgen. *Paederidus* italiani (Col. Staphylinidae). Mem. Soc. Ent. Ital. 36: 65-77.

Fourcroy, A.F. de

1795. Entomologia Parisiensis; sive catalogus Insectorum quae in agro Parisiensi reperiuntur; secundum methodum Geoffraeanam in sectiones genera & species distributus: cui addita sunt nomina trivalia & fere trecentae novae species. Pars prima, viii + 544 p. Aedibus Serpentineis; Paris.

Frank, J.H. & K. Kanamitsu.

1987. *Paederus* sensu lato (Coleoptera: Staphylinidae): Natural history and medical importance. J. Med. Ent. 24: 155-191.

Gaedike, H.

1981. Katalog der in den Sammlungen der Abteilung Taxonomie der Insekten des Institutes für Pflanzenschutzforschung, Bereich Eberswalde (ehemals Deutsches Entomologisches

Institut), aufbewahrten Typen - XIX. Beitr. Ent., Berlin 31: 175-232.

Gahan, C.J.

1900. Staphylinidae [p. 89-90, in] C.W. Andrews (ed.) A monograph of Christmas Island (Indian Ocean), physical features and geology, with descriptions of the fauna and flora by numerous contributors. xiii + 337 p. + 22 pl. British Museum (Natural History), Dept. Geology; London.

Ganglbauer, I.

1895. Die Käfer von Mitteleuropa. Die Käfer der österreichisch-ungarischen Monarchie, Deutschlands, der Schweiz, sowie des französischen und italienischen Alpengebietes. Vol. 2. Familienreihe Staphylinidae. 1. Theil: Staphylinidae, Pselaphidae. vi + 881 p. Carl Gerold's Sohn; Vienna.

Gautier des Cottés, C.

- 1861a. Caractères différentiels de trois espèces nouvelles de *Paederus* propres à la faune française. Ann. Soc. Ent. France (sér. 4) 1: 393-394.
- 1861b. [Genre nouveau de staphylinien et deux nouvelles espèces de *Paederus*]. Bull. Soc. Ent. France (sér. 4) 1: xxxvi.
1862. Genre nouveau de staphylinien et description de nouvelles espèces de coléoptères de Syrie et d'Europe. Ann. Soc. Ent. France (sér. 4) 2: 75-78.
1866. Descriptions d'espèces nouvelles de carabiques d'Europe, suivies d'observations synonymiques sur deux *Paederus* décrits par moi dans les an. de la S. Ent. de France. Mitt. Schweiz. Ent. Ges. 2: 110-114.

Gemminger, M. & E. von Harold

1868. Catalogus Coleopterorum hucusque descriptorum synonymicus et systematicus. Vol. 2, p. 425-752. Gummi; Munich.

Germar, E.F.

1848. Beiträge zur Insectenfauna von Adelaide. Linn. Ent. 3: 153-247.

Gerstaecker, A.

1867. Beitrag zur Insekten-Fauna von Zanzibar, nach dem während der Expedition des Baron v. d. Decken gesammelten Material zusammengestellt. Arch. Naturges. 33: 1-49.
1873. Gliederthiere (Insekten, Arachniden, Myriopoden und Isopoden) [Pt. 2, xvi + 542 p. + 18 pl., in] Baron Carl Claus von der Decken's Reisen in Ost-Afrika in den Jahren 1859-1865. Vol. 3: Wissenschaftlichen Ergebnisse. Winter; Leipzig.

- Gistel, J.N.F.X.
1848. Naturgeschichte des Thierreichs für höhere Schulen. xvi + 216 + 4 p. + 32 pl. Hoffman; Stuttgart.
- Goeze, J.A.E.
1777. Entomologische Beyträge zu des Ritter Linne zwölfsten Ausgabe des Natursystems. Vol. 1, 736 p. Weidmanns, Erben & Reich; Leipzig.
- Göldi, E. A.
1913. Die sanitärisch-pathologische Bedeutung der Insekten und verwandten Gliedertiere, namentlich als Krankheits-Erreger und Krankheits-Überträger. 155 p. Friedländer; Berlin.
- Gravenhorst, J.L.C.
1802. Coleoptera microptera Brunsvicensia nec non exoticorum quotquot exstant in collectionibus entomologorum Brunsvicensium in genera familias et species distribuit. lxxvi + 206 p. Reichard; Brunswick.
1806. Monographia Coleopterorum Micropterorum. xvi + 236 + 12 p. + 1 pl. Dieterich; Göttingen.
- Guerin, F.-E.
1830. Insectes [p. 57-302, pl. 1-22, in] L.I. Duperrey. Voyage autour du monde, exécuté par ordre du roi, sur la corvette de sa majesté, La Coquille, pendant les années 1822, 1823, 1824 et 1825, sous la ministère et conformément aux instructions de S.E.M. le Marquis de Clermont-Tonnerre, ministre de la marine; et publié sous les auspices de son excellence Mgr. le Cte de Chabrol, ministre de la marine et des colonies par M. L.I. Duperrey, capitaine de frégate, chevalier de Saint-Louis et membre de la Legion d'Honneur, commandant de l'expédition. Zoologie, Vol. 2, Pars 2, Div. 1, 319 p. and Atlas, Insectes, pl. 1-22. Bertrand; Paris.
- Gyllenhal, L.
1810-1827. Insecta Suecica descripta a Leonardo Gyllenhal. Classis I. Coleoptera sive Eleuterata. Vol.1, pars 2, xx + 660 p. (1810) Leverentz; Mannheim, Vol. 1, pars 4, x + 762 p. (1827) Fleischer; Mannheim.
- Hammond, P.M.
1977. Staphylinidae [p. 22-42, in] R.D. Pope (ed.) A check list of British insects. Part 3, Coleoptera and Strepsiptera. Handbk. Indent. British Insects (R. Ent. Soc. Lond.) 11(3): i-xiv, 1-105.
- Hansen, V.
1951. Biller XV Rovbiller 1. Del. Danmarks Fauna Vol. 57, 274 p. Dansk Naturhist. Foren.; Copenhagen.
- Harrer, G.A.
1784. Beschreibung derjenigen Insecten, welche Herr D. Jacob Christoph Schäffer in CCLXXX ausgemahlten Kupfertafeln unter dem titel Icones Insectorum circa Ratisbonam indigenorum ehemals in drey Theilen herausgegeben hat. I. Theil. Hartschalige Insecten. [32 +] 328 p. Keyser; Regensburg.
- Hatch, M.H.
1957. The beetles of the Pacific Northwest. Part 2: Staphyliniformia. ix + 384 p. Univ. Washington Press; Seattle.
- Heer, O.
1839-1841. Fauna Coleopterorum Helvetica. Pars 1, fasc. 2, p. 145-360 (1839), Pars 1, fasc. 3, p. 361-652 (1841). Orellius & Fuesslin; Zurich.
- Heller, K.M.
1898. Neue Käfer von Celebes III. Abh. Ber. Königl. Zool. Anthropol. Ethnogr. Mus. Dresden 7(3): 1-42.
- Herbst, J.F.W.
1784. Kritisches Verzeichniss meiner Insecten-Sammlung. Fuessly Arch. Insectengesch. 5: 73-151, pl. 24-30.
1786. Erste Mantissee zum Verzeichniss der ersten Klasse meiner Insectensammlung. Fuessly Arch. Insectengesch. 7-8: 153-182, pl. 43-48.
- Herman, L.H.
1981. Revision of the subtribe Dolicoonina of the New World, with discussions of phylogeny and the Old World genera (Staphylinidae, Paederinae). Bull. American Mus. Nat. Hist. 167: 327-520.
- ICZN.
1964. International code of zoological nomenclature adopted by the XV International Congress of Zoology. xix + 176 p. Int. Trust Zool. Nomencl.; London.
1985. International code of zoological nomenclature third edition adopted by the XX General Assembly of the International Union of Biological Sciences. xx + 338 p. Int. Trust Zool. Nomencl.; London.
- Illiger, J.C.W.
1807. Literatur. Mag. Insektenk. 6: 350-361.
- Jarrige, J.
1944. Les *Paederus* de France. Entomologiste 1: 5-9.
1970. Contribution à l'étude des Coleoptera Brachelytra du Massif du Tsaratanana (Madagascar nord). Mém. ORSTOM 37: 31-63.
1971. Contribution à la faune de l'Iran. 21. Coléoptères Brachelytra. Ann. Soc. Ent. France (n.s.) 7: 483-502.

- Jarrige, J. (continued)
 1978. Contribution à l'étude des Coleoptera Brachelytra Staphylinidae du Massif de l'Andringitra (Madagascar centre). Bull. Mus. Natn. Hist. Nat. Paris (Zool.) 353: 245-296.
- Jeannel, R. & J. Jarrige.
 1949. Biospéologica LXVIII. Coléoptères staphylinides (première série). Arch. Zool. Exp. Gén. 86: 255-392.
- Kirschenblatt, Y.D.
 1932. Review of the beetles of the genus *Paederus* Fabr. occurring in the territories of the USSR [in Russian, part in German]. Parazit. Sborn. Zool. Inst. Akad. Nauk USSR 3: 215-222.
- Klug, J.C.F.
 1834. Bericht über eine auf Madagascar veranstaltete Sammlung von Insecten aus der Ordnung Coleoptera. Abh. Akad. Wiss. Berlin (1832) (1): 91-223, pl. 1-5.
 1855. Diagnosen neuen Coleoptera aus Mossambique. Ber. Verhandl. Akad. Wiss. Berlin (1855): 643-666.
- Koch, C.
 1934. Wissenschaftliche Ergebnisse der entomologischen Expeditionen seiner Durchlaucht des Fuersten Alessandro della Torre e Tasso nach Aegypten und auf die Halbinsel Sinai. IV. Staphylinidae. (Coleoptera). Bull. Soc. R. Ent. Egypte (1934): 33-91.
 1937. Diagnoses préliminaires de quelques nouveaux staphylinides de France et du Maroc. Rev. Française Ent. 4: 24-31.
 1938. Über neue und wenig bekannte paläarktische Paederinae. Ent. Bl. 34: 103-116.
 1941. Récoltes de R. Paulian & A. Villiers dans le Haut-Atlas Marocain, 1938 (XI note) Coléoptères Staphylinides. Ann. Soc. Ent. France 105: 53-70.
- Kolbe, H.
 1883. Neue Coleoptera von Westafrika. Berliner Ent. Ztschr. 27: 15-36.
 1886. Beiträge zur Kenntnis der Coleopteren-Fauna Koreas, bearbeitet auf Grund der von Herrn Dr. C. Gottsche während der Jahre 1883 und 1884 in Korea veranstalteten Sammlung; nebst Bemerkungen über die zoo-geographischen Verhältnisse dieses Faunengebietes und Untersuchungen über einen Sinnesapparat im Gaumen von *Misolampidius morio*. Arch. Naturges. 52: 139-240, pl. 10-11.
- Korge, H.
 1969. Über zwei Arten der Gattung *Paederus* aus pontischen Gebirgen (Col. Staphylinidae). Mitt. Deutsch. Ent. Ges. 28: 60-62.
- Kraatz, G.
 1857. Naturgeschichte der Insecten Deutschlands, begonnen von Dr. H. Schaum, G. Kraatz und H. v. Kiesenwetter. p. 377-768. Nicolai; Berlin.
 1859a. Die Staphylinen-Fauna von Ost-Indien, insbesondere der Insel Ceylan. Arch. Naturges. 25: 1-198.
 1859b. Zeitschriftschau: Bulletin de la Société Impériale des Naturalistes de Moscou, publié sous la redaction du Docteur Renard. 1858, I, II. Berliner Ent. Ztschr. (1859): xxxi-xxxii.
 1867. Synonymische Bemerkungen. Deutsche Ent. Ztschr. 10: 413-418.
- Lacordaire, J.T.
 1835. [in] J.B.A. Boisduval & J.T. Lacordaire. Faune entomologique des environs de Paris; ou species general des insectes qui se trouvent dans un rayon de quinze à vingt lieues aux alentours de Paris. Vol. 1, 17 + 696 p. + 3 pl. Méquignon-Marvis; Paris.
- Laporte de Castelnau, F.L.N.
 1835. Etudes entomologiques, ou description d'insectes nouveaux, et observations sur la synonymie. Pt. 1, p. 95-159, pl. 1-4, Carnassiers. Méquignon-Marvis; Paris.
 1840. Histoire naturelle des insectes coléoptères avec une introduction renfermant l'anatomie et la physiologie des animaux articulés par M. Brulle. Vol. 1, cxxv p. + 24 pl., 325 p. + 19 pl. Duménil; Paris.
- Last, H.R.
 1950. New species of Coleoptera (Staphylinidae) from South Africa. Entomologists's Mon. Mag. 86: 136-137.
 1952. New species of Staphylinidae (Col.) from Africa. Entomologists's Mon. Mag. 88: 89-92.
 1966. Coleoptera from southeast Asia (IV). Family Staphylinidae. Mem. Fac. Lib. Arts Educ. Kagawa Univ. Pt. 2, no. 135: 1-10.
 1979. Records of the genus *Paederus* taken by Dr. R. Hornabrook in New Guinea with descriptions of new species (Coleoptera, Staphylinidae). Faun. Abh. Staatl. Mus. Tierk. Dresden 7: 123-138.
 1980. Records of New Guinea Staphylinidae (Coleoptera) in the Hungarian Natural History Museum. Ann. Hist. Nat. Mus. Natn. Hungarici 72: 139-161.
 1984. Recorded and new species of Coleoptera (Staphylinidae, Paederinae) in Papua New-Guinea. Folia Ent. Hungarica 45: 109-126.
 1985. Species of New Guinea *Paederus* (Coleoptera):

- Staphylinidae) in the collection of Bishop Museum. *Internat. J. Ent.* 27: 326-337.
- Latreille, P.A.
 1804. Histoire naturelle, générale et particulière, des crustacées et des insectes. Vol. 9, 416 p. + pl. 74-80. Dufart; Paris.
 1806. Genera Crustaceorum et Insectorum secundum ordinem naturalem in familias disposita, iconibus exemplisque plurimis explicata. Vol. 1, xviii p. + xvi pl. + 302 p. Koenig; Paris.
 1810. Considerations générales sur l'ordre naturel des animaux composant les classes des crustacés, des arachnides et des insectes; avec un tableau méthodique de leurs genres, disposés en familles. 444 p. Schoell; Paris.
- Lea, A.M.
 1904. Descriptions of new species of Australian Coleoptera. Part VII. *Proc. Linn. Soc. New South Wales* 29: 60-107, pl. 4.
 1923. On Australian Staphylinidae (Coleoptera). *Trans. R. Soc. South Australia* 47: 1-53.
 1931. Australasian Coleoptera. *Rec. S. Australian Mus.* 4: 365-408.
- Leconte, J.L.
 1858. Description of new species of Coleoptera, chiefly collected by the United States and Mexican Boundary Commission, under Major W.H. Emory, U.S.A. *Proc. Acad. Nat. Sci. Philadelphia* 10: 58-89.
 1863. New species of North American Coleoptera. Part 1. *Smithsonian Misc. Colln.* 6(167): 1-86.
 1878. [p. 373-434, 470-472, *in*] E.A. Schwarz. The Coleoptera of Florida. *Proc. American Philos. Soc.* 17: 353-472.
- Legosz-Owsianna, D.
 1963. Przegląd krajowych gatunków z rodzaju *Paederus* Fabricius (Coleoptera, Staphylinidae). *Fragm. Faun.* 10: 317-359.
- Leng, C.W.
 1920. Catalogue of the Coleoptera of America, north of Mexico. x + 470 p. Sherman; Mount Vernon, New York.
- Leng, C.W. & A.J. Mutchler.
 1914. A preliminary list of the Coleoptera of the West Indies as recorded to January 1, 1914. *Bull. American Mus. Nat. Hist.* 33: 391-493.
 1917. Supplement to preliminary list of the Coleoptera of the West Indies. *Bull. American Mus. Nat. Hist.* 37: 191-220.
- Lentz, F.L.
 1856. Kugelannsche Rathsel für Entomologen. *Preuss. Provinzialbl.* 56: 49-62.
- Linnaeus, C.
 1758. *Systema naturae per regna tria naturae, secundum classes, ordines, genera, species, cum characteribus, differentiis, synonymis, locis.* 10th edn. Vol. 1, 823 p. Salvius; Stockholm.
 1767. *Caroli a Linné systema naturae.* 12th edn. Vol. 1, pt. 2, p. 533-1327 + 37 p. Salvius; Stockholm.
 1789. *Entomologia, faunae Suecicae descriptionibus aucta; DD. Scopoli, Geoffroy, de Geer, Fabricii, Schrank, &c. speciebus vel in systemate non enumeratis, vel nuperrime detectis, vel speciebus Galliae Australis locupletata, generum specierumque rariorum iconibus ornata; curante & augmente Carolo de Villers.* Vol. 1, 4 + xvi + 766 p. + 3 pl. Piestre & Delamollière; Lyon.
- Lohse, G.A.
 1964. Staphylinidae. I. Die Käfer Mitteleuropas. Vol. 4, 264 p. Goecke & Evers; Krefeld.
- Lucas, P.H.
 1849. Histoire naturelle des animaux articulés. Troisième famille. Les brachélytres [p. 99-128, *in*] *Exploration scientifique de l'Algérie pendant les années 1840, 1841,*
 1842. *Sciences physiques. Zoologie* 2, 590 p. Bertrand; Paris.
 1857. *Entomologie* [Vol. 3, Pt. 1, 204 p. + 14 pl., *in*] *Animaux nouveaux ou rares recueillis pendant l'expédition dans les parties centrales de l'Amérique du Sud, de Rio de Janeiro à Lima, et de Lima au Para; exécutée par ordre du gouvernement français pendant les années 1843 à 1847, sous la direction du comte Francis de Castelnau.* Bertrand; Paris.
- Lynch Arribalzaga, F.E.
 1884. Los estafilinos de Buenos Aires. *Bol. Acad. Nac. Cienc., Cordoba* 7: 6-392.
- MacLeay, W.J.
 1871. Notes on a collection of insects from Gayndah. *Trans. Ent. Soc. New South Wales* 2: 79-205.
- Mannerheim, C.G. von
 1831. Précis d'un nouvel arrangement de la famille des brachélytres de l'ordre des insectes coléoptères. *Mem. Acad. Imp. Sci. St. Petersburg* 1: 415-501.
- Marcuzzi, G.
 1977. Stafilinidi (Col. Staphyl.) raccolti da G. Marcuzzi e collaboratori in Venezuela negli anni 1948-50. *Quaderni Ecol. Anim.* 12: 1-20.
- Marseul, S.A. de
 1866. [Synonymies nouvelles]. *Bull. Soc. Ent. France* (1866): v-viii.

- Marshall, T.A.
1878. Notes on the entomology of the Windward Islands. Trans. Ent. Soc. London (1878): xxvii-xxxviii.
- Miller, L.
1852. [Drei neue Staphyliniden der Wiener Gegend]. Verh. Zool. Bot. Ver. Wien 2: 26-28.
1868. Eine entomologische Reise in die ostgalizischen Karpathen. Verh. Zool. Bot. Ges. Wien 18: 3-34.
- Moore, I. & E.F. Legner.
1975. A catalogue of the Staphylinidae of America north of Mexico (Coleoptera). Univ. California Divn. Agric. Sci., Spec. Publ. 3015: 1-514.
- Motschulsky, T.V. von
1849. Coléoptères reçus d'un voyage de M. Handschuh dans le midi de l'Espagne énumérés et suivis de notes. Bull. Soc. Imp. Nat. Moscou 22: 52-163.
1858. Énumération des nouvelles espèces de coléoptères rapportés de ses voyages. Bull. Soc. Imp. Nat. Moscou 31(2): 634-670.
1860a. Insectes des Indes orientales, et de contrées analogues. 2. de série. Etudes Ent. 8: 25-118.
1860b. Énumération des nouvelles espèces de coléoptères rapportées de ses voyages. Bull. Soc. Imp. Nat. Moscou 33(2): 539-588.
- Mulsant, E. & C. Rey.
1878. Histoire naturelle des coléoptères de France. Brévipennes: Pédériens-Evesthétiens. 354 p. + 6 pl. Deyrolle; Paris.
- Muona, J.
1979. Staphylinidae [p. 14-28, in] H. Silfverberg (ed.) Enumeratio Coleopterorum Fennoscandiae et Daniae. Helsingin Hyonteisvaihtoyhdistys; Helsinki.
- Olivier, A.G.
1795. *Paederus* [Coléoptères, Vol. 3, pt. 44, p. 1-7, in] Entomologie ou histoire naturelle des insectes avec leurs caractères génériques et spécifiques, leur synonymie, et leur figure enluminée. Lanneau; Paris.
1812. Insectes [Vol. 8, pt. 2, p. 361-722, in] Encyclopédie méthodique. Histoire naturelle. Agasse; Paris.
- Outereho, R.
1980. *Parameropaederus carpeticus* n. sp. (Staphylinidae: Paederinae). Nouv. Rev. Ent. 10: 31-35.
- Panzer, G.W.F.
1793-1809. Faunae Insectorum Germanicae initia, oder Deutschlands Insecten gesammelt und herausgegeben. Hft. 9: 1-24 (1793a), Hft. 11: 1-24 (1793b), Hft. 27: 1-24 (1796a), Hft. 36: 1-24 (1796b), Hft. 43: 1-24 (1797), Hft. 104: 1-24 (1809). Felsecker; Nuremberg.
- Paulian, R.
1947. Observations sur la faune du Pic Bintmane (Sierra Leone) d'après les récoltes de M. P. Jaeger. Notes Africaines, I.F.A.N., Dakar 34: 27-28.
- Paykull, G. von
1789. Monographia Staphylinorum Sueciae. 82 p. Edman; Uppsala.
1790. Monographia Caraborum Sueciae. 138 p. Edman; Uppsala.
1800. Fauna Suecica. Insecta. Vol. 3, 459 p. Edman; Uppsala.
- Perroud, B.-P.
1865. [Staphyliniens, p. 83-89, in] B.-P. Perroud & A. Montrouzier. Essai sur la faune entomologique de Kanala (Nouvelle-Calédonie) et description de quelques espèces nouvelles ou peu connues. Ann. Soc. Linn. Lyon (sér. 2) 11: 46-257.
- Perty, J.A.M.
1830. Insecta Brasiliensia, p. 1-60, pl. 1-12, in C.F.P. de Martius (ed.) Delectus animalium articulatum, quae in itinere per Brasiliam annis MDCCCXVII-MDCCCXX jussu et auspiciis Maximiliani Josephi I. Bavariae regis augustissimi peracto collegerunt Dr. J.B. de Spix et Dr. C.F.P. de Martius. Fasc. 1. Imp. Edit.; Munich.
- Peyerimhoff, P. de
1931. Mission scientifique du Hoggar envoyée de février à mai 1928 par M. Pierre Bordes Gouverneur Général de l'Algérie. Coléoptères. Mem. Soc. Hist. Nat. Afrique Nord 2: 1-173, pl. 1-3, 2 maps.
1949. Note sur quelques staphylinides marocains. Bull. Soc. Ent. France (1949): 104-110.
- Peyron, E.
1858. Catalogue des coléoptères des environs de Tarsous (Caramanie), avec la description des espèces nouvelles. Ann. Soc. Ent. France (ser. 3) 6: 353-434.
- Plavilstschikov, N.N.
1929. Les staphylinides récoltés par M. Nicolas Iconicov dans le Pérou. Ent. Nachrichtenbl. 3: 29-30.
- Puthz, V.
1969. Staphylinidae (Coleoptera) from New Guinea. Pacific Insects 11: 529-534.
- Quedenfeldt, M.
1881. Vier neue Staphylinen-Arten aus dem Mittelmeer-Faunengebiet. Berliner Ent. Ztschr. (1881): 291-293.

Reiche, L.

1847. Entomologie [Vol. 3, p. 257-471, et Atlas, pt. 1, pl. 16-33, in] P.V.A. Ferret et J.G. Galinier. Voyage en Abyssinie dans les provinces du Tigre, du Samen et de l'Amhara. Paulin; Paris.

Reitter, E.

1884. *Paederus pelikani*, eine neue Art von den jonischen Inseln. Wiener Ent. Ztg. 3: 43-45.
 1885. Bemerkungen zur den Catalogs-Berichtigungen des Herrn. A. Fauvel in der Revue d'Entomologie (1884). Wiener Ent. Ztg. 4: 151-156.
 1889. Uebersicht der bekannten Arten der Coleopteren-Gattung *Paederus*. Ent. Nachrichten 15: 169-171.
 1890. Coleopterologische Notizen XXXIV. Wiener Ent. Ztg. 9: 13-15.

Rossi, P.

1790. Fauna Etrusca sistens Insecta quae in provinciis Florentina et Pisana praesertim collegit. Vol. 1, 272 p. Masi; Livorno.

Roubal, J.

1932. Sur quelques coléoptères des environs de Bagdad. Bull. Soc. Ent. France 37: 59-64.

Sahlberg, J.R.

1876. Enumeratio Coleopterorum Brachelytrorum Fenniae I. Staphylinidae. Acta Soc. Faun. Flor. Fennica 1: 1-247.

Samouelle, G.

1819. The entomologist's useful compendium; or introduction to the knowledge of British insects, containing the best means of obtaining and preserving them, and a description of the apparatus generally used; together with the genera of Linné, and the modern method of arranging the classes Crustacea, Myriapoda, spiders, mites and insects, from the affinities and structure, according to the views of Dr. Leach. 496 p. Boys; London.

Sanderson, M.W.

1967. Revision of the Greater Antillean *Paederus* (Coleoptera: Staphylinidae). Coleopt. Bull. 21: 59-62.

Saulcy, F.H.C. de

1862. [Sur trois *Paederus* décrits par M. Gautier]. Bull. Soc. Ent. France (sér. 4) 2: vi-vii.
 1864. Descriptions des espèces nouvelles de coléoptères recueillis en Syrie, en Egypte et en Palestine pendant les mois d'Octobre 1863 à Janvier 1864, par M. de Saulcy. 2e partie. Ann. Soc. Ent. France (sér. 4) 4: 629-660.

Say, T.

1823. Descriptions of coleopterous insects collected in the late expedition to the Rocky Mountains,

performed by order of Mr. Calhoun, Secretary of War, under the command of Major Long. J. Acad. Nat. Sci. Philadelphia 3: 139-216.

1834. Descriptions of new North American insects and observations on some already described. Trans. American Philos. Soc. 4: 409-470.

Schaeffer, J.C.

1779. Icones Insectorum circa Ratisbonam indigenorum coloribus naturam referentibus expressae. Vol. 1, pt. 1, pl. 1-50. Breitfeld; Regensburg.

Scheerpeltz, O.

1933. Staphylinidae II [in] Coleopterorum Catalogus (W. Junk, Berlin) 129: 989-1500.

1935. Staphyliniden aus dem Sundainseln und Nordaustralien. Rev. Suisse Zool. 42: 593-659.

- 1957a. Vorläufige Diagnosen einiger neuen palaarktischen Arten und Formen der Gattungen *Paederidus* Muls. Rey, *Paederus* Fabr. (mit den neuen Untergattungen *Eopaederus*, *Paederus* s. str. nov., *Heteropaederus*, *Dioncopaederus* und *Oedopaederus*), *Parameropaederus* nov. gen., *Lobopaederus* nov. gen. und *Megalopaederus* nov. gen. Mem. Mus. Stor. Nat. Venezia Tridentina 11: 447-475.

- 1957b. Staphylinidae (Col.) von Sumba und Flores (4. Beitrag zur Kenntnis der Staphyliniden der orientalischen Region). Verh. Naturf. Ges. Basel 68: 217-357.

1960. Zur Kenntnis neotropischer Staphyliniden (Col.). Beitr. Neotrop. Fauna 2: 65-141.

1963. Coleoptera, Staphylinidae. 116. Beitrag zur Kenntnis der paläarktischen Staphyliniden. Lunds Univ. Arsskr. (N.F. 2) 58: 1-38.

- 1965a. Die bis jetzt aus Venezuela bekannt gewordenen Arten der Gattung *Paederus* Fabricius. Acta Biol. Venezuel. 4: 213-227.

- 1965b. Wissenschaftliche Ergebnisse der Schwedischen Expedition 1934 nach Indien und Burma. Coleoptera Staphylinidae (except. Megalopsidiinae et Steninae). Ark. Zool. 17: 93-371.

1966. Ergebnisse der Österreichischen Neukaledonien-Expedition

1965. Staphylinidae. Ann. Naturhist. Mus. Wien 69: 393-418.

1968. Coleoptera - Staphylinidae [pt. xvfa, p. 1-279, in] Catalogus faunae Austriae. Österreichische Staatsdruckerei; Vienna.

1976. Wissenschaftliche Ergebnisse entomologischer Aufsammlungen in Nepal (Col. Staphylinidae). Khumbu Himal 5: 77-173.

1978. Wissenschaftliche Ergebnisse der von Dr. Ra-

- Scheerpeltz, O. (continued)
 ul do Nascimento Ferreira-Coimbra, Portugal während seines Aufenthaltes von Ende des Jahres 1964 bis zum Anfang des Jahres 1966 auf der Insel Timor, der östlichsten der kleinen Sundainseln, durchgeführten Aufsammlungen von Staphyliniden (Coleoptera). *Eos, Madrid* 52: 185-232.
- Schrank, F. von Paula von
 1781. *Enumeratio Insectorum Austriae indigenorum*. [22 +] 548 [+ 4] p. + 4 pl. Eberhard, Klet & Franck; Augsburg.
- Shubert, K.
 1906. Beitrag zur Staphylinidenfauna Deutsch-Ostafrika. *Deutsche Ent. Ztschr.* (1906): 371-377.
- Scopoli, J.A.
 1763. *Entomologia Carniolica exhibens Insecta Carnioliae indigena et distributa in ordines, genera, species, varietates. Methodo Linnaeana*. 40 + 421 p. + 43 pl. Trattner; Vienna.
- Sharp, D.
 1874. The Staphylinidae of Japan. *Trans. Ent. Soc. London* (1874): 1-103.
 1876a. Contributions to an insect fauna of the Amazon Valley: Coleoptera - Staphylinidae. *Trans. Ent. Soc. London* (1876): 27-424.
 1876b. Description of a new genus, and some new species, of Staphylinidae from Mexico and Central America. *Trans. Ent. Soc. London* (1876): 425-432.
 1886. Staphylinidae [Vol. 1, pt. 2, p. 537-672, *in*] *Coleoptera [in] Insecta [in] Biologia Centrali-Americana*.
 1891. Coleoptera - (continued). [p. 40-44, *in*] E. Whymper (ed.) *Supplementary appendix to travels amongst the great Andes of the Equator*. xxii + 147 p. John Murray; London.
- Shibata, Y.
 1973. Preliminary check list of the family Staphylinidae of Taiwan (Insecta: Coleoptera). *Annl. Bull. Nichidai Sanko* 16: 21-88.
 1977. Provisional check list of the family Staphylinidae of Japan. II (Insecta: Coleoptera). *Annl. Bull. Nichidai Sanko* 20: 16-83.
- Smetana, A.
 1959. Zur Kenntnis der Staphyliniden-Fauna Albanien (Col., Staphylinidae). *Acta Ent. Mus. Natn. Pragae* 33: 195-218, pl. 10-17.
 1962. Zwei neue Arten der Untergattung *Harpopaederus* des Gattung *Paederus*. *Reichenbachia* 8: 45-50.
 1964. Systematische und faunistische Beiträge zur Kenntnis der Staphyliniden-Fauna der Tschechoslowakei VI (Col., Staph.). *Cas. Csl. Spol. Ent.* 61: 162-172.
 1979. "*Staphylinus fulgidus*" as the type species of several staphylinid genera (Insecta, Coleoptera, Staphylinidae) Z.N.(S)2221. *Bull. Zool. Nomencl.* 36: 44-52.
- Solsky, S.M. de
 1866. Deux staphylins nouveaux du Mexique. *Hor. Soc. Ent. Rossicae* 4: 105-107.
 1871. Prémices d'une faune entomologique de la vallée de Zaravschén, dans l'Asie centrale. *Hor. Soc. Ent. Rossicae* 8: 133-165, pl. 5.
- Steel, W.O.
 1955. New species of Staphylinidae from Australia. *Proc. Linn. Soc. New South Wales* 80: 180-185.
- Stephens, J.F.
 1829. A systematic catalogue of British insects; being an attempt to arrange all the hitherto discovered indigenous insects in accordance with their natural affinities. Pt. 1, xxxv + 416 p. Baldwin & Cradock; London.
 1833. Illustrations of British entomology; or, a synopsis of indigenous insects: containing their generic and specific distinctions; with an account of their metamorphoses, times of appearance, localities, food, and economy, as far as practicable. *Mandibulata*, Vol. 5, p. 241-304, pl. 27. Baldwin & Cradock; London.
- Szujecki, A.
 1965a. Some remarks on the palearctic species of Paederinae (Coleoptera Staphylinidae). *Bull. Acad. Polonaise Sci. Cl. II, Ser. Sci. Biol.* 13: 83-85.
 1965b. Klucze do oznaczania owadów polski. *Polski Zwiasek Entomologiczny* 19(24c): 1-74.
- Thomson, C.G.
 1860. Skandinaviens Coleoptera, synoptisk bearbetade. Vol. 2, 304 p. Berlinoska; Lund.
- Tikhomirova, A.I.
 Morpho-ecological characteristics and phylogeny of Staphylinidae (with a catalogue of the fauna of the U.S.S.R.). 191 p. Nauka; Moscow. [in Russian].
- Tottenham, C.E.
 1953. Staphylinidae from Kilimanjaro, Mt. Kenya and the Ruwenzori. *Ann. Mag. Nat. Hist. (ser. 12)* 6: 481-512.
- Tronquet, M.
 1981. Staphylinidae d'Afghanistan (Coleoptera). *Rev. Française Ent. (n.s.)* 3: 69-83.
- Walker, F.
 1858. Characters of some apparently undescribed

- Ceylon insects. Ann. Mag. Nat. Hist. (ser. 3) 2: 202-209.
- Waltl, J.
1838. Beiträge zur nahern naturhistorischen Kenntnis des Unterdonaukreises in Bayern. Isis Oken 4: column 250-273.
- Wasmann, E.
1911. Zur Kenntnis der Termiten und Termitengäste vom Belgischen Congo. Rev. Zool. Afr. 1: 91-117, pl. 3-5.
- Watanabe, Y.
1986. A revisional study of the Japanese species of the genus *Megalopaederus* (Coleoptera, Staphylinidae). Ent. Papers Presented Kurosawa, Tokyo: 159-169.
- Weise, J.
1877. Zwei neue japanische Staphylinen. Deutsche Ent. Ztschr. (1877): 367-368.
- Wendeler, H.
1926a. Neue exotische Staphylinidae (Col.). (7. Beitrag zur Kenntnis der Staphyliniden). Deutsche Ent. Ztschr. (1926): 71-75.
1926b. Neue exotische *Paederus*-Arten (Col. Staph.). (8. Beitrag zur Kenntnis der Staphyliniden). Deutsche Ent. Ztschr. (1926): 199-203.
1926c. Ein neuer *Paederus* aus Rumänien (Col. Staph.). (9. Beitrag zur Kenntnis der Staphyliniden). Deutsche Ent. Ztschr. (1926): 204.
1926d. Fauna Buruana. Genus *Paederus* Fabricius (Col. Staph.). (6. Beitrag zur Kenntnis der Staphyliniden). Treubia 7: 328-330.
1927. Ein neues Subgenus und eine neue Species des Genus *Paederus* (Col., Staphyl.). (15. Beitrag zur Kenntnis der Staphyliniden). Neue Beitr. Syst. Insektenk. 4: 1-2.
1928a. *Paederognathus* nom. nov. (Col. Staphylin.). Deutsche Ent. Ztschr. (1928): 37.
1928b. Zwei neue paläarktische Staphyliniden (Col.). (14. Beitrag zur Kenntnis der Staphyliniden). Col. Centralbl. 2: 263-265.
1928c. Neue exotische Staphyliniden (Coleoptera). (16. Beitrag zur Kenntnis der Staphyliniden). Neue Beitr. Syst. Insektenk. 4: 32-35.
1930. Neue exotische Staphyliniden (Coleoptera). (17. Beitrag zur Kenntnis der Staphyliniden). Neue Beitr. Syst. Insektenk. 4: 181-192, 248-252.
1931. Eine Reihe neuer *Paederus*-Arten aus verschiedenen exotischen Faunengebieten (Col. Staph.). (12. Beitrag zur Kenntnis der Staphyliniden). Deutsche Ent. Ztschr. (1931): 37-48.
1933. Sechs neue *Paederus*. (18. Beitrag zur Kenntnis der Staphyliniden) (Col.). Mitt. Deutsche Ent. Ges. 4: 55-58.
1942. Drei neue *Paederus*-Arten des Deutschen Entomologischen Instituts aus dem tropischen Amerika (Coleoptera: Staphylinidae). (19. Beitrag zur Kenntnis der Staphyliniden). Arb. Morph. Taxon. Ent. 9: 102-103.
- Wickham, H.F.
1913. Fossil Coleoptera from the Wilson ranch near Florissant, Colorado. Univ. Iowa Bull. Lab. Nat. Hist. 6(4): 3-29, pl. 1-7.
- Wiedemann, C.R.W.
1823. Zweihundert neue Käfer von Java, Bengalen und dem Vorgebirge der guten Hoffnung. Zool. Mag. 2(1): 1-135.
- Wollaston, T.V.
1867. Coleoptera Hesperidum, being an account of the coleopterous insects of the Cape Verde archipelago [with appendix containing addenda etc. to the Coleoptera Atlantidum]. xxxix + 285 p. + 1 map. Van Voorst; London.
- Wu, C.-F.
1937. Catalogus Insectorum Sinensium (Catalogue of Chinese insects). Vol. 3, x + 1312 p. Fan Memorial Inst. Biol.; Beijing.
- Zetterstedt, J.W.
1838. Coleoptera [columns 9-240, in] Insecta Lapponica descripta. Voss; Leipzig.