

Acta Crystallographica Section E

Structure Reports

Online

ISSN 1600-5368

Ethyl 4-[3-(2-methylbenzoyl)thioureido]-benzoate

Aamer Saeed,^{a*} Hummera Rafique,^a Amara Mumtaz^a and Michael Bolte^b

^aDepartment of Chemistry, Quaid-i-Azam University, Islamabad 45320, Pakistan., and ^bInstitut für Anorganische Chemie, J. W. Goethe-Universität Frankfurt, Max-von-Laue-Str. 7, 60438 Frankfurt/Main, Germany.

Correspondence e-mail: aamersaeed@yahoo.com


Received 20 September 2009; accepted 13 October 2009

Key indicators: single-crystal X-ray study; $T = 173$ K; mean $\sigma(\text{C}-\text{C}) = 0.002$ Å; disorder in main residue; R factor = 0.037; wR factor = 0.100; data-to-parameter ratio = 18.9.

The molecular conformation of the title compound, $\text{C}_{18}\text{H}_{18}\text{N}_2\text{O}_3\text{S}$, is stabilized by an intramolecular $\text{N}-\text{H}\cdots\text{O}$ hydrogen bond. The crystal packing shows centrosymmetric dimers connected by $\text{N}-\text{H}\cdots\text{S}$ hydrogen bonds. The terminal ethoxy substituents are statistically disordered [occupancy ratio 0.527 (5):0.473 (5)].

Related literature

For the use of thiourea derivatives in organic synthesis and analysis, see: Eynde & Watte (2003); Fu *et al.* (1999); Rashdan *et al.* (2006); Maryanoff *et al.* (1986); Wang *et al.* (2005); Saeed *et al.* (2008); Koch, (2001). For their bioactivity and pharmaceutical applications, see: Upadhyaya & Srivastava (1982); Ramadas *et al.* (1998); Blum & Hayes (1979); DeBeer *et al.* (1936). For related structures, see: Saeed & Flörke (2007a,b); Saeed *et al.* (2009).



Experimental

Crystal data

$\text{C}_{18}\text{H}_{18}\text{N}_2\text{O}_3\text{S}$
 $M_r = 342.40$
 Triclinic, $P\bar{1}$
 $a = 7.4555$ (3) Å

$b = 7.6311$ (4) Å
 $c = 15.2468$ (8) Å
 $\alpha = 96.456$ (4)°
 $\beta = 103.860$ (5)°

$\gamma = 92.908$ (4)°
 $V = 834.13$ (7) Å³
 $Z = 2$
 Mo $K\alpha$ radiation

$\mu = 0.21$ mm⁻¹
 $T = 173$ K
 $0.33 \times 0.32 \times 0.28$ mm

Data collection

Stoe IPDS II two-circle-diffractometer
 Absorption correction: multi-scan (MULABS; Spek, 2009; Blessing, 1995)
 $T_{\min} = 0.933$, $T_{\max} = 0.943$
 22798 measured reflections
 4659 independent reflections
 4311 reflections with $I > 2\sigma(I)$
 $R_{\text{int}} = 0.057$

Refinement

$R[F^2 > 2\sigma(F^2)] = 0.037$
 $wR(F^2) = 0.100$
 $S = 1.04$
 4659 reflections
 246 parameters
 H-atom parameters constrained
 $\Delta\rho_{\text{max}} = 0.28$ e Å⁻³
 $\Delta\rho_{\text{min}} = -0.37$ e Å⁻³

Table 1

Hydrogen-bond geometry (Å, °).

$D-H\cdots A$	$D-H$	$H\cdots A$	$D\cdots A$	$D-H\cdots A$
$\text{N1}-\text{H1}\cdots\text{O1}$	0.80	2.01	2.669 (1)	139
$\text{N2}-\text{H2}\cdots\text{S1}^i$	0.89	2.67	3.5551 (9)	170

Symmetry code: (i) $-x + 2, -y, -z + 1$.

Data collection: *X-AREA* (Stoe & Cie, 2001); cell refinement: *X-AREA*; data reduction: *X-AREA*; program(s) used to solve structure: *SHELXS97* (Sheldrick, 2008); program(s) used to refine structure: *SHELXL97* (Sheldrick, 2008); molecular graphics: *XP* in *SHELXTL-Plus* (Sheldrick, 2008); software used to prepare material for publication: *SHELXL97*.

Supplementary data and figures for this paper are available from the IUCr electronic archives (Reference: IM2145).

References

- Blessing, R. H. (1995). *Acta Cryst.* **A51**, 33–38.
 Blum, J. J. & Hayes, A. (1979). *J. Supramol. Struct.* **12**, 23–34.
 DeBeer, E. J., Buck, J. S., Ide, W. S. & Hjort, A. M. (1936). *J. Pharmacol.* **57**, 19–33.
 Eynde, J. J. V. & Watte, O. (2003). *Arkivoc*, **iv**, 93–101.
 Fu, M., Fernandez, M., Smith, M. L. & Flygaa, J. A. (1999). *Org. Lett.* **1**, 1351–1353.
 Koch, K. R. (2001). *Coord. Chem. Rev.* **216–217**, 473–488.
 Maryanoff, C. A., Stanzione, R. C., Plampin, J. N. & Mills, J. E. (1986). *J. Org. Chem.* **51**, 1882–1884.
 Ramadas, K., Suresh, G., Janarthanan, N. & Masilamani, S. (1998). *Pestic. Sci.* **52**, 145–151.
 Rashdan, S., Light, M. E. & Kilburn, J. D. (2006). *Chem. Commun.* pp. 4578–4580.
 Saeed, A. & Flörke, U. (2007a). *Acta Cryst.* **E63**, o4259.
 Saeed, A. & Flörke, U. (2007b). *Acta Cryst.* **E63**, o4614.
 Saeed, A., Khera, R. A., Simpson, J. & Stanley, R. G. (2009). *Acta Cryst.* **E65**, o1735–o1736.
 Saeed, A., Zaman, S. & Bolte, M. (2008). *Synth. Commun.* **38**, 2185–2199.
 Sheldrick, G. M. (2008). *Acta Cryst.* **A64**, 112–122.
 Spek, A. L. (2009). *Acta Cryst.* **D65**, 148–155.
 Stoe & Cie (2001). *X-AREA*. Stoe & Cie, Darmstadt, Germany.
 Upadhyaya, J. S. & Srivastava, P. K. (1982). *J. Indian Chem. Soc.* **59**, 767–769.
 Wang, X.-C., Wang, F., Quan, Z.-J., Wang, M.-G. & Li, Z. (2005). *J. Chem. Res.* **61**, 689–690.

supplementary materials

Acta Cryst. (2009). E65, o2774 [doi:10.1107/S160053680904183X]

Ethyl 4-[3-(2-methylbenzoyl)thioureido]benzoate

A. Saeed, H. Rafique, A. Mumtaz and M. Bolte

Comment

The background of this study has been described in our earlier paper concerning the crystal structure of 1-(2-Chloro-5-nitrophenyl)-3-(2,2-dimethylpropionyl)thiourea (Saeed *et al.*, 2009). As part of our work on the structure of thioureas, we report here the structure of the title derivative, I, Fig 1.

The molecular conformation of the title compound, C₁₈H₁₈N₂O₃S, is stabilized by intramolecular N—H···O hydrogen bonds. The crystal packing shows centrosymmetric dimers connected by N—H···S hydrogen bonds (Table 1). Terminal ethoxy substituents are statistically disordered.

Experimental

A solution of 2-methylbenzoyl chloride (10 mmol) in acetone (50 ml) was added dropwise to a suspension of potassium thiocyanate (10 mmol) in acetone (30 ml) and the reaction mixture was refluxed for 30 min. After cooling to room temperature, a solution of 4-aminobenzoic acid ethyl ester (10 mmol) in acetone (10 ml) was added and the resulting mixture refluxed for 3 h. The reaction mixture was poured into cold water and the precipitated thiourea was recrystallized from aqueous ethanol. Anal. calcd. for C₁₈H₁₈N₂O₃S: C, 63.14; H, 5.30; N, 8.18; S, 9.36% found: C, 63.26; H, 5.34; N, 8.21; S, 9.27%;

Refinement

H atoms were positioned geometrically and refined using a riding model with fixed individual displacement parameters [$U(H) = 1.2 U_{eq}(C,N)$ or $U(H) = 1.5 U_{eq}(C_{methyl})$] using a riding model with $C—H(\text{aromatic}) = 0.95 \text{ \AA}$, $C—H(\text{methyl}) = 0.98 \text{ \AA}$, or $C—H(\text{methylene}) = 0.99 \text{ \AA}$, respectively. H atoms bonded to N were set to the position where they were found in the difference map. The ethoxy group is disordered over two positions with a site occupation factor of 0.527 (5) for the major occupied site.

Figures




Fig. 1. Perspective view of the title compound. The disordered atoms of the minor occupied site have been omitted for clarity. Displacement ellipsoids are shown at the 50 % probability level.




Fig. 2. Packing diagram of the title compound. Hydrogen bonds shown as dashed lines. The minor occupied sites are omitted for clarity.

Ethyl 4-[3-(2-methylbenzoyl)thioureido]benzoate

Crystal data

$C_{18}H_{18}N_2O_3S$

$M_r = 342.40$

Triclinic, $P\bar{1}$

Hall symbol: -P 1

$a = 7.4555$ (3) Å

$b = 7.6311$ (4) Å

$c = 15.2468$ (8) Å

$\alpha = 96.456$ (4)°

$\beta = 103.860$ (5)°

$\gamma = 92.908$ (4)°

$V = 834.13$ (7) Å³

$Z = 2$

$F_{000} = 360$

$D_x = 1.363$ Mg m⁻³

Mo $K\alpha$ radiation, $\lambda = 0.71073$ Å

Cell parameters from 42517 reflections

$\theta = 3.4$ – 29.9 °

$\mu = 0.21$ mm⁻¹

$T = 173$ K

Block, colourless

$0.33 \times 0.32 \times 0.28$ mm

Data collection

Stoe IPDS II two-circle-diffractometer

Radiation source: fine-focus sealed tube

Monochromator: graphite

$T = 173$ K

ω scans

Absorption correction: multi-scan (MULABS; Spek, 2009; Blessing, 1995)

$T_{\min} = 0.933$, $T_{\max} = 0.943$

22798 measured reflections

4659 independent reflections

4311 reflections with $I > 2\sigma(I)$

$R_{\text{int}} = 0.057$

$\theta_{\max} = 29.6$ °

$\theta_{\min} = 3.4$ °

$h = -10 \rightarrow 10$

$k = -10 \rightarrow 10$

$l = -21 \rightarrow 21$

Refinement

Refinement on F^2

Least-squares matrix: full

Secondary atom site location: difference Fourier map

Hydrogen site location: inferred from neighbouring sites

$$R[F^2 > 2\sigma(F^2)] = 0.037$$

$$wR(F^2) = 0.100$$

$$S = 1.04$$

4659 reflections

246 parameters

Primary atom site location: structure-invariant direct methods

H-atom parameters constrained

$$w = 1/[\sigma^2(F_o^2) + (0.0517P)^2 + 0.2098P]$$

$$\text{where } P = (F_o^2 + 2F_c^2)/3$$

$$(\Delta/\sigma)_{\max} = 0.002$$

$$\Delta\rho_{\max} = 0.28 \text{ e } \text{\AA}^{-3}$$

$$\Delta\rho_{\min} = -0.37 \text{ e } \text{\AA}^{-3}$$

Extinction correction: none

Special details

Geometry. All esds (except the esd in the dihedral angle between two l.s. planes) are estimated using the full covariance matrix. The cell esds are taken into account individually in the estimation of esds in distances, angles and torsion angles; correlations between esds in cell parameters are only used when they are defined by crystal symmetry. An approximate (isotropic) treatment of cell esds is used for estimating esds involving l.s. planes.

Refinement. Refinement of F^2 against ALL reflections. The weighted R -factor wR and goodness of fit S are based on F^2 , conventional R -factors R are based on F , with F set to zero for negative F^2 . The threshold expression of $F^2 > \sigma(F^2)$ is used only for calculating R -factors(gt) *etc.* and is not relevant to the choice of reflections for refinement. R -factors based on F^2 are statistically about twice as large as those based on F , and R -factors based on ALL data will be even larger.

Fractional atomic coordinates and isotropic or equivalent isotropic displacement parameters (\AA^2)

	<i>x</i>	<i>y</i>	<i>z</i>	$U_{\text{iso}}^*/U_{\text{eq}}$	Occ. (<1)
S1	1.09603 (4)	0.22920 (4)	0.599200 (16)	0.02926 (8)	
O1	0.54635 (11)	0.37544 (11)	0.42973 (5)	0.03282 (18)	
O2	1.1748 (4)	1.1151 (2)	0.86115 (13)	0.0302 (5)	0.527 (5)
O2A	1.0912 (4)	1.1206 (2)	0.88161 (14)	0.0309 (6)	0.473 (5)
O3	1.23261 (17)	0.91072 (12)	0.96012 (6)	0.0521 (3)	
N1	0.82261 (12)	0.44800 (11)	0.57980 (5)	0.02358 (17)	
H1	0.7282	0.4728	0.5478	0.028*	
N2	0.77873 (12)	0.19546 (11)	0.47488 (6)	0.02420 (17)	
H2	0.8242	0.0956	0.4570	0.029*	
C1	0.89035 (13)	0.29850 (13)	0.55169 (6)	0.02227 (18)	
C2	0.61524 (13)	0.23496 (13)	0.41838 (6)	0.02368 (18)	
C11	0.90732 (12)	0.57062 (12)	0.65792 (6)	0.02124 (17)	
C12	0.98398 (14)	0.51635 (13)	0.74226 (7)	0.02457 (19)	
H12	0.9840	0.3941	0.7490	0.029*	
C13	1.06022 (14)	0.64342 (13)	0.81625 (7)	0.02587 (19)	
H13	1.1131	0.6075	0.8739	0.031*	
C14	1.06025 (15)	0.82288 (13)	0.80705 (7)	0.0273 (2)	
C15	0.97895 (15)	0.87601 (13)	0.72331 (7)	0.0286 (2)	
H15	0.9755	0.9983	0.7170	0.034*	
C16	0.90262 (14)	0.74958 (13)	0.64881 (6)	0.02458 (19)	
H16	0.8471	0.7856	0.5915	0.029*	
C17	1.1522 (2)	0.95152 (16)	0.88788 (9)	0.0456 (3)	
C18	1.2785 (4)	1.2514 (3)	0.93166 (16)	0.0336 (6)	0.527 (5)
H18A	1.3222	1.3504	0.9033	0.040*	0.527 (5)

supplementary materials

H18B	1.3886	1.2024	0.9681	0.040*	0.527 (5)
C19	1.1602 (4)	1.3183 (4)	0.9924 (2)	0.0404 (6)	0.527 (5)
H19A	1.2319	1.4107	1.0394	0.061*	0.527 (5)
H19B	1.1193	1.2206	1.0215	0.061*	0.527 (5)
H19C	1.0518	1.3673	0.9564	0.061*	0.527 (5)
C18A	1.1603 (4)	1.2562 (4)	0.9587 (2)	0.0324 (6)	0.473 (5)
H18C	1.1639	1.2048	1.0158	0.039*	0.473 (5)
H18D	1.0751	1.3519	0.9553	0.039*	0.473 (5)
C19A	1.3507 (5)	1.3318 (4)	0.9602 (2)	0.0419 (8)	0.473 (5)
H19D	1.3942	1.4226	1.0131	0.063*	0.473 (5)
H19E	1.3469	1.3849	0.9043	0.063*	0.473 (5)
H19F	1.4356	1.2375	0.9644	0.063*	0.473 (5)
C21	0.53091 (13)	0.08983 (13)	0.34332 (7)	0.02406 (19)	
C22	0.50526 (15)	-0.08026 (14)	0.36571 (8)	0.0294 (2)	
H22	0.5425	-0.1013	0.4275	0.035*	
C23	0.42550 (17)	-0.21929 (16)	0.29831 (9)	0.0373 (3)	
H23	0.4064	-0.3349	0.3136	0.045*	
C24	0.37444 (18)	-0.18683 (18)	0.20872 (9)	0.0419 (3)	
H24	0.3212	-0.2812	0.1621	0.050*	
C25	0.40002 (17)	-0.01808 (17)	0.18616 (8)	0.0369 (3)	
H25	0.3647	0.0011	0.1241	0.044*	
C26	0.47678 (14)	0.12474 (15)	0.25293 (7)	0.0277 (2)	
C27	0.49622 (18)	0.30658 (17)	0.22573 (8)	0.0363 (2)	
H27A	0.3852	0.3678	0.2291	0.054*	
H27B	0.5109	0.2969	0.1632	0.054*	
H27C	0.6053	0.3737	0.2671	0.054*	

Atomic displacement parameters (\AA^2)

	U^{11}	U^{22}	U^{33}	U^{12}	U^{13}	U^{23}
S1	0.02783 (13)	0.03461 (14)	0.02109 (12)	0.01339 (10)	-0.00154 (9)	-0.00282 (9)
O1	0.0292 (4)	0.0317 (4)	0.0296 (4)	0.0119 (3)	-0.0037 (3)	-0.0098 (3)
O2	0.0369 (11)	0.0226 (7)	0.0255 (8)	-0.0031 (6)	0.0009 (7)	-0.0036 (6)
O2A	0.0409 (14)	0.0227 (8)	0.0229 (8)	0.0024 (7)	-0.0005 (8)	-0.0048 (6)
O3	0.0761 (7)	0.0317 (4)	0.0293 (4)	-0.0019 (4)	-0.0204 (4)	-0.0010 (3)
N1	0.0232 (4)	0.0242 (4)	0.0187 (3)	0.0062 (3)	-0.0018 (3)	-0.0032 (3)
N2	0.0249 (4)	0.0245 (4)	0.0195 (4)	0.0070 (3)	0.0007 (3)	-0.0045 (3)
C1	0.0242 (4)	0.0250 (4)	0.0161 (4)	0.0044 (3)	0.0029 (3)	-0.0004 (3)
C2	0.0224 (4)	0.0264 (4)	0.0199 (4)	0.0046 (3)	0.0031 (3)	-0.0032 (3)
C11	0.0198 (4)	0.0235 (4)	0.0180 (4)	0.0019 (3)	0.0021 (3)	-0.0020 (3)
C12	0.0290 (4)	0.0217 (4)	0.0205 (4)	0.0023 (3)	0.0022 (3)	0.0007 (3)
C13	0.0305 (5)	0.0248 (4)	0.0188 (4)	0.0020 (4)	-0.0003 (3)	0.0017 (3)
C14	0.0304 (5)	0.0234 (4)	0.0219 (4)	-0.0016 (4)	-0.0035 (4)	-0.0004 (3)
C15	0.0337 (5)	0.0215 (4)	0.0252 (5)	-0.0009 (4)	-0.0020 (4)	0.0024 (3)
C16	0.0262 (4)	0.0253 (4)	0.0193 (4)	0.0023 (3)	0.0002 (3)	0.0025 (3)
C17	0.0634 (8)	0.0239 (5)	0.0322 (6)	-0.0038 (5)	-0.0185 (6)	-0.0001 (4)
C18	0.0361 (12)	0.0261 (11)	0.0317 (11)	-0.0045 (10)	0.0014 (9)	-0.0073 (9)
C19	0.0509 (14)	0.0331 (13)	0.0341 (14)	0.0081 (11)	0.0081 (11)	-0.0053 (11)

C18A	0.0436 (14)	0.0228 (12)	0.0262 (13)	0.0009 (10)	0.0060 (10)	-0.0101 (10)
C19A	0.0430 (15)	0.0352 (15)	0.0415 (15)	-0.0034 (13)	0.0032 (12)	-0.0028 (12)
C21	0.0200 (4)	0.0269 (4)	0.0222 (4)	0.0034 (3)	0.0035 (3)	-0.0065 (3)
C22	0.0272 (5)	0.0292 (5)	0.0310 (5)	0.0026 (4)	0.0085 (4)	-0.0031 (4)
C23	0.0334 (5)	0.0283 (5)	0.0479 (7)	-0.0032 (4)	0.0129 (5)	-0.0090 (5)
C24	0.0357 (6)	0.0405 (6)	0.0406 (6)	-0.0035 (5)	0.0054 (5)	-0.0208 (5)
C25	0.0342 (5)	0.0455 (6)	0.0244 (5)	0.0044 (5)	0.0018 (4)	-0.0121 (4)
C26	0.0243 (4)	0.0336 (5)	0.0222 (4)	0.0049 (4)	0.0036 (3)	-0.0053 (4)
C27	0.0398 (6)	0.0395 (6)	0.0283 (5)	0.0073 (5)	0.0056 (4)	0.0034 (4)

Geometric parameters (Å, °)

S1—C1	1.6709 (10)	C18—H18A	0.9900
O1—C2	1.2247 (12)	C18—H18B	0.9900
O2—C17	1.370 (2)	C19—H19A	0.9800
O2—C18	1.451 (3)	C19—H19B	0.9800
O2A—C17	1.395 (2)	C19—H19C	0.9800
O2A—C18A	1.449 (4)	C18A—C19A	1.499 (4)
O3—C17	1.2038 (15)	C18A—H18C	0.9900
N1—C1	1.3388 (12)	C18A—H18D	0.9900
N1—C11	1.4220 (11)	C19A—H19D	0.9800
N1—H1	0.7998	C19A—H19E	0.9800
N2—C2	1.3824 (12)	C19A—H19F	0.9800
N2—C1	1.3936 (12)	C21—C22	1.3941 (15)
N2—H2	0.8911	C21—C26	1.4001 (14)
C2—C21	1.4937 (13)	C22—C23	1.3904 (15)
C11—C16	1.3895 (13)	C22—H22	0.9500
C11—C12	1.3941 (13)	C23—C24	1.382 (2)
C12—C13	1.3881 (13)	C23—H23	0.9500
C12—H12	0.9500	C24—C25	1.385 (2)
C13—C14	1.3923 (14)	C24—H24	0.9500
C13—H13	0.9500	C25—C26	1.4000 (14)
C14—C15	1.3900 (14)	C25—H25	0.9500
C14—C17	1.4858 (14)	C26—C27	1.5021 (17)
C15—C16	1.3904 (13)	C27—H27A	0.9800
C15—H15	0.9500	C27—H27B	0.9800
C16—H16	0.9500	C27—H27C	0.9800
C18—C19	1.493 (4)		
C17—O2—C18	115.67 (16)	C18—C19—H19A	109.5
C17—O2A—C18A	118.32 (17)	C18—C19—H19B	109.5
C1—N1—C11	126.33 (8)	H19A—C19—H19B	109.5
C1—N1—H1	116.3	C18—C19—H19C	109.5
C11—N1—H1	117.2	H19A—C19—H19C	109.5
C2—N2—C1	128.53 (8)	H19B—C19—H19C	109.5
C2—N2—H2	115.7	O2A—C18A—C19A	111.2 (4)
C1—N2—H2	115.5	O2A—C18A—H18C	109.4
N1—C1—N2	116.03 (8)	C19A—C18A—H18C	109.4
N1—C1—S1	125.74 (7)	O2A—C18A—H18D	109.4
N2—C1—S1	118.21 (7)	C19A—C18A—H18D	109.4

supplementary materials

O1—C2—N2	122.79 (9)	H18C—C18A—H18D	108.0
O1—C2—C21	123.60 (9)	C18A—C19A—H19D	109.5
N2—C2—C21	113.61 (8)	C18A—C19A—H19E	109.5
C16—C11—C12	120.35 (8)	H19D—C19A—H19E	109.5
C16—C11—N1	117.45 (8)	C18A—C19A—H19F	109.5
C12—C11—N1	122.11 (9)	H19D—C19A—H19F	109.5
C13—C12—C11	119.08 (9)	H19E—C19A—H19F	109.5
C13—C12—H12	120.5	C22—C21—C26	120.98 (9)
C11—C12—H12	120.5	C22—C21—C2	118.37 (9)
C12—C13—C14	120.85 (9)	C26—C21—C2	120.65 (9)
C12—C13—H13	119.6	C23—C22—C21	120.47 (11)
C14—C13—H13	119.6	C23—C22—H22	119.8
C15—C14—C13	119.70 (9)	C21—C22—H22	119.8
C15—C14—C17	122.23 (10)	C24—C23—C22	119.01 (12)
C13—C14—C17	118.05 (9)	C24—C23—H23	120.5
C14—C15—C16	119.80 (9)	C22—C23—H23	120.5
C14—C15—H15	120.1	C23—C24—C25	120.70 (10)
C16—C15—H15	120.1	C23—C24—H24	119.7
C11—C16—C15	120.18 (9)	C25—C24—H24	119.7
C11—C16—H16	119.9	C24—C25—C26	121.39 (11)
C15—C16—H16	119.9	C24—C25—H25	119.3
O3—C17—O2	124.17 (12)	C26—C25—H25	119.3
O3—C17—O2A	120.61 (13)	C25—C26—C21	117.44 (11)
O3—C17—C14	124.28 (11)	C25—C26—C27	119.59 (10)
O2—C17—C14	109.38 (11)	C21—C26—C27	122.97 (9)
O2A—C17—C14	112.60 (11)	C26—C27—H27A	109.5
O2—C18—C19	110.5 (3)	C26—C27—H27B	109.5
O2—C18—H18A	109.6	H27A—C27—H27B	109.5
C19—C18—H18A	109.6	C26—C27—H27C	109.5
O2—C18—H18B	109.6	H27A—C27—H27C	109.5
C19—C18—H18B	109.6	H27B—C27—H27C	109.5
H18A—C18—H18B	108.1		
C11—N1—C1—N2	177.55 (9)	C15—C14—C17—O3	-175.89 (16)
C11—N1—C1—S1	-3.80 (15)	C13—C14—C17—O3	2.8 (2)
C2—N2—C1—N1	6.94 (15)	C15—C14—C17—O2	-12.1 (2)
C2—N2—C1—S1	-171.82 (9)	C13—C14—C17—O2	166.59 (17)
C1—N2—C2—O1	0.83 (17)	C15—C14—C17—O2A	22.0 (2)
C1—N2—C2—C21	-178.64 (9)	C13—C14—C17—O2A	-159.3 (2)
C1—N1—C11—C16	137.72 (11)	C17—O2—C18—C19	-77.4 (3)
C1—N1—C11—C12	-45.70 (15)	C17—O2A—C18A—C19A	79.8 (3)
C16—C11—C12—C13	-1.85 (15)	O1—C2—C21—C22	-128.25 (12)
N1—C11—C12—C13	-178.34 (9)	N2—C2—C21—C22	51.21 (12)
C11—C12—C13—C14	0.19 (16)	O1—C2—C21—C26	50.88 (15)
C12—C13—C14—C15	1.59 (17)	N2—C2—C21—C26	-129.66 (10)
C12—C13—C14—C17	-177.10 (12)	C26—C21—C22—C23	0.19 (16)
C13—C14—C15—C16	-1.71 (17)	C2—C21—C22—C23	179.32 (9)
C17—C14—C15—C16	176.93 (12)	C21—C22—C23—C24	0.89 (17)
C12—C11—C16—C15	1.74 (15)	C22—C23—C24—C25	-0.76 (19)
N1—C11—C16—C15	178.39 (9)	C23—C24—C25—C26	-0.44 (19)

C14—C15—C16—C11	0.06 (17)	C24—C25—C26—C21	1.48 (17)
C18—O2—C17—O3	-10.6 (4)	C24—C25—C26—C27	-177.99 (11)
C18—O2—C17—O2A	83.6 (3)	C22—C21—C26—C25	-1.35 (15)
C18—O2—C17—C14	-174.5 (2)	C2—C21—C26—C25	179.54 (9)
C18A—O2A—C17—O3	13.0 (4)	C22—C21—C26—C27	178.11 (10)
C18A—O2A—C17—O2	-93.5 (4)	C2—C21—C26—C27	-1.00 (15)
C18A—O2A—C17—C14	175.8 (3)		

Hydrogen-bond geometry (Å, °)

<i>D</i> —H \cdots <i>A</i>	<i>D</i> —H	H \cdots <i>A</i>	<i>D</i> \cdots <i>A</i>	<i>D</i> —H \cdots <i>A</i>
N1—H1 \cdots O1	0.80	2.01	2.669 (1)	139
N2—H2 \cdots S1 ⁱ	0.89	2.67	3.5551 (9)	170

Symmetry codes: (i) $-x+2, -y, -z+1$.

Fig. 1




Fig. 2

