

Scale Insects (Hemiptera: Coccoidea) of ornamental plants from São Carlos, São Paulo, Brazil

A. L. B. G. Peronti

Department of Ecology and Evolutionary Biology
Federal University of São Carlos
C. P. 676, 13565-905 São Carlos
São Paulo, Brazil

D. R. Miller

Systematic Entomology Laboratory
Plant Sciences Institute
Agricultural Research Service, USDA
Beltsville, Maryland 20705

C. R. Sousa-Silva

Department of Ecology and Evolutionary Biology
Federal University of São Carlos
C. P. 676, 13565-905 São Carlos
São Paulo, Brazil

Abstract. A list of 35 scale insects collected from 72 ornamental plant species in São Carlos, São Paulo, Brazil is provided. Regarding host specificity, 30 scale insects were polyphagous, 4 oligophagous, and 1 monophagous. A total of 102 coccoid/plant associations are recorded, 29 of which are new host records for the species; 60 are new host records for the species in Brazil. *Pulvinaria urbicola* Cockerell, 1893 (Coccidae), *Phenacoccus similis* Granara de Willink, 1983 (Pseudococcidae), and *Orthezia molinarii* (Morrison, 1952) (Ortheziidae) are recorded for the first time in Brazil. In addition, we describe the injury caused by scale insects on ornamental plants.

Key words. Scale insect diversity, Scale/plant associations, Floricultural pests.

Introduction

Floriculture, which consists of the cultivation of ornamental plants used for cut flowers, flowering and non-flowering potted plants, and the production of seeds, bulbs, and large trees, is an important part of Brazilian agriculture, with annual gross sales of approximately one billion Brazilian reals (currently about half a billion U. S. dollars). The state of São Paulo holds a 70% share of this market, and specialists in floriculture estimate an annual growth of 20% in the next few years (Arruda et al., 1996).

Scale insects (Superfamily Coccoidea) stand out as important pests among insects that feed on ornamental plants (Arruda et al., 1996). They cause damage to plants directly by sucking their sap, and indirectly by injecting toxic salivary secretions, transmitting pathogens, attracting ants, and encouraging the development of sooty-mold (Lara, 1992; Zucchi et al., 1993). Besides their impact on

the commercial value of ornamental plants, these insects also affect urban landscapes. Infested containerized plants become disseminators of harmful insects when they are transported to regions or countries far from their place of origin or production.

Despite the increase of ornamental plant production in Brazil, very few studies have been undertaken on the insects associated with plants in gardens. Recent publications include Bergmann & Alexandre (1995) and Imenes and Alexandre (1996); they discuss the occurrence of pests, diseases, and forms of control in roses (*Rosa* sp.) and chrysanthemums (*Chrysanthemum* sp.), respectively. Favero (1996) describes some pests of flowers, ornamental foliage, and orchids and gives methods of control, and Imenes and Bergmann (2000) characterize the principal pests of ornamental plants including some scale insect-plant interactions and provide information on control strategies.

TABLE 1. Scale insects and their ornamental host plants collected in São Carlos – SP from August 1997 to April 1999.

Scale insects collected	Host plant species/Common names in Portuguese shown in quotes	Plant family	Plant Origin	Plant Form
MONOPHAGOUS				
Diaspididae				
<i>Kuwaniaspis bambusicola</i> (Cockerell, 1899)(vrw)	<i>Bambusa gracilis</i> Hort. “bambu-de-jardim”	Poaceae	exotic	shrub
OLIGOPHAGOUS				
Diaspididae				
<i>Acutaspis oliverai</i> (Lepage & Gianotti, 1942)(vrw)	** <i>Hedera canariensis</i> Willd. “hera-da-algéria”	Araliaceae	exotic	vine
Eriococcidae				
<i>Eriococcus campinensis</i> Hempel, 1937 (drm)	** <i>Mimosa caesalpiniæfolia</i> Benth. “sansão-do-campo”	Fabaceae	native	tree
Ortheziidae				
<i>Orthezia molinarii</i> Morrison, 1952 (drm)	** <i>Bougainvillea spectabilis</i> Willd. “primavera” ** <i>Cordilyne terminalis</i> Kunth. “dracena-vermelha”	Nyctaginaceae	native	shrub
Pseudococcidae				
<i>Phenacoccus similis</i> Granara de Willink, 1983 (drm)	** <i>Saintpaulia ionantha</i> Wendl. “violeta-africana”	Gesneriaceae	exotic	herb
POLYPHAGOUS				
Coccidae				
<i>Ceroplastes cirripediformis</i> Comstock, 1881 (mlw)	* <i>Duranta repens</i> var. <i>aurea</i> L. Hort. “pingo-de-ouro”	Verbenaceae	native	shrub
<i>Ceroplastes floridensis</i> Comstock, 1881 (mlw)	<i>Hedera canariensis</i> “hera-da-algéria” * <i>Schefflera actinophylla</i> Harms “cheflera” * <i>Schefflera arboricola</i> (Hay.) Merr. “cheflera-pequena”	Araliaceae	exotic	vine
		Araliaceae	exotic	shrub
		Araliaceae	exotic	shrub

The objective of this study was to survey scale insect-plant interactions in ornamental plants in the city of São Carlos, state of São Paulo (SP), Brazil. In addition, we classified the species of scale insects according to their host specificity and characterized host-plant injury caused by scale insects.

Materials and Methods

The buds, branches, and leaves of ornamental plants infested by scale insects were collected in public and private gardens in the city of São Carlos, SP, Brazil and taken to the laboratory of the Department of Ecology and Evolutionary Biology of the Federal University of São Carlos (UFSCar).

Collections were made from August 1997 to April 1999. In the laboratory, scale insects were collected from host plants with brushes and were stored in vials filled with 70% alcohol. They were mounted on microscope slides according to the technique described by Granara de Willink (1990). Scale insects were identified with a compound light microscope using keys Granara de Willink (1999) and Hodgson (1994) for Coccidae; Ferris (1937, 1938, 1941, 1942) and Lepage and Gianotti (1942) for Diaspididae; Morrison (1925, 1952) for Ortheziidae; and Williams and Granara de Willink (1992) for Pseudococcidae. Most of the scale insects were sent to specialists for confirmation of final identifications. Categorization of scale insect host-plant specificity was based in large part on classic defini-

TABLE 1. Continued.

Scale insects collected	Host plant species/Common names In Portuguese shown in quotes	Plant family	Plant origin	Plant form
<i>Coccus hesperidum</i> Linnaeus, 1758 (drm, mlw)	** <i>Davallia fejeensis</i> Hook. “renda-portuguesa” * <i>Platycerium bifurcatum</i> (Cav.) Chr. “chifre-de-veado” * <i>Schefflera actinophylla</i> “cheflera”	Davalliaceae	exotic	herb
		Polypodiaceae	exotic	herb
		Araliaceae	exotic	shrub
<i>Coccus viridis</i> (Green, 1889) (drm, mlw)	* <i>Murraya exotica</i> Jack “falsa-murta” <i>Gardenia jasminoides</i> Ellis “jasmin-do-cabo” * <i>Polyscias fruticosa</i> N.E.Br. “árvore-da-felicidade-fêmea” * <i>Ixora coccinea</i> L. var. <i>compacta</i> Hort. “ixora-compacta”	Rutaceae	exotic	tree
		Rubiaceae	exotic	shrub
		Araliaceae	exotic	shrub
		Rubiaceae	exotic	shrub
<i>Coccus</i> sp. (alp)	<i>Ixora chinensis</i> Lam. “ixora-chinesa”	Rubiaceae	exotic	shrub
<i>Parasaissetia nigra</i> (Nietner, 1861) (drm, mlw)	* <i>Schinus molle</i> L. “aoeira-salsa” * <i>Euphorbia fulgens</i> Karw ex Kl. “chiquita-bacana”	Anacardiaceae	native	tree
		Euphorbiaceae	exotic	shrub
<i>Parasaissetia</i> sp. (drm)	<i>Camellia japonica</i> L. “camélia”	Theaceae	exotic	shrub
<i>Protopulvinaria pyriformis</i> (Cockerell, 1894) (drm)	<i>Schefflera arboricola</i> “cheflera-pequena”	Araliaceae	exotic	shrub
	<i>Bauhinia variegata</i> Linn. “pata-de-vaca”	Fabaceae	native	tree
<i>Pulvinaria urbicola</i> Cockerell, 1893 (drm)	** <i>Schinus molle</i> “aoeira-salsa”	Anacardiaceae	native	tree

tions, in which “polyphagous” is used for species that live on a variety of non-related plants, i.e., plants from different families and, “oligophagous” is defined as species that live on a reduced number of non-related host plants. We defined “monophagous” as species that feed on one or more plants of the same family. Scale insect specimens are deposited in the Insect Collection of the Department of Ecology and Evolutionary Biology (Colecoc/DEBE) at UFSCar and in the Smithsonian’s National Coccoidea Collection, Beltsville, Maryland, USA. Ornamental plants were identified according to Lorenzi (1992) and Lorenzi and Souza (1995) or sent to specialists. Determination of new host records are based on a search of the literature or ScaleNet (Ben-

Dov et al. 2001). Determinations of scale insects were made by Douglass R. Miller (drm), Michael L. Williams (mlw), Vera R. dos Santos Wolff (vrw), and Ana Lúcia B. G. Peronti (alp) and are so indicated in the accompanying table. New records for Brazil are indicated with an asterisk (*); new host records for a species of scale are indicated with a double asterisk (**).

Results and Discussion

A total of 184 samples of scale insects were collected from 72 ornamental species (8 trees, 39 shrubs, 23 herbaceous plants, and 2 vines) distributed among 37 botanical families. Approximately

TABLE 1. Continued.

Scale insects collected	Host plant species/Common names In Portuguese shown in quotes	Plant family	Plant origin	Plant Form
<i>Saissetia coffeae</i> (Walker, 1852) (drm, mlw)	* <i>Ervatamia coronaria</i> Stapf. "jasmim-café" * <i>Pseuderanthemum atropurpureum</i> Radlk. "pseudoerântemo" ** <i>Nephrolepis</i> sp. "samambaia" * <i>Asparagus africanus</i> "aspargo"	Apocynaceae	exotic	shrub
Diaspididae				
<i>Abgrallaspis cyanophylli</i> (Signoret, 1869) (vrw)	* <i>Philodendron</i> sp. "filodendro"	Araceae	native	shrub
<i>Acutaspis perseae</i> (Comstock, 1881) (vrw)	** <i>Dieteris iridioides</i> Sweet "moréia"	Iridaceae	exotic	herb
<i>Chrysomphalus aonidum</i> (Linnaeus, 1758) (vrw)	<i>Chrysalidocarpus luteus</i> Wendl. "areca-bambu" * <i>Dracaena marginata</i> Lam. "dracena-de-madagascar" "orquídea"	Palmae	exotic	palm tree shrub
<i>Chrysomphalus dictyospermi</i> (Morgan, 1889) (vrw)	<i>Rosa</i> sp. "roseira"	Rosaceae	exotic	shrub
<i>Diaspis boisduvalii</i> Signoret, 1869 (drm, vrw)	* <i>Oncidium varicosum</i> Lindl. "chuva-de-ouro" <i>Cattleya walkueriana</i> var. <i>princeps</i> Barb. Rodr. "orquídea"	Orchidaceae	native	herb
<i>Fiorinia fioriniae</i> (Targioni-Tozzetti, 1867) (drm, vrw)	* <i>Camellia japonica</i> "camélia" <i>Podocarpus lambertii</i> Klotz. "podocarpo"	Theaceae	exotic	shrub
<i>Hemiberlesia rapax</i> (Comstock, 1881) (vrw)	* <i>Chrysalidocarpus luteus</i> "areca-bambu"	Palmae	exotic	palm tree

74% of host plants were exotics, and all were perennials. One hundred and two coccoid-plant associations were recorded; 26 of them were new for the scale species and 60 were new for Brazil. Thirty-five species of scale insects were identified, including 18 diaspidids (armored scales), 8 coccids (soft scales), 6 pseudococcids (mealybugs), 2 ortheziids (ensign scales), and 1 eriococcid (felt scale). The *Icerya* sp. (Margarodidae) found on *Pittosporum tobira* (Thunb.) Ait. (Pittosporaceae) is probably *Icerya purchasi* Maskell (Table 1).

Pulvinaria urbicola Cockerell, 1893 (Coccidae), *Phenacoccus similis* Granara de Willink, 1983

(Pseudococcidae), and *Orthezia molinarii* (Morrison, 1952) (Ortheziidae) were recorded for the first time in Brazil.

Based on host-plant information provided in Morrison (1952), Vernalha (1953), Silva et al. (1968), Ben-Dov (1993, 1994), Ben-Dov et al. 2001, and Williams and Granara de Willink (1992), 30 of the scale-insect species are polyphagous, 4 are oligophagous, and 1 is monophagous (Table 1).

Among the polyphagous species, the most commonly encountered were *Coccus viridis* and *Saissetia coffeae* (Coccidae); *Ischnaspis longirostris*, *Pinnaspis strachani* and *Parlatoria proteus* (Diaspid-

TABLE 1. Continued.

Scale insects collected	Host plant species/Common names In Portuguese shown in quotes	Plant family	Plant origin	Plant Form
<i>Howardia biclavis</i> (Comstock, 1833) (vrw)	** <i>Ervatamia coronaria</i> “jasmim-café”	Apocynaceae	exotic	shrub
<i>Ischnaspis longirostris</i> (Signoret, 1882) (drm, vrw)	<i>Chrysalidocarpus lutezens</i> “areca-bambu” ** <i>Ctenanthe oppenheimiana</i> Schum. “maranta-variegata” ** <i>Dietetes iridioides</i> “moréia” <i>Ficus benjamina</i> L. “ficus” * <i>Heliconia rostrata</i> Ruiz et Pav. “helicônia” <i>Ixora coccinea</i> L. “ixora-coral” <i>Phoenix roebelinii</i> O’ Brien. “tamareira-de-jardim”	Palmae Marantaceae Iridaceae Moraceae Musaceae Rubiaceae Palmae	exotic native exotic	palm tree herb tree shrub shrub shrub palm tree
<i>Parlatoria proteus</i> (Curtis, 1843) (vrw)	* <i>Monstera</i> sp. “monstera” * <i>Schefflera arboricola</i> “cheflera-pequena” ** <i>Spathiphyllum wallissii</i> Regel “lírio-da-paz” * <i>Yucca elephantipes</i> Hort. ex Regel “yuca-elefante” “orquídea”	Araceae Araliaceae Araceae Liliaceae Orchidaceae	_____	herb shrub herb shrub herb
<i>Pinnaspis aspidistrae</i> (Signoret, 1869) (drm, vrw)	* <i>Platycerium bifurcatum</i> (Cav.) Chr. “chifre-de-veado”	Polypodiaceae	exotic	herb

idae); *Phenacoccus madeirensis* and *Pseudococcus longispinus* (Pseudococcidae); and *Orthezia insignis* (Ortheziidae). These eight species occurred on 42% of the scale-infested hosts.

Eriococcus campinensis (Eriococcidae), described by Hempel (1937), and *Acutaspis oliverai* (Diaspididae), described by Lepage and Giannotti (1942), are recorded from Brazil only, and *Orthezia molinarii* (Ortheziidae) and *Phenacoccus similis* (Pseudococcidae), previously were recorded by Morrison (1952) and Williams and Granara de Willink (1992), respectively, only in Argentina. It is possible that additional records will place these species in the polyphagous category.

Kuwanaspis bambusicola (Diaspididae), collected from *Bambusa gracilis*, is a monophagous species. According to Ben-Dov (1990), this species has been reported from species in only two genera of bamboo, *Bambusa* and *Dendrocalamus*.

Most samples of scale insects were collected from plants growing in pots or narrow flower beds in the shade (indoors) or semi-shade (under nursery netting, trees, or open porches). Infestations of the armored scale *Ischnaspis longirostris* on *Ficus benjamina* and of the mealybug *Pseudococcus longispinus* on *Cordyline terminalis* were found only on potted plants, with the former interaction occurring both in shade and in bright sunlight and the latter only indoors. In larger flowerbeds where these pests were found, they appeared in small numbers.

Dense scale-insect populations were observed on the majority of host plants. Most infested plants showed symptoms such as dried or shriveled leaves; in a small number of cases flowers or fruit were aborted and leaves were deformed. Chlorosis of leaf tissue was detected mainly on hosts infested by species of Diaspididae.

TABLE 1. Continued.

Scale insects collected	Host plant species/Common names In Portuguese shown in quotes	Plant family	Plant Origin	Plant form
<i>Pinnaspis strachani</i> (Cooley, 1899) (vrw)	* <i>Asparagus densiflorus</i> (Kunth.) Jess. var. <i>sprengeri</i> Hort. “aspargo-pendente” * <i>Bauhinia variegata</i> Linn. “pata-de-vaca” <i>Hibiscus rosa-sinensis</i> L. “hibisco” * <i>Murraya exotica</i> “falsa-murta” ** <i>Nephrolepis exalta</i> Schot var. <i>forida-ruffle</i> “samambaia-crespa”	Liliaceae	exotic	herb
<i>Pinnaspis</i> sp. (vrw)	<i>Asplenium nidus</i> L. “asplênio”	Aspleniaceae	exotic	herb
<i>Pseudaonidia trilobitiformis</i> (Green, 1896) (drm, vrw)	* <i>Nerium oleander</i> L. “espirradeira” * <i>Ficus pumila</i> L. “unha-de-gato”	Apocynaceae	exotic	shrub
<i>Pseudaulacaspis pentagona</i> (Targioni -Tozzetti, 1886) (vrw)	sp. 2 “orquídea” * <i>Ligustrum sinense</i> Lour. “ligusto-chinês”	Oleaceae	exotic	shrub
<i>Pseudischnaspis bowreyi</i> (Cockerell, 1893) (vrw)	** <i>Chrysalidocarpus luteus</i> “areca-bambu”	Palmae	exotic	palm tree
<i>Selenaspis articulatus</i> (Morgan, 1889) (vrw)	** <i>Ervatamia coronaria</i> “jasmim-café” * <i>Murraya exotica</i> “falsa-murta”	Apocynaceae	exotic	shrub
Margarodidae: <i>Icerya</i> sp. (drm)	<i>Pittosporum tobira</i> (Thunb.) Ait. “pitosporo-japonês”	Pittosporaceae	exotic	shrub

Sooty-mold frequently was found on plants infested by species of Ortheziidae, Pseudococcidae, and Coccidae. A large amount of sooty mold was found on *Nectandra megapotamica* infested by *Nipaecoccus nipae* (Pseudococcidae) and on *Schefflera arboricola* and *S. actinophylla* infested by *Orthezia* sp. (Ortheziidae).

Withered and dead plants of *Cordyline terminalis* were observed infested by *P. longispinus* (Pseudococcidae) as were plants of *Asparagus densiflorus* and *Murraya exotica* infested by *Pinnaspis strachani* (Diaspididae), and *Saintpaulia ionantha* infested by *Phenacoccus similis* (Pseudococcidae). These plants were apparently negatively impacted

because of direct and indirect damage caused by these insects.

In most cases, it was not possible to determine the extent of the ultimate destruction caused by these insects because most infested hosts were treated in some way to protect them from extensive damage.

Acknowledgments

The authors gratefully acknowledge the help of Dra. Vera R. dos Santos Wolff (Fundação Estadual de Pesquisa Agropecuária, FEPAGRO, Porto Alegre,

TABLE 1. Continued.

Scale insects collected	Host plant species/Common names in Portuguese shown in quotes	Plant family	Plant Origin	Plant form
Ortheziidae				
<i>Orthezia insignis</i> Browne, 1887 (drm, alp)	<i>Ardisia crenata</i> Sims. “ardísia” * <i>Iresine herbstii</i> Hook. “iresínea” * <i>Duranta repens</i> L. var. <i>aurea</i> Hort. “pingo-de-ouro” ** <i>Pseuderanthemum atropurpureum</i> “pseudoerântemo” * <i>Pachystachys lutea</i> Nees. “camarão-amarelo” <i>Solenostemon scutellarioides</i> (L.) Lodd. “cóleus”	Myrsinaceae	exotic	shrub
<i>Orthezia sp</i> (drm, alp).	<i>Brunfelsia uniflora</i> D. Don. “manacá-de-cheiro” <i>Codiaeum variegatum</i> Blume. “crótão” <i>Dichorisandra thyrsiflora</i> Mik. “mariarinha” <i>Duranta repens</i> L. “violeta-ira” <i>Euphorbia milii</i> des Moulins var. <i>mili</i> “coroa-de-cristo” <i>Euphorbia milii</i> des Moulins var. <i>breonii</i> Hort. “coroa-de-cristo” <i>Hibiscus syriacus</i> L. “hibisco-colunar” <i>Lantana camara</i> L. “camarazinho” <i>Schefflera actinophylla</i> “cheflera” <i>Schefflera arboricola</i> “cheflera-pequena” <i>Syngonium podopylum</i> Schott. “singônio”	Solanaceae	native	shrub
		Euphorbiaceae	exotic	shrub
		Comeliaceae	native	shrub
		Verbenaceae	native	shrub
		Euphorbiaceae	exotic	shrub
		Euphorbiaceae	exotic	shrub
		Malvaceae	exotic	shrub
		Verbenaceae	native	shrub
		Araliaceae	exotic	shrub
		Araliaceae	exotic	shrub
		Araceae	exotic	herb

Brazil) and Dr. Michael L. Williams (Entomology Department, Auburn University, Auburn, Alabama, USA) for the identification of a large part of the scale insects; Dra. Maria Inês Salgueiro Lima (Departamento de Botânica, Universidade Federal de São Carlos, UFSCar, São Carlos, Brazil) for the identification of some plant species; and CNPq (Conselho Nacional de Desenvolvimento Científico e Tecnológico, Brazil) for their financial support. We are grateful to Drs. Manya B. Stoetzel and John W. Brown, Systematic Entomology Laboratory, US Department of Agriculture and Avas B. Hamon, Division of Plant Industry, Florida State Collection of Arthropods, Gainesville, Florida, for their comments and criticisms of this manuscript.

References

- Arruda, S. T., M. P. A. Olivette, and C. E. F. Castro. 1996. Diagnóstico da Floricultura do Estado de São Paulo. Revista Brasileira Horticultura Ornamental, Campinas, 2:1-18.
- Ben-Dov, Y. 1990. Bamboo. In: David Rosen (ed.), Armored scale insects. Their biology natural enemies and control. World Crop Pests 4B:655-660. Elsevier Science Publisher B.V., Amsterdam. 687 p.
- Ben-Dov, Y. 1993. A Systematic Catalogue of the Soft Scale Insects of the World. Flora & Fauna Handbook Number 9, Sandhill Crane Press, Incorporated, Gainesville, Florida. 536 p.

TABLE 1. Continued.

Scale insects collected	Host plant species/Common names in Portuguese shown in quotes	Plant family	Plant Origin	Plant form
Pseudococcidae				
<i>Ferrisia virgata</i> (Cockerell, 1893) (drm)	<i>Codiaeum variegatum</i> “crótão” ** <i>Rhipsalis baccifera</i> (Mill.)Stearn “cacto-macarrão”	Euphorbiaceae	exotic	shrub
<i>Nipaecoccus nipae</i> (Maskell, 1893) (drm)	** <i>Nectandra megapotamica</i> Spreng. “canelinha”	Lauraceae	native	tree
<i>Phenacoccus madeiriensis</i> Green, 1923 (drm)	* <i>Pachystachys lutea</i> “camarão-amarelo” * <i>Salvia splendens</i> Ker-Gawl. “sangue-de-adão” <i>Hibiscus rosa-sinensis</i> “hibisco” <i>Acalypha wilkesiana</i> M. Arg. “acalifa” <i>Lantana camara</i> “camarazinho”	Acanthaceae	exotic	shrub
<i>Planococcus</i> sp. (drm)	<i>Oncidium varicosum</i> Lindl. “chuva-de-ouro”	Orchidaceae	native	herb
<i>Planococcus citri</i> (Risso, 1813) (drm)	** <i>Kalanchoe blossfeldiana</i> v. Poelln. “calançoê”	Crassulaceae	exotic	shrub
<i>Pseudococcus longispinus</i> (Targioni-Tozzetti, 1867) (drm)	* <i>Cycas circinalis</i> Roxb. “cicas” ** <i>Cordyline terminalis</i> “dracena-vermelha” ** <i>Dracaena surculosa</i> Lindl. “dracena-bambu” ** <i>Syngonium podophyllum</i> “singônio” ** <i>Rosa</i> sp. “roseira”	Cycadaceae	exotic	shrub
<i>Pseudococcus</i> sp. (drm, alp)	<i>Davallia fejeensis</i> “renda-portuguesa” <i>Dieffenbachia amoena</i> Hort. ex Gent. “comigo-niguém-pode”	Davalliaceae	exotic	herb
		Araceae	exotic	herb

Ben-Dov, Y. 1994. A Systematic Catalogue of the Mealybugs of the World (Insecta: Homoptera: Coccoidea: Pseudococcidae: Putoidae). Intercept Ltd., Andover, UK, 686 p.

Ben-Dov, Y., D. R. Miller and G. A. P. Gibson. 2001. ScaleNet, Systematic Entomology Laboratory, Maryland, USDA, <http://www.se.barc.usda.gov/scalenet>. Consulted February 22, 2001.

Bergmann, E. C. and M. A. V. Alexandre. 1995. Aspectos fitossanitários da roseira. Boletim Técnico do Instituto Biológico, São Paulo, 2:5-51.

Cockerell, T. D. A. 1893. Two new species of *Pulvinaria* from Jamaica. Transactions of the Entomological Society of London 1893:159-163.

Favero, S. 1996. Pragas de Plantas Ornamentais. Universidade Estadual do Norte Fluminense. Centro de Ciências e Tecnologias Agropecuárias.

- Campos dos Goyatacazes, Rio de Janeiro, Boletim Técnico. 3. 16p.
- Ferris, G. F.** 1937. Atlas of the scale insects of North America. Series I. Stanford University Press, Palo Alto, California, 275 p.
- Ferris, G. F.** 1938. Atlas of the scale insects of North America. Series II. Stanford University Press, Palo Alto, California, 264 p.
- Ferris, G. F.** 1941. Atlas of the scale insects of North America. Series III. Stanford University Press, Palo Alto, California, 230 p.
- Ferris, G. F.** 1942. Atlas of the scale insects of North America. Series IV. Stanford University Press, Palo Alto, California, 243 p.
- Granara de Willink, M. C.** 1983. Tres especies de *Phenacoccus* de la Provincia de Tucumán, Argentina (Hom. Pseudococcidae). (In Spanish; Summary In English). *Acta Zoologica Lilloana* 37: 59-64.
- Granara de Willink, M. C.** 1990. Conociendo nuestra fauna I. Superfamilia Coccoidea (Homoptera: Sternorrhyncha). Facultad de Ciencias Naturales e Instituto Miguel Lillo, Universidad Nacional de Tucumán, Argentina, Serie Monográfica y Didáctica vol 6., 43 p.
- Granara de Willink, M. C.** 1999. Las cochinillas blandas de la República Argentina (Homoptera: Coccoidea: Coccidae). Contributions on Entomology, International 3 (1):1-183.
- Hempel, A.** 1937. Novas espécies de coccídeos (Homoptera) do Brasil. Archivos do Instituto Biológico. São Paulo, 8: 5-36.
- Hodgson, C. J.** 1994. The scale insect family Coccidae: an identification manual to genera. CAB. International, Wallingford, Oxon, U. K. 638 p.
- Imenes, S. D. L. and M. A. V. Alexandre.** 1996. Aspectos fitossanitários do crisântemo. . Boletim Técnico do Instituto Biológico, São Paulo, 5:5-47.
- Imenes, S. D. L. and E. C. Bergmann.** 2000. Reconhecimento e controle de pragas de plantas ornamentais. Vetores & Pragas, 7:8-15.
- Lara, F. M.** 1992. Principios de Entomologia. Editora Ícone Ltda. 331 p.
- Lepage, H. S. and O. Giannotti.** 1942. Contribuição para o levantamento fitossanitário do nordeste Brasileiro. Coccoideos assinalados nos estados do Ceará, Paraíba do Norte e Rio Grande do Norte. Boletim da Sociedade Brasileira de Agronomia. 5:444-458.
- Lorenzi, H.** 1992. Árvores Brasileiras Manual de identificação e cultivo de plantas arbóreas nativas do Brasil. Editora Plantarum LTDA, Nova Odessa, São Paulo. 368 p.
- Lorenzi, H. and H. M. Souza.** 1995. Plantas ornamentais no Brasil: arbustivas, herbáceas e trepadeiras. Editora Plantarum LTDA, Nova Odessa, São Paulo. 736 p.
- Morrison, H.** 1925. Classification of Scale Insects of the Subfamily Ortheziinae. Journal of Agricultural Research., Washington, D. C., 30: 97-154.
- Morrison, H.** 1952. Classification of the Ortheziidae. Supplement to classification of scale insects of the subfamily Ortheziinae. United States Department of Agriculture, Technical Bulletin, 1050 Washington, D.C. 80 p.
- Silva, A. G. d'A. G. R. Gonçalves, D. M. Galvão, A. J. L. Gonçalves, J. Gomes, M. N. Silva and L. Simoni.** 1968. Quarto Catálogo dos Insetos que vivem nas plantas do Brasil, seus parasitos e predadores. Rio de Janeiro, Ministério da Agricultura. Parte II, 1º tomo. 622 p.
- Vernalha, M. M.** 1953. Coccídeos da coleção I. B. P. T. Arquivos de Biologia e Tecnologia, Curitiba, 8: 11-304.
- Williams, D. J. and M. C. Granara de Willink.** 1992. Mealybugs of Central and South America. CAB. International. Wallingford, Oxon, U. K. 634 p.
- Zucchi, R. A., S. S. Neto and O. Nakano.** 1993. Guia de Identificação de Pragas Agrícolas. Fundação de Estudos Agrários Luiz de Queiros (FEALQ), Piracicaba. 139 p.