

Title: **Single-molecule photobleaching reveals MET receptor dimerization after ligand binding on intact cells**

Running title: Single-molecule photobleaching of MET dimers

Authors: Marina S. Dietz¹, Daniel Haße², Davide M. Ferraris³, Antonia Göhler⁴, Hartmut H. Niemann², and Mike Heilemann¹

Addresses: ¹Institute of Physical and Theoretical Chemistry, Johann Wolfgang Goethe-University, Max-von-Laue-Str. 7, 60438 Frankfurt, Germany

²Structural Biochemistry, Department of Chemistry, Bielefeld University, Universitaetsstrasse 25, 33615 Bielefeld, Germany

³Department of Pharmaceutical Science, University of Eastern Piedmont "Amedeo Avogadro", Via Bovio 6, 28100 Novara, Italy

⁴Department of Biotechnology and Biophysics, Julius-Maximilians University, Am Hubland, Biozentrum, 97074 Wuerzburg, Germany

Correspondence: Mike Heilemann, Hartmut Niemann

Telephone: +49 69 798 29736 (MH); +49 521 106 6154 (HN)


Fax: +49 69 798 29560 (MH); +49 521 106 6146 (HN)

E Mail: heilemann@chemie.uni-frankfurt.de ; hartmut.niemann@uni-bielefeld.de

SUPPLEMENTARY INFORMATION

SUPPLEMENTAL FIGURES

Figure S1



dSTORM imaging of MET receptor in fixed HeLa cells.

MET was immunostained with Alexa Fluor 647. (A) Comparison of wide field and super-resolution dSTORM images (scale bar 2 μm). (B) Enlarged section ($2 \times 2 \mu\text{m}^2$) of the inset in (A), (C) dSTORM image of the section in (B).