

The electronic publication

The Boscaglia Vegetation Complex in Southern Somalia

(Pignatti and Warfa 1983)

has been archived at <http://publikationen.ub.uni-frankfurt.de/> (repository of University Library Frankfurt, Germany).

Please include its persistent identifier [urn:nbn:de:hebis:30:3-369855](http://nbn-resolving.org/urn:nbn:de:hebis:30:3-369855) whenever you cite this electronic publication.

The screenshot shows the digital library interface for the University of Frankfurt. At the top, there is a blue navigation bar with the university logo, the text 'Elektronische Dokumente Universitätsbibliothek UB', and navigation icons for 'Startseite', 'Suchen', 'Browsen', 'Veröffentlichen', and 'FAQ'. A 'Login' and 'Englisch' link is also present.

The main content area displays the title 'The Boscaglia Vegetation Complex in Southern Somalia' by Sandro Pignatti and Ahmed Mumin Warfa. A brief abstract is provided: 'The boscaglia vegetation complex in Southern Somalia on fossil dunes is composed by 3 associations: (a) the Gisekio-Tephrosietum, a xerophilous pasture with dominance of paleotropical weeds; (b) the Acacietum tortil-busseii, a thorny woodland with several Acacia-species and (c) the Justioio-Asystasiatum gangeticum with sciophilous species. Each of them is described in its floristical characters and dynamism.'

Options for 'VOLLTEXT DATEIEN HERUNTERLADEN' (Download full text files) and 'METADATEN EXPORTIEREN' (Export metadata) are visible. The 'WEITERE DIENSTE' (Further services) section includes icons for Twitter and Google Scholar.

The 'Metadaten' (Metadata) section is presented as a table:

Verfasserangaben:	Sandro Pignatti, Ahmed Mumin Warfa
URN:	urn:nbn:de:hebis:30:3-369855
ISSN:	0722-494X
Titel des übergeordneten Werkes (Mehrsprachig):	Tuexenia : Mitteilungen der Floristisch-Soziologischen Arbeitsgemeinschaft
Dokumentart:	Wissenschaftlicher Artikel
Sprache:	Englisch
Datum der Veröffentlichung (online):	08.02.2015
Jahr der Erstveröffentlichung:	1983
Veröffentlichende Institution:	Univ.-Bibliothek Frankfurt am Main
Datum der Freischaltung:	08.02.2015
Jahrgang:	3
Erste Seite:	159
Letzte Seite:	168
DDC-Klassifikation:	580 Pflanzen (Botanik)
Sammlungen:	Sondersammelgebiets-Volltexte
Zeitschriften / Jahresberichte:	Tuexenia : Mitteilungen der Floristisch-Soziologischen Arbeitsgemeinschaft, Band 3 (1983)
Zeitschrift:	Dazugehörige Zeitschrift anzeigen
Lizenz (Deutsch):	Veröffentlichungsvertrag für Publikationen

The Boscaglia Vegetation Complex in Southern Somalia

- Sandro Pignatti and Ahmed Mumin Warfa' -

SUMMARY

The boscaglia vegetation complex in Southern Somalia on fossil dunes is composed by 3 associations: (a) the *Gisekio-Tephrosietum*, a xerophilous pasture with dominance of paleotropical weeds; (b) the *Acacietum tortili-bussei*, a thorny woodland with several *Acacia*-species and (c) the *Justicio-Asystasiatum gangeticae* with sciaphilous species. Each of them is described in its floristical characters and dynamism.

ZUSAMMENFASSUNG

Als Boscaglia wird ein Vegetationskomplex bezeichnet, der in Süd-Somalia auf fossilen Dünen vorkommt und aus 3 Assoziationen besteht: (a) das *Gisekio-Tephrosietum*, eine xerophile Weidengesellschaft mit Dominanz von paleotropischen Krautarten; (b) das *Acacietum tortili-bussei*, ein stacheliges Gebüsch mit mehreren *Acacia*-Arten und (c) das *Justicio-Asystasiatum* mit einli- gen Schattenpflanzen. Jede Gesellschaft wird in ihren floristischen Merkmalen und ihrer Dynamik beschrieben.

INTRODUCTION

Mogadishu, the capital of Somalia, is surrounded by a system of sand dunes. This subdesertic landscape is covered by an open community with dominance of several thorny *Acacia*-species alternating with xerophilous pastures. This vegetation (locally called "boscaglia") is broadly widespread in Southern Somalia and reaches up to the suburbia of the town. During the summer semester 1981, the first of us was appointed as professor of systematic botany at the Agriculture Faculty of the Somali National University, and an investigation of this vegetation was carried out¹⁾.

The first botanical survey of Southern Somalia was published by ROBECCHI-BRICHETTI (1899), and a more detailed study by PAOLI (1916). In these works the name "boscaglia" appears; it comes from the Italian language and was introduced by the explorers during the colonial period. CHIOVENDA (1929), summarized the informations on this vegetation (further additions in SENNI, 1935) and gave a definition of boscaglia as "the more or less dense shrub vegetation covering a broad belt of the interior plain". At the present time, boscaglia is the only word used to designate the thorny woodland in Somalia, even with derivatives (boscagliosi = the nomads living in the shrubs). The Somali language, very rich in precise denominations for the different plant species (and particularly for the Acacias, does not seem to possess an equivalent word for the thorny woodland. A survey of the (mostly physiognomic) literature on this vegetation in Southern Somalia is given in PICHI SERMOLLI (1957, p. 39-49). No phytosociological investigations have been carried out up to the present on the boscaglia. Other vegetation types of the territory are described in CIFERRI (1939) and RAIMONDO & WARFA' (1980).

The investigated area corresponds to the former Benadir in Southern Somalia, i.e. the region surrounding Mogadishu from the coast to the river Shabelle. The whole area is flat, without reliefs of any importance; the only permanent water is the river Shabelle. The area is rather densely populated, chiefly in the irriguouse belt along the river; the major urban area is Mogadishu, large agricultural settlements are also Afgoy and Balad.

METHODS

The analysis of the vegetation was carried out in accord with the principles of the phytosociological method (BRAUN-BLANQUET 1964, WESTHOFF & VAN DER MAAREL 1973). In the herbaceous vegetation the usual phytosociological techniques have been used, with relevés of mostly 100 m² (exceptionally 50-250 m²)

¹⁾ We are indebted to Dr. KAZMI (The National Herbarium-Mogadishu), for facilitation in consulting the collections, to Prof. SHIRWA (Agricultural Faculty-Mogadishu) for unpublished climatic data, and to Dr. P. NIMIS (Istituto Botanico - Trieste) for a careful lecture and discussion of the manuscript.

in almost omotonous areas, largely exceeding the minimal area of the association. The woody vegetation, on the contrary, required some methodological modifications because of their general discontinuity: the relevés were begun in a single dense shrub of 50-70 m² area (hardly corresponding to the minimal area of this vegetation), and then continued on surrounding shrub up to a complete inventory of the woody component, i.e. 200-800 m²; cover values have been estimated in the first area investigated, the border of shrub (mostly infiltrated with herbaceous vegetation), has been neglected. Floristic nomenclature follows CUFODONTIS (1953-1972) with some recent additions. Ordination of phytosociological tables with the WILDI and ORLOCI software (1980). Type relevés labelled with (*).

CLIMATE

The mean yearly temperature in Southern Somalia is lower along the coasts (from 26° to 27°C) and slowly increases in the internal plain, up to 28.7° in Bulo Burti, but the differences are feeble. Even seasonal differences are small and temperature remains remarkably constant all over the year. On the contrary, strong differences can be observed in rainfall. In the coastal region only one rainy period occurs, in correspondence with the SW-Monsun, i.e. from April to July, the completely dry period lasts 4 months, from December to March; total precipitations vary from 250 to 400 mm in the year. In the internal plain, even at only 30-40 km distance from the sea, rainfall shows an almost bimodal distribution, with two maxima, in April-May and in October-November; the dry period is mostly reduced to 3 months (January to March); total precipitation reaches 400 to 500 mm. These features correspond to a semi-arid climate of the II(III)2 type following WALTER & LIETH. With increasing distance from the coast a second period of drought (from June to September) appears, mean yearly temperature increases up to 30°, total precipitation decreases under 250 mm and the climate can be regarded as semi-desertic; further informations in PRETO (1976).

Mean values of yearly rainfall: Mogadishu 483 mm
Afgoy 534 mm
Bur Acaba 559 mm

GEOMORPHOLOGY AND SUBSTRATE

The investigated vegetation grows on the big Somali dune (cfr. PERISOTTO 1978). This is a system of dunes, parallel to the coast of the Indian Ocean, extending from Northern Somalia to the mouth of the river Juba, next to the Kenya frontier. The big Somali dune is about 1200 km long and mostly 20-30 km wide, and was built in geological times; actually this dune system may be regarded as a fossil one. The dune system mostly begins some hundred meters from the sea shore (which partly consists of sand of recent coralline origin and partly of sedimentary rocks) and reaches the maximum elevation (60 to 120 metres above sea level) some km far from the sea; more inwards, the dune is flattened and partly eroded and gently declines to the alluvial plain of the river Shabelle.

The dune is composed by a typical reddish sand rich in silicates and poor in cations. The superficial sand of the dune is completely incoherent and can be easily blown away by the Monsun wind. In places where vegetation is scarce or lacking imponent phenomena of aeolian erosion (on the coastal side), wandering dunes and overlay with sand of the agricultural land have been observed. Under the vegetation cover, the dune is almost stable, and in 50-100 cm depth, the sand appears compact and coherent (even in dry state).

BIOCLIMATICAL ZONES

According to the scheme proposed by CHEVALIER (1933) and recently reviewed by SCHNELL (1976), Southern Somalia (with yearly rainfall of 350-600 mm) can be regarded as belonging to the "Sahelian zone" of vegetation. The climate of Northern Somalia on the contrary is of the desertic type (rainfall lower than 350 mm). In the southernmost part of the Somali Republic near to the frontier of Kenya, annual rainfall reaches up to 600-700 mm and this territory possibly belongs to the "Sudanian zone". All relevés of this study have been carried out within the Sahelian zone of Somalia. The territory is flat and therefore it is not possible to distinguish vegetation belts or any other altitudinal difference in vegetation. A vegetation map in scale 1:500 000 was carried out by DE MARCO & FAGOTTO (1978) with quite accurate cartography of the boscaglia limits in the territory between Brava and Merca.

VEGETATION

1. Pastures: *Gisekio-Tephrosietum pumilae* (tab. 1)

1.1 Description

The vegetation of the pastures on the dunes surrounding Mogadishu and in the neighbouring area, up to Balad (in the North) and to Afgoy (in the West) can be indicated as *Gisekio-Tephrosietum pumilae*. In general this association occurs in mosaic with the xerophilous thorny woodland, characterized by *Acacia* species.

Tephrosia pumila (Lam.) Pers. is the dominant species in these pastures. It is a small perennial herb (fam. *Papilionaceae*) with 5-15 cm long creeping branches and woody base; the root system is well developed and contributes to the consolidation of sand. Nodules with nitrifying bacteria are frequent; the herb seems to be of low pubular value. Other high-frequency species are *Ipomoea garckeana*, *Commelina forskahlei*, *Gisekia pharnacoides*, *Latipes senegalensis*, *Pedaliium murex*, *Psilotrichum gnaphalobryum* and *Eragrostis ciliaris*.

Facies with dominance of *Tephrosia/Ipomoea* or with dominance of *Commelina/Latipes* occur frequently; vegetational aspects with dominance of *Pedaliium* or *Gisekia* are also diffused. A quite unusual facies with abundant *Oxygonum* is widespread near Balad. In the coastal belt near the airport there is a facies with *Aristolochia rigida* (not recorded in table 1). Seedlings of woody plants

Tab. 1 - GISEKIO-TEPHROSIIETUM PUMILAE

Number of relevés	1	2	3	4	5	6	7	8	9*	10	11	12	13	14	15	16	17			
Elevation in m	120	120	90	90	90	40	90	10	60	100	90	90	110	90	90	90	90			
Slope in %	-	-	-	-	-	-	0-5	-	2	-	1	-	-	-	-	-	-			
Exposure	95	-	-	-	-	-	-	-	N	-	N	-	-	-	-	-	-			
Cover in %	95	95	70	80	70	70	85	80	80	60	75	85	70	95	80	85	85			
Surface in m ²	150	150	75	70	100	50	100	50	100	250	100	100	150	100	100	100	100			
Character-species of the association																				
<i>Tephrosia pumila</i> (Lam.) Pers.	+	+	3,1	1,1	+	2,2	2,2	2,3	3,3	+	3,2	2,2	3,3	+	3,3	3,2	3,3	100		
<i>Gisekia pharnacoides</i> L.	+	+	2,2	+	+	+	+	+	+	1,1	+	+	+	+	+	1,2	2,1	+	88	
<i>Sesilla aemulans</i> Baker	(+)	+	+	+	+	+	+	+	(+)										41	
Differential species of an East-African syntom																				
<i>Ipomoea garckeana</i> Vathe	3,3	+	1,1	2,1	1,1	1,1	1,1	1,1	+	+	+	+	+	1,1	+	1,1	1,1	1,1	100	
<i>Psilotrichum gnaphalobryum</i> (Hochst.) Schinz	+	1,1	+	1,1	2,1	+	1,2	1,1	1,2	+	+	1,1	+	+	+	+	1,1	+	1,1	100
<i>Indigofera ferric</i> Chiov.	+	+	+	1,1	+	+	+	+	+	+	+	+	+	(+)	+	+	+	+	+	65
<i>Cissampelos strigosa</i> (Kojer) Oliver							1,1	+	+	+	+	1,1	(+)	+	+	1,1	1,1	53	35	
<i>Dignatia nitida</i> Stapf	+2	1,2					+	+	+	+										24
<i>Indigofera tomponkiana</i> Baker		1,1																		
Palaeotropical weeds (ch.-species of the class ?)																				
<i>Commelina forskahlei</i> Vahl	1,1	1,1	1,1	1,1	2,2	1,1	2,2	(+)	1,1	1,1	+	1,1	+	+	2,2	1,2	1,1	100	100	
<i>Pedaliium murex</i> L.	r	r	+2	2,1	2,1	1,1	2,1	(+)	(+)	+	1,1	2,2	+	+	1,2	2,1	+	100	100	
<i>Latipes senegalensis</i> Kunth	1,2	1,2	+	1,2	+	2,3	+	2,2	2,2	+	+	2,2	+	+	1,1	1,2	+2	94	94	
<i>Eragrostis ciliaris</i> (L.) K.Brown	1,1	1,1	1,1	+	1,1	1,1	+	+	+	+	+	1,1	1,1	+	1,1	1,1	+	1,1	94	
<i>Oxygonum striplifolium</i> (Meisn.) Martelli	+	+	+	+	+	+	+	(+)	3,4	1,1	1,1	+	+	1,1	+	1,1	+	1,1	94	
<i>Borreria sodiba</i> (Schum. et Th.) K.Schum.	1,1	1,1	+	+	+	+	2,3	1,1	+	+	+	+	+	1,1	+	+	+	+	88	
<i>Digitaria miltaniensis</i> Stapf	+	2,3	+	+	2,2	+	(+)	+	+	1,1	+	+	+	1,1	+	+	+	+	82	
<i>Zornia glochidiata</i> Rech.	+	+	+	+	+	+	+	r	1,1	1,1	1,1	1,1	+2	1,1	1,2	65	65	65		
<i>Portulaca quadrifida</i> L.	+	1,1	+	+	+	+	+	+	+	+	+	(+)	+	+	+	+	+	+	53	
<i>Heliotropium subulatum</i> (DC.) Vathe	+	+	+	+	+	1,1	+	+	+	+	+	+	+	+	+	+	+	+	53	
<i>Hibiscus cranesburianus</i> Hochst.	+	+	+	+	+	+	+	1,1	+	+	+	+	+	+	+	+	+	+	41	
<i>Tephrosia multiflora</i> Baker	+	+	+	+	+	+	+	+	+	+	+	+	(+)	+	+	+	+	+	67	
<i>Cuscuta dipetala</i> Ehrenberg	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+	41	
<i>Cenchrus ciliaris</i> L.	2,2													1,1	+	+	+	+	67	
<i>Echaulia nasayitana</i> Schindl.	+													(+)	+	+	+	+	35	
<i>Indigofera schimper</i> Jaub. et Spach	+								(+)	+	+	+	+	+	+	+	+	+	35	
<i>Phyllanthus rotundifolius</i> Klein	+													+	+	1,1	+	+	29	
<i>Dactyloctenium aegyptium</i> Neesler	+	1,2	+											+	+	+	1,1	+	29	
<i>Scleraria adalis</i> (Forsk.) Pers.	1,1	1,1						2,2	1,1										24	
<i>Urochloa panicoides</i> P.Beauv.	+	1,2																	18	
<i>Tephrosia quinqueana</i> Cuf.	(+)																		18	
Companions																				
<i>Aerva persica</i> (Burm.f.) Merrill	1,1	+	+	1,1	+	+	+	+	(+)	+	+	+	+	+	1,1	+	+	+	96	
<i>Cenchrus ciliaris</i> Meisner	+	+	+	+	+	+	+	+	+	1,1	+	+	+	+	+	+	+	+	35	
<i>Cyperus amblyanthus</i> L.	+	+2	+2	1,1	1,2								(+)	+	+	+	+2	+2	53	
<i>Acacia senegal</i> (L.) Willd. (young plants)																			53	
<i>Jussiaea flava</i> (Vahl) Vahl		1,1	+											1,2				1,1	35	
<i>Cenchrus biflorus</i> Roxburgh	2,2	+																	35	
<i>Acacia tortilis</i> (Forsk.) Hayne (young plants)	+	+																	35	
<i>Dactyloctenium aegyptium</i> (L.) Richter	+	+																	35	
<i>Furcraea</i> sp.	+	+																	29	
<i>Oxymon hederifera</i> Forsk.	+	+								+									24	
<i>Achyrocline satureioides</i> (L.) Anders.		1,1								1,1				+					24	
<i>Burkera waggana</i> Rendle	+2	+	+							1,2									24	
<i>Acacia bussei</i> Harms (young plants)	+	+												+					24	
<i>Solanum jubae</i> Bitter	+	+																	18	
Sporadic species (under 20 %)																				
	4	2	2	1	3	1	3	4	6	2	1	3	2	5	-	-	-	-	-	

(*Acacia tortilis*, *A. bussei*, *A. senegal*) mostly 1-3 dm high are very scattered but nearly always present; they show the trend of the vegetational succession.

The majority of the more diffused species are perennial herbs with prostrate stems or rhizomes. Consequently the vegetation hardly reaches 20 cm in height. Only a few species with low cover values form an upper layer 30-60 cm high; they are small shrubs with erect habitus and woody branches (*Aerva*) or at least with woody base (*Cleome*, *Indigofera*).

The seasonal development of the *Gisekio-Tephrosietum* begins in June with the rainfall period and lasts about three months. In this short period most species accomplish their vegetative growth, produce flowers, fruits and seeds. During August or in the first decade of September leaves and other aerial parts progressively dry up and the vegetation turns in a condition of vegetative dormancy. Perennial herbs survive up to the next year chiefly with their root systems or with subterranean rhizomes. Reproduction by seeds seems relatively infrequent and only few seedlings can be observed at the beginning of the rainfall period. Even therophytes are scarce.

1.2 Minimal area

The extension of the minimal area was calculated for 6 relevés of tab. 1; the results are given in tab. 2. The number of observed species increases very steeply from the first m² up to 10 m². After that, the increase is more slow. In 10 m² 17 species are present (mean value) i.e. nearly 70% of the complete assemblage of the association. In 20 m² more as 19 species are present (78%). Further broadening of the investigated area brings very few new species. The flexus of the curve seems to be near 20 m². Therefore, it may be assumed that relevés of 100 m² can retain the complete floristical information of this vegetation type.

TAB. 2 - Minimal area

relevés	1	2	3	4	5	6	observations (total)	mean	sum
1 m ²	8	8	11	9	8	8	52	8.66	8.66
2	3	1	1	2	2	3	12	2.0	10.66
5	5	3	5	3	1	1	18	3.0	13.66
10	2	4	4	3	4	3	20	3.33	17.00
20	1	4	1		5	2	13	2.16	19.16
30	1	1	2	1	1	1	7	1.16	20.32
40	1			3	1	1	6	1.00	21.32
50	1				1	1	3	0.50	21.82
100	3		3	2	2	5	15	2.50	24.32
200					1		1	0.16	24.48

1.3 Human activity

The pastures of the *Gisekio-Tephrosietum pumilae* type are submitted to intense grazing by goat, camels, zebu and sheep. The available biomass is of 100-250 g/m² of dry fodder (average value of 7 determinations), i.e. very low. The main fodder species seem to be *Commelina*, *Borreria*, *Psilotrichum* and the grasses; *Tephrosia* on the contrary, seems to be avoided by the cattle. Poisonous plants (*Euphorbia*, *Solanum*) occur in single individuals and with very low frequency. Spiny plants are almost absent from the herbaceous layer (on the contrary, they dominate the woody vegetation). The discontinuous vegetation prevents the expansion of fire and consequently burning does not seem to have a significant importance for this vegetation type.

The traditional agricultural exploitation is based on the conversion of open pastures into closed fields (shambas). Small depressions between the dunes are fenced with branches of spiny *Acacia*, the woody vegetation is cut and burned and the herbaceous layer can massively develop because of the higher humidity, the presence of ash in the soil and even of a weak manuring. In this way soil conditions improve and the herbaceous vegetation can reach a cover of nearly 100%, with height of 2-4 dm and biomass of 500 g/m² and more. The fodder is harvested in July-September and sold in the town. After harvesting, cattle is introduced in the shamba with some restrictions as far as grazing is concer-

ned. Unregulated grazing is almost avoided. A single surface can be used as shamba for many years (up to 50 years and more) and after exhaustion of the residual fertility is abandoned again. Differential species of the shambas aspects are *Blepharis edulis*, *Dactyloctenium aegyptium*, *Dignatia* and others.

1.4 Syntaxonomy

The *Gisekio-Tephrosietum* shows close relationships to the anthropogenous vegetation of the neighborhood of Mogadishu described by RAIMONDO & WARFA' (1980). The floristic affinity is particularly high with the *Aristolochia rigida* community, which is surely a distinct association, but possibly belonging together with the *Gisekio-Tephrosietum* to a single alliance. Unfortunately RAIMONDO & WARFA' do not indicate which higher syntaxa their associations can be referred to.

An association with *Tephrosia pumila* as a character species is described by JENIK & HALL (1976) from the pastures of the Accra Plains (Ghana); this community belongs to the class *Vetiverietaea*. No other species are in common with the *Gisekio-Tephrosietum* of Somalia and even the bioclimatic features of the Accra Plains are quite different (rainfall of 700-1100 mm, i.e. typical for the Sudanian zone). Consequently it is impossible to include the *Gisekio-Tephrosietum* into the *Vetiverietaea*.

Some pasture communities from the Sahara described by QUEZEL (1965) seem to be related to the *Gisekio-Tephrosietum*, namely the ass. of *Lotus glinoides* and *Matthiola livida* (from the Hoggar in 1000-1500 m altitude), the ass. of *Fuellia patula* and *Cadaba farinosa* and the ass. of *Aristida funiculata* and *Indigofera sessiliflora* (both from the Emi Koussi in 1260-1550 m and 500-1200 m altitude respectively). These associations have a group of species in common with the *Gisekio-Tephrosietum* (*Aerva persica*, *Grewia tenax*, *Gisekia pharnacioides*, *Blepharis edulis*, *Cenchrus biflorus*) but the saharo-sindian and submediterranean species are dominating. Consequently it is not clear, if these associations belong to the same class as the *Gisekio-Tephrosietum*. Anyway, they represent the extreme penetration in northern direction of equatorial elements, living in the middle of the Saharian zone. They are limited to the montane habitats (extrazonal) and therefore they cannot be regarded as particularly significant.

Closer floristic affinity shows the community of *Schmittia pappophoroides* and *Panicum turgidum* described by QUEZEL (1969) in the Darfur area (Sudan Republic). Species in common with the *Gisekio-Tephrosietum* are *Cenchrus biflorus*, *Gisekia pharnacioides*, *Zornia glochidiata*, *Latispes senegalensis*, *Dactyloctenium aegyptium*, *Aerva persica*, *Oxygonum atriplicifolium*. This community has been observed on sandy soils at 750-1100 m altitude in a territory with 300-350 mm rainfall (i.e. probably belonging to the Sahelian zone). The higher syntaxonomical units for this vegetation are not indicated.

In conclusion, it seems likely, that at least the *Gisekio-Tephrosietum*, the vegetation described by RAIMONDO & WARFA' near Mogadishu and the Darfur community belong to a unique vegetation class. As character species for this class several paleotropical weeds with sahelian distribution (from West Africa to Arabia and Sindh) could be proposed. Consequently, the class seems to have a broad distributional range. Otherwise, the communities described in Somalia (both by RAIMONDO & WARFA' and by us) are characterized by several East African species, not occurring elsewhere; they may probably belong to a particular syntaxon. The species in Tab. 1 have been ordered following these assumptions, but we think that the present knowledge of this vegetation types is too fragmentary and not sufficient for proposing a new syntaxonomical scheme.

2. The open thorny woodland: *Acacietum tortillibussei* (tab. 3)

2.1 Description

The *Acacia* community is widespread on the sand-dune near Afgoy, between Mogadishu and Afgoy and up to Gesira and Balad. A similar community has been physiognomically recognized also along the dune system from Gandersher to Merca and Brava. Therefore, this association is spread at least over 200 km along the coast.

The dominating element of this association is given by the *Acacia* species:

- A. tortilis* subsp. *spirocarpa* - small tree or gregarious shrub
- A. senegal* - gregarious shrub
- A. bussei* - gregarious shrub
- A. misera* - scattered shrub
- A. nilotica* - shrub, + accidental
- A. mellifera* - shrub, ± accidental

Other widespread shrubs are *Solanum jubae*, many *Capparidaceae* (*Courbonia*, *Cadaba*, *Maerua*, *Boscia*), *Commiphora*, *Uvaria dehnhartiana*, *Anisotes involucreatus* etc.

The following vegetation strata can be identified:

A₂ - Umbrelliform trees-layer (5-8 m high) chiefly composed by *Acacia tortilis* and single individuals of other *Acacia* species, *Balanites* and *Dobera*. The arboreal *Acacia* species cover a very small surface (mostly not more than 5-20%).

B₁ - Higher shrubs-layer (2-6 m height) a compact shrub of different *Acacia* species, in association with other shrubs. This vegetation covers about 80-95% of the surface.

B₂ - Lower shrubs-layer (5-20 dm); sporadically occurs at the edges of *Acacia* thickets covering no more than 5%.

C - Herbs-layer (5-80 cm), particularly abundant under the shrubs, composed by species of the pasture like *Commelina forskahlei* and *Ipomoea garrckeana* or by shadow elements like *Justicia flava*. This layer covers about 20-40%.

D - Climbers, very scattered and with low cover (1% or less).




Fig. 1: Vegetation profile of the *Acacietaum tortili-bussei*.

- | | |
|-----------------------------|--------------------------------|
| 1: <i>Acacia tortilis</i> | 6: <i>Uvaria denhardtiana</i> |
| 2: <i>Acacia senegal</i> | 7: <i>Grewia tenax</i> |
| 3: <i>Acacia misera</i> | 8: <i>Cucumis dipsaceus</i> |
| 4: <i>Solanum jubae</i> | 9: <i>Justicia flava</i> |
| 5: <i>Maerua angolensis</i> | 10: <i>Momordica balsamina</i> |

The photosynthetic activity in the thorny woodland lasts for the whole year, with maxima in the humid periods but without a complete interruption in the dry season. The single species seem to produce leaves and flowers independently from one another and even single individuals of the same species show quite different phenology. In general, plants produce leaves and flowers during a short period (sometimes only 2-3 weeks). We observed leaves and flowers in *Acacia tortilis* (in VII), in *A. bussei* and *A. senegal* (in VIII), in *Courbonia* and *Cadaba* (in IX) and in several *Commiphora* (in X). The wood of *Acacia* shows irregular growth rings, which may be assumed to be annual.

The minimal area of the *Acacietaum tortili-bussei* was not directly investigated, but seems to reach some 100 m².

2.2 Syntaxonomy

Communities of thorny woodland with *Acacia* (*A. tortilis* and other species), *Boscia*, *Balanites*, *Dobera*, *Cadaba*, *Courbonia* etc. are widespread all over the Sahelian zone from Senegal to Eastern Africa and even to Arabia and SW Asia.

TAB. 3 - ACACIETUM TORTILI-BUSSEI

Number of relevés	18	19	20	21	22	23*	24	25	26
Elevation in m	120	120	90	90	90	120	90	90	90
Slope in %	-	-	-	-	-	2	-	-	-
Exposure	-	-	-	-	-	W	-	-	-
Surface in m ²	400	500	300	600	500	800	200	300	400
A₂ - Umbrelliform trees (5-8 m) %	5	20	15	30	2	5	20	5	1
<i>Acacia tortilis</i> (Forak.) Hayne	1.1	1.1	1.1	2.3	+	(+)	1.2	1.1	(+)
<i>Balanites aegyptiaca</i> (L.) Del.	+	(+)				1.1			
<i>Acacia misera</i> Vatke				1.1	+				
<i>Acacia senegal</i> (L.) Willd.		+							(+)
<i>Dobera glabra</i> (Forak.) Juss.							1.1		
<i>Acacia bussei</i> Harms								1.1	
B₁ - Higher shrubs (2-6 m) %	80	98	80	80	85	80	90	70	95
I - thorny Acacia									
<i>Acacia tortilis</i> (Forak.) Hayne	3.3	2.2	4.5	2.3	2.2	1.1	3.4	+	1.1
<i>Acacia senegal</i> (L.) Willd.	3.3	2.2	1.1	1.1	2.2	3.4	2.1	2.2	1.1
<i>Acacia bussei</i> Harms	+	1.1	1.1	1.1	2.2	+	1.1	2.2	4.4
<i>Acacia misera</i> Vatke	+		+		+	3.4			(+)
<i>Acacia nilotica</i> (L.) Del.							1.1	+	
<i>Acacia mellifera</i> (Vahl) Benth.			1.1						
II - others									
<i>Solanum jubas</i> Bitter	1.1	+	+	1.1	+	+	+	1.1	+
<i>Urtica denhartwegii</i> Engl. et Diels	+		1.1	1.1	+	1.1	+	1.1	
<i>Anlaetas umolucatus</i> Fiori	+	(+)	+	2.3	+	(+)	+	+	1.1
<i>Commiphora ellenbachii</i> Engler		(+)			+	(+)	+	+	
<i>Maera angolensis</i> DC.			1.1		2.2	1.1	+	1.1	1.1
<i>Courbonia ramularifolia</i> Mattei			2.2	+	(+)		(+)	1.1	+
<i>Balanites aegyptiaca</i> (L.) Del.	1.1	+			(+)				(+)
<i>Solanum</i> cfr. <i>somalense</i> Franchet	1.1	(+)			1.1	+			
<i>Commiphora africana</i> (Rich.) Engler			+				(+)	1.1	
<i>Combretum oconstrictum</i> (Benth.) Lawson				+	+				(+)
<i>Chordia amalaensis</i> Baker	+	+	(+)	(+)					
<i>Theopatia dante</i> Oliver	1.2	2.2					+		
<i>Cadaba farinosa</i> Forak.	+	+							(+)
<i>Bonasia oortiana</i> Pax	1.1			1.1					
<i>Maera oblongifolia</i> (Forak.) Rich.				+				1.1	2.2
<i>Commiphora</i> cfr. <i>lughensis</i> Chiov.				(+)					+
<i>Cassia longaracemosa</i> Vatke							1.1		
<i>Dobera glabra</i> (Forak.) Juss.									(+)
<i>Chordia ghazaf</i> (Forak.) Aschers.									+
<i>Grewia villosa</i> Willd.									(+)
<i>Dioscorea glomerata</i> (Forak.) Chiov.	+								
<i>Grewia verrucifera</i> Chiov.						+			
<i>Allophylus rubifolius</i> (Hochst.) Engler									+
<i>Hildenbrandtia africana</i> Vatke			+						
B₂ - Lower shrubs (5-20 dm) %	20	5	5	0	5	5	1	5	5
<i>Indigofera sohimperi</i> Jamb. et Spach	+	1.1			1.1	1.1	+	1.1	
<i>Barleria waggana</i> Rendle		1.1	1.1		+2	+2		+	1.2
<i>Grewia tenax</i> (Forak.) Fiori	+	+	+2	+	+	+			
<i>Hibiscus arvensis</i> Hochst.	1.1	+			+				+
<i>Aerva persica</i> (Burm.f.) Merrill							2.3	+	
<i>Pavonia</i> sp.	1.1			+					
<i>Corchorus erinaceus</i> Weimarck					+	+			+
C - Herbs (5-80 cm) %	10	40	20	30	25	25	40	30	40
<i>Commelina forskahlei</i> Vahl	1.1		1.1	2.1	1.1	1.1	1.2	2.3	2.3
<i>Justicia flava</i> (Vahl) Vahl	+	1.2	(+)	+	1.1	1.1	2.3	+	
<i>Ipomoea garckeana</i> Vatke	1.1	+		1.1	+	1.1	+		
<i>Asystasia gargetica</i> (L.) Anderss.	+	+			+	1.1	1.2		
<i>Urochloa panicoides</i> P.Beauv.		2.2	+		+			+	+
<i>Pennisetum graphalobryum</i> (Hochst.) Schinz	+	+	+	+					+
<i>Ottima hadiana</i> Forak.		+	1.1					1.2	1.2
<i>Laticipes senegalensis</i> Kunth				+2	+	+			+
D - Climbers %	10	1	1	1	2	5	5	1	1
<i>Momordica balsamina</i> L.	(+)	1.1			(+)	1.1	1.1	+	+
<i>Combretum oconstrictum</i> (Benth.) Lawson			+	(+)	+	(+)	+		+
<i>Pentstemon spiralis</i> (Forak.) Decne			+	+	+	+			66
<i>Cucumis dipsacous</i> Ehrenb.	+	+			+			+	+
<i>Coccoloba grandis</i> (L.) Voigt						(+)			(+)
<i>Triumfetta astinocarpa</i> Moore								+	
<i>Cissampelos digitata</i> (Forak.) Lam.					+				
<i>Sarcostemma viminale</i> (L.) R.Br.				(+)					
<i>Ipomoea garckeana</i> Vatke									+
<i>Pergularia damia</i> (Forak.) Chiov.									(+)
Sporadic species (under 20 %)	2	7	4	1	1	3	-	-	6

For a survey cfr. SCHNELL (1976). Phytosociological tables have been published by QUEZEL (1969) from the Darfur area in Sudan (the *Acacia mellifera*-, the *Acacia nubica*- and the *Digera-Aeschynomene* communities). This vegetation is probably related to the microphyllous thorny woodland of the Transvaal Bushveld (South Africa), which shows (VAN DER MEULEN, 1978) *Acacia tortilis*, *A. misera*, *A. nilotica* and other species present in Southern Somalia and close vicarism at the genus level. It seems likely, that a "Sahelian" vegetation class with continental distribution still exists, but the informations deriving from scattered territories are up to now too fragmentary for a more precise definition.

3. The vegetation of shady places: *Justicia - Asystasiaetum gangeticae* (tab. 4)

Acacias produce a feeble shadow, particularly when covered by climbers and a dense community of *Justicia flava* and *Asystasia gangetica* regularly occurs on these shady niches. In most cases there are only fragments of 1-5 m² which are hardly sufficient for relevés. Several species of the pastures and of the thorny woodland are also present. Therefore the community can be identified only in a very confuse form. A similar vegetation is described by QUEZEL for the Darfur area (Sudan) as meso-sciaphilous community of *Craterostigma plantagineum* and *Kalanchoe modesta*. In this community some species occurring also in Somalia have been found, namely *Achyranthes aspera*, *Ocimum haldiense* and another *Justicia* (*J. galeopsis*). The syntaxonomical position of this vegetation is quite obscure.

TAB. 4 - JUSTICIO-ASYSTASIIETUM GANGETICAE

Number of relevés	27	28	29	30	31*	32	33
Elevation in m	120	90	90	90	90	90	75
Slope in %	-	2	2	-	-	-	-
Exposure	-	N	N	-	-	-	-
Cover in %	100	100	100	70	60	95	100
Surface in m ²	6	8	4	4	10	25	8
Character-species of ass. and higher syntaxa							
<i>Asystasia gangetica</i> (L.) Anders.	2.2	2.2	(+)	1.1	1.1	2.2	1.1
<i>Justicia flava</i> (Vahl) Vahl	+	1.2	(+)	2.2	2.3	+	3.3
<i>Achyranthes aspera</i> L.	(+)	1.2				1.1	
<i>Ocimum haldiense</i> Forsk.			2.2		(+)		
Climbers							
<i>Ficus laevis</i> (Forsk.) Chiov.		1.1			+	+	+
<i>Cucumis dipsacoides</i> Ehrenb.		+				+	
<i>Momordica balsatina</i> L.			+				
<i>Flauropentaria revoluta</i> Franchet	+						
<i>Coccinia grandis</i> (L.) Voigt	+						
Paleotropical weeds							
<i>Commelina forskahlei</i> Vahl	2.2	3.3	2.2	1.1	1.1	1.1	+
<i>Urochloa panicoides</i> P.Beauv.	1.2		1.2	+		2.2	+
<i>Cenchrus ciliaris</i> L.						+	2.2
<i>Indigofera ciferrii</i> Chiov.	+		+				
<i>Borreria arabica</i> (Schum. et Th.) K.Schum.			1.1				
Companions							
<i>Momordica debilis</i> (Forsk.) Nees	2.2				(+)	1.1	+
<i>Ipomoea garckeana</i> Vatke		1.1	+	+			+
<i>Indigofera arvensis</i> Hochst.				1.1	+		+
<i>Pavonia</i> sp.	+					+	
<i>Tridax procumbens</i> L.	+		1.1		1.1		
<i>Ptilorhizum graphalobryum</i> (Hochst.) Schinz					1.1	+	+
<i>Solanum</i> cfr. <i>somalense</i> Franchet	1.1		+				
<i>Borreria laggardii</i> Rendle	1.2		+	2			
<i>Cassia longicaespita</i> Vatke						+	
<i>Withania somnifera</i> (L.) Dunal		+				(+)	
<i>Boerhaavia diffusa</i> L.					+	+	
<i>Solanum arundo</i> Mattei						+	
<i>Acacia tortilis</i> (Forsk.) Hayne (young plants)						3.3	
Sporadic species (under 20 %)	5	1	1	2	1	2	

LANDSCAPE ECOLOGY

The Boscaglia is the vegetation complex consisting of the *Gisekio-Tephrosietum*, *Acacietum tortili-bussei* and *Justicio-Asystasiatum*. In most places the three vegetation types occur together, often with mosaic-like patterns. The thorny woodland dominates the landscape; pastures are also frequent and mostly cover 20-30% of the area, whereas the vegetation of shady places is quite rare and without an appreciable cover.

Natural growth, pasture and human activity are responsible of successional cycles in the boscaglia. At least 3 phases can be distinguished:

I) young phase - only higher shrubs, mostly with discontinuous cover. Dominant species are *Acacia senegal* and *A. bussei*; elements of the *Gisekio-Tephrosietum* are almost abundant; pasture intense.

II) optimal phase - higher shrubs compose a dense and almost continuous vegetation 2-6 m high (*Acacia senegal*, *A. bussei* and *A. tortilis* are co-dominant) with single umbrelliform trees of *A. tortilis* reaching 6-8 m in height; herbs became very rare; pasture activity is scarce and limited to the marginal zones.

III) senescence phase - umbrelliform trees became more frequent, higher and with denser canopy. In their shadow the shrubs show reduced vitality or disappear; such places are often occupied by nomads as temporary refuge, lower shrubs are cut, climbers (sometimes favoured by man) and nitrophilous species became more frequent; elements of the *Justicio-Asystasiatum* often spread in more shady places.

The phase III allows more or less frequent passage of humans and cattle, intensive pasture and cutting of the woody species. When the umbrelliform trees disappear (cut for wood or naturally dead) direct sunlight can penetrate up to the soil surface and the vegetation turns back to phase I. Then, the cycle is complete.

In present time this dynamism is chiefly caused by human activity and can be interpreted as completely anthropogenous. Despite this, vegetation seems to be in a semi-natural condition. It is a matter of fact, that pasture intensity in this ecosystem was always high: presently by cattle, but before the settlement of man by wild species: dig-dig, antelopes, giraffes, donkeys and other big mammals of the savanna- and sahelian-environment. Human action substituted the action of wildlife without producing major changes in the vegetation. Plant growth, natural decay of organic matter and pasture compose in Southern Somalia a well integrated ecosystem which works since immemorial times in a condition of steady state.

LITERATURE

- BRAUN-BLANQUET, J. (1964): Pflanzensozioologie 3. Aufl. - Wien (Springer): 865 pp.
- CHEVALIER, A. (1933): Le territoire géo-botanique de l'Afrique tropicale Nord-occidentale et ses subdivisions. - Bull. Soc. Bot. Fr. 80: 4-26.
- CHIOVENDA, E. (1929): Flora Somala. - Roma (S.I.Arta Grafiche): 436 pp.
- CIFERRI, R. (1939): Le associazioni del litorale marino della Somalia meridionale. - Riv. Biol. Colon. 2: 5-42.
- CUFODONTIS, G. (1953-72): Enumeratio plantarum Aethiopiae Spermatophyta. - Bull. Jard. Bot. Etat Bruxelles 23-42.
- DE MARCO, G., FAGOTTO, F. (1978): Geobotanical and faunistic map of the lower Scebeli river (Somalia). - Pubbl. Centro Cartogr. Ecol. Appl. Univ. Roma.
- JENIK, J., HALL, J.B. (1976): Plant communities of the Accra Plains, Ghana. - Folia Geobot. Phytotax. 11: 163-212. Praha.
- PAOLI, G. (1916): Risultati botanici delle missioni scientifiche Stefanini-Paoli nella Somalia Italiana meridionale. - Mem. Accad. Lincei Cl. Sc. Fis. Mat. Sez. 5, vol. 11, fasc. 10.
- PERISOTTO, A. (1978): Appunti di geomorfologia della Somalia. - Quad. Geol. Somalia 2: 45-64.
- PICHI-SERMOLLI, R.E. (1952): Una carta geobotanica dell'Africa Orientale (Eritrea, Etiopia, Somalia). - Webbia 13: 15-132.
- QUEZEL, P. (1965): La végétation du Sahara, du Tschad à la Mauritanie. - Stuttgart (G. Fischer): 333 pp.
- (1969): Flore et végétation des plateaux du Darfur Nord-occidentale et du jebel Gorgeuil. - C.N.R.S. Dossiers R.C.P. n° 45: 145 pp.

- RAIMONDO, F.M., WARFA', A.M. (1980): Preliminary phytosociological research on synanthropical vegetation in Southern Somalia. - Not. Fitosoc. 15: 189-206.
- ROBECCHI-BRICCHETTI, L. (1899): Somalia e Benadir (in append. PIROTTA, R.: Notizie sulle collezioni botaniche etc.). Milano, pp.701-726.
- SCHNELL, R. (1976): La flore et la végétation de l'Afrique tropicale. - Paris (Gauthier-Villars): 459 pp.
- SENNI, L. (1935): Gli alberi e le formazioni legnose della Somalia. - Ist. Agric. Coloniale It. Firenze: 305 pp.
- VAN DER MEULEN, F. (1979): Plant sociology of the western Transvaal Bushveld, South Africa. - Diss. Bot. 49. Vaduz (Cramer): 191 pp.
- WESTHOFF, V., VAN DER MAAREL, E. (1973): The Braun-Blanquet approach. - In: Handb. of Veget. Science V: WHITTAKER, R.H.: Ordination and classification of vegetation: 619-726. The Hague (Junk).

Anschriften der Verfasser:

Prof. Dr. Sandro Pignatti
Istituto Botanico - Città Universitaria
I - 00100 Roma (Italy)

Ahmed Mumin WARFA'
Università Nazionale Somala
Facoltà di Agraria
p.o. box 801
Mogadishu (Somalia)