



First report of *Culex (Culex) tritaeniorhynchus* Giles, 1901 (Diptera: Culicidae) in the Cape Verde Islands

Joana Alves^{1,2}, Adilson de Pina¹, Mawlouth Diallo³ & Ibrahima Dia³

Keywords: Culicidae, *Culex tritaeniorhynchus*, Cape Verde Islands, first occurrence.

ABSTRACT

During an entomological survey in Santiago Island, Cape Verde Islands, in November-December 2011 in order to study the bio-ecology and susceptibility to insecticides of *Anopheles arabiensis* Patton, 1905, *Culex tritaeniorhynchus* Giles, 1901 was found to be present in the Santa Cruz District. Both adult and immature specimens were collected and a description of both is given. Further confirmation of the taxonomic identity of the specimens was obtained from studying the male genitalia. This is the first known occurrence of *Culex tritaeniorhynchus* (a primary vector of Japanese encephalitis in eastern and southern Asia and a potential vector of West Nile Virus, Sindbis and Rift Valley Fever Virus) in the Cape Verde Islands.

RESUMO

Durante um levantamento entomológico realizado na ilha de Santiago, arquipélago de Cabo Verde, em Novembro-Dezembro de 2011, com o objectivo de estudar a bioecologia e susceptibilidade de *Anopheles arabiensis* Paton, 1905, foi encontrado *Culex tritaeniorhynchus* Giles, 1901, no Concelho de Santa Cruz. Foram recolhidos, identificados e descritos espécimes quer no estado adulto quer na forma do quarto estadio larvar. A identificação taxonómica foi posteriormente confirmada através do estudo da genitália de machos. Relata-se pela primeira vez, nas ilhas de Cabo Verde, a ocorrência de *Culex tritaeniorhynchus*, vector primário do vírus da Encefalite Japonesa na Ásia Oriental e do Sul e vector potencial dos seguintes vírus: o da febre do Nilo Ocidental, Sindbis e febre do Vale do Rift.

¹ Direcção Nacional da Saúde, Ministério da Saúde, Palácio da Várzea, Praia, Republic of Cape Verde; Joana.Alves@cnds.gov.cv

² Centro de Malária e Outras Doenças Tropicais, Universidade Nova de Lisboa, Portugal

³ Unité d'entomologie médicale, Institut Pasteur de Dakar, BP 220, Dakar, Sénégal

INTRODUCTION

Updating knowledge of the culicidean fauna of the Cape Verde Islands has been a priority for many years (e.g. Ribeiro *et al.* 1980, Cambournac *et al.* 1982, Alves *et al.* 2010). During an entomological survey, conducted between November and December 2011, in order to study the bio-ecology

and susceptibility to insecticides of *Anopheles arabiensis* Patton, 1905 in Santiago Island, *Culex (Culex) tritaeniorhynchus* Giles, 1901 was found to be present in the Santa Cruz district (Fig. 1). We describe the site and provide descriptions of both adult and larval stages.




Fig. 1. Maps of Santiago Island and the Santa Cruz district, indicating (*) the locality where specimens of *Culex tritaeniorhynchus* were found.

The habitat consisted of a natural basin of rain water drainage in a confluent zone between a lagoon and a beach in a coastal region (Fig. 2). The breeding site was exposed to sunlight, characterized by the presence of emergent and fluctuant aquatic vegetation and the water was slightly brackish, but within the limits of being considered freshwater. So far, only *Culex ethiopicus* Edwards, 1912 had been found breeding at this and similar sites in the area. *Culex*

tritaeniorhynchus is considered a primary vector of Japanese encephalitis in eastern and southern Asia (van den Hurk *et al.* 2009). It is also a potential vector of West Nile Virus (WNV), Sindbis and Rift Valley Fever Virus (RVFV) (Wills *et al.* 1985, Kheir *et al.* 2010). The presence of *C. tritaeniorhynchus* in this area underlines the need for further investigations into its significance for public health and epidemiological issues.

MATERIAL AND METHODS

We visited localities in five districts of Santiago Island to collect immature stages of mosquitoes at several breeding sites, including ponds, pools, pits

and tanks. Upon collection, all immature mosquitoes were placed in containers and transported to the laboratory, where they were reared until

they emerged as adults. In addition, adult mosquitoes were collected with CDC light traps (placed in local people's bedrooms and in sheds for cattle, goats and sheep) and by using pyrethrum spray in human dwellings. After

collection, mosquito specimens were sorted, counted and identified morphologically using a stereomicroscope and according to the identification keys provided by Ribeiro & Ramos (1995).



Fig. 2. Partial view of the breeding site of *Culex tritaeniorhynchus*, Santa Cruz district, Santiago Island, Cape Verde Islands, where specimens were collected (photo by Ibrahima Dia; © Joana Alves).

RESULTS

A total of 2,192 mosquito specimens were identified. The distribution by district and capture method is given in Table 1. A large majority of specimens were immatures (94.1%), followed by resting females from human dwellings. Specimens from CDC light traps in bedrooms and sheds were the least abundant (0.4% and 0.1% respectively). From the total of emerged specimens, *Culex* spp. averaged 7%, while in the district of Santa Cruz the average of emerged *Culex* spp. was 27%. In most samples, *Culex* spp. were found in association with *Anopheles gambiae* s.l. Giles, 1902 and *A. pretoriensis* (Theobald, 1903).

While identifying the collected mosquitoes, we found specimens with a median pale ring in the proboscis. Using the key of Ribeiro & Ramos (1995), 136 specimens were identified as *C. tritaeniorhynchus*. The identification of 4th instar larvae confirmed this. Identification was also

confirmed by the morphology of the male genitalia (Fig. 3) (cf. Ribeiro 1966, Harbach 1988).

ADULTS Pale ring of female proboscis extending just to the base, absence of low mesepimeral setae, absence of pale scales on femora; scutum dark brown with pale scales almost completely confined to near scutellum; abdominal tergites with basal pale bands; wings without pale disseminated scales.

LARVAE Mentum with well developed teeth; comb of the abdominal segment VIII composed entirely by typical scales without spine or spiniform scales in number between 30-45; syphon light brown to nearly colorless; pecten of the syphon simple at tip; subventral tufts of the syphon smaller than the diameter, usually with 3-4 branches, rarely 2; head setae 5, 3-branched; pre-clipeal spines brown and strong; anal brush without median tufts.



Fig. 3. Left: coxite of male terminalia of *Culex tritaeniorhynchus* from Santa Cruz, Santiago, Cape Verde Islands (magnification x1000). Right: Basal parts (frontal view) of the male terminalia of *Culex tritaeniorhynchus* from Santa Cruz, Santiago, Cape Verde Islands (magnification x1000).

Species	Collecting method	Praia	Santa Cruz	Santa Catarina	São Miguel	Tarrafal
<i>Anopheles pretoriensis</i>	immature stages	894	241	60	250	30
<i>Anopheles gambiae</i>		12	36	5	45	21
<i>Culex pipiens</i>		21	86	8	15	41
<i>Aedes aegypti</i>		89	0	0	0	43
<i>Culex tigripes</i>		0	4	0	7	5
<i>Culex tritaeniorhynchus</i>		0	136	0	0	0
<i>Culex ethiopicus</i>		0	0	0	3	0
<i>Aedes caspius</i>		0	0	0	10	1
Subtotal		1016	503	73	330	141
<i>Culex pipiens</i>	spray catches	11	53	0	54	0
Subtotal		11	53	0	54	0
<i>Culex pipiens</i>	CDC in bedroom	7	1	0	0	0
Subtotal		7	1	0	0	0
<i>Anopheles pretoriensis</i>	CDC in cowshed	0	0	0	0	2
<i>Culex pipiens</i>		0	0	0	0	1
Subtotal		0	0	0	0	3
TOTAL		1034	557	72	584	144

Table 1. Distribution of mosquito specimens by collecting method in the five districts of Santiago Island prospected.

DISCUSSION

Culex tritaeniorhynchus, a mosquito with a cosmopolitan distribution, is considered a primary vector of JEV (Flavivirus: Flaviviridae) (Self *et al.* 1973, Takahashi 1977, van den Hurk *et al.* 2009), a disease that mainly affects children in the Asian region, with a lethality rate of around 30% (Nett *et al.* 2009). It is also a vector of WNV (Flavivirus: Flaviviridae), considered an emerging vector-borne disease in some parts of the world due to the increased circulation of people and trade goods as a consequence of globalization, while migratory birds may also play a role (Kanojia & Geevarghese 2004, Medlock *et al.* 2005), as well as SINV (Alfavirus: Togaviridae) and the Rift Valley fever virus (Plebovirus: Bunyaviridae) (Kheir *et al.* 2010). The occurrence of *Culex tritaeniorhynchus* in the Cape Verde Islands prompts the need for focused

follow-up investigations in order to evaluate its possible implication in the transmission of vector-borne pathogens in the archipelago. The presence of this species, with its large trophic range (cattle, pigs, birds, humans) (Bram 1967) and as a potential transmitter of Japanese encephalitis, may introduce yet another vector-borne emerging disease and calls for close monitoring. Therefore, practical measures for its surveillance and control, including pathogen isolation, must be taken. At the same time, investigations on other arboviruses, such as WNV, Sindbis and Valley Rift fever virus, transmitted by *Culex pipiens pipiens* L., 1758 and *Culex perexiguus* Theobald, 1903 (Jupp *et al.* 1986, Turell *et al.* 1996, 2002, Jupp 2001, Orshan *et al.* 2008), species both known to occur in Cape Verde, should not be neglected.

ACKNOWLEDGEMENTS

The authors wish to thank the Ministry of Health of Cape Verde for the opportunity to participate in the study and the WHO Office in Cape Verde for support. Thanks also to the delegates, technicians and sanitary agents from the health districts investigated for their assistance in field work, to the Centro Nacional de Desenvolvimento

Sanitário for providing working space and conditions, to the inhabitants of the different localities for their help and cooperation and to Marga Miquel for her help in constructing Fig. 1. We also thank two anonymous reviewers for their helpful comments on earlier drafts of the manuscript.

REFERENCES

- Alves, J., B. Gomes, R. Rodrigues, J. Silva, A.P. Arez, J. Pinto & C.A. Sousa, 2010. Mosquito fauna on the Cape Verde Islands (West Africa): an update on species distribution and a new finding. *Journal of Vector Ecology* 35: 307-312.
- Bram, R.A., 1967. Contributions to the mosquito fauna of South Asia. II. The genus *Culex* in Thailand (Diptera: Culicidae). *Contributions of the American Entomological Institute* 2: 1-296.
- Cambournac, F.J.C., M.C. Oliveira, A. Correia, M.A. Coutinho, J. Torinho & A.B. Soares, 1984. *Culex (Lutzia) tigripes* (Grandpré); mais uma espécie nova para Cabo Verde. *Anais do Instituto de Higiene e Medicina Tropical* 10: 41-46.
- Harbach, R.E., 1988. The mosquitoes of the subgenus *Culex* in southwestern Asia and Egypt (Diptera: Culicidae). *Contributions of the American Entomological Institute* 24: 1-240.
- Jupp, P.G., 2001. The ecology of the West Nile Virus in South Africa and the occurrence of outbreaks in humans. *Annals of the New York Academy of Sciences* 951: 143-152.
- Jupp, P.G., B.M. McIntosh & N.K. Blackburn, 1986. Experimental assessment of the vector competence of *Culex (Culex) neavei* Theobald with West Nile and Sindbis viruses in South Africa. *Transactions of the Royal Society of Tropical Medicine and Hygiene* 80: 226-230.
- Kanojia, P.C. & G. Geevarghese, 2004. First report on high-degree endophilism in *Culex tritaeniorhynchus* (Diptera: Culicidae) in an area endemic for Japanese encephalitis. *Journal of Medical Entomology* 41: 994-996.
- Kheir, S.M., A.M. Alahmed, M.A. Al Kuriji & S.F. Zubyani, 2010. Distribution and seasonal activity of mosquitoes in al Madinah Al

- Munwwrah, Saudi Arabia. *Journal of the Egyptian Society of Parasitology* 40: 215-227.
- Medlock, J.M., K.R. Snow & S. Leach, 2005. Potential transmission of West Nile virus in the British Isles: an ecological review of candidate mosquito bridge vectors. *Medical and Veterinary Entomology* 19: 2-21.
- Nett, R.J., G.L. Campbell & W.K. Reisen, 2009. Potential for the emergence of Japanese encephalitis virus in California. *Vector-Borne Zoonotic Diseases* 9: 511-517.
- Orshan, L., H. Bin, H. Schnur, A. Kaufman, A. Valinsky, L. Shulman, L. Weiss, E. Mendelson & H. Pener, 2008. Mosquito vectors of West Nile Fever in Israel. *Journal of Medical Entomology* 45: 939-947.
- Ribeiro, H., 1966. Research on the mosquitoes of Angola (Diptera: Culicidae). II - Some new culicine records. *Anais do Instituto de Medicina Tropical* 23: 163-166.
- Ribeiro, H. & H.C. Ramos, 1995. Guia ilustrado para a identificação dos mosquitos de Angola. *Boletim da Sociedade Portuguesa de Entomologia*, Suplemento 4, 287 pp.
- Ribeiro, H., H.C. Ramos, R.A. Capela & C.A. Pires, 1980. Os mosquitos de Cabo Verde (Diptera: Culicidae). Sistemática, distribuição, bioecologia e importância médica. *Estudos, Ensaios e Documentos (Junta de Investigações do Ultramar)* 135: 1-141.
- Self, L.S., H.K. Shin, K.H. Kim, K.W. Lee, C.Y. Chow & H.K. Hong, 1973. Ecological studies on *Culex tritaeniorhynchus* as a vector of Japanese encephalitis. *Bulletin of the World Health Organization* 49: 42-47.
- Takahashi, M., 1977. The effects of environmental and physiological conditions of *Culex tritaeniorhynchus* on the pattern of transmission of Japanese encephalitis virus. *Journal of Medical Entomology* 13: 275-284.
- Turell, M.J., S.M. Presley, A.M. Gad, S.E. Cope, D.J. Dohm, J.C. Morrill & R.R. Arthur, 1996. Vector competence of Egyptian mosquitoes for Rift Valley fever virus. *American Journal of Tropical Medicine and Hygiene* 54: 136-139.
- Turell, M.J., J.C. Morrill, C.A. Rossi, A.M. Gad, S.E. Cope, T.L. Clements, R.R. Arthur, L.P. Wasieloski, D.J. Dohm, D. Nash, M.M. Hassan, A.N. Hassan, Z.S. Morsy & S.M. Presley, 2002. Isolation of West Nile and Sindbis viruses from mosquitoes collected in the Nile Valley of Egypt during an outbreak of Rift Valley Fever. *Journal of Medical Entomology* 39: 248-250.
- van den Hurk, A.F., S.A. Ritchie & J.S. Mackenzie, 2009. Ecology and geographical expansion of Japanese Encephalitis Virus. *Annual Review of Entomology* 54: 17-35.
- Wills, W.M., W.L. Jakob, D.J.B. Farancy, R.E. Oerthey, E. Anami, C.H. Callsher & T.P. Monath, 1985. Sindbis virus isolations from Saudi Arabian mosquitoes. *Transactions of the Royal Society of Tropical Medicine and Hygiene* 79: 63-66.

Received 13 March 2013

Revision received 15 May 2014

Accepted 25 August 2014