


## Supplementary Figure 6



**Uncropped silver-stained gel and Western blot of figure 4D.** The nitrocellulose membrane was scanned by a GS-710 Imaging Densitometer (Bio-Rad) and for image processing the Quantity One 4.2.1 software (Bio-Rad) was used. The general settings were as follows:

Left panel; silver stained gel (A):


- Application: Gel
- Filter: Silver stain
- Resolution: 42,3 \* 42,3 microns =High for small gels with tiny features
- 

Transform operations of the silver stained gel: high: 4000; low: 1486; gamma: 4.00

Right panel, nitrocellulose membrane (B):

- Application: Blot; HRP-Substrate (DAB)
- Filter: Green
- Resolution: 42,3 \* 42,3 microns =High for small gels with tiny features

Transform operations: high: 4095; low; gamma: 1.00



**Uncropped gel and Western blot of figure 4E.** Nitrocellulose membranes were scanned by a GS-710 Imaging Densitometer (Bio-Rad) and for image processing the Quantity One 4.2.1 software (Bio-Rad) was used. The general settings were as follows:

- Application: Blot; HRP-Substrate (DAB)
- Filter: Green
- Resolution: 42,3 \* 42,3 microns =High for small gels with tiny features

Transform operations: high: 4095; low; gamma: 1.00