

Table S1. Median (IQR) intra-patient TE, pSWE, APRI, FIB-4 and Fibrotest values changes at different time points relative in chronic HBV patients with at least two determinations of non-invasive methods over time.

Time Point	Patients	TE (kPa)	pSWE (m/s)		APRI		FIB-4		Fibrotest		
FU1	<i>n</i> = 255	-0.10	<i>p</i> = 0.82	0.00	<i>p</i> = 0.65	+0.01	<i>p</i> = 0.67	+0.05	<i>p</i> = 0.00	+0.01	<i>p</i> = 0.16
FU2	<i>n</i> = 156	-0.01	<i>p</i> = 0.88	-0.02	<i>p</i> = 0.62	-0.01	<i>p</i> = 0.82	-0.01	<i>p</i> = 0.01	-0.03	<i>p</i> = 0.26
FU3	<i>n</i> = 107	-0.06	<i>p</i> = 0.91	-0.01	<i>p</i> = 0.66	-0.01	<i>p</i> = 0.73	+0.05	<i>p</i> = 0.05	-0.02	<i>p</i> = 0.17
FU4	<i>n</i> = 71	-0.20	<i>p</i> = 0.72	0.00	<i>p</i> = 0.85	-0.01	<i>p</i> = 1.0	+0.02	<i>p</i> = 0.12	-0.03	<i>p</i> = 0.16
FU5	<i>n</i> = 30	-0.09	<i>p</i> = 0.77	-0.07	<i>p</i> = 0.09	-0.03	<i>p</i> = 0.03	+0.02	<i>p</i> = 0.22	+0.03	<i>p</i> = 0.00
FU6	<i>n</i> = 6	-0.51	<i>p</i> = 0.27	-0.04	<i>p</i> = 0.05	-0.05	<i>p</i> = 0.00	-0.04	<i>p</i> = 0.30	+0.01	<i>p</i> = 0.00

APRI, aspartate to platelet ratio index; FIB-4, fibrosis index based on four factors; FU, Follow up; pSWE, point shearwave elastography; TE, transient elastography.