

## Codebook Stangier et al. (2021) - Metta-based Therapy

| Variable name/name component | Meaning                                      | Values / Range  |
|------------------------------|--|---|
|                              |  |   |
| PR                           | pre/baseline measures                        | -   |
| IM                           | intermediate measures                        | -   |
| PO                           | post measures                                | -   |
| FU                           | 6-month-follow-up measures                   | -   |
| PART_NO                      | Participant number                           | -   |
| GROUP                        | Participant group                            | 0 = Metta-based Therapy<br>1 = Wait-list control group              |
| COMPL_DO                     | Completers and drop-outs                     | 0 = completers<br>1 = drop-outs                                     |
| ANTDM_PR                     | antidepressant medication PR                 | 0 = no medication<br>1 = antidepressant medication at baseline      |
| ANTDM_PO                     | antidepressant medication PO                 | 0 = no medication<br>1 = antidepressant medication at post-measures |
| QIDS_RESP_PO                 | response at the QIDS at post-measures        | 0 = no response<br>1 = response                                     |
| QIDS_RESP_FU                 | response at the QIDS at 6-month-follow-up    | 0 = no response<br>1 = response                                     |
| BDI_RESP_PO                  | response at the BDI-II at post-measures      | 0 = no response<br>1 = response                                     |
| BDI_RESP_FU                  | response at the QIDS at 6-month-follow-up    | 0 = no response<br>1 = response                                     |
| CHANGE_ANTDM_PR_PO           | Change in antidepressive medication pre-post | 0 = no change<br>1 = discontinued                                   |

|          |  |  |
|----------|--|--|
|          |  | 2 = reduced<br>3 = increased<br>4 = new drug   |
| CHH_TR   | Childhood trauma   | 0 = no scale of the CTQ has severe or extreme values<br>1 = At least 1 scale of the CTQ has severe or extreme values |
| CTQ_TOT  | Childhood Trauma Questionnaire - total score               | 25-125   |
| QIDS_TOT | Quick Inventory of Depressive Symptomatology - total score | 0-27   |
| BDI_TOT  | Beck Depression Inventory II - total score                 | 0-63   |
| BADS_TOT | Behavioral Activation for Depression Scale - total score   | 0-150  |
| CLS_MEAN | Compassionate Love Scale - mean score                      | 1-7  |
| FFMQ_TOT | Five Facet Mindfulness Questionnaire - total score         | 39-195   |
| RSQ_SYM  | Response Styles Questionnaire - symptom-focused rumination | 8-32   |
| SASS_TOT | Social Adaptation Self-evaluation Scale - total score      | 0-60   |
| OFD_TOT  | Operationalized Skills Assessment - total score            | 25-125   |
|          |  |  |