

Oligonucleotides suppress PKB/Akt and act as superinductors of apoptosis in human keratinocytes

Stefan Kippenberger^{1,*}, Jutta Müller¹, Maike Schultz¹, Annette Dorn¹,
Andreas Bock², Hüseyin Aygün³, Diamant Thaçi¹, Matthias Hofmann¹,
Roland Kaufmann¹ and August Bernd¹

¹Department of Dermatology and Venerology, J.W. Goethe-University, 60590 Frankfurt/Main, ²Henkel AG & Co. KGaA, Henkelstr. 67, 40191 Düsseldorf and ³BioSpring GmbH, Alt Fechenheim 34, 60386 Frankfurt, Germany

Received January 28, 2009; Revised and Accepted April 3, 2009

ABSTRACT

DNA oligonucleotides (ODN) applied to an organism are known to modulate the innate and adaptive immune system. Previous studies showed that a CpG-containing ODN (CpG-1-PTO) and interestingly, also a non-CpG-containing ODN (nCpG-5-PTO) suppress inflammatory markers in skin. In the present study it was investigated whether these molecules also influence cell apoptosis. Here we show that CpG-1-PTO, nCpG-5-PTO, and also natural DNA suppress the phosphorylation of PKB/Akt in a cell-type-specific manner. Interestingly, only epithelial cells of the skin (normal human keratinocytes, HaCaT and A-431) show a suppression of PKB/Akt. This suppressive effect depends from ODN lengths, sequence and backbone. Moreover, it was found that TGF α -induced levels of PKB/Akt and EGFR were suppressed by the ODN tested. We hypothesize that this suppression might facilitate programmed cell death. By testing this hypothesis we found an increase of apoptosis markers (caspase 3/7, 8, 9, cytosolic cytochrome c, histone associated DNA fragments, apoptotic bodies) when cells were treated with ODN in combination with low doses of staurosporin, a well-known pro-apoptotic stimulus. In summary the present data demonstrate DNA as a modulator of apoptosis which specifically targets skin epithelial cells.

INTRODUCTION

Redundancy seems to be a basic principle of living matter providing an economic handling with limited resources.

An example of this is the desoxyribonucleic acid (DNA) that was for a long time only considered as carrier of genetic information. It is more than one hundred years ago since William Coley found that a preparation of heat inactivated streptococci induces some immunological response in cancer patients (1,2). It took until 1984 since bacterial DNA itself was identified to convey this effect (3). This observation demonstrated that information within the DNA molecule is not restricted to the genetic code. Later it was found that a specific sequence, a motif displaying an unmethylated cytidine-phosphate-guanosine (CpG) is able to activate the immune system (4). The fact that the CpG motif has a much lower incidence in human DNA than predicted by random base utilization and the increased amount of methylated cytidine residues makes this motif an ideal pathogen-associated molecular pattern (PAMP) recognized by the endosomal toll-like receptor-9 (5–8). Clinically, CpG and also some non-CpG oligonucleotides (ODN) are currently tested as immunomodulators for immune protection, allergic response, vaccination booster and in antitumor therapy (9). For these applications much efforts have been placed on the development of structurally modified ODN. Particularly, the substitution of one of the nonbridging oxygen atoms bound to phosphorus by sulfur is very commonly used. These phosphorothioates offer an increased serum stability and resistance against nucleases (10). Moreover, phosphorothioates can form mRNA–DNA duplexes which elicit RNase H activity triggering the degradation of mRNA (11). This mechanism is considered to play an important role in most antisense-based applications.

Both the presence of CpG-motifs and the modification of the DNA-backbone with phosphorothioates are described to stimulate pro-inflammatory responses (4,12–14). Therefore, it was a surprise that a CpG-ODN and a non-CpG-ODN both with phosphorothioate backbones suppressed pro-inflammatory markers in skin

*To whom correspondence should be addressed. Tel: +49 69 6301 7734; Fax: +49 69 6301 6466; Email: kippenberger@em.uni-frankfurt.de

keratinocytes and attenuated the cutaneous delayed-type hypersensitivity response in a mouse model (15). These findings speak for a tissue-specific mode of action.

As the duration of inflammation is frequently regulated by apoptosis of the effector cells (16), we investigated the impact of CpG-ODN, and non-CpG-ODN with and without phosphorothiate backbones on apoptosis in a variety of epithelial and non-epithelial cells. Of note, it was found that epithelial cells of the skin namely, HaCaT cells, A-431 cells and normal human keratinocytes show PKB/Akt suppression and facilitated apoptosis in response to ODN. This effect was modulated by ODN backbone, sequence and length. These findings suggest that ODN can act as modulators of skin homeostasis.

MATERIALS AND METHODS

Reagents

ODN with phosphorothioate backbone were synthesized and purified by BioSpring GmbH (Frankfurt/Main, Germany), reconstituted in water and stored at -20°C . ODN were given to the cells at the indicated concentration without DNA complexing reagents. Hybridization of CpG-1-PTO and its reverse strand CpG-1-PTO-rev was performed as described (15). DNA from *Escherichia coli*, *Clostridium perfringens* and DNA from Salmon sperm were purchased from ICN Pharmaceuticals (Heidelberg, Germany) and dissolved in water. Staurosporin (STS) was purchased from Sigma (Taufkirchen, Germany). TGF α was purchased from PeproTech (Hamburg, Germany).

Cell culture

Spontaneously immortalized human keratinocyte cell line (HaCaT) (a generous gift by Prof. Fusenig, German Cancer Research Institute, Heidelberg, Germany) was cultured in carbonate buffered Hank's medium with 5% fetal calf serum, 0.35 g/l glutamine (Invitrogen, Paisley, UK) and 1% penicillin-streptomycin solution (Biochrom KG, Berlin, Germany) at 37°C in 5% CO_2 atmosphere. The human epidermoid carcinoma cell line A-431, the human epithelial kidney line HEK-293, the kidney fibroblast cell line Cos-7 (all purchased from the American Type Culture Collection, ATCC), the human sebocyte cell line SZ95 (kindly provided by Christos Zouboulis, Universitätsklinikum Benjamin Franklin, Berlin) were cultured in Dulbecco's modified Eagle's medium (DMEM, Gibco, Karlsruhe, Germany) containing 1% penicillin-streptomycin solution and 10% FCS. Normal human keratinocytes (NHK) and fibroblasts were isolated from skin specimen of cosmetic surgeries. NHK were propagated in serum-free keratinocyte medium (Gibco) and fibroblasts were held in RPMI 1640 medium (Biochrom, Berlin, Germany) with 10% FCS and 1% penicillin-streptomycin solution. Likewise, G-361 melanoma cells (ATCC), derived from a 31-year-old Caucasian male were cultured under the same conditions as fibroblasts. Human umbilical vein endothelial cells (HUVEC) were purchased from Promocell (Heidelberg, Germany) and propagated in endothelial growth medium (EGM,

Lonza, Wuppertal, Germany). The medium was renewed twice a week. All experiments were done in agreement with the local ethics commission.

Plasmids and transfection

The human dominant-negative TLR-9 plasmid was kindly provided by Hermann Wagner (Technische Universität München, München, Germany) (17). The dominant-negative flag-tagged MyD88 was kindly provided from Tularik (San Francisco, USA) (17). Empty pcDNA-3 served as control vector. Transient transfection was performed using A-431 cells plated in 6-well multidishes at a confluence of 70% using Lipofectamine 2000 reagent (Invitrogen) according to the manufacturers' instructions. The following day, cells were treated with $4\mu\text{M}$ CpG-1-PTO or nCpG-5-PTO. After 30 min proteins were extracted as described.

Mechanical stimulation

Stretching of *in vitro* cultured cells was performed as described previously (18). Briefly, silicone elastomer MED-4011 (Armando Medizin Technik, Düsseldorf, Germany) was stirred, poured into preformed teflon matrices and allowed to polymerize. For allowing cell attachment silicone dishes were treated with 5.7% KOH in methanol for 5 min in order to neutralize the polymerization-derived HCl. After washing with double-distilled water, silicone dishes were coated with 2% arginine for 2 h to facilitate cell attachment (19) and afterwards rinsed with PBS. Subsequently, the dishes were incubated with fetal calf serum for 2 h. After withdrawal of the serum, cells were plated in flexible silicone chambers and incubated for 24 h under regular conditions. Prior to the application of mechanical stretch, the cells were held for 24 h under serum-free conditions. For the last hour $4\mu\text{M}$ ODN were added to the cells. Then the silicone chambers were extended to 10% for 5 min. After stretch stimulation, protein samples were prepared at the indicated time intervals.

Immunoblotting

For detection of PKB/Akt and caspase 3 and 8 (antibodies from Cell Signaling Technology, Frankfurt, Germany), cells were lysed in $100\mu\text{l}$ SDS sample buffer [62.5 mM Tris-HCl (pH 6.8), 2% SDS, 10% glycerol, 50 mM DTT, 0.1% bromophenol blue], sonicated and boiled for 5 min, and separated on SDS-polyacrylamide gels. For detection of EGFR (Cell Signaling Technology), cells were scraped into lysis buffer [20 mM Tris (pH 7.4), 150 mM NaCl, 1 mM EDTA, 1 mM EGTA, 1% Triton, 2.5 mM sodium pyrophosphate, 1 mM β -glycerolphosphate, 1 mM Na_3VO_4 , $1\mu\text{g/ml}$ leupeptin, 1 mM PMSF], sonicated and centrifuged. Protein concentration of the supernatant was determined (Biorad DC Protein Assay Kit, Biorad, München, Germany) and standardized using bovine serum albumin. Twenty micrograms of protein were mixed with SDS sample buffer and run on SDS-polyacrylamide gels. Consecutively, proteins were immunoblotted to a PVDF membrane. The membrane was blocked in blocking

buffer [TBS (pH 7.6), 0.1% Tween-20, 5% nonfat dry milk] for at least 3 h at 4°C followed by incubation with the primary antibody in TBS (pH 7.6), 0.05% Tween-20 and 5% BSA. The bound primary antibodies were detected using anti-mouse IgG-horseradish peroxidase conjugate and visualized with the ECL detection system (Amersham).

Cytosolic cytochrome *c*

In order to detect the amount of cytosolic cytochrome *c*, cells were fractionated as described (20). Briefly, 2×10^6 cells were trypsinated, washed and solubilized in 250 μ l phosphate buffered saline (PBS). A mild lysis of the cell membrane was performed by adding 250 μ l digitonin-solution (80 μ g/ml sucrose solution, 500 mM). After vigorously vortexing for 10 s, cell extracts were centrifuged (1 min, 14 000g, 4°C) and the protein content of the supernatant was determined as described under *Immunoblotting*. Thirty micrograms protein of the supernatant was applied to a commercial human cytochrome *c* immunoassay (R&D Systems, Wiesbaden, Germany).

Histone-associated DNA fragments

Apoptosis was quantified on the basis of cytoplasmic histone-associated DNA fragments using the Cell Death Detection ELISA (Roche, Mannheim, Germany) according to the manufacturer's manual. In brief, cells were cultured in microwell plates (2×10^4 cells per 0.33 cm²) and treated with increasing amounts of staurosporin (STS) in combination with different ODN. After 24 h cytosolic fraction (200g supernatant) was used as antigen source in a sandwich enzyme-linked immunosorbent assay with primary anti-histone antibody coated to a microtiter plate and secondary anti-DNA antibody coupled to peroxidase. Optical density was measured at

530 nm in an ELISA reader (MR 5000, Dynatech, Guernsey, UK).

Caspase 3/7 and 9 activity

Activity of caspase 3/7 and 9 was quantitatively assessed by using commercial assays (Caspase-Glo 3/7, Caspase-Glo 9, Promega, Mannheim, Germany). Cells were seeded in microplate wells and treated with staurosporin (STS) and ODN as aforementioned. After 24 h cells were lysed and luminogenic substrates specific for the different caspase species were added. Light emission was measured in a luminometer (Berthold, Bad Wildbad, Germany).

Statistical analysis

The grey levels of western blot bands were measured using ImageJ (Bethesda, USA). Data from western blots, histone associated DNA fragments, cytosolic cytochrome *c* and caspase 3/7 and 9 activity are given as means \pm SD. Each column represents at least three independent experiments. Statistical analysis was performed using the Wilcoxon–Mann–Whitney U-test (BIAS, Frankfurt, Germany). Differences were considered significant at $P < 0.05$ indicated by asterisks. Figure legends indicate the data columns which were compared.

RESULTS

CpG-1-PTO and non-CpG-5-PTO suppress PKB/Akt

HaCaT keratinocytes were exposed to increasing amounts (0.5, 1, 2, 4 μ M) of CpG-1-PTO or nCpG-5-PTO for 30 min. Protein extracts were separated by SDS–PAGE and probed against phospho-PKB/Akt and total PKB/Akt for loading control, respectively. Figure 1A shows a dose-dependent suppression of basal PKB/Akt activation




Figure 1. Oligonucleotides suppress basal PKB/Akt activation. Protein extracts derived from HaCaT cells exposed to CpG-1-PTO and nCpG-5-PTO were separated by SDS–PAGE. The blotted proteins were probed with anti-PKB/Akt-Ser473, anti-PKB/Akt-Thr308 and phospho-non-specific anti-PKB/Akt. (A) Concentration dependent effect of CpG-1-PTO and nCpG-5-PTO. Cells were exposed to 0.5, 1, 2, 4 μ M of CpG-1-PTO and nCpG-5-PTO respectively for 30 min. (B) Statistical analysis of (A). Each column represents at least three independent experiments; standard deviations are indicated. Data were compared to untreated controls. * $P < 0.05$; solid bars, serine; white bars, threonine.




Figure 2. Time-dependent suppression of basal PKB/Akt by CpG-1-PTO and nCpG-5-PTO. HaCaT cells were exposed for the indicated times to 4 μ M (A) CpG-1-PTO and (B) nCpG-5-PTO. (C) and (D) represent the statistical analysis of (A) and (B), respectively. Each column represents at least three independent experiments; standard deviations are indicated. Data were compared to untreated controls. * $P < 0.05$; solid bars, serine; white bars, threonine.

at both phosphorylation sites. Already a concentration of 0.5 μ M nCpG-5-PTO led to a significant suppression of PKB/Akt (see statistical analysis in Figure 1B). In order to test the dynamic of this suppression HaCaT cells were exposed to 4 μ M CpG-1-PTO (Figure 2A) and nCpG-5-PTO (Figure 2B), respectively for different times ranging from 5 min to 24 h. Statistical analysis in Figure 2C and D shows a rapid and robust downregulation by both oligos. Of note, the suppression of PKB/Akt phosphorylation lasted for at least 24 h. CpG-1-PTO and nCpG-5-PTO have a length of 20 nucleotides. In order to test if the lengths of the ODN molecules have an impact on PKB/Akt suppression different deletion mutants were tested (Table 1, Figure 3A–D). Deletion of flanking nucleotides from both ends of CpG-1-PTO reversed the

PKB/Akt suppression in a length dependent manner. The hexamer CpG-6-PTO which still bears a complete CpG-motif shows no suppressive effect on PKB/Akt at 4 μ M after incubation for 30 min. Vice versa nucleotide deletion of nCpG-5-PTO showed also a length dependent suppression of PKB/Akt (Figure 3B and D). ODN molecules with lengths from 20 down to 16 nucleotides showed a significant suppression of basal PKB/Akt phosphorylation.

Natural DNA suppress PKB/Akt

The aforementioned data show that both, CpG- and nCpG-ODN suppress basal PKB/Akt phosphorylation. In order to test if DNA in general conveys this effect,

Table 1. Oligonucleotides

CpG-1-PTO	5'-TCC ATG <u>ACG</u> TTC CTG <u>ACG</u> TT-3'
CpG-18-PTO	5'-TCC ATG <u>ACG</u> TTC CTG <u>ACG</u> -3'
CpG-16-PTO	5'-TCC ATG <u>ACG</u> TTC CTG A-3'
CpG-14-PTO	5'-CC ATG <u>ACG</u> TTC CTG-3'
CpG-12-PTO	5'-CAT GAC <u>GTT</u> CCT-3'
CpG-6-PTO	5'-GAC <u>GTT</u> -3'
nCpG-5-PTO	5'-CCC <u>CCC</u> CCC CCC CCC CC-3'
nCpG-18-PTO	5'-CCC CCC CCC CCC CCC CC-3'
nCpG-16-PTO	5'-CCC CCC CCC CCC CC-3'
nCpG-14-PTO	5'-CCC CCC CCC CC-3'
nCpG-12-PTO	5'-CCC CCC CC-3'
nCpG-6-PTO	5'-CCC CC-3'
Scramb	5'-CTC TAG GAC TCT CTG GAC TT-3'
CpG-1-PTO-rev	5'-AAC GTC AGG AAC <u>GTC</u> ATG GA-3'
nCpG-5-PDE	5'-ccc ccc ccc ccc cc-3'
CpG-1-PDE	5'-tcc atg <u>acg</u> ttc ctg <u>acg</u> tt-3'

Phosphorothioates (PTO) in capital letters, phosphodiester (PDE) in small letters. CpG-motifs are underlined.

we treated HaCaT keratinocytes with different natural occurring DNA derived from prokaryotes and eukaryotes. Figure 4A and B show that DNA from *Escherichia coli*, *Clostridium perfringens* and DNA derived from Salmon sperm suppress the basal phosphorylation of PKB/Akt. The observed effects are distinct but not as strong as for synthetic CpG-1-PTO. Furthermore, the natural DNA species offered no significant toxic effect on HaCaT cell in the tested concentrations as detected by the release of lactate dehydrogenase (LDH) (Figure S1, Supplementary Data). Figure 4C shows ethidiumbromide stained DNA samples separated by agarose gel electrophoresis. DNA derived from *Escherichia coli* gives a smear over the whole range and a slightly condensed section at approximately 10 kbp. This accumulation was more prominent for DNA derived from *Clostridium perfringens* and Salmon sperm. In order to test if shearing of DNA has an




Figure 3. Lengths dependent suppression of basal PKB/Akt activation by CpG-PTO and nCpG-PTO. HaCaT cells were treated with different deletion mutants (4 μ M) of CpG-1-PTO and nCpG-5-PTO for 30 min. Then protein extracts were utilized for detection of PKB/Akt activation. (A) Deletion mutants of CpG-1-PTO ranging from 6 (CpG-6-PTO) to 20 nucleotides (CpG-1-PTO). (B) Deletion mutants of nCpG-5-PTO ranging from 6 (nCpG-6-PTO) to 20 nucleotides (nCpG-5-PTO). (C) and (D) represent the statistical analysis of (A) and (B), respectively. Each column represents at least three independent experiments; standard deviations are indicated. Data were compared to untreated controls. * $P < 0.05$; solid bars, serine; white bars, threonine.




Figure 4. Natural DNA suppresses basal PKB/Akt activation. (A) HaCaT cells were treated with different concentrations of DNA derived from natural sources for 60 min. Then protein extracts were utilized for detection of PKB/Akt activation. CpG-1-PTO served as positive control. *E. coli*, *Escherichia coli*; *C. perfr.*, *Clostridium perfringens*; Salm. sp., Salmon sperm. (B) represents the statistical analysis of (A). Each column represents at least three independent experiments; standard deviations are indicated. Data were compared to untreated controls. * $P < 0.05$; solid bars, serine; white bars, threonine. (C) Agarose gel electrophoresis with DNA derived from *E. coli*, *C. perfringens* and Salmon sperm.

impact on the observed PKB/Akt suppression preparations of all three DNA species were treated with ultrasound. The effect of this treatment was monitored by agarose gel electrophoresis as aforementioned (Figure S2A, Supplementary Data). No changed effect on PKB/Akt suppression was detected suggesting that the molecular size is not a significant determinant in this case (Figure S2B, Supplementary Data).

PKB/Akt suppression depends from DNA sequence and backbone

In order to test what distinct information conveys the suppressive effect on PKB/Akt different synthetic 20-mer were tested (Figure 5A–C). It was found that both, the sequence and backbone are important for the observed effect. A CpG-1-molecule with phosphodiester bonds (CpG-1-PDE) offered no significant suppressive effect on PKB/Akt, whereas its phosphorothiotate counterpart

shows the already noted strong inhibition. A scrambled sequence of CpG-1-PTO (Scramb) also shows strong inhibition. nCpG-5-PDE, which is a phosphodiester of nCpG-5-PTO, suppresses PKB/Akt in a moderate manner. Furthermore, the effect of a DNA hybrid was tested (Figure 5B). At first it was found that 4 μM of the reverse strand of CpG-1-PTO (CpG-1-PTO-rev) offered a slightly stronger suppressive effect than CpG-1-PTO indicating sequence specificity. This strong effect was maintained for the hybrid of CpG-1-PTO and CpG-1-PTO-rev, indicating that a single strand is sufficient to reduce PKB/Akt phosphorylation significantly.

Cell-specific suppression of PKB/Akt by CpG-1-PTO and nCpG-5-PTO

The aforementioned data were derived from experiments with HaCaT keratinocytes. In order to test if the suppression of PKB/Akt by CpG-1-PTO and nCpG-5-PTO is




Figure 5. Suppression of PKB/Akt depends on DNA-backbone and sequence. HaCaT cells were treated with (A) different concentrations of CpG-1-PTO (with phosphorothioate backbone), CpG-1-PDE (with phosphodiester backbone), a scrambled sequence of CpG-1-PTO (Scramb) and with nCpG-5-PDE a phosphodiester of nCpG-5-PTO for 30 min. Then protein extracts were utilized for detection of PKB/Akt activation. (B) 4 μM CpG-1-PTO, CpG-1-PTO-rev, the reverse strand, and the hybrid of both strands were given to the cells for 30 min. Then proteins were analyzed as aforementioned. (C) represents the statistical analysis of (A) and (B), respectively. Each column represents at least three independent experiments; standard deviations are indicated. Data were compared to untreated controls. * $P < 0.05$; solid bars, serine; white bars, threonine.

cell specific also other cell species from different tissues were tested. Figure 6A shows the effect of CpG-1-PTO or nCpG-5-PTO on basal PKB/Akt phosphorylation on epithelial cells (A-431, HaCaT, NHK, HEK293, SZ95), fibroblasts (skin fibroblasts, Cos-7), melanoma cells (G361) and endothelial cells (HUVEC) after incubation for 60 min. The western blot analysis demonstrates a suppression of PKB/Akt phosphorylation in A-431, NHK and HaCaT cell as reference. Sebocytes (SZ95), fibroblasts derived from human skin, melanoma cells (G361), endothelial cells (HUVEC) and fibroblasts from kidney tissue (Cos-7) show no shift in PKB/Akt phosphorylation in response to ODN treatment. In order to test if classical CpG-effects are induced by CpG-1-PTO the murine macrophage cell line Raw264.7 was utilized (Figure 6B). It was found that treatment with 4 μM CpG-1-PTO leads to a time-dependent activation of PKB/Akt. This effect is well described leading to an anti-apoptotic response mediated via TLR-9, the

CpG-receptor (21). A control experiment using nCpG-5-PTO devoid of a classical CpG-motif shows no activation of PKB/Akt in these cells. In sum, the results presented show a heterogenic response to CpG-1-PTO and nCpG-5-PTO depending on the cell species used. Of note, in epithelial cells derived from skin (HaCaT, NHK) and squamous epithelium (A-431) the ODN tested offer a clear suppression of basal PKB/Akt phosphorylation at both phosphorylation sites.

CpG-1-PTO and nCpG-5-PTO suppress induced levels of PKB/Akt and EGFR

The aforementioned data show that CpG-1-PTO and nCpG-5-PTO suppress the basal phosphorylation of PKB/Akt. Figure 7A shows a concentration dependent increase of PKB/Akt phosphorylation in response to TGFα. This phosphorylation was attenuated by a preincubation with CpG-1-PTO or nCpG-5-PTO at a concentration of 4 μM for 30 min. As TGFα induces




Figure 6. Cell-specific suppression of PKB/Akt by CpG-1-PTO and nCpG-5-PTO. (A) A-431, HaCaT (human keratinocyte cell line), NHK (normal human keratinocytes), SZ95 (human sebocyte line), Fib (human skin fibroblasts), G-361 (human melanoma line), Cos-7 (monkey kidney fibroblast line), HUVEC (human endothelial cells) and HEK-293 (human kidney epithelial line) were treated with 4 μ M CpG-1-PTO or nCpG-5-PTO for 60 min. Then protein extracts were utilized for detection of PKB/Akt activation. (B) The murine macrophage cell line Raw267.7 known to activate PKB/Akt in response to CpG-DNA was treated with 4 μ M CpG-1-PTO or nCpG-5-PTO for the indicated time intervals. Proteins were extracted and analyzed as aforementioned. The blots show representative results.

PKB/Akt phosphorylation via EGFR we investigated the effect of ODN on EGFR activation (Figure 7B). As expected, TGF α causes a dose dependent phosphorylation of EGFR. Interestingly, a pretreatment with CpG-1-PTO and particularly nCpG-5-PTO reversed this induction indicating a suppressive effect already at the surface receptor level. A statistical analysis performed for data given in Figure 7A and B shows significant reductions of PKB/Akt and EGFR in cells treated with TGF α (exemplary tested for 10 ng/ml) in the presence of CpG-1-PTO or nCpG-5-PTO, respectively (Figure 7C and D).

No impact of dominant-negative expression of TLR-9 and MyD88 on PKB/Akt suppression

CpG-DNA is recognized by a specific receptor, the toll-like receptor 9 (8) which mediates its signal via the adaptor protein MyD88 (22). In order to test if the here presented suppression of PKB/Akt by CpG-ODN and also

non-CpG-ODN utilizes this signalling pathway A-431 cells were transfected with dominant-negative constructs for TLR-9 and MyD88 (Figure 8). Data given in Figure 8A and B show that mock transfected cells (pcDNA3) show a similar suppression of PKB/Akt as cells transfected with dominant-negative constructs for TLR-9 (Δ TLR-9) and MyD88 (Δ MyD88). These results demonstrate that the observed effects are rather independent from the TLR-9 signalling pathway.

CpG-1-PTO and nCpG-5-PTO suppress PKB/Akt activation in response to stretch

In the previous experiments ODN suppressed basal levels of PKB/Akt and after stimulation with a prototypical soluble mediator (TGF α). In the following a complete different stimulus, mechanical stretch, was applied to HaCaT cells (see 'Materials and Methods' section). Published data from our group already identified




Figure 7. Partial suppression of TGF α -induced activation of PKB/Akt and EGF-R by CpG-1-PTO and nCpG-5-PTO. HaCaT cells were treated with 4 μ M ODN for 30 min, followed by the addition of TGF α at the indicated concentrations. After 5 min proteins were extracted and analyzed for (A) PKB/Akt and (B) EGF-R activation. (C) and (D) represent the statistical analysis of (A) and (B) in the presence or absence of 10 ng/ml TGF. Each column represents at least three independent experiments; standard deviations are indicated. Data were compared to TGF α controls. * $P < 0.05$; in (C) solid bars, serine; white bars, threonine, in (D) solid bars, EGFR phosphorylation.

mechanical stretch as strong inducer of PKB/Akt phosphorylation (23). Figure 9A shows that application of a stretching stimulus for 5 min induces PKB/Akt phosphorylation at both phosphorylation sites. A preincubation with 4 μ M CpG-1-PTO or nCpG-5-PTO for 30 min suppressed the stretch-induced phosphorylation. Of note, PKB/Akt induction by mechanical stretch shows some variation in the time kinetic resulting in a higher standard deviation (Figure 9B). Nevertheless, treatment with CpG-1-PTO or nCpG-5-PTO suppressed significantly the stretch-induced PKB/Akt levels at all tested time intervals.

CpG-1-PTO and nCpG-5-PTO sensitize for apoptosis

In the present study we have shown that treatment with CpG-1-PTO and nCpG-5-PTO suppresses basal and induced PKB/Akt—one of the key regulators in apoptosis. From this it could be construed that the ODN tested might change the sensitivity of cells against proapoptotic stimulation. This issue was addressed by using a combination of ODN and staurosporin—a well-known inducer of apoptosis. Data given in Figures 10 and 11 show that CpG-1-PTO and nCpG-5-PTO increase the pro-apoptotic effects staurosporin in HaCaT cells. In Figure 10A the

application of staurosporin for 24 h led to a significant increase of cleaved caspase 8 and 3 fragments in a dose dependent manner. When staurosporin and 8 μ M ODN were applied at the same time the appearance of cleaved caspases shifted to lower staurosporine concentrations indicating an accelerated onset of apoptosis. By using a quantitative detection kit for caspase 3/7 activity these results were confirmed: 8 μ M CpG-1-PTO and nCpG-5-PTO significantly amplified the effect of staurosporin (Figure 10B). These results were corroborated by measuring the cytosolic levels of cytochrome *c* (Figure 10C). A treatment with increasing concentrations of staurosporin alone for 14 h led to a concentration dependent increase of cytosolic cytochrome *c* as measured by ELISA. In the presence of 8 μ M CpG-1-PTO the amount of cytosolic cytochrome *c* at 0.5 μ M staurosporin almost doubles. This effect was even more pronounced for 8 μ M nCpG-5-PTO: at 0.1 and 0.5 μ M staurosporin a significant increase of cytochrome *c* was measured. At 1.0 μ M staurosporin the measured levels of cytochrome *c* decreased again indicating advanced apoptosis. On the morphological level treatment with 1 μ M staurosporin for 8 h induced the enhanced formation of apoptotic bodies as visualized by bisbenzimid staining (1 μ g/ml) (Figure 10D). In the




Figure 8. No effect of Δ TLR-9 and Δ MyD88 on PKB/Akt suppression by CpG-1-PTO and nCpG-5-PTO. (A) A-431 cells were transfected with pcDNA3 and dominant-negative constructs of TLR-9 (Δ TLR-9) and MyD88 (Δ MyD88), respectively. After 24h cells were treated with 4 μ M CpG-1-PTO and nCpG-5-PTO for 30 min. Then protein extracts were utilized for detection of PKB/Akt activation. (B) represents the statistical analysis of (A). Each column represents at least three independent experiments; standard deviations are indicated. Data of each set were compared to transfection controls. * $P < 0.05$; solid bars, serine; white bars, threonine.

presence of 8 μ M CpG-1-PTO this effect was considerably enhanced (Figure 10D).

The release of cytochrome *c* from mitochondria marks the onset of the intrinsic apoptosis triggering initiator caspase 9 activation. Quantitative detection of caspase 9 activity showed a significant amplification of the staurosporin effect (Figure 11A). These data are coherent in respect to results shown in Figure 10C documenting the activation of caspase 3/7 which are downstream from caspase 9. A late event in apoptosis is the degradation of DNA which can be quantitatively measure by the formation of histone associated DNA fragments. Figure 11B shows that CpG-1-PTO and nCpG-5-PTO enhance the pro-apoptotic effect of staurosporin while the hexamers

of both ODN (CpG-6-PTO, nCpG-6-PTO) do not indicating that the apoptosis sensitizing effect of ODN depends on the molecule length. Finally, the effect of ODN on apoptosis was tested in the murine macrophage cell line RAW264.7 (Figure 11C). In contrast to skin epithelial cells CpG-1-PTO protect RAW264.7 cell from staurosporin induced apoptosis, an effect already documented (21).

DISCUSSION

In the present study we show that CpG-ODN, nCpG-ODN and also natural DNA suppress PKB/Akt




Figure 9. Stretch-induced PKB/Akt activation is suppressed by CpG-1-PTO and nCpG-5-PTO. (A) HaCaT cells were seeded in flexible silicon dishes, treated with 4 μ M CpG-1-PTO or nCpG-5-PTO for 60 min and then mechanically stretched for 5 min as described under 'Materials and Methods' section. Proteins were extracted after 0, 10 and 30 min and analyzed for PKB/Akt activation. (B) represents the statistical analysis of (A). Each column represents at least three independent experiments; standard deviations are indicated. Data were compared to the referring controls. * $P < 0.05$; solid bars, serine; white bars, threonine.

phosphorylation in epithelial skin cells. This is interesting as a recent study shows that a CpG-ODN from the same subclass (CpG-B) activates PKB/Akt in the mouse macrophage cell line RAW264.7 with an anti-apoptotic effect (21). This experiment was confirmed by us with using CpG-1-PTO. Likewise to this experiment others have found that CpG-ODN delay the onset of apoptosis in neutrophil granulocytes (24). Surprisingly, in our experimental setting using epithelial cells we found that both, CpG-ODN and also non-CpG-ODN sensitize cells for apoptosis. These results support current data showing that ODN trigger unexpected responses in epithelial skin cells and epithelial tissues (15). In this article it was demonstrated that the same ODN as used in the present study offer anti-inflammatory properties.

Besides the huge amount of data documenting the effect of CpG ODN on human cells there is increasing evidence that also some non-CpG ODN have physiological significance (9). Similarly, to CpG ODN also non-CpG ODN can exert an immune stimulatory effect in human leukocytes (25). Interestingly, the mode of stimulation differs between both ODN entities, and in contrast to CpG ODN non-CpG ODN support a Th2-biased response (26,27). Of note, the signaling of certain non-CpG-ODN require the presence of functional TLR-9 (27), which is surprising as previous work suggested that only CpG ODN, but not non-CpG ODN, trigger TLR9-mediated signaling (17,28,29). In contrast our data show that suppression of PKB/Akt in skin keratinocytes by CpG ODN, and non-CpG ODN seems to




Figure 10. CpG-1-PTO and nCpG-5-PTO amplify staurosporin-induced caspase 3, 7, 8 activity, cytochrome *c* liberation and formation of apoptotic bodies. HaCaT cells were treated with $8\mu\text{M}$ CpG-1-PTO or nCpG-5-PTO in the presence of 0.1, 0.5 and $1\mu\text{M}$ staurosporin (STS). (A) After 24 h proteins were extracted and analyzed for degradation of caspase 3 and 8. The blot shows representative results. (B) After 24 h caspase 3/7 activity was assessed using a commercial assay as described under 'Materials and Methods' section. Activity of untreated cells was set to 100%. Each bar represents the mean of four parallel experiments; the standard deviations are indicated ($*P < 0.05$). (C) After 14 h a cytosolic extract was analyzed for cytochrome *c*. Each bar represents the mean of four parallel experiments; the standard deviations are indicated ($*P < 0.05$) (D) Cells were treated with $8\mu\text{M}$ CpG-1-PTO and $1\mu\text{M}$ STS for 8 h. Then cells were fixed and apoptotic bodies were stained using bisbenzimidazole ($1\mu\text{g/ml}$). Experiments were repeated with similar results.

be independent from TLR-9 signaling as displayed by dominant-negative expression of TLR-9 and the adaptor protein MyD88. Instead a dependency from ODN lengths, sequence and backbone was demonstrated. The relevance of these parameters was also shown for immunostimulation by non-CpG ODN (25). Similar to our findings phosphorothioates with lengths longer than 18 nucleotides offer high stimulatory effects on B-cells. Interestingly, these authors found thymidine-rich sequences more effective than sequences with high amounts of cytosine. Our data regarding PKB/Akt suppression show that a sequence solely consisting of cytosines (nCpG-5-PDE) is more effective than CpG-1-PDE with an amount of almost 30% thymidines. Although, ODN with

phosphodiester bondings and also naturally occurring DNA derived from bacteria and salmon sperm have the potential to suppress PKB/Akt the presence of phosphorothioates amplified the efficacy.

The kinase PKB/Akt plays an important role in the suppression of apoptotic cell death (30,31). Proximal from surface receptors, the phosphoinositide 3-OH kinase (PI3K) conveys activation of PKB/Akt via phosphoinositide-dependent kinases (PDK). It has been demonstrated that PDK-1 phosphorylates PKB/Akt at threonine-308 (32), whereas the mechanism of the serine-473 phosphorylation is still under debate (33,34). Treatment with ODN as performed in this study causes suppression at both relevant phosphorylation sites,




Figure 11. CpG-1-PTO and nCpG-5-PTO amplify staurosporin-induced caspase 9 activity and cytoplasmic histone-associated DNA fragments in HaCaT cells—CpG-1-PTO protects RAW264.7 cells from apoptosis. HaCaT cells were treated with 8 μM CpG-1-PTO or nCpG-5-PTO in the presence of 0.1, 0.5 and 1 μM staurosporin (STS). (A) After 24 h caspase 9 activity was assessed using a commercial assay as described under ‘Materials and Methods’ section. Each bar represents the mean of four parallel experiments; the standard deviations are indicated (**P* < 0.05). (B) After 24 h cytoplasmic histone associated DNA fragments were assessed as described under ‘Materials and methods’ section. The effect of 20-mers (CpG-1-PTO, nCpG-5-PTO) was compared to hexamers (CpG-6-PTO, nCpG-6-PTO). Each bar represents the mean of four parallel experiments; the standard deviations are indicated (**P* < 0.05). (C) The murine macrophage cell line Raw267.7 was treated with STS and different amounts of CpG-1-PTO. After 24 h cytoplasmic histone associated DNA fragments were assessed as aforementioned. Each bar represents the mean of four parallel experiments; the standard deviations are indicated (**P* < 0.05).

although the lack of only one of these sites abrogates PKB/Akt signalling (35). PKB/Akt is known to control a plethora of molecules involved in apoptosis (31,36). Interestingly, treatment with ODN in the absence of an additional pro-apoptotic stimulus is not sufficient to trigger apoptosis. Moreover, in the presence of staurosporin the onset of apoptosis becomes facilitated. Our data provide evidence for an activation of the mitochondrial apoptosis machinery as documented by elevated cytochrome *c* levels in response to ODN treatment. Suppression of PKB/Akt leads to activation of the BCL-2 homology-3 domain-only proteins BIM and BAD which in turn mediate the release of cytochrome *c* (37,38). The release of cytochrome *c* from mitochondria marks the onset of the intrinsic apoptosis with subsequent activation of caspase-9 and caspase-3 (39). Furthermore, the activation of caspase-8, the increase of histone associated DNA fragments and the presence of apoptotic bodies indicate a massive induction of apoptosis at different stages.

More distal we found a suppression of the epidermal growth factor receptor (EGFR) by CpG-1-PTO, and particularly nCpG-5-PTO after stimulation with TGF α . The EGFR is characterized as a key element in the downstream signaling to PKB/Akt (40–42). Rockwell *et al.* (43) showed that the polyanionic nature of phosphorothioates enables interaction with proteins of the cell surface. Particularly, a specific inhibition of protein tyrosine receptors including EGFR was found. We speculate that the high amount of EGFR on epithelial cells may contribute to the selective action of ODN presented in this article.

Our findings suggest that ODN and particularly phosphorothioates are interesting pharmacological compounds in the treatment of hyperproliferative skin diseases such as psoriasis or actinic keratosis. It could be speculated that ODN containing preparations help to lower the concentration of pro-apoptotic compounds with toxic side effects maintaining similar clinical effects.

SUPPLEMENTARY DATA

Supplementary Data are available at NAR Online.

ACKNOWLEDGEMENTS

We are particularly thankful to Heike Beschmann for excellent assistance providing the statistical analysis.

FUNDING

Federal Ministry of Education and Research (BMBF), Germany (BioChancePLUS, Number 0313642B).

Conflict of interest statement. None declared.

REFERENCES

- Coley, W.B. (1893) The treatment of malignant tumors by repeated inoculations with of erysipelas with a report of ten original cases. *Am. J. Med. Sci.*, **105**, 487–511.
- Coley, W.B. (1884) Treatment of inoperable malignant tumors with the toxins of erysipelas and the bacillus prodigiosus. *Am. J. Med. Sci.*, **108**, 183–212.
- Tokunaga, T., Yamamoto, H., Shimada, S., Abe, H., Fukuda, T., Fujisawa, Y., Furutani, Y., Yano, O., Kataoka, T. and Sudo, T. (1984) Antitumor activity of deoxyribonucleic acid fraction from *Mycobacterium bovis* BCG. I. Isolation, physicochemical characterization and antitumor activity. *J. Natl Cancer Inst.*, **72**, 955–962.
- Krieg, A.M., Yi, A.-K., Matson, S., Waldschmidt, T.J., Bishop, G.A., Teasdale, R., Koretzky, G.A. and Klinman, D.M. (1995) CpG motifs in bacterial DNA trigger direct B-cell activation. *Nature*, **374**, 546–549.
- Bird, A.P. (1985) CpG islands and genes. *Curr. Opin. Genet.*, **5**, 309–314.
- Bird, A.P. (1987) CpG islands as gene markers in the vertebrate nucleus. *Trends Genet.*, **3**, 342–347.
- Hergersberg, M. (1991) Biological aspects of cytosine methylation in eukaryotic cells. *Experientia*, **47**, 1171–1185.
- Hemmi, H., Takeuchi, O., Kawai, T., Kaisho, T., Sato, S., Sanjo, H., Matsumoto, M., Hoshino, K., Wagner, H., Takeda, K. and Akira, S. (2000) A Toll-like receptor recognizes bacterial DNA. *Nature*, **408**, 740–745.
- Dorn, A. and Kippenberger, S. (2008) Clinical application of CpG-, non-CpG-, and antisense oligodeoxynucleotides as immunomodulators. *Curr. Opin. Mol. Ther.*, **10**, 10–20.
- Stein, C.A., Subasinghe, C., Shinozuka, K. and Cohen, J.S. (1988) Physicochemical properties of phosphorothioate oligodeoxynucleotides. *Nucleic Acids Res.*, **16**, 3209–3221.
- Matteucci, M. (1997) Oligonucleotide analogues: an overview. *Ciba Found Symp.*, **209**, 5–14.
- Tokunaga, T., Yano, O., Kuramoto, E., Kimura, Y., Yamamoto, T., Kataoka, T. and Yamamoto, S. (1992) Synthetic oligonucleotides with particular base sequences from the cDNA encoding proteins of *Mycobacterium bovis* BCG induce interferons and activate natural killer cells. *Microbiol. Immunol.*, **36**, 55–66.
- Yamamoto, T., Yamamoto, S., Kataoka, T. and Tokunaga, T. (1994) Lipofection of synthetic oligodeoxyribonucleotide having a palindromic sequence of AACGTT to murine splenocytes enhances interferon production and natural killer activity. *Microbiol. Immunol.*, **38**, 831–836.
- Liang, H., Nishioka, Y., Reich, C.F., Pisetsky, D.S. and Lipsky, P.E. (1996) Activation of human B cells by phosphorothioate oligodeoxynucleotides. *J. Clin. Invest.*, **98**, 1119–1129.
- Dorn, A., Ludwig, R.J., Bock, A., Thaci, D., Hardt, K., Bereiter-Hahn, J., Kaufmann, R., Bernd, A. and Kippenberger, S. (2007) Oligonucleotides suppress IL-8 in skin keratinocytes in vitro and offer anti-inflammatory properties in vivo. *J. Invest. Dermatol.*, **127**, 846–854.
- Luo, H.R. and Loison, F. (2008) Constitutive neutrophil apoptosis: Mechanisms and regulation. *Am. J. Hematol.*, **83**, 288–295.
- Bauer, S., Kirschning, C.J., Hacker, H., Redecke, V., Hausmann, S., Akira, S., Wagner, H. and Lipford, G.B. (2001) Human TLR9 confers responsiveness to bacterial DNA via species-specific CpG motif recognition. *Proc. Natl Acad. Sci. USA*, **98**, 9237–9242.
- Kippenberger, S., Bernd, A., Loitsch, S., Guschel, M., Müller, J., Bereiter-Hahn, J. and Kaufmann, R. (2000) Signaling of mechanical stretch in human keratinocytes via MAP kinases. *J. Invest. Dermatol.*, **114**, 408–412.
- Karasek, M.A. (1980) Effect of chemical modification of keratinocytes and collagen in keratinocyte-collagen interaction. *Curr. Probl. Dermatol.*, **10**, 143–158.
- Leist, M., Single, B., Naumann, H., Fava, E., Simon, B., Kuhnle, S. and Nicotera, P. (1999) Nitric oxide inhibits execution of apoptosis at two distinct ATP-dependent steps upstream and downstream of mitochondrial cytochrome *c* release. *Biochem. Biophys. Res. Commun.*, **258**, 215–221.
- Kuo, C.C., Liang, C.M., Lai, C.Y. and Liang, S.M. (2007) Involvement of heat shock protein (Hsp)90 beta but not Hsp90 alpha in antiapoptotic effect of CpG-B oligodeoxynucleotide. *J. Immunol.*, **178**, 6100–6108.
- Medzhitov, R., Preston-Hurlburt, P., Kopp, E., Stadler, A., Chen, C., Ghosh, S. and Janeway, C.A. Jr (1998) MyD88 is an adaptor

- protein in the hToll/IL-1 receptor family signaling pathways. *Mol. Cell*, **2**, 253–258.
23. Kippenberger, S., Loitsch, S., Guschel, M., Muller, J., Knies, Y., Kaufmann, R. and Bernd, A. (2005) Mechanical stretch stimulates protein kinase B/Akt phosphorylation in epidermal cells via angiotensin II type I receptor and epidermal growth factor receptor. *J. Biol. Chem.*, **280**, 3060–3067.
 24. Jozsef, L., Khreiss, T. and Filep, J.G. (2004) CpG motifs in bacterial DNA delay apoptosis of neutrophil granulocytes. *FASEB J.*, **18**, 1776–1778.
 25. Vollmer, J., Janosch, A., Laucht, M., Ballas, Z.K., Schetter, C. and Krieg, A.M. (2002) Highly immunostimulatory CpG-free oligodeoxynucleotides for activation of human leukocytes. *Antisense Nucleic Acid Drug Dev.*, **12**, 165–175.
 26. Sano, K., Shirota, H., Terui, T., Hattori, T. and Tamura, G. (2003) Oligodeoxynucleotides without CpG motifs work as adjuvant for the induction of Th2 differentiation in a sequence-independent manner. *J. Immunol.*, **170**, 2367–2373.
 27. Vollmer, J., Weeratna, R.D., Jurk, M., Samulowicz, U., McCluskie, M.J., Payette, P., Davis, H.L., Schetter, C. and Krieg, A.M. (2004) Oligodeoxynucleotides lacking CpG dinucleotides mediate Toll-like receptor 9 dependent T helper type 2 biased immune stimulation. *Immunology*, **113**, 212–223.
 28. Takeshita, F., Leifer, C.A., Gursel, I., Ishii, K.J., Takeshita, S., Gursel, M. and Klinman, D.M. (2001) Cutting edge. Role of toll-like receptor 9 in CpG DNA-induced activation of human cells. *J. Immunol.*, **167**, 3555–3558.
 29. Chuang, T.H., Lee, J., Kline, L., Mathison, J.C. and Ulevitch, R.J. (2002) Toll-like receptor 9 mediates CpG-DNA signaling. *J. Leukoc. Biol.*, **71**, 538–544.
 30. Franke, T.F., Kaplan, D.R., Cantley, L.C. and Tokar, A. (1997) Direct regulation of the Akt proto-oncogene product by phosphatidylinositol-3,4-bisphosphate. *Science*, **275**, 665–668.
 31. Kennedy, S.G., Wagner, A.J., Conzen, S.D., Jordan, J., Bellacosa, A., Tschlis, P.N. and Hay, N. (1997) The PI 3-kinase/Akt signaling pathway delivers an anti-apoptotic signal. *Genes Dev.*, **11**, 701–713.
 32. Alessi, D.R., Deak, M., Casamayor, A., Caudwell, F.B., Morrice, N., Norman, D.G., Gaffney, P., Reese, C.B., MacDougall, C.N., Harbison, D. *et al.* (1997) 3-Phosphoinositide-dependent protein kinase-1 (PDK1): structural and functional homology with the Drosophila DSTPK61 kinase. *Curr. Biol.*, **7**, 776–789.
 33. Toker, A. and Newton, A.C. (2000) Akt/protein kinase B is regulated by autophosphorylation at the hypothetical PDK-2 site. *J. Biol. Chem.*, **275**, 8271–8274.
 34. Attwell, S., Roskelley, C. and Dedhar, S. (2000) The integrin-linked kinase (ILK) suppresses anoikis. *Oncogene*, **19**, 3811–3815.
 35. Alessi, D.R., Andjelkovic, M., Caudwell, B., Cron, P., Morrice, N., Cohen, P. and Hemmings, B.A. (1996) Mechanism of activation of protein kinase B by insulin and IGF-1. *EMBO J.*, **15**, 6541–6551.
 36. Kandel, E.S. and Hay, N. (1999) The regulation and activities of the multifunctional serine/threonine kinase Akt/PKB. *Exp. Cell Res.*, **253**, 210–229.
 37. Khwaja, A. (1999) Akt is more than just a Bad kinase. *Nature*, **401**, 33–34.
 38. Rusinol, A.E., Thewke, D., Liu, J., Freeman, N., Panini, S.R. and Sinensky, M.S. (2004) AKT/protein kinase B regulation of BCL family members during oxysterol-induced apoptosis. *J. Biol. Chem.*, **279**, 1392–1399.
 39. Slee, E.A., Harte, M.T., Kluck, R.M., Wolf, B.B., Casiano, C.A., Newmeyer, D.D., Wang, H.G., Reed, J.C., Nicholson, D.W., Alnemri, E.S. *et al.* (1999) Ordering the cytochrome c-initiated caspase cascade: hierarchical activation of caspases-2, -3, -6, -7, -8, and -10 in a caspase-9-dependent manner. *J. Cell Biol.*, **144**, 281–292.
 40. Burgering, B.M. and Coffey, P.J. (1995) Protein kinase B (c-Akt) in phosphatidylinositol-3-OH kinase signal transduction. *Nature*, **376**, 599–602.
 41. Franke, T.F., Yang, S.I., Chan, T.O., Datta, K., Kazlauskas, A., Morrison, D.K., Kaplan, D.R. and Tschlis, P.N. (1995) The protein kinase encoded by the Akt proto-oncogene is a target of the PDGF-activated phosphatidylinositol 3-kinase. *Cell*, **81**, 727–736.
 42. Kohn, A.D., Kovacina, K.S. and Roth, R.A. (1995) Insulin stimulates the kinase activity of RAC-PK, a pleckstrin homology domain containing ser/thr kinase. *EMBO J.*, **14**, 4288–4295.
 43. Rockwell, P., O'Connor, W.J., King, K., Goldstein, N.I., Zhang, L.M. and Stein, C.A. (1997) Cell-surface perturbations of the epidermal growth factor and vascular endothelial growth factor receptors by phosphorothioate oligodeoxynucleotides. *Proc. Natl Acad. Sci. USA*, **94**, 6523–6528.