

Supporting Information

The resolution revolution in cryoEM requires new sample preparation procedures: A rapid pipeline to high resolution maps of yeast FAS

Mirko Joppe^{1#}, Edoardo D'Imprima^{2,3#}, Nina Salustros^{2,4}, Karthik S. Paithankar¹, Janet Vonck², Martin Grininger^{1*}, Werner Kühlbrandt^{2*}

¹Buchmann Institute for Molecular Life Sciences, Goethe University Frankfurt, Institute of Organic Chemistry and Chemical Biology, Frankfurt am Main, Max von Laue Strasse 15, 60438, Germany

²Max Planck Institute of Biophysics, Department of Structural Biology, Frankfurt am Main, Max von Laue Strasse 3, 60438, Germany

³Current Address: Structural and Computational Biology Unit, European Molecular Biology Laboratory, Meyerhofstrasse 1, 69117 Heidelberg, Germany

⁴Current Address: University of Copenhagen, Department of Biomedical Sciences, Nørre Allé 14, DK-2200 Copenhagen, Denmark

#These authors contributed equally

*Correspondence should be addressed to: grininger@chemie.uni-frankfurt.de, werner.kuehlbrandt@biophys.mpg.de

File(s):

Figure S1. CryoEM reconstruction of yeast FAS colored by local resolution.




Figure S1. CryoEM reconstruction of yeast FAS colored by local resolution. Yeast FAS is shown in top view (A), side view (B) and as cross-section along the threefold axis (C).