

A journal of world insect systematics

INSECTA MUNDI

1059

Anthrenus (Nathrenus) peacockae (Coleoptera:
Dermestidae: Megatominae), a new species from Pakistan

Graham J. Holloway

Cole Museum of Zoology, School of Biological Sciences, Health and Life Sciences Building,
The University of Reading, Whiteknights, Reading, Berkshire, RG6 6EX, UK

Date of issue: June 28, 2024

Center for Systematic Entomology, Inc., Gainesville, FL

Holloway GJ. 2024. *Anthrenus (Nathrenus) peacockae* (Coleoptera: Dermestidae: Megatominae), a new species from Pakistan. *Insecta Mundi* 1059: 1–5.

Published on June 28, 2024 by
Center for Systematic Entomology, Inc.
P.O. Box 141874
Gainesville, FL 32614-1874 USA
<http://centerforsystematicentomology.org/>

INSECTA MUNDI is a journal primarily devoted to insect systematics, but articles can be published on any non-marine arthropod. Topics considered for publication include systematics, taxonomy, nomenclature, checklists, faunal works, and natural history. *Insecta Mundi* will not consider works in the applied sciences (i.e. medical entomology, pest control research, etc.), and no longer publishes book reviews or editorials. *Insecta Mundi* publishes original research or discoveries in an inexpensive and timely manner, distributing them free via open access on the internet on the date of publication.

Insecta Mundi is referenced or abstracted by several sources, including the Zoological Record and CAB Abstracts. *Insecta Mundi* is published irregularly throughout the year, with completed manuscripts assigned an individual number. Manuscripts must be peer reviewed prior to submission, after which they are reviewed by the editorial board to ensure quality. One author of each submitted manuscript must be a current member of the Center for Systematic Entomology.

Guidelines and requirements for the preparation of manuscripts are available on the *Insecta Mundi* website at <http://centerforsystematicentomology.org/insectamundi/>

Chief Editor: David Plotkin, insectamundi@gmail.com

Assistant Editor: Paul E. Skelley, insectamundi@gmail.com

Layout Editor: Robert G. Forsyth

Editorial Board: Davide Dal Pos, M. J. Paulsen, Felipe Soto-Adames

Founding Editors: Ross H. Arnett, Jr., J. H. Frank, Virendra Gupta, John B. Heppner, Lionel A. Stange, Michael C. Thomas, Robert E. Woodruff

Review Editors: Listed on the *Insecta Mundi* webpage

Printed copies (ISSN 0749-6737) annually deposited in libraries

Florida Department of Agriculture and Consumer Services, Gainesville, FL, USA

The Natural History Museum, London, UK

National Museum of Natural History, Smithsonian Institution, Washington, DC, USA

Zoological Institute of Russian Academy of Sciences, Saint-Petersburg, Russia

Electronic copies (online ISSN 1942-1354) in PDF format

Archived digitally by Portico.

Florida Virtual Campus: <http://purl.fcla.edu/fcla/insectamundi>

University of Nebraska-Lincoln, Digital Commons: <http://digitalcommons.unl.edu/insectamundi/>

Goethe-Universität, Frankfurt am Main: <http://nbn-resolving.de/urn/resolver.pl?urn:nbn:de:hebis:30:3-135240>

This is an open access article distributed under the terms of the Creative Commons, Attribution Non-Commercial License, which permits unrestricted non-commercial use, distribution, and reproduction in any medium, provided the original author(s) and source are credited.
<https://creativecommons.org/licenses/by-nc/3.0/>

Anthrenus (Nathrenus) peacockae (Coleoptera: Dermestidae: Megatominae), a new species from Pakistan

Graham J. Holloway

Cole Museum of Zoology, School of Biological Sciences, Health and Life Sciences Building,
The University of Reading, Whiteknights, Reading, Berkshire, RG6 6EX, UK
g.j.holloway@reading.ac.uk

🌐 <https://orcid.org/0000-0003-0495-0313>

Abstract. A new species, *Anthrenus (Nathrenus) peacockae* Holloway (Coleoptera: Dermestidae: Megatominae) from Pakistan is described and compared with *A. narani* Háva and Ahmed, 2014, the most similar species. Images of the habitus, ventrites, antenna, aedeagus, sternites VIII and IX are presented. *Anthrenus peacockae* is only the third species in the subgenus *Nathrenus* Casey, 1900 to be recorded from Pakistan.

Key words. Carpet beetle, *Anthrenus narani*, aedeagus, median lobe, sternite, taxonomy, distribution, taxonomy.

ZooBank registration. urn:lsid:zoobank.org:pub:F68C53E2-3BCF-4F6B-A838-A14EE2FA4B66

Introduction

Dermestidae Latreille, 1804 (Coleoptera) contains over 1900 species (Háva 2023). One of the largest genera within the family is *Anthrenus* Geoffroy, 1762 with nearly 300 species, split into 10 subgenera (Háva 2023). Research on larval features suggests that *Anthrenus* is polytypic (Kadej 2018), so it is possible the taxonomy of *Anthrenus* might be revised. For the moment, taxonomy of *Anthrenus* is based largely on antennal structure (Peacock 1993).

Despite being a large genus, *Anthrenus* continues to become more speciose as workers turn their attention to dissection and genital structure to differentiate species. This approach was largely pioneered by Beal (1998) but first applied to Palaearctic *Anthrenus* in a focused manner by Kadej et al. (2007). Kadej et al. (2007) described three new species, with additional new *Anthrenus* species discovered through dissection by Kadej and Háva (2011), Holloway (2019, 2020, 2021, 2023), and Holloway et al. (2023b) mostly by reviewing material in museum collections.

The Natural History Museum, London (NHML) contains a good collection of Dermestidae. Whilst examining the collection of *Anthrenus* species at NHML, the author came across nine specimens of an unfamiliar species. One of the specimens had been labelled by Enid Peacock as '*Anthrenus (Nathrenus)* sp. n.?. Here *Anthrenus (Nathrenus) peacockae*, a new species from Pakistan, is described.

Materials and Methods

The dry, carded specimens were borrowed from NHML for study. All specimens were macerated in a solution of 2% acetic acid for five days to soften and allow removal from staging prior to dissection. Dissection was carried out under a Brunel BMSL zoom stereo LED microscope and involved detaching the abdomen from the rest of the insect using two entomological pins. The soft tergites were then peeled away from the harder ventrites to expose the genitalia. Sternite VIII was detached from tergite IX, and the aedeagus and sternite IX were detached from the ring sclerite. Habitus images, both dorsal and ventral sides, were captured at $\times 20$ magnification using a Canon EOS 2000D camera mounted on the BMSL microscope. Images of the antenna, aedeagus, and sternite IX were captured at $\times 100$ magnification using a Canon EOS 1300D camera mounted on a Brunel monocular SP28 microscope. Images of sternite VIII were captured via the Canon EOS 1300D at $\times 50$. After dissection, all body parts were mounted on card. All images were fed through Helicon Focus Pro version 8.2.2 focus-stacking software. All measurements were made using DsCap.Ink software version 3.90. Measurements taken:

- Body length (BL): distance from anterior margin of pronotum to the apex of the elytra.
- Body width (BW): maximum distance across the elytra.
- Antennal club length (AL): length of the last three antennomeres.
- Antennal club width (AW): maximum width across the terminal antennomere.
- Paramere length (PL): distance from the anterior end of the parameres to the apex of the parameres.
- Sternite IX length (SL): distance from the tip of one anterior horn to the tip of the posterior lobe.

The data for the distribution map (Shorthouse 2010) were derived from the data labels.

Results

Anthrenus (Nathrenus) peacockae Holloway, new species

(Fig. 1, 2)

Type specimens. Holotype male. Hazara, Lower Kagan Valley, Pakistan (32.795°N, 74.285°E), 6 vi 1927, H.G. Champion leg. NHML.

Paratypes. Eight males, NHML, same data as holotype.

Description, external characteristics. Holotype *Anthrenus (Nathrenus) peacockae* (Fig. 1A) (BL = 4.1 mm, BW = 2.65 mm) has a single orange ocellus on head between the eyes but below the level of the top of the eyes. The inner margins of the eyes smoothly curved with no notch, typical of the subgenus *Nathrenus* Casey, 1900 (Peacock 1993). The head and prothorax are a very dark brown; the elytra are a pale chestnut brown. The pronotum and elytra are covered in loosely packed pale cream and very dark brown scales. The scales are slim and sharply pointed at both ends. The pale scales are clustered in the basal ¼ of each elytron and the posterior lateral corners of the pronotum. Elsewhere, the pale scales are scattered along the elytral suture and form a loose fascia beyond the midpoint with patches at the lateral elytral margins just beyond the midpoint and sub-apically. The ocellus is




Figure 1. *Anthrenus peacockae* holotype. **A)** Habitus dorsal aspect (scale bar = 1 mm). **B)** Abdominal ventrites (scale bar = 1 mm). **C)** Antenna (scale bar = 100 μ m).




Figure 2. *Anthrenus peacockae* holotype. **A)** Aedeagus dorsal aspect. **B)** Sternite VIII dorsal aspect. **C)** Sternite IX dorsal aspect. All scale bars = 100 μm .

surrounded by dark brown scales and pale cream scales line the inner margins of each eye. Palps orange (evident in some paratypes).

The ventrites (Fig. 1B) are mid-brown and covered in pale cream scales. The lateral margins of each ventrite carry large spots of dark brown scales, largest on ventrite I and becoming progressively smaller towards ventrite IV. There are two spots of brown scales either side of the midline on ventrites II, III, and IV. The dark spot of scales on ventrite I is not entirely bounded by pale scales along the anterior edge. The tergites are a pale yellowish brown.

The 11-segmented antenna (Fig. 1C) (AL = 263 μm , AW = 107 μm) is slim and entirely pale reddish brown. Antennomeres 1 and 2 are large and round, antennomeres 3–8 are smaller and squarish, and antennomeres 9–11 form a well-defined elongated club. All components of the legs are similarly pale reddish brown, contrasting with the dark brown of the thorax. The anterior faces of the femora are covered in pale scales, the tibiae and tarsi lack scales.

Description, internal characteristics. Figure 2A shows the ventral surface of the holotype male aedeagus. The parameres (PL = 530 μm) are narrow, but deeper dorsoventrally. The lateral margins are mostly parallel to each other with only the slightest inwards curve at the apex. The inner margins of the parameres are slightly concave and covered in long, fine hairs. The apices of the parameres are adorned with long, fine, straight hairs.

The median lobe is very broad, parallel sided from the base to the midway point, apically the lateral margins converge to a rounded, blunt apex that falls short of the tips of the parameres. Along the ventral surface of the median lobe and projecting beyond the apex is a membranous tube.

Sternite VIII (Fig. 2B) is an ornate structure with several broad spines or spicules emerging from each curved sclerotized rib and pointing obliquely anterior. Most spicules with apex obliquely flattened. The anterior halves of both sides of sternite VIII consist of a lattice of sclerotized rods.

Sternite IX (Fig. 2C, SL = 585 μm) is a very thin membranous structure that expands from the base to about the basal $\frac{1}{3}$ and from there the lateral margins converge to the apical $\frac{1}{5}$, where margins become parallel to a rounded apex. The margins from the apex to about midway are covered in many long hairs, which are longest and densest around the apex; hairs hook inwards apically. There are a few hairs on the central part of sternite IX, but the hairs are concentrated on the outer sections of the sternite.

Morphometrics. Mean BL = 3.94 ± 0.22 (standard deviation) mm, mean BW/BL = 0.65 ± 0.01 , $n = 9$ in both cases. The data indicate that 95% BL of *A. peacockae* would lie within 3.43–4.46 mm. The smallest individual found in the sample studied was 3.45 mm, the largest 4.2 mm.

Distribution. The location of collection of the holotype and all paratypes is shown in Fig 3.

Etymology. *Anthrenus peacockae* is named after Enid Peacock, author of the Royal Entomological Society Handbook on Dermestidae and Derodontidae in the UK, who noted the *A. peacockae* specimens studied here were a new species.

Differential diagnosis. Initially, *A. peacockae* was thought to be *A. narani* Háva and Ahmed, 2014, but the internal features illustrated by Háva and Ahmed (2014) (Fig. 4) bear little resemblance to *A. peacockae*. Externally, *A. peacockae* and *A. narani* are similar.

The antenna illustrated for *Anthrenus narani* by Háva and Ahmed (2014) (Fig. 4A) has a truncated apex to the terminal antennomere (*A. peacockae* is rounded, Fig. 1C), only antennomere 1 is large and rounded in *A. narani* (both antennomeres 1 and 2 are large and rounded in *A. peacockae*), antennomeres 2–5 are similar sized and quadrate, antennomeres 6–8 are much broader than long (antennomeres 3–8 are all quadrate in *A. peacockae*).

According to Háva and Ahmed (2014), the aedeagus of *A. narani* (Fig. 4B) has strongly curved parameres with much smaller tufts of hairs at the apex than *A. peacockae* (Fig. 2A). The illustration in Háva and Ahmed (2014) suggests that only the apex of each paramere has hairs. The paramere apices almost touch and are distinctly longer than the median lobe. Otherwise, the median lobe of *A. narani* illustrated by Háva and Ahmed (2014) is similar to *A. peacockae* (Fig. 2A).

Sternite IX illustrated by Háva and Ahmed (2014) (Fig. 4C) has a truncated posterior apex (*A. peacockae* is rounded, Fig. 2C) with only short hairs (*A. peacockae* sternite IX has many long hairs apically and along the sides). Háva and Ahmed (2014) illustrate a sternite IX with posterior margins converging all the way to the posterior apex (sternite IX margins for the final 1/5 are parallel sided in *A. peacockae*). Háva and Ahmed (2014) did not illustrate sternite VIII.




Figure 3. Collection location of holotype and all paratype *Anthrenus peacockae* from Pakistan.




Figure 4. *Anthrenus narani* as illustrated by Háva and Ahmed (2014). **A)** Antenna. **B)** Aedeagus. **C)** Sternite IX.

Discussion

Anthrenus peacockae represents only the third *Nathrenus* species known from Pakistan, the others being *A. narani* and *A. verbasci* Linnaeus, 1767. Háva and Ahmed (2014) report that *A. narani* is large with the holotype BL = 4.3 mm. *Anthrenus peacockae* is also large with average BL = 3.94 mm. One of the few other *Nathrenus* species with some specimens approaching 4 mm is *A. transcaspicus* Mroczkowski, 1960 (Herrmann 2023), which is known from Iran and Turkmenistan, the same region of Asia as Pakistan.

Anthrenus narani and *A. peacockae* are externally similar, both in terms of size and scale pattern. The present study illustrates the necessity to dissect genitalia to establish species identity for sound taxonomic work and to determine distribution (Holloway et al. 2023a). From the information provided by Háva and Ahmed (2014) and the information presented here, it is clear that *A. peacockae* and *A. narani* are different species.

Acknowledgments

The author is very grateful to Dr. Chris Foster, Andreas Herrmann and Dr. Paul Skelley for reviewing the manuscript so efficiently, and to Max Barclay and the Coleoptera curatorial team at NHML for maintaining and making available specimens for comparison.

Literature Cited

- Beal RS Jr. 1998.** Taxonomy and biology of Nearctic species of *Anthrenus* (Coleoptera: Dermestidae). Transactions of the American Entomological Society 124: 271–332. <https://www.jstor.org/stable/25078667>
- Háva J 2023.** Dermestidae World (Coleoptera). Available at <http://dermestidae.wz.cz/world-dermestidae> (Last accessed 12 December 2023.)
- Háva J, Ahmed Z. 2014.** A new species of the genus *Anthrenus* from Pakistan (Coleoptera: Dermestidae: Megatominae: Anthrenini). Studies and Reports Taxonomical Series 10 (1): 89–92.

- Herrmann A. 2023.** Dermestidae (Coleoptera) of the World. <https://www.dermestidae.com> (Last accessed 12 December 2023).
- Holloway GJ. 2019.** *Anthrenus* (s. str.) *amandae* (Coleoptera: Dermestidae): a new species from Mallorca, Spain. *Zootaxa* 4543 (4): 595–599. <http://doi.org/10.11646/zootaxa.4543.4.9>
- Holloway GJ. 2020.** *Anthrenus* (s. str.) *chikatunovi* (Coleoptera: Dermestidae): a new species from southern France. *Israel Journal of Entomology* 50: 69–75. <http://doi.org/10.5281/zenodo.4088743>
- Holloway GJ. 2021.** *Anthrenus* (s. str.) *corona* (Coleoptera, Dermestidae, Anthrenini): a new species in the *A. pimpinellae* (Fabricius, 1775) complex from Turkey. *Zootaxa* 4991(3): 555–560. <https://doi.org/10.11646/zootaxa.4991.3.7>
- Holloway GJ. 2023.** *Anthrenus* (*Anthrenus*) *mumbaiensis* sp. nov. from India and a morphometric examination of *Anthrenus* (*Anthrenus*) *festivus* (Coleoptera, Dermestidae, Anthrenini). *Zootaxa* 5306(3): 377–384. <https://doi.org/10.11646/zootaxa.5306.3.5>
- Holloway GJ, McLure CJ, Foster CW. 2023a.** Palaearctic distributions of *Anthrenus pimpinellae* (Fabricius) and *Anthrenus isabellinus* Küster (Coleoptera: Dermestidae). *Entomologist's Monthly Magazine* 159: 239–244. <https://doi.org/10.31184/m00138908.1594.4207>
- Holloway GJ, Thanasoulis G, Herrmann A. 2023b.** A new species *Anthrenus bakaloudisi* sp. nov. (Coleoptera, Dermestidae, Megatominae) from Macedonia, Greece and comparison with *Anthrenus pfefferi* Kalík, 1954, *Anthrenus delicatus* Kiesenwetter, 1851, and *Anthrenus warchalowskii* Kadej, Háva & Kalík, 2007. *Baltic Journal of Coleopterology* 23(1): 41–50. [https://doi.org/10.59893/bjc.23\(1\).003](https://doi.org/10.59893/bjc.23(1).003)
- Kadej M. 2018.** Contribution to knowledge of the immature stages of Dermestidae with special emphasis on the larval morphology of the genus *Anthrenus* Geoffroy, 1762 (Megatominae, Anthrenini). *Polish Entomological Monographs* No. 16. Polish Entomological Society; Poznań, Poland. 180 p.
- Kadej M, Háva J. 2011.** Three new species of *Anthrenus pimpinellae* species group from Palaearctic Region (Coleoptera: Dermestidae: Megatominae: Anthrenini). *Studies and Reports, Taxonomical Series* 7: 241–248.
- Kadej M, Háva J, Kalík V. 2007.** Review of the *Anthrenus pimpinellae* species group from Palaearctic region (Coleoptera: Dermestidae: Anthrenini). *Genus* 18(4): 721–750.
- Peacock ER. 1993.** Adults and larvae of hide, larder and carpet beetles and their relatives (Coleoptera: Dermestidae) and of derodontid beetles (Coleoptera: Derodontidae). *Handbooks for the identification of British insects*, vol. 5, no. 3. Natural History Museum; London, UK. 144 p.
- Shorthouse DP. 2010.** SimpleMappr, an online tool to produce publication-quality point maps. Available at <https://www.simplemappr.net> (Last accessed 11 December 2023.)

Received February 16, 2024; accepted June 3, 2024.

Review editor Paul E. Skelley.