



Anisotropic flow of charged hadrons, pions and (anti-)protons measured at high transverse momentum in Pb–Pb collisions at $\sqrt{s_{NN}} = 2.76$ TeV [☆]

ALICE Collaboration

ARTICLE INFO

Article history:

Received 28 May 2012

Received in revised form 28 November 2012

Accepted 28 December 2012

Available online 4 January 2013

Editor: V. Metag

ABSTRACT

The elliptic, v_2 , triangular, v_3 , and quadrangular, v_4 , azimuthal anisotropic flow coefficients are measured for unidentified charged particles, pions, and (anti-)protons in Pb–Pb collisions at $\sqrt{s_{NN}} = 2.76$ TeV with the ALICE detector at the Large Hadron Collider. Results obtained with the event plane and four-particle cumulant methods are reported for the pseudo-rapidity range $|\eta| < 0.8$ at different collision centralities and as a function of transverse momentum, p_T , out to $p_T = 20$ GeV/c. The observed non-zero elliptic and triangular flow depends only weakly on transverse momentum for $p_T > 8$ GeV/c. The small p_T dependence of the difference between elliptic flow results obtained from the event plane and four-particle cumulant methods suggests a common origin of flow fluctuations up to $p_T = 8$ GeV/c. The magnitude of the (anti-)proton elliptic and triangular flow is larger than that of pions out to at least $p_T = 8$ GeV/c indicating that the particle type dependence persists out to high p_T .

© 2013 CERN. Published by Elsevier B.V. Open access under [CC BY-NC-ND license](http://creativecommons.org/licenses/by-nc-nd/3.0/).

The goal of ultra-relativistic nucleus–nucleus collisions is to study nuclear matter under extreme conditions. For non-central collisions, in the plane perpendicular to the beam direction, the geometrical overlap region, where the highly Lorentz contracted nuclei intersect and where the initial interactions occur, is azimuthally anisotropic. This initial spatial asymmetry is converted via interactions into an anisotropy in momentum space, a phenomenon referred to as transverse anisotropic flow (for a review see [1]). Anisotropic flow has become a key observable for the characterization of the properties and the evolution of the system created in a nucleus–nucleus collision.

Identified particle anisotropic flow provides valuable information on the particle production mechanism in different transverse momentum, p_T , regions [1]. For $p_T < 2$ –3 GeV/c, the flow pattern of different particle species is qualitatively described by hydrodynamic model calculations [2]. At intermediate p_T , $3 < p_T < 6$ GeV/c, the observed flow of the baryons is larger than that of the mesons [3,4]. For $p_T \gtrsim 8$ GeV/c, the fragmentation of high-energy partons, resulting from initial hard scatterings, is expected to play the dominant role. While traversing the hot and dense matter these partons experience collisional and radiative energy loss [5,6], which are strongly dependent on the thickness of the created medium [7]. In the azimuthally asymmetric system, the energy loss depends on the azimuthal emission angle of the parton, which leads to an azimuthal anisotropy in particle production at high p_T [8,9].

The magnitude of the anisotropic flow is characterized by the coefficients in the Fourier expansion of the azimuthal distribution of particles with respect to the collision symmetry plane [10,11]:

$$v_n(p_T, \eta) = \langle \cos[n(\phi - \Psi_n)] \rangle, \quad (1)$$

where p_T , η , and ϕ are the particle's transverse momentum, pseudo-rapidity, and the azimuthal angle, respectively, and Ψ_n is the n -th harmonic symmetry plane angle. For a smooth matter distribution in the colliding nuclei, the symmetry planes of all harmonics coincide with the reaction plane defined by the beam direction and the impact parameter, the vector connecting the centers of the two colliding nuclei at closest approach. In this case, for particles produced at midrapidity, all odd Fourier coefficients are zero by symmetry. Due to event-by-event fluctuations of the positions of the participating nucleons inside the nuclei, the shape of the initial energy density of the heavy-ion collision in general is not symmetric with respect to the reaction plane, and the Ψ_n may deviate from the reaction plane. This gives rise to non-zero odd harmonic coefficients [12–18], and contributes to the difference in flow coefficients calculated from two- or multi-particle azimuthal correlations, and also to the difference in v_n measured with respect to different harmonic symmetry planes.

Large elliptic flow, v_2 , and significant triangular flow, v_3 , were observed at the Relativistic Heavy Ion Collider (RHIC) [19–21] and at the Large Hadron Collider (LHC) [22–28]. In this Letter we present the measurement of unidentified charged particle anisotropic flow out to $p_T = 20$ GeV/c, and for protons and

charged pions¹ out to $p_T = 16$ GeV/c. We also present unidentified charged particle quadrangular flow, v_4 , measured with respect to the second (Ψ_2) and fourth (Ψ_4) harmonic symmetry planes.

The data sample recorded by ALICE during the 2010 heavy-ion run at the LHC is used for this analysis. Detailed descriptions of the ALICE detector can be found in [29–31]. The Time Projection Chamber (TPC) was used to reconstruct charged particle tracks and measure their momenta with full azimuthal coverage in the pseudo-rapidity range $|\eta| < 0.8$, and for particle identification via the specific ionization energy loss, dE/dx , in the transverse momentum region $p_T > 3$ GeV/c. Two scintillator arrays (VZERO) which cover the pseudo-rapidity ranges $-3.7 < \eta < -1.7$ and $2.8 < \eta < 5.1$ were used for triggering, and the determination of centrality [32] and symmetry planes. The trigger conditions and the event selection criteria are identical to those described in [22,23,32]. Approximately 10^7 minimum-bias Pb-Pb events with a reconstructed primary vertex within ± 10 cm from the nominal interaction point in the beam direction are used for this analysis. Charged particles reconstructed in the TPC in $|\eta| < 0.8$ and $0.2 < p_T < 20$ GeV/c were selected. The charged track quality cuts described in [22] were applied to minimize contamination from secondary charged particles and fake tracks. The charged particle track reconstruction efficiency and contamination were estimated from HIJING Monte Carlo simulations [33] combined with a GEANT3 [34] detector model, and found to be independent of the collision centrality. The reconstruction efficiency, which may bias the determination of the p_T averaged flow, increases from 70% to 80% for particles with $0.2 < p_T < 1$ GeV/c and remains constant at $80 \pm 5\%$ for $p_T > 1$ GeV/c. The estimated contamination by secondary charged particles from weak decays and photon conversions is less than 6% at $p_T = 0.2$ GeV/c and falls below 1% for $p_T > 1$ GeV/c.

The selection of pions and protons at $p_T > 3$ GeV/c is based on the measurement of the dE/dx in the TPC, following the procedure described in [35]. Enriched pion (proton) samples are obtained by selecting tracks from the upper (lower) part of the expected pion (proton) dE/dx distribution. For example, protons were typically selected, depending on their momentum, in the range from 0 to -3σ or from -1.5σ to -4.5σ around their nominal value in dE/dx , where σ is the energy loss resolution. Note that dE/dx of pions is larger than that of protons in the p_T range used for this study. The track selection criteria have been adjusted to keep the contamination by other particle species below 1% for pions and below 15% for protons. The pion and proton v_2 and v_3 are not corrected for this contamination. The systematic uncertainties in v_2 and v_3 related to the purity of the pion and proton samples are 2% for $p_T < 8$ GeV/c and 10% for $p_T \geq 8$ GeV/c.

The flow coefficients v_n are measured using the event plane method ($v_n\{\text{EP}\}$ [1]) and the four-particle cumulant technique ($v_n\{4\}$ [36]), which have different sensitivity to flow fluctuations and correlations unrelated to the azimuthal asymmetry in the initial geometry (“non-flow”). The non-flow contribution to $v_n\{4\}$ is estimated to be negligible from analytic calculations and Monte Carlo simulations [37–39]. The contribution from flow fluctuations was shown to be negative for $v_n\{4\}$ and positive for $v_n\{\text{EP}\}$ [1].

The orientation of the symmetry planes Ψ_n is estimated with the event plane angle determined from the azimuthal distribution of hits measured by the VZERO scintillators. The corresponding event plane resolution is estimated from correlations between event planes determined in the TPC and the two VZERO detectors. The large gap in pseudo-rapidity between the charged particles in the TPC and those in the VZERO detectors greatly suppresses non-

flow contributions, which largely come from the inter-jet correlations and resonance decays and are narrow in rapidity. An estimate of the remaining non-flow contributions is obtained by rescaling the correlation measured in pp collisions under the assumption that it scales inversely proportional to the total multiplicity. It was observed that the two-particle azimuthal correlations in pp and the most peripheral Au–Au collisions at $\sqrt{s_{\text{NN}}} = 0.2$ TeV are very similar [40], which suggests that non-flow dominates correlations in the centrality range 80–90%. The systematic uncertainty from the remaining non-flow, δ_n^{cent} , in the measured $v_n\{\text{EP}\}$ coefficients was estimated based on the equation:

$$\delta_n^{\text{cent}} = v_n^{80-90\%} \sqrt{\frac{M^{80-90\%}}{M^{\text{cent}}}}, \quad (2)$$

where $v_n^{80-90\%}$ and $M^{80-90\%}$ are the magnitude of v_n and average multiplicity for the centrality range 80–90%, respectively, and M^{cent} is the average multiplicity in a given centrality class. The non-flow increases with p_T and from central to peripheral collisions. For example, the non-flow contributions to v_2 in 5–10% (40–50%) most central collisions are about 1% (2%) at $p_T = 1$ GeV/c and reach up to 10% (12%) for $p_T > 10$ GeV/c. Other sources of systematic uncertainties were evaluated from the variation of the results with different cuts on the reconstructed collision vertex and the centrality estimated from the charged particle multiplicity measured in the TPC and VZERO detectors. Changes due to variations of the track selection criteria and the difference of the results obtained using only positively or negatively charged particles were considered as a part of the systematic error. The difference in the extracted coefficients using one or the other of the two VZERO detectors was found to be below 1% for v_2 and v_3 , and below 5% for v_4 over the measured region of transverse momentum. The combined results from correlations with both VZERO detectors are denoted as $v_n\{\text{EP}, |\Delta\eta| > 2.0\}$ in the following. The contributions from all sources were added in quadrature as an estimate of the total systematic uncertainty. The resulting systematic uncertainties in v_2 are 3% for $0.9 < p_T < 1$ GeV/c and ${}_{-11}^{+3}\%$ (${}_{-12}^{+3}\%$) for $9 < p_T < 10$ GeV/c in the 5–10% (40–50%) centrality class. The resulting systematic uncertainties in v_3 are 3% for $0.9 < p_T < 1$ GeV/c and increase to 6% (10%) for $7 < p_T < 9$ GeV/c for centrality 5–10% (40–50%). We assign an 8% (16%) systematic uncertainty to v_4 for $0.9 < p_T < 1$ GeV/c in the 5–10% (40–50%) centrality class, while for $p_T > 6$ GeV/c the systematic uncertainty is dominated by non-flow contributions.

Fig. 1 shows unidentified charged particle v_2 , v_3 , and v_4 as a function of transverse momentum for different centrality classes. The difference between $v_2\{\text{EP}\}$ and $v_2\{4\}$ for $p_T < 7$ GeV/c is predominantly due to flow fluctuations. The measured v_2 at $p_T > 8$ GeV/c is non-zero, positive and approximately constant, while its value increases from central to mid-peripheral collisions. In the 20–50% centrality range, the observed $v_2\{\text{EP}\}$ at $p_T > 10$ GeV/c is fairly well described by extrapolation to the LHC energy [41] of the WHDG model calculations [42] for v_2 of neutral pions including collisional and radiative energy loss of partons in a Bjorken-expanding medium [43]. The coefficient v_3 exhibits a weak centrality dependence with a magnitude significantly smaller than that of v_2 , except for the most central collisions. Unlike v_3 , which originates entirely from fluctuations of the initial geometry of the system, v_4 has two contributions, which are probed by correlations with the Ψ_2 and Ψ_4 symmetry planes. The measured $v_{4/\Psi_4}\{\text{EP}\}$ does not depend strongly on the collision centrality which points to a strong contribution from flow fluctuations. In contrast, $v_{4/\Psi_2}\{\text{EP}\}$ shows a strong centrality dependence which is typical for correlations with respect to the true reaction

¹ In this analysis we do not differentiate between particle and antiparticle.




Fig. 1. (Color online.) v_2 , v_3 , and v_4 measured for unidentified charged particles as a function of transverse momentum for various centrality classes. The dashed line represents the WHDG model calculations for neutral pions v_2 [43] extrapolated to the LHC collision energy. For clarity, the markers for v_3 and v_4/ψ_4 results are slightly shifted along the horizontal axis. Note that the highest p_T data point for v_4/ψ_4 in 5–10% centrality is out of the plotting range. Error bars (shaded boxes) represent the statistical (systematic) uncertainties.




Fig. 2. (Color online.) Comparison of the ALICE results on $v_n(p_T)$ obtained with the event plane method to the analogous measurements from ATLAS [26] and CMS [27] collaborations, as well as v_2 measurements by STAR [44]. Only statistical errors are shown.

plane. The difference between the two, indicative of flow fluctuations, persists at least up to $p_T = 8$ GeV/c.

Fig. 2 compares our results obtained with the event plane method for 30–40% centrality to the analogous measurements by ATLAS [26] and CMS [27] collaborations, and results obtained at RHIC by the STAR Collaboration [44]. An excellent agreement is observed between results from all three LHC experiments. $v_2(p_T)$ at top RHIC energy has a peak value about 10% lower than at LHC although it is very similar in shape.

To investigate further the role of flow fluctuations at different transverse momenta we study the relative difference between $v_2\{EP\}$ and $v_2\{4\}$, $[(v_2\{EP\}^2 - v_2\{4\}^2)/(v_2\{EP\}^2 + v_2\{4\}^2)]^{1/2}$, which for small non-flow is proportional to the relative flow fluctua-

tions $\sigma_{v_2}/\langle v_2 \rangle$ [1]. Fig. 3 presents this quantity as a function of transverse momentum for various centrality classes. The relative flow fluctuations are minimal for mid-central collisions and become larger for peripheral and central collisions, similar to those observed at RHIC energies [1]. It is remarkable that in the 5–30% centrality range, relative flow fluctuations are within errors independent of momentum up to $p_T \sim 8$ GeV/c, far beyond the region where the flow magnitude is well described by hydrodynamic models ($p_T < 2-3$ GeV/c). This indicates a common origin for flow fluctuations, which are usually associated with fluctuations of the initial collision geometry, at least up to the regime where hard scattering and jet energy loss are expected to dominate. The ratio develops a momentum dependence, starting to




Fig. 3. (Color online.) Relative event-by-event elliptic flow fluctuations for unidentified charged particles versus transverse momentum for different centrality classes. For clarity, the markers for centrality classes $\geq 10\%$ are slightly shifted along the horizontal axis. Error bars (shaded boxes) represent the statistical (systematic) uncertainties.




Fig. 4. (Color online.) Unidentified charged particle v_2 , v_3 , and v_4 integrated over the transverse momentum range $10 < p_T < 20$ GeV/c as a function of collision centrality, with the more central (peripheral) collisions shown on the left-(right)-hand side, respectively. The dashed line represents the WHDG model calculations for neutral pions [43] extrapolated to the LHC collision energy. Error bars (shaded boxes) represent the statistical (systematic) uncertainties.

increase at $p_T \sim 1.5$ GeV/c, for more peripheral collisions (30–50%), and in most central collisions (0–5%), where it is most pronounced. In both cases, the relative contribution of non-flow effects is expected to be the largest.

Fig. 4 shows unidentified charged particle v_2 , v_3 , and v_4 averaged over $10 < p_T < 20$ GeV/c as a function of centrality. v_2 increases from central to peripheral collisions. No significant difference between $v_2\{EP\}$ and $v_2\{4\}$ results is observed, which might indicate that the fluctuations of the initial collision geometry become unimportant for $p_T > 10$ GeV/c. The centrality dependence of v_3 differs significantly from that of v_2 . v_4 measured with respect to the second and fourth harmonic symmetry planes is consistent with zero within relatively large uncertainties. All these observations indicate that for $p_T > 10$ GeV/c the effect of fluctuations of the initial collision geometry on the final momentum anisotropy might be very different compared to that at low and intermediate p_T .

Fig. 5 presents charged pion and proton v_2 and v_3 as a function of p_T in the 10–50% centrality range from the event plane method. The proton v_2 and v_3 are higher than that of pions out to $p_T = 8$ GeV/c where the uncertainties become large. This behavior is qualitatively consistent with a picture where particle production in this intermediate p_T region includes interaction of jet fragments with bulk matter, e.g. as in model [45]. The magnitude of the measured charged pion elliptic flow for $p_T > 8$ GeV/c is compatible with that for unidentified charged particles, and π^0 measured by PHENIX [46] in Au–Au collisions at $\sqrt{s_{NN}} = 0.2$ TeV, and reproduced by the WHDG model calculations for v_2 of neutral pions [43]. The unidentified charged particle, pion, and proton v_3 are the same within uncertainties for $p_T > 8$ GeV/c.

In summary, we have presented elliptic, triangular, and quadrangular flow coefficients measured by the ALICE Collaboration in Pb–Pb collisions at $\sqrt{s_{NN}} = 2.76$ TeV over a broad range of




Fig. 5. (Color online.) v_2 (top) and v_3 (bottom) of charged pion and proton as a function of transverse momentum for 10–50% centrality class compared to unidentified charged particles results from the event plane method. For clarity, the markers for v_2 and v_3 at $p_T > 8$ GeV/c are slightly shifted along the horizontal axis. PHENIX π^0 v_2 measurements [46] are also shown. The dashed line represents the WHDG model calculations for neutral pions [43] extrapolated to the LHC collision energy for the 20–50% centrality range. Error bars (shaded boxes) represent the statistical (systematic) uncertainties.

transverse momentum. For $p_T > 8$ GeV/c, we find that the unidentified charged particle v_2 in 0–70% and v_3 in 0–20% centrality ranges are finite, positive and only weakly dependent on transverse momentum, while v_3 for 20–50% and v_4 for 5–50% centrality are consistent with zero within rather large statistical and systematic uncertainties. The observed difference in the centrality dependence of v_4/ψ_4 and v_4/ψ_2 , and the results on v_2 obtained with the event plane and four-particle cumulant methods indicate that the effect of flow fluctuations extends at least up to $p_T = 8$ GeV/c and does not change significantly in magnitude, except for very central collisions. It shows that the effect of fluctuations of the initial collision geometry on particle production is similar at low and intermediate p_T regions, which are considered to be dominated by hydrodynamical flow and quark coalescence, respectively. For $p_T > 10$ GeV/c, where particle production is dominated by fragmentation of hard partons, the response to fluctuations of the initial collision geometry might be different, but more data is needed to study this regime in more detail. The pion v_2 at LHC energies is very close to that measured at RHIC out to $p_T = 16$ GeV/c and is reproduced by WHDG model calculations for $p_T > 8$ GeV/c. The proton v_2 and v_3 are finite, positive, and have a larger magnitude than that of the pion for $p_T < 8$ GeV/c, indicating that the particle type dependence, which is typical at low p_T , persists out to intermediate transverse momenta. The pion

and proton v_3 are consistent with zero within uncertainties for $p_T > 8$ GeV/c.

Acknowledgements

The ALICE Collaboration would like to thank all its engineers and technicians for their invaluable contributions to the construction of the experiment and the CERN accelerator teams for the outstanding performance of the LHC complex.

The ALICE Collaboration acknowledges the following funding agencies for their support in building and running the ALICE detector:

Calouste Gulbenkian Foundation from Lisbon and Swiss Fonds Kidagan, Armenia;

Conselho Nacional de Desenvolvimento Científico e Tecnológico (CNPq), Financiadora de Estudos e Projetos (FINEP), Fundação de Amparo à Pesquisa do Estado de São Paulo (FAPESP);

National Natural Science Foundation of China (NSFC), the Chinese Ministry of Education (CMOE) and the Ministry of Science and Technology of China (MSTC);

Ministry of Education and Youth of the Czech Republic;

Danish Natural Science Research Council, the Carlsberg Foundation and the Danish National Research Foundation;

The European Research Council under the European Community's Seventh Framework Programme;

Helsinki Institute of Physics and the Academy of Finland;

French CNRS-IN2P3, the 'Region Pays de Loire', 'Region Alsace', 'Region Auvergne' and CEA, France;

German BMBF and the Helmholtz Association;

General Secretariat for Research and Technology, Ministry of Development, Greece;

Hungarian OTKA and National Office for Research and Technology (NKTH);

Department of Atomic Energy and Department of Science and Technology of the Government of India;

Istituto Nazionale di Fisica Nucleare (INFN) of Italy;

MEXT Grant-in-Aid for Specially Promoted Research, Japan;

Joint Institute for Nuclear Research, Dubna;

National Research Foundation of Korea (NRF);

CONACYT, DGAPA, México, ALFA-EC and the HELEN Program (High-Energy physics Latin-American-European Network);

Stichting voor Fundamenteel Onderzoek der Materie (FOM) and the Nederlandse Organisatie voor Wetenschappelijk Onderzoek (NWO), Netherlands;

Research Council of Norway (NFR);

Polish Ministry of Science and Higher Education;

National Authority for Scientific Research – NASR (Autoritatea Națională pentru Cercetare Științifică – ANCS);

Federal Agency of Science of the Ministry of Education and Science of Russian Federation, International Science and Technology Center, Russian Academy of Sciences, Russian Federal Agency of Atomic Energy, Russian Federal Agency for Science and Innovations and CERN-INTAS;

Ministry of Education of Slovakia;

Department of Science and Technology, South Africa;

CIEMAT, EELA, Ministerio de Educación y Ciencia of Spain, Xunta de Galicia (Consellería de Educación), CEADEN, Cubaenergía, Cuba, and IAEA (International Atomic Energy Agency);

Swedish Research Council (VR) and Knut & Alice Wallenberg Foundation (KAW);

Ukraine Ministry of Education and Science;

United Kingdom Science and Technology Facilities Council (STFC);

The United States Department of Energy, the United States National Science Foundation, the State of Texas, and the State of Ohio.

Open access

This article is published Open Access at sciencedirect.com. It is distributed under the terms of the Creative Commons Attribution License 3.0, which permits unrestricted use, distribution, and reproduction in any medium, provided the original authors and source are credited.

References

- [1] S.A. Voloshin, A.M. Poskanzer, R. Snellings, in: *Relativistic Heavy Ion Physics*, in: Landolt-Boernstein, vol. 1/23, Springer-Verlag, 2010, pp. 5–54.
- [2] R. Snellings, STAR Collaboration, ALICE Collaboration, *Eur. Phys. J. C* 49 (2007) 87.
- [3] B.I. Abelev, et al., STAR Collaboration, *Phys. Rev. Lett.* 99 (2007) 112301.
- [4] A. Adare, et al., PHENIX Collaboration, arXiv:1203.2644 [nucl-ex].
- [5] M. Gyulassy, M. Plumer, *Phys. Lett. B* 243 (1990) 432.
- [6] X.N. Wang, M. Gyulassy, *Phys. Rev. Lett.* 68 (1992) 1480.
- [7] M. Gyulassy, X.N. Wang, *Nucl. Phys. B* 420 (1994) 583.
- [8] R.J.M. Snellings, A.M. Poskanzer, S.A. Voloshin, arXiv:nucl-ex/9904003.
- [9] X.N. Wang, *Phys. Rev. C* 63 (2001) 054902.
- [10] S. Voloshin, Y. Zhang, *Z. Phys. C* 70 (1996) 665.
- [11] A.M. Poskanzer, S.A. Voloshin, *Phys. Rev. C* 58 (1998) 1671.
- [12] A.P. Mishra, R.K. Mohapatra, P.S. Saumia, A.M. Srivastava, *Phys. Rev. C* 77 (2008) 064902.
- [13] A.P. Mishra, R.K. Mohapatra, P.S. Saumia, A.M. Srivastava, *Phys. Rev. C* 81 (2010) 034903.
- [14] J. Takahashi, et al., *Phys. Rev. Lett.* 103 (2009) 242301.
- [15] B. Alver, G. Roland, *Phys. Rev. C* 81 (2010) 054905;
B. Alver, G. Roland, *Phys. Rev. C* 82 (2010) 039903 (Erratum).
- [16] B.H. Alver, C. Gombaud, M. Luzum, J.Y. Ollitrault, *Phys. Rev. C* 82 (2010) 034913.
- [17] D. Teaney, L. Yan, arXiv:1010.1876 [nucl-th].
- [18] M. Luzum, *Phys. Lett. B* 696 (2011) 499.
- [19] K.H. Ackermann, et al., STAR Collaboration, *Phys. Rev. Lett.* 86 (2001) 402.
- [20] S.S. Adler, et al., PHENIX Collaboration, *Phys. Rev. Lett.* 91 (2003) 182301.
- [21] A. Adare, et al., PHENIX Collaboration, *Phys. Rev. Lett.* 107 (2011) 252301.
- [22] K. Aamodt, et al., ALICE Collaboration, *Phys. Rev. Lett.* 105 (2010) 252302.
- [23] K. Aamodt, et al., ALICE Collaboration, *Phys. Rev. Lett.* 107 (2011) 032301.
- [24] G. Aad, et al., ATLAS Collaboration, *Phys. Lett. B* 707 (2012) 330.
- [25] S. Chatrchyan, et al., CMS Collaboration, arXiv:1201.3158 [nucl-ex].
- [26] G. Aad, et al., ATLAS Collaboration, arXiv:1203.3087 [hep-ex].
- [27] S. Chatrchyan, et al., CMS Collaboration, arXiv:1204.1409 [nucl-ex].
- [28] S. Chatrchyan, et al., CMS Collaboration, arXiv:1204.1850 [nucl-ex].
- [29] ALICE Collaboration, *JINST* 3 (2008) S08002.
- [30] ALICE Collaboration, *J. Phys. G* 30 (2004) 1517.
- [31] ALICE Collaboration, *J. Phys. G* 32 (2006) 1295.
- [32] K. Aamodt, et al., ALICE Collaboration, *Phys. Rev. Lett.* 106 (2011) 032301.
- [33] M. Gyulassy, X.-N. Wang, *Comput. Phys. Commun.* 83 (1994) 307;
X.N. Wang, M. Gyulassy, *Phys. Rev. D* 44 (1991) 3501.
- [34] R. Brun, et al., CERN Program Library Long Write-up, W5013, GEANT Detector Description and Simulation Tool, 1994.
- [35] K. Aamodt, et al., ALICE Collaboration, *Eur. Phys. J. C* 71 (2011) 1655.
- [36] A. Bilandzic, R. Snellings, S. Voloshin, *Phys. Rev. C* 83 (2011) 044913.
- [37] C. Adler, et al., STAR Collaboration, *Phys. Rev. C* 66 (2002) 034904.
- [38] N.K. Pruthi, STAR Collaboration, arXiv:1111.5637 [nucl-ex].
- [39] A. Bilandzic, Ph.D. thesis, Utrecht University, 2012, available at <http://aliweb.cern.ch/node/21417>.
- [40] J. Adams, et al., STAR Collaboration, *Phys. Rev. Lett.* 93 (2004) 252301.
- [41] W.A. Horowitz, M. Gyulassy, *Nucl. Phys. A* 872 (2011) 265.
- [42] S. Wicks, W. Horowitz, M. Djordjevic, M. Gyulassy, *Nucl. Phys. A* 784 (2007) 426.
- [43] W.A. Horowitz, M. Gyulassy, *J. Phys. G* 38 (2011) 124114.
- [44] J. Adams, et al., STAR Collaboration, *Phys. Rev. C* 72 (2005) 014904.
- [45] K. Werner, I. Karpenko, M. Bleicher, T. Pierog, S. Porteboeuf-Houssais, arXiv:1203.5704 [nucl-th];
K. Werner, arXiv:1205.3379 [nucl-th].
- [46] A. Adare, et al., PHENIX Collaboration, *Phys. Rev. Lett.* 105 (2010) 142301.

ALICE Collaboration

B. Abelev¹²⁵, J. Adam⁷⁵, D. Adamová¹⁰², A.M. Adare¹¹⁹, M.M. Aggarwal⁸⁷, G. Aglieri Rinella²⁵, A.G. Agocs¹²⁶, A. Agostinelli⁴⁷, S. Aguilar Salazar¹⁰³, Z. Ahammed⁵², A. Ahmad Masoodi²⁶, N. Ahmad²⁶, S.U. Ahn⁵⁴, A. Akindinov²⁸, D. Aleksandrov¹¹⁵, B. Alessandro⁹², R. Alfaro Molina¹⁰³, A. Alici^{60,i}, A. Alkin¹⁰⁵, E. Almaráz Aviña¹⁰³, J. Alme¹⁸, T. Alt⁵¹, V. Altini⁹⁷, S. Altinpinar⁸³, I. Altsybeev⁵⁸, C. Andrei⁴⁹, A. Andronic⁷³, V. Anguelov¹¹, J. Anielski⁹⁹, T. Antičić¹⁵, F. Antinori¹⁰⁴, P. Antonioli¹¹⁷, L. Aphecetche⁷⁸, H. Appelshäuser²⁰, N. Arbor⁵⁶, S. Arcelli⁴⁷, N. Armesto¹⁰⁶, R. Arnaldi⁹², T. Aronsson¹¹⁹, I.C. Arsene⁷³, M. Arslandok²⁰, A. Augustinus²⁵, R. Averbeck⁷³, T.C. Awes¹¹¹, J. Äystö²⁹, M.D. Azmi²⁶, M. Bach⁵¹, A. Badalà⁷⁴, Y.W. Baek^{5,ii}, R. Bailhache²⁰, R. Bala⁹², R. Baldini Ferroli⁶⁰, A. Baldisseri¹²³, A. Baldit⁵, F. Baltasar Dos Santos Pedrosa²⁵, J. Bán⁸⁴, R.C. Baral⁹¹, R. Barbera⁸¹, F. Barile⁹⁷, G.G. Barnaföldi¹²⁶, L.S. Barnby³¹, V. Barret⁵, J. Bartke¹⁴, M. Basile⁴⁷, N. Bastid⁵, S. Basu⁵², B. Bathen⁹⁹, G. Batigne⁷⁸, B. Batyunya³⁸, C. Baumann²⁰, I.G. Bearden¹¹⁶, H. Beck²⁰, I. Belikov⁹⁸, F. Bellini⁴⁷, R. Bellwied⁶⁹, E. Belmont-Moreno¹⁰³, G. Bencedi¹²⁶, S. Beole²⁴, I. Berceau⁴⁹, A. Bercuci⁴⁹, Y. Berdnikov⁸⁸, D. Berenyi¹²⁶, A.A.E. Bergognon⁷⁸, D. Berzano⁹², L. Betev²⁵, A. Bhasin³⁷, A.K. Bhati⁸⁷, J. Bhom⁶², L. Bianchi²⁴, N. Bianchi⁹, C. Bianchin³, J. Bielčik⁷⁵, J. Bielčíková¹⁰², A. Bilandzic¹¹⁶, S. Bjelogrić², F. Blanco⁵⁵, F. Blanco⁶⁹, D. Blau¹¹⁵, C. Blume²⁰, N. Bock⁵⁹, S. Böttger⁵⁷, A. Bogdanov¹⁶, H. Bøggild¹¹⁶, M. Bogolyubsky⁴⁵, L. Boldizsár¹²⁶, M. Bombara¹³, J. Book²⁰, H. Borel¹²³, A. Borissov⁹⁴, S. Bose¹², F. Bossú²⁴, M. Botje³², B. Boyer³⁵, E. Braidot¹⁰, P. Braun-Munzinger⁷³, M. Bregant⁷⁸, T. Breitner⁵⁷, T.A. Browning⁹⁶, M. Broz⁶³, R. Brun²⁵, E. Bruna⁹², G.E. Bruno⁹⁷, D. Budnikov⁹³, H. Buesching²⁰, S. Bufalino⁹², K. Bugaiev¹⁰⁵, O. Busch¹¹, Z. Buthelezi⁸⁹, D. Caffarri³, X. Cai⁶⁶, H. Caines¹¹⁹, E. Calvo Villar³³, P. Camerini³⁰, V. Canoa Roman⁴⁸, G. Cara Romeo¹¹⁷, F. Carena²⁵, W. Carena²⁵, F. Carminati²⁵, A. Casanova Díaz⁹, J. Castillo Castellanos¹²³, E.A.R. Casula⁶¹, V. Catanescu⁴⁹, C. Cavicchioli²⁵, C. Ceballos Sanchez⁷¹, J. Cepila⁷⁵, P. Cerello⁹², B. Chang²⁹, S. Chapeland²⁵, J.L. Charvet¹²³, S. Chattopadhyay¹², S. Chattopadhyay⁵², I. Chawla⁸⁷, M. Cherney¹¹², C. Cheshkov³⁹, B. Cheynis³⁹, E. Chiavassa⁹², V. Chibante Barroso²⁵, D.D. Chinellato¹¹³, P. Chochula²⁵, M. Chojnacki², S. Choudhury⁵², P. Christakoglou³², C.H. Christensen¹¹⁶, P. Christiansen¹⁰⁷, T. Chujo⁶², S.U. Chung⁷⁶, C. Cicalo²⁷, L. Cifarelli⁴⁷, F. Cindolo¹¹⁷, J. Cleymans⁸⁹, F. Coccetti⁶⁰, F. Colamaria⁹⁷,

D. Colella⁹⁷, G. Conesa Balbastre⁵⁶, Z. Conesa del Valle²⁵, P. Constantin¹¹, G. Contin³⁰, J.G. Contreras⁴⁸,
 T.M. Cormier⁹⁴, Y. Corrales Morales²⁴, I. Cortés Maldonado⁷⁷, P. Cortese¹¹⁸, M.R. Cosentino¹⁰,
 F. Costa²⁵, M.E. Cotallo⁵⁵, P. Crochet⁵, E. Cruz Alaniz¹⁰³, E. Cuautle⁷², L. Cunqueiro⁹, G. D’Erasmus⁹⁷,
 A. Dainese¹⁰⁴, H.H. Dalsgaard¹¹⁶, A. Danu³⁶, D. Das¹², I. Das³⁵, K. Das¹², A. Dash¹¹³, S. Dash¹²⁴,
 S. De⁵², G.O.V. de Barros⁴⁶, A. De Caro^{60,iii}, G. de Cataldo⁹⁰, J. de Cuveland⁵¹, A. De Falco⁶¹,
 D. De Gruttola⁴², N. De Marco⁹², S. De Pasquale⁴², R. de Rooij², H. Delagrange⁷⁸, A. Deloff⁸²,
 V. Demanov⁹³, E. Dénes¹²⁶, A. Deppman⁴⁶, D. Di Bari⁹⁷, C. Di Giglio⁹⁷, S. Di Liberto¹⁰⁰, A. Di Mauro²⁵,
 P. Di Nezza⁹, M.A. Diaz Corchero⁵⁵, T. Dietel⁹⁹, R. Divià²⁵, Ø. Djuvslund⁸³, A. Dobrin^{94,*},
 T. Dobrowolski⁸², I. Domínguez⁷², B. Dönigus⁷³, O. Dordic²¹, O. Driga⁷⁸, A.K. Dubey⁵², L. Ducroux³⁹,
 P. Dupieux⁵, A.K. Dutta Majumdar¹², M.R. Dutta Majumdar⁵², D. Elia⁹⁰, D. Emschermann⁹⁹, H. Engel⁵⁷,
 H.A. Erdal¹⁸, B. Espagnon³⁵, M. Estienne⁷⁸, S. Esumi⁶², D. Evans³¹, G. Eyyubova²¹, D. Fabris¹⁰⁴,
 J. Faivre⁵⁶, D. Falchieri⁴⁷, A. Fantoni⁹, M. Fasel⁷³, R. Fearick⁸⁹, A. Fedunov³⁸, D. Fehlker⁸³,
 L. Feldkamp⁹⁹, D. Felea³⁶, B. Fenton-Olsen¹⁰, G. Feofilov⁵⁸, A. Fernández Téllez⁷⁷, A. Ferretti²⁴,
 R. Ferretti¹¹⁸, J. Figiel¹⁴, M.A.S. Figueredo⁴⁶, S. Filchagin⁹³, D. Finogeev⁸⁰, F.M. Fionda⁹⁷, E.M. Fiore⁹⁷,
 M. Floris²⁵, S. Foertsch⁸⁹, P. Foka⁷³, S. Fokin¹¹⁵, E. Fragiaco¹⁰⁸, U. Frankenfeld⁷³, U. Fuchs²⁵,
 C. Furget⁵⁶, M. Fusco Girard⁴², J.J. Gaardhøje¹¹⁶, M. Gagliardi²⁴, A. Gago³³, M. Gallio²⁴,
 D.R. Gangadharan⁵⁹, P. Ganoti¹¹¹, C. Garabatos⁷³, E. Garcia-Solis¹²⁰, I. Garishvili¹²⁵, J. Gerhard⁵¹,
 M. Germain⁷⁸, C. Geuna¹²³, A. Gheata²⁵, M. Gheata^{25,iv}, B. Ghidini⁹⁷, P. Ghosh⁵², P. Gianotti⁹,
 M.R. Girard⁹⁵, P. Giubellino²⁵, E. Gladysz-Dziadus¹⁴, P. Glässel¹¹, R. Gomez⁶⁷, A. Gonschior⁷³,
 E.G. Ferreira¹⁰⁶, L.H. González-Trueba¹⁰³, P. González-Zamora⁵⁵, S. Gorbunov⁵¹, A. Goswami⁵³,
 S. Gotovac¹¹⁴, V. Grabski¹⁰³, L.K. Graczykowski⁹⁵, R. Grajcarek¹¹, A. Grelli², A. Grigoras²⁵,
 C. Grigoras²⁵, V. Grigoriev¹⁶, A. Grigoryan⁷, S. Grigoryan³⁸, B. Grinyov¹⁰⁵, N. Grion¹⁰⁸,
 J.F. Grosse-Oetringhaus²⁵, J.-Y. Grossiord³⁹, R. Grosso²⁵, F. Guber⁸⁰, R. Guernane⁵⁶,
 C. Guerra Gutierrez³³, B. Guerzoni⁴⁷, M. Guilbaud³⁹, K. Gulbrandsen¹¹⁶, T. Gunji²², A. Gupta³⁷,
 R. Gupta³⁷, H. Gutbrod⁷³, Ø. Haaland⁸³, C. Hadjidakis³⁵, M. Haiduc³⁶, H. Hamagaki²², G. Hamar¹²⁶,
 L.D. Hanratty³¹, A. Hansen¹¹⁶, Z. Harmanova¹³, J.W. Harris¹¹⁹, M. Hartig²⁰, D. Hasegan³⁶,
 D. Hatzifotiadou¹¹⁷, A. Hayrapetyan^{7,v}, S.T. Heckel²⁰, M. Heide⁹⁹, H. Helstrup¹⁸, A. Herghelegiu⁴⁹,
 G. Herrera Corral⁴⁸, N. Herrmann¹¹, B.A. Hess¹, K.F. Hetland¹⁸, B. Hicks¹¹⁹, P.T. Hille¹¹⁹, B. Hippolyte⁹⁸,
 T. Horaguchi⁶², Y. Hori²², P. Hristov²⁵, I. Hřivnáčová³⁵, M. Huang⁸³, T.J. Humanic⁵⁹, D.S. Hwang⁶⁵,
 R. Ichou⁵, R. Ilkaev⁹³, I. Ilkiv⁸², M. Inaba⁶², E. Incani⁶¹, G.M. Innocenti²⁴, M. Ippolitov¹¹⁵,
 M. Irfan²⁶, C. Ivan⁷³, A. Ivanov⁵⁸, M. Ivanov⁷³, V. Ivanov⁸⁸, O. Ivanytskyi¹⁰⁵, A. Jachołkowski²⁵,
 P.M. Jacobs¹⁰, S. Jangal⁹⁸, M.A. Janik⁹⁵, R. Janik⁶³, P.H.S.Y. Jayarathna⁶⁹, S. Jena¹²⁴, D.M. Jha⁹⁴,
 R.T. Jimenez Bustamante⁷², L. Jiriden²⁵, P.G. Jones³¹, H. Jung⁵⁴, A. Jusko³¹, V. Kakoyan⁷, S. Kalcher⁵¹,
 P. Kaliňák⁸⁴, T. Kalliokoski²⁹, A. Kalweit⁸⁶, K. Kanaki⁸³, J.H. Kang⁶, V. Kaplin¹⁶, A. Karasu Uysal²⁵,
 O. Karavichev⁸⁰, T. Karavicheva⁸⁰, E. Karpechev⁸⁰, A. Kazantsev¹¹⁵, U. Kebschull⁵⁷, R. Keidel⁶⁸,
 M.M. Khan²⁶, P. Khan¹², S.A. Khan⁵², A. Khanzadeev⁸⁸, Y. Kharlov⁴⁵, B. Kileng¹⁸, B. Kim⁶, D.J. Kim²⁹,
 D.W. Kim⁵⁴, J.H. Kim⁶⁵, J.S. Kim⁵⁴, M. Kim⁶, M. Kim⁵⁴, S. Kim⁶⁵, S.H. Kim⁵⁴, T. Kim⁶, S. Kirsch⁵¹,
 I. Kisel⁵¹, S. Kiselev²⁸, A. Kisiel⁹⁵, J.L. Klay⁴⁰, J. Klein¹¹, C. Klein-Bösing⁹⁹, A. Kluge²⁵, M.L. Knichel⁷³,
 A.G. Knospe⁷⁹, K. Koch¹¹, M.K. Köhler⁷³, A. Kolojvari⁵⁸, V. Kondratiev⁵⁸, N. Kondratyeva¹⁶,
 A. Konevskikh⁸⁰, A. Korneev⁹³, R. Kour³¹, M. Kowalski¹⁴, S. Kox⁵⁶, G. Koyithatta Meethalevedu¹²⁴,
 J. Kral²⁹, I. Králik⁸⁴, F. Kramer²⁰, I. Kraus⁷³, T. Krawutschke^{11,vi}, M. Krelina⁷⁵, M. Kretz⁵¹,
 M. Krivda^{31,vii}, F. Krizek²⁹, M. Krus⁷⁵, E. Kryshen⁸⁸, M. Krzewicki⁷³, Y. Kucheriaev¹¹⁵, C. Kuhn⁹⁸,
 P.G. Kuijer³², I. Kulakov²⁰, P. Kurashvili⁸², A. Kurepin⁸⁰, A.B. Kurepin⁸⁰, A. Kuryakin⁹³, S. Kuschpil¹⁰²,
 V. Kuschpil¹⁰², M.J. Kweon¹¹, Y. Kwon⁶, S.L. La Pointe², P. La Rocca⁸¹, P. Ladrón de Guevara⁷²,
 I. Lakomov³⁵, R. Langoy⁸³, C. Lara⁵⁷, A. Lardeux⁷⁸, C. Lazzeroni³¹, Y. Le Bornec³⁵, R. Lea³⁰,
 M. Lechman²⁵, G.R. Lee³¹, K.S. Lee⁵⁴, S.C. Lee⁵⁴, F. Lefèvre⁷⁸, J. Lehnert²⁰, L. Leistam²⁵,
 R.C. Lemmon¹⁹, M. Lenhardt⁷⁸, V. Lenti⁹⁰, I. León Monzón⁶⁷, H. León Vargas²⁰, M. Leoncino⁹²,
 P. Lévai¹²⁶, J. Lien⁸³, R. Lietava³¹, S. Lindal²¹, V. Lindenstruth⁵¹, C. Lippmann⁷³, M.A. Lisa⁵⁹, L. Liu⁸³,
 P.I. Loenne⁸³, V.R. Loggins⁹⁴, V. Loginov¹⁶, S. Lohn²⁵, D. Lohner¹¹, C. Loizides¹⁰, K.K. Loo²⁹, X. Lopez⁵,
 E. López Torres⁷¹, G. Løvhøiden²¹, X.-G. Lu¹¹, P. Luettig²⁰, M. Lunardon³, J. Luo⁶⁶, G. Luparello²,
 L. Luquin⁷⁸, C. Luzzi²⁵, R. Ma¹¹⁹, A. Maevskaya⁸⁰, M. Mager²⁵, D.P. Mahapatra⁹¹, A. Maire¹¹,
 D. Mal’Kevich²⁸, M. Malaev⁸⁸, I. Maldonado Cervantes⁷², L. Malinina³⁸, P. Malzacher⁷³, A. Mamonov⁹³,

L. Manceau⁹², V. Manko¹¹⁵, F. Manso⁵, V. Manzari⁹⁰, Y. Mao⁶⁶, M. Marchisone^{24,viii}, J. Mareš²³, G.V. Margagliotti³⁰, A. Margotti¹¹⁷, A. Marín⁷³, C.A. Marin Tobon²⁵, C. Markert⁷⁹, I. Martashvili⁴⁴, P. Martinengo²⁵, M.I. Martínez⁷⁷, A. Martínez Davalos¹⁰³, G. Martínez García⁷⁸, Y. Martynov¹⁰⁵, A. Mas⁷⁸, S. Masciocchi⁷³, M. Maserà²⁴, A. Masoni²⁷, M. Mastromarco⁹⁰, A. Mastroserio⁹⁷, Z.L. Matthews³¹, A. Matyja^{78,ix}, D. Mayani⁷², C. Mayer¹⁴, J. Mazer⁴⁴, M.A. Mazzoni¹⁰⁰, F. Meddi⁴³, A. Menchaca-Rocha¹⁰³, J. Mercado Pérez¹¹, M. Meres⁶³, Y. Miake⁶², L. Milano²⁴, J. Milosevic²¹, A. Mischke², A.N. Mishra⁵³, D. Miśkowiec^{25,x}, C. Mitu³⁶, J. Mlynarz⁹⁴, A.K. Mohanty²⁵, B. Mohanty⁵², L. Molnar²⁵, L. Montaño Zetina⁴⁸, M. Monteno⁹², E. Montes⁵⁵, T. Moon⁶, M. Morando³, D.A. Moreira De Godoy⁴⁶, S. Moretto³, A. Morsch²⁵, V. Muccifora⁹, E. Mudnic¹¹⁴, S. Muhuri⁵², M. Mukherjee⁵², H. Müller²⁵, M.G. Munhoz⁴⁶, L. Musa²⁵, A. Musso⁹², B.K. Nandi¹²⁴, R. Nania¹¹⁷, E. Nappi⁹⁰, C. Natrass⁴⁴, N.P. Naumov⁹³, S. Navin³¹, T.K. Nayak⁵², S. Nazarenko⁹³, G. Nazarov⁹³, A. Nedosekin²⁸, M. Nicassio⁹⁷, M. Niculescu^{36,v}, B.S. Nielsen¹¹⁶, T. Niida⁶², S. Nikolaev¹¹⁵, V. Nikolic¹⁵, S. Nikulin¹¹⁵, V. Nikulin⁸⁸, B.S. Nilsen¹¹², M.S. Nilsson²¹, F. Noferini^{60,i}, P. Nomokonov³⁸, G. Nooren², N. Novitzky²⁹, A. Nyanin¹¹⁵, A. Nyatha¹²⁴, C. Nygaard¹¹⁶, J. Nystrand⁸³, H. Oeschler⁸⁶, S. Oh¹¹⁹, S.K. Oh⁵⁴, J. Oleniacz⁹⁵, C. Oppedisano⁹², G. Ortona²⁴, A. Oskarsson¹⁰⁷, J. Otwinowski⁷³, K. Oyama¹¹, Y. Pachmayer¹¹, M. Pachr⁷⁵, F. Padilla²⁴, P. Pagano⁴², G. Paić⁷², F. Painke⁵¹, C. Pajares¹⁰⁶, S. Pal¹²³, S.K. Pal⁵², A. Palaha³¹, A. Palmeri⁷⁴, V. Papikyan⁷, G.S. Pappalardo⁷⁴, W.J. Park⁷³, A. Passfeld⁹⁹, D.I. Patalakha⁴⁵, V. Paticchio⁹⁰, A. Pavlinov⁹⁴, T. Pawlak⁹⁵, T. Peitzmann², H. Pereira Da Costa¹²³, E. Pereira De Oliveira Filho⁴⁶, D. Peresunko¹¹⁵, C.E. Pérez Lara³², E. Perez Lezama⁷², D. Perini²⁵, D. Perrino⁹⁷, W. Peryt⁹⁵, A. Pesci¹¹⁷, V. Peskov²⁵, Y. Pestov⁴¹, V. Petráček⁷⁵, M. Petran⁷⁵, M. Petris⁴⁹, P. Petrov³¹, M. Petrovici⁴⁹, C. Petta⁸¹, S. Piano¹⁰⁸, A. Piccotti⁹², M. Pikna⁶³, P. Pillot⁷⁸, O. Pinazza²⁵, L. Pinsky⁶⁹, N. Pitz²⁰, F. Piuze²⁵, D.B. Piyarathna⁶⁹, M. Płoskoń¹⁰, J. Pluta⁹⁵, S. Pochybova¹²⁶, P.L.M. Podesta-Lerma⁶⁷, M.G. Poghosyan^{24,v}, B. Polichtchouk⁴⁵, A. Pop⁴⁹, S. Porteboeuf-Houssais⁵, V. Pospíšil⁷⁵, B. Potukuchi³⁷, S.K. Prasad⁹⁴, R. Preghenella^{60,i}, F. Prino⁹², C.A. Pruneau⁹⁴, I. Pshenichnov⁸⁰, S. Puchagin⁹³, G. Puddu⁶¹, P. Pujahari¹²⁴, J. Pujol Teixido⁵⁷, A. Pulvirenti⁸¹, V. Punin⁹³, M. Putiš¹³, J. Putschke⁹⁴, E. Quercigh²⁵, H. Qvigstad²¹, A. Rachevski¹⁰⁸, A. Rademakers²⁵, S. Radomski¹¹, T.S. Rähä²⁹, J. Rak²⁹, A. Rakotozafindrabe¹²³, L. Ramello¹¹⁸, A. Ramírez Reyes⁴⁸, R. Raniwala⁵³, S. Raniwala⁵³, S.S. Räsänen²⁹, B.T. Rascanu²⁰, D. Rathee⁸⁷, K.F. Read⁴⁴, J.S. Real⁵⁶, K. Redlich⁸², P. Reichelt²⁰, M. Reicher², R. Renfordt²⁰, A.R. Reolon⁹, A. Reshetin⁸⁰, F. Rettig⁵¹, J.-P. Revol²⁵, K. Reygers¹¹, L. Riccati⁹², R.A. Ricci¹⁷, T. Richert¹⁰⁷, M. Richter²¹, P. Riedler²⁵, W. Riegler²⁵, F. Riggi⁸¹, B. Rodrigues Fernandes Rabacal²⁵, M. Rodríguez Cahuantzi⁷⁷, A. Rodriguez Manso³², K. Røed⁸³, D. Rohr⁵¹, D. Röhrich⁸³, R. Romita⁷³, F. Ronchetti⁹, P. Rosnet⁵, S. Rossegger²⁵, A. Rossi^{3,v}, C. Roy⁹⁸, P. Roy¹², A.J. Rubio Montero⁵⁵, R. Rui³⁰, R. Russo²⁴, E. Ryabinkin¹¹⁵, A. Rybicki¹⁴, S. Sadovsky⁴⁵, K. Šafařík²⁵, R. Sahoo⁷⁰, P.K. Sahu⁹¹, J. Saini⁵², H. Sakaguchi¹⁰⁹, S. Sakai¹⁰, D. Sakata⁶², C.A. Salgado¹⁰⁶, J. Salzwedel⁵⁹, S. Sambyal³⁷, V. Samsonov⁸⁸, X. Sanchez Castro⁹⁸, L. Šándor⁸⁴, A. Sandoval¹⁰³, M. Sano⁶², S. Sano²², R. Santo⁹⁹, R. Santoro^{25,xi}, J. Sarkamo²⁹, E. Scapparone¹¹⁷, F. Scarlassara³, R.P. Scharenberg⁹⁶, C. Schiaua⁴⁹, R. Schicker¹¹, C. Schmidt⁷³, H.R. Schmidt¹, S. Schreiner²⁵, S. Schuchmann²⁰, J. Schukraft²⁵, Y. Schutz^{78,v}, K. Schwarz⁷³, K. Schweda⁷³, G. Scioli⁴⁷, E. Scapparini⁹², P.A. Scott³¹, R. Scott⁴⁴, G. Segato³, I. Selyuzhenkov⁷³, S. Senyukov¹¹⁸, J. Seo⁷⁶, S. Serci⁶¹, E. Serradilla^{103,xii}, A. Sevcenco³⁶, A. Shabetai⁷⁸, G. Shabratova³⁸, R. Shahoyan²⁵, N. Sharma⁸⁷, S. Sharma³⁷, K. Shigaki¹⁰⁹, M. Shimomura⁶², K. Shtejer⁷¹, Y. Sibiriyak¹¹⁵, M. Siciliano²⁴, E. Sicking²⁵, S. Siddhanta²⁷, T. Siemiarzczuk⁸², D. Silvermyr¹¹¹, C. Silvestre⁵⁶, G. Simatovic¹⁵, G. Simonetti²⁵, R. Singaraju⁵², R. Singh³⁷, S. Singha⁵², V. Singhal⁵², B.C. Sinha⁵², T. Sinha¹², B. Sitar⁶³, M. Sitta¹¹⁸, T.B. Skaali²¹, K. Skjerdal⁸³, R. Smakal⁷⁵, N. Smirnov¹¹⁹, R.J.M. Snellings², C. Søgaard¹¹⁶, R. Soltz¹²⁵, H. Son⁶⁵, J. Song⁷⁶, M. Song⁶, C. Soos²⁵, F. Soramel³, I. Sputowska¹⁴, M. Spyropoulou-Stassinaki¹⁰¹, B.K. Srivastava⁹⁶, J. Stachel¹¹, I. Stan³⁶, G. Stefanek⁸², G. Stefanini²⁵, T. Steinbeck⁵¹, M. Steinpreis⁵⁹, E. Stenlund¹⁰⁷, G. Steyn⁸⁹, J.H. Stiller¹¹, D. Stocco⁷⁸, M. Stolpovskiy⁴⁵, K. Strabykin⁹³, P. Strmen⁶³, A.A.P. Suaide⁴⁶, M.A. Subieta Vásquez²⁴, T. Sugitate¹⁰⁹, C. Suire³⁵, M. Sukhorukov⁹³, R. Sultanov²⁸, M. Šumbera¹⁰², T. Susa¹⁵, A. Szanto de Toledo⁴⁶, I. Szarka⁶³, A. Szczepankiewicz¹⁴, A. Szostak⁸³, M. Szymanski⁹⁵, J. Takahashi¹¹³, J.D. Tapia Takaki³⁵, A. Tauro²⁵, G. Tejeda Muñoz⁷⁷, A. Telesca²⁵, C. Terrevoli⁹⁷, J. Thäder⁷³, D. Thomas², R. Tieulent³⁹, A.R. Timmins⁶⁹, A. Toia⁵¹, H. Torii²², F. Tosello⁹², W.H. Trzaska²⁹, T. Tsuji²²,

A. Tumkin⁹³, R. Turrisi¹⁰⁴, T.S. Tveter²¹, J. Ulery²⁰, K. Ullaland⁸³, J. Ulrich⁵⁷, A. Uras³⁹, J. Urbán¹³, G.M. Urciuoli¹⁰⁰, G.L. Usai⁶¹, M. Vajzer¹⁰², M. Vala^{38,vii}, L. Valencia Palomo³⁵, S. Vallero¹¹, N. van der Kolk³², M. van Leeuwen², P. Vande Vyvre²⁵, L. Vannucci¹⁷, A. Vargas⁷⁷, R. Varma¹²⁴, M. Vasileiou¹⁰¹, A. Vasiliev¹¹⁵, V. Vechernin⁵⁸, M. Veldhoen², M. Venaruzzo³⁰, E. Vercellin²⁴, S. Vergara⁷⁷, R. Vernet⁵⁰, M. Verweij², L. Vickovic¹¹⁴, G. Viesti³, O. Vikhlyantsev⁹³, Z. Vilakazi⁸⁹, O. Villalobos Baillie³¹, A. Vinogradov¹¹⁵, L. Vinogradov⁵⁸, Y. Vinogradov⁹³, T. Virgili⁴², Y.P. Viyogi⁵², A. Vodopyanov³⁸, K. Voloshin²⁸, S. Voloshin⁹⁴, G. Volpe²⁵, B. von Haller²⁵, D. Vranic⁷³, G. Øvrebekk⁸³, J. Vrláková¹³, B. Vulpescu⁵, A. Vyushin⁹³, B. Wagner⁸³, V. Wagner⁷⁵, R. Wan⁶⁶, D. Wang⁶⁶, M. Wang⁶⁶, Y. Wang¹¹, Y. Wang⁶⁶, K. Watanabe⁶², M. Weber⁶⁹, J.P. Wessels^{25,xiii}, U. Westerhoff⁹⁹, J. Wiechula¹, J. Wikne²¹, M. Wilde⁹⁹, A. Wilk⁹⁹, G. Wilk⁸², M.C.S. Williams¹¹⁷, B. Windelband¹¹, L. Xaplanteris Karampatsos⁷⁹, C.G. Yaldo⁹⁴, Y. Yamaguchi²², H. Yang¹²³, S. Yang⁸³, S. Yasnopolskiy¹¹⁵, J. Yi⁷⁶, Z. Yin⁶⁶, I.-K. Yoo⁷⁶, J. Yoon⁶, W. Yu²⁰, X. Yuan⁶⁶, I. Yushmanov¹¹⁵, C. Zach⁷⁵, C. Zampolli¹¹⁷, S. Zaporozhets³⁸, A. Zarochentsev⁵⁸, P. Závada²³, N. Zaviyalov⁹³, H. Zbroszczyk⁹⁵, P. Zelnicek⁵⁷, I.S. Zgura³⁶, M. Zhalov⁸⁸, H. Zhang⁶⁶, X. Zhang^{66,viii}, D. Zhou⁶⁶, F. Zhou⁶⁶, Y. Zhou², J. Zhu⁶⁶, X. Zhu⁶⁶, A. Zichichi^{47,xi}, A. Zimmermann¹¹, G. Zinovjev¹⁰⁵, Y. Zoccarato³⁹, M. Zynovyev¹⁰⁵, M. Zyzak²⁰

¹ Eberhard Karls Universität Tübingen, Tübingen, Germany

² Nikhef, National Institute for Subatomic Physics and Institute for Subatomic Physics of Utrecht University, Utrecht, Netherlands

³ Dipartimento di Fisica dell'Università and Sezione INFN, Padova, Italy

⁴ COMSATS Institute of Information Technology (CIIT), Islamabad, Pakistan

⁵ Laboratoire de Physique Corpusculaire (LPC), Clermont Université, Université Blaise Pascal, CNRS-IN2P3, Clermont-Ferrand, France

⁶ Yonsei University, Seoul, South Korea

⁷ Yerevan Physics Institute, Yerevan, Armenia

⁸ Scientific Research Technological Institute of Instrument Engineering, Kharkov, Ukraine

⁹ Laboratori Nazionali di Frascati, INFN, Frascati, Italy

¹⁰ Lawrence Berkeley National Laboratory, Berkeley, CA, United States

¹¹ Physikalisches Institut, Ruprecht-Karls-Universität Heidelberg, Heidelberg, Germany

¹² Saha Institute of Nuclear Physics, Kolkata, India

¹³ Faculty of Science, P.J. Šafárik University, Košice, Slovakia

¹⁴ The Henryk Niewodniczanski Institute of Nuclear Physics, Polish Academy of Sciences, Cracow, Poland

¹⁵ Rudjer Bošković Institute, Zagreb, Croatia

¹⁶ Moscow Engineering Physics Institute, Moscow, Russia

¹⁷ Laboratori Nazionali di Legnano, INFN, Legnano, Italy

¹⁸ Faculty of Engineering, Bergen University College, Bergen, Norway

¹⁹ Nuclear Physics Group, STFC Daresbury Laboratory, Daresbury, United Kingdom

²⁰ Institut für Kernphysik, Johann Wolfgang Goethe-Universität Frankfurt, Frankfurt, Germany

²¹ Department of Physics, University of Oslo, Oslo, Norway

²² University of Tokyo, Tokyo, Japan

²³ Institute of Physics, Academy of Sciences of the Czech Republic, Prague, Czech Republic

²⁴ Dipartimento di Fisica Sperimentale dell'Università and Sezione INFN, Turin, Italy

²⁵ European Organization for Nuclear Research (CERN), Geneva, Switzerland

²⁶ Department of Physics, Aligarh Muslim University, Aligarh, India

²⁷ Sezione INFN, Cagliari, Italy

²⁸ Institute for Theoretical and Experimental Physics, Moscow, Russia

²⁹ Helsinki Institute of Physics (HIP) and University of Jyväskylä, Jyväskylä, Finland

³⁰ Dipartimento di Fisica dell'Università and Sezione INFN, Trieste, Italy

³¹ School of Physics and Astronomy, University of Birmingham, Birmingham, United Kingdom

³² Nikhef, National Institute for Subatomic Physics, Amsterdam, Netherlands

³³ Sección Física, Departamento de Ciencias, Pontificia Universidad Católica del Perú, Lima, Peru

³⁴ Gauhati University, Department of Physics, Guwahati, India

³⁵ Institut de Physique Nucléaire d'Orsay (IPNO), Université Paris-Sud, CNRS-IN2P3, Orsay, France

³⁶ Institute of Space Sciences (ISS), Bucharest, Romania

³⁷ Physics Department, University of Jammu, Jammu, India

³⁸ Joint Institute for Nuclear Research (JINR), Dubna, Russia

³⁹ Université de Lyon, Université Lyon 1, CNRS/IN2P3, IPN-Lyon, Villeurbanne, France

⁴⁰ California Polytechnic State University, San Luis Obispo, CA, United States

⁴¹ Budker Institute for Nuclear Physics, Novosibirsk, Russia

⁴² Dipartimento di Fisica 'E.R. Caianiello' dell'Università and Gruppo Collegato INFN, Salerno, Italy

⁴³ Dipartimento di Fisica dell'Università 'La Sapienza' and Sezione INFN, Rome, Italy

⁴⁴ University of Tennessee, Knoxville, TN, United States

⁴⁵ Institute for High Energy Physics, Protvino, Russia

⁴⁶ Universidade de São Paulo (USP), São Paulo, Brazil

⁴⁷ Dipartimento di Fisica dell'Università and Sezione INFN, Bologna, Italy

⁴⁸ Centro de Investigación y de Estudios Avanzados (CINVESTAV), Mexico City and Mérida, Mexico

⁴⁹ National Institute for Physics and Nuclear Engineering, Bucharest, Romania

⁵⁰ Centre de Calcul de l'IN2P3, Villeurbanne, France

⁵¹ Frankfurt Institute for Advanced Studies, Johann Wolfgang Goethe-Universität Frankfurt, Frankfurt, Germany

⁵² Variable Energy Cyclotron Centre, Kolkata, India

⁵³ Physics Department, University of Rajasthan, Jaipur, India

⁵⁴ Gangneung-Wonju National University, Gangneung, South Korea

⁵⁵ Centro de Investigaciones Energéticas Medioambientales y Tecnológicas (CIEMAT), Madrid, Spain

⁵⁶ Laboratoire de Physique Subatomique et de Cosmologie (LPSC), Université Joseph Fourier, CNRS-IN2P3, Institut Polytechnique de Grenoble, Grenoble, France

- 57 Institut für Informatik, Johann Wolfgang Goethe-Universität Frankfurt, Frankfurt, Germany
- 58 V. Fock Institute for Physics, St. Petersburg State University, St. Petersburg, Russia
- 59 Department of Physics, Ohio State University, Columbus, OH, United States
- 60 Centro Fermi – Centro Studi e Ricerche e Museo Storico della Fisica “Enrico Fermi”, Rome, Italy
- 61 Dipartimento di Fisica dell’Università and Sezione INFN, Cagliari, Italy
- 62 University of Tsukuba, Tsukuba, Japan
- 63 Faculty of Mathematics, Physics and Informatics, Comenius University, Bratislava, Slovakia
- 64 Kirchoff-Institut für Physik, Ruprecht-Karls-Universität Heidelberg, Heidelberg, Germany
- 65 Department of Physics, Sejong University, Seoul, South Korea
- 66 Hua-Zhong Normal University, Wuhan, China
- 67 Universidad Autónoma de Sinaloa, Culiacán, Mexico
- 68 Zentrum für Technologietransfer und Telekommunikation (ZTT), Fachhochschule Worms, Worms, Germany
- 69 University of Houston, Houston, TX, United States
- 70 Indian Institute of Technology Indore (IIT), Indore, India
- 71 Centro de Aplicaciones Tecnológicas y Desarrollo Nuclear (CEADEN), Havana, Cuba
- 72 Instituto de Ciencias Nucleares, Universidad Nacional Autónoma de México, Mexico City, Mexico
- 73 Research Division and ExtreMe Matter Institute EMMI, GSI Helmholtzzentrum für Schwerionenforschung, Darmstadt, Germany
- 74 Sezione INFN, Catania, Italy
- 75 Faculty of Nuclear Sciences and Physical Engineering, Czech Technical University in Prague, Prague, Czech Republic
- 76 Pusan National University, Pusan, South Korea
- 77 Benemérita Universidad Autónoma de Puebla, Puebla, Mexico
- 78 SUBATECH, Ecole des Mines de Nantes, Université de Nantes, CNRS-IN2P3, Nantes, France
- 79 The University of Texas at Austin, Physics Department, Austin, TX, United States
- 80 Institute for Nuclear Research, Academy of Sciences, Moscow, Russia
- 81 Dipartimento di Fisica e Astronomia dell’Università and Sezione INFN, Catania, Italy
- 82 Soltan Institute for Nuclear Studies, Warsaw, Poland
- 83 Department of Physics and Technology, University of Bergen, Bergen, Norway
- 84 Institute of Experimental Physics, Slovak Academy of Sciences, Košice, Slovakia
- 85 Korea Institute of Science and Technology Information, Daejeon, South Korea
- 86 Institut für Kernphysik, Technische Universität Darmstadt, Darmstadt, Germany
- 87 Physics Department, Panjab University, Chandigarh, India
- 88 Petersburg Nuclear Physics Institute, Gatchina, Russia
- 89 Physics Department, University of Cape Town, iThemba LABS, Cape Town, South Africa
- 90 Sezione INFN, Bari, Italy
- 91 Institute of Physics, Bhubaneswar, India
- 92 Sezione INFN, Turin, Italy
- 93 Russian Federal Nuclear Center (VNIIEF), Sarov, Russia
- 94 Wayne State University, Detroit, MI, United States
- 95 Warsaw University of Technology, Warsaw, Poland
- 96 Purdue University, West Lafayette, IN, United States
- 97 Dipartimento Interateneo di Fisica ‘M. Merlin’ and Sezione INFN, Bari, Italy
- 98 Institut Pluridisciplinaire Hubert Curien (IPHC), Université de Strasbourg, CNRS-IN2P3, Strasbourg, France
- 99 Institut für Kernphysik, Westfälische Wilhelms-Universität Münster, Münster, Germany
- 100 Sezione INFN, Rome, Italy
- 101 Physics Department, University of Athens, Athens, Greece
- 102 Nuclear Physics Institute, Academy of Sciences of the Czech Republic, Řež u Prahy, Czech Republic
- 103 Instituto de Física, Universidad Nacional Autónoma de México, Mexico City, Mexico
- 104 Sezione INFN, Padova, Italy
- 105 Bogolyubov Institute for Theoretical Physics, Kiev, Ukraine
- 106 Departamento de Física de Partículas and IGFAE, Universidad de Santiago de Compostela, Santiago de Compostela, Spain
- 107 Division of Experimental High Energy Physics, University of Lund, Lund, Sweden
- 108 Sezione INFN, Trieste, Italy
- 109 Hiroshima University, Hiroshima, Japan
- 110 Bose Institute, Department of Physics and Centre for Astroparticle Physics and Space Science (CAPSS), Kolkata, India
- 111 Oak Ridge National Laboratory, Oak Ridge, TN, United States
- 112 Physics Department, Creighton University, Omaha, NE, United States
- 113 Universidade Estadual de Campinas (UNICAMP), Campinas, Brazil
- 114 Technical University of Split FESB, Split, Croatia
- 115 Russian Research Centre Kurchatov Institute, Moscow, Russia
- 116 Niels Bohr Institute, University of Copenhagen, Copenhagen, Denmark
- 117 Sezione INFN, Bologna, Italy
- 118 Dipartimento di Scienze e Tecnologie Avanzate dell’Università del Piemonte Orientale and Gruppo Collegato INFN, Alessandria, Italy
- 119 Yale University, New Haven, CT, United States
- 120 Chicago State University, Chicago, IL, United States
- 121 Fachhochschule Köln, Köln, Germany
- 122 China Institute of Atomic Energy, Beijing, China
- 123 Commissariat à l’Energie Atomique, IRFU, Saclay, France
- 124 Indian Institute of Technology, Mumbai, India
- 125 Lawrence Livermore National Laboratory, Livermore, CA, United States
- 126 KFKI Research Institute for Particle and Nuclear Physics, Hungarian Academy of Sciences, Budapest, Hungary

* Corresponding author.

E-mail address: alexandru.florin.dobrin@cern.ch (A. Dobrin).

i Also at: Sezione INFN, Bologna, Italy.

ii Also at: Gangneung-Wonju National University, Gangneung, South Korea.

iii Also at: Dipartimento di Fisica ‘E.R. Caianiello’ dell’Università and Gruppo Collegato INFN, Salerno, Italy.

iv Also at: Institute of Space Sciences (ISS), Bucharest, Romania.

v Also at: European Organization for Nuclear Research (CERN), Geneva, Switzerland.

- vi Now at: Fachhochschule Köln, Köln, Germany.
- vii Also at: Institute of Experimental Physics, Slovak Academy of Sciences, Košice, Slovakia.
- viii Also at: Laboratoire de Physique Corpusculaire (LPC), Clermont Université, Université Blaise Pascal, CNRS-IN2P3, Clermont-Ferrand, France.
- ix Now at: The Henryk Niewodniczanski Institute of Nuclear Physics, Polish Academy of Sciences, Cracow, Poland.
- x Now at: Research Division and ExtreMe Matter Institute EMMI, GSI Helmholtzzentrum für Schwerionenforschung, Darmstadt, Germany.
- xi Also at: Centro Fermi – Centro Studi e Ricerche e Museo Storico della Fisica “Enrico Fermi”, Rome, Italy.
- xii Also at: Centro de Investigaciones Energéticas Medioambientales y Tecnológicas (CIEMAT), Madrid, Spain.
- xiii Now at: Institut für Kernphysik, Westfälische Wilhelms-Universität Münster, Münster, Germany.