

Pion, kaon, and (anti) proton production in U+U collisions at $\sqrt{s_{NN}} = 193$ GeV measured with the STAR detector

M. S. Abdallah,¹ B. E. Aboona,² J. Adam,³ J. R. Adams,⁴ J. K. Adkins,⁵ G. Agakishiev,⁶ I. Aggarwal,⁷ M. M. Aggarwal,⁷ Z. Ahammed,⁸ A. Aitbaev,⁶ I. Alekseev,^{9,16} D. M. Anderson,² A. Aparin,⁶ J. Atchison,¹⁰ G. S. Averichev,⁶ V. Bairathi,¹¹ W. Baker,¹² J. G. Ball Cap,¹³ K. Barish,¹² P. Bhagat,¹⁴ A. Bhasin,¹⁴ S. Bhatta,¹⁵ I. G. Bordyuzhin,⁹ J. D. Brandenburg,³ A. V. Brandin,¹⁶ X. Z. Cai,¹⁷ H. Caines,¹⁸ M. Calderón de la Barca Sánchez,¹⁹ D. Cebra,¹⁹ I. Chakaberia,²⁰ B. K. Chan,²¹ Z. Chang,²² S. Chattopadhyay,⁸ D. Chen,¹² J. Chen,²³ J. H. Chen,²⁴ X. Chen,²⁵ Z. Chen,²³ J. Cheng,²⁶ S. Choudhury,²⁴ W. Christie,³ X. Chu,³ H. J. Crawford,²⁷ M. Daugherty,¹⁰ T. G. Dedovich,⁶ I. M. Deppner,²⁸ A. A. Derevschikov,²⁹ A. Dhamija,⁷ L. Di Carlo,³⁰ L. Didenko,³ P. Dixit,³¹ X. Dong,²⁰ J. L. Drachenberg,¹⁰ E. Duckworth,³² J. C. Dunlop,³ J. Engelage,²⁷ G. Eppley,³³ S. Esumi,³⁴ O. Evdokimov,³⁵ A. Ewigleben,³⁶ O. Eyser,³ R. Fatemi,⁵ F. M. Fawzi,¹ S. Fazio,³⁷ C. J. Feng,³⁸ Y. Feng,³⁹ E. Finch,⁴⁰ Y. Fisyak,³ A. Francisco,¹⁸ C. Fu,⁴¹ F. Geurts,³³ N. Ghimire,⁴² A. Gibson,⁴³ K. Gopal,⁴⁴ X. Gou,²³ D. Grosnick,⁴³ A. Gupta,¹⁴ A. Hamed,¹ Y. Han,³³ M. D. Harasty,¹⁹ J. W. Harris,¹⁸ H. Harrison,⁵ S. He,⁴¹ W. He,²⁴ X. H. He,⁴⁵ Y. He,²³ S. Heppelmann,¹⁹ E. Hoffman,¹³ C. Hu,⁴⁵ Q. Hu,⁴⁵ Y. Hu,²⁰ H. Huang,³⁸ H. Z. Huang,²¹ S. L. Huang,¹⁵ T. Huang,³⁸ X. Huang,²⁶ Y. Huang,²⁶ T. J. Humanic,⁴ D. Isenhower,¹⁰ M. Isshiki,³⁴ W. W. Jacobs,²² C. Jena,⁴⁴ Y. Ji,²⁰ J. Jia,^{3,15} K. Jiang,²⁵ C. Jin,³³ X. Ju,²⁵ E. G. Judd,²⁷ S. Kabana,¹¹ M. L. Kabir,¹² D. Kalinkin,^{22,3} K. Kang,²⁶ D. Kapukchyan,¹² K. Kauder,³ H. W. Ke,³ D. Keane,³² A. Kechechyan,⁶ M. Kelsey,³⁰ B. Kimelman,¹⁹ A. Kiselev,³ A. G. Knospe,³⁶ H. S. Ko,²⁰ L. Kochenda,¹⁶ A. A. Korobitsin,⁶ P. Kravtsov,¹⁶ L. Kumar,⁷ S. Kumar,⁴⁵ R. Kunnawalkam Elayavalli,¹⁸ J. H. Kwasizur,²² R. Lacey,¹⁵ S. Lan,⁴¹ J. M. Landgraf,³ A. Lebedev,³ R. Lednicky,⁶ J. H. Lee,³ Y. H. Leung,²⁰ N. Lewis,³ C. Li,²³ C. Li,²⁵ W. Li,³³ W. Li,¹⁷ X. Li,²⁵ Y. Li,²⁵ Y. Li,²⁶ Z. Li,²⁵ X. Liang,¹² Y. Liang,³² T. Lin,²³ Y. Lin,⁴¹ F. Liu,⁴¹ H. Liu,²² H. Liu,⁴¹ T. Liu,¹⁸ X. Liu,⁴ Y. Liu,² T. Ljubicic,³ W. J. Llope,³⁰ R. S. Longacre,³ E. Loyd,¹² T. Lu,⁴⁵ N. S. Lukow,⁴² X. F. Luo,⁴¹ L. Ma,²⁴ R. Ma,³ Y. G. Ma,²⁴ N. Magdy,³⁵ D. Mallick,⁴⁶ S. L. Manukhov,⁶ S. Margetis,³² H. S. Matis,²⁰ J. A. Mazer,⁴⁷ G. McNamara,³⁰ N. G. Minaev,²⁹ D. Mishra,⁴⁶ B. Mohanty,⁴⁶ M. M. Mondal,⁴⁶ I. Mooney,¹⁸ D. A. Morozov,²⁹ M. I. Nagy,⁴⁸ A. S. Nain,⁷ J. D. Nam,⁴² Md. Nasim,³¹ K. Nayak,⁴⁴ D. Neff,²¹ J. M. Nelson,²⁷ D. B. Nemes,¹⁸ M. Nie,²³ G. Nigmatkulov,¹⁶ T. Niida,³⁴ R. Nishitani,³⁴ L. V. Nogach,²⁹ T. Nonaka,³⁴ A. S. Nunes,³ G. Odyniec,²⁰ A. Ogawa,³ S. Oh,²⁰ V. A. Okorokov,¹⁶ K. Okubo,³⁴ B. S. Page,³ R. Pak,³ J. Pan,² A. Pandav,⁴⁶ A. K. Pandey,³⁴ Y. Panebratsev,⁶ P. Parfenov,¹⁶ A. Paul,¹² C. Perkins,²⁷ B. R. Pokhrel,⁴² J. Porter,²⁰ M. Posik,⁴² N. K. Pruthi,⁷ J. Putschke,³⁰ Z. Qin,²⁶ H. Qiu,⁴⁵ A. Quintero,⁴² C. Racz,¹² S. K. Radhakrishnan,³² N. Raha,³⁰ R. L. Ray,⁴⁹ H. G. Ritter,²⁰ O. V. Rogachevsky,⁶ J. L. Romero,¹⁹ D. Roy,⁴⁷ P. Roy Chowdhury,⁵⁰ L. Ruan,³ A. K. Sahoo,³¹ N. R. Sahoo,²³ H. Sako,³⁴ S. Salur,⁴⁷ E. Samigullin,⁹ S. Sato,³⁴ W. B. Schmidke,³ N. Schmitz,⁵¹ J. Seger,⁵² M. Sergeeva,²¹ R. Seto,¹² P. Seyboth,⁵¹ N. Shah,⁵³ E. Shabaliev,⁶ P. V. Shanmuganathan,³ M. Shao,²⁵ T. Shao,²⁴ R. Sharma,⁴⁴ A. I. Sheikh,³² D. Y. Shen,²⁴ K. Shen,²⁵ S. S. Shi,⁴¹ Y. Shi,²³ Q. Y. Shou,²⁴ E. P. Sichtermann,²⁰ J. Singh,⁷ S. Singha,⁴⁵ P. Sinha,⁴⁴ M. J. Skoby,^{54,39} Y. Söhngen,²⁸ W. Solyst,²² Y. Song,¹⁸ B. Srivastava,³⁹ T. D. S. Stanislaus,⁴³ D. J. Stewart,³⁰ M. Strikhanov,¹⁶ B. Stringfellow,³⁹ C. Sun,¹⁵ X. M. Sun,⁴¹ X. Sun,⁴⁵ Y. Sun,²⁵ Y. Sun,⁵⁵ B. Surrow,⁴² D. N. Svirida,⁹ Z. W. Sweger,¹⁹ A. H. Tang,³ Z. Tang,²⁵ A. Taranenko,¹⁶ T. Tarnowsky,⁵⁶ J. H. Thomas,²⁰ D. Tlusty,⁵² T. Todoroki,³⁴ M. Tokarev,⁶ C. A. Tomkiel,³⁶ S. Trentalange,²¹ R. E. Tribble,² P. Tribedy,³ S. K. Tripathy,⁴⁸ O. D. Tsai,²¹ C. Y. Tsang,^{32,3} Z. Tu,³ T. Ullrich,³ D. G. Underwood,^{57,43} I. Upsal,³³ G. Van Buren,³ A. N. Vasiliev,^{29,16} V. Verkest,³⁰ F. Videbæk,³ S. Vokal,⁶ S. A. Voloshin,³⁰ F. Wang,³⁹ G. Wang,²¹ J. S. Wang,⁵⁵ P. Wang,²⁵ X. Wang,²³ Y. Wang,⁴¹ Y. Wang,²⁶ Z. Wang,²³ J. C. Webb,³ P. C. Weidenkaff,²⁸ G. D. Westfall,⁵⁶ D. Wielanek,⁵⁰ H. Wieman,²⁰ S. W. Wissink,²² J. Wu,⁴¹ J. Wu,⁴⁵ Y. Wu,¹² B. Xi,¹⁷ Z. G. Xiao,²⁶ G. Xie,²⁰ W. Xie,³⁹ H. Xu,⁵⁵ N. Xu,²⁰ Q. H. Xu,²³ Y. Xu,²³ Z. Xu,³ Z. Xu,²¹ G. Yan,²³ Z. Yan,¹⁵ C. Yang,²³ Q. Yang,²³ S. Yang,⁵⁸ Y. Yang,³⁸ Z. Ye,³³ Z. Ye,³⁵ L. Yi,²³ K. Yip,³ Y. Yu,²³ W. Zha,²⁵ C. Zhang,¹⁵ D. Zhang,⁴¹ J. Zhang,²³ S. Zhang,²⁵ S. Zhang,²⁴ Y. Zhang,⁴⁵ Y. Zhang,²⁵ Y. Zhang,⁴¹ Z. J. Zhang,³⁸ Z. Zhang,³ Z. Zhang,³⁵ F. Zhao,⁴⁵ J. Zhao,²⁴ M. Zhao,³ C. Zhou,²⁴ J. Zhou,²⁵ Y. Zhou,⁴¹ X. Zhu,²⁶ M. Zurek,⁵⁷ and M. Zyzak⁵⁹

(STAR Collaboration)

¹American University of Cairo, New Cairo 11835, New Cairo, Egypt

²Texas A&M University, College Station, Texas 77843

³Brookhaven National Laboratory, Upton, New York 11973

⁴Ohio State University, Columbus, Ohio 43210

⁵University of Kentucky, Lexington, Kentucky 40506-0055

⁶Joint Institute for Nuclear Research, Dubna 141 980

⁷Panjab University, Chandigarh 160014, India

⁸Variable Energy Cyclotron Centre, Kolkata 700064, India

⁹Alikhanov Institute for Theoretical and Experimental Physics NRC "Kurchatov Institute", Moscow 117218

¹⁰Abilene Christian University, Abilene, Texas 79699

¹¹Instituto de Alta Investigación, Universidad de Tarapacá, Arica 1000000, Chile

¹²University of California, Riverside, California 92521

- ¹³University of Houston, Houston, Texas 77204
¹⁴University of Jammu, Jammu 180001, India
¹⁵State University of New York, Stony Brook, New York 11794
¹⁶National Research Nuclear University MEPhI, Moscow 115409
¹⁷Shanghai Institute of Applied Physics, Chinese Academy of Sciences, Shanghai 201800
¹⁸Yale University, New Haven, Connecticut 06520
¹⁹University of California, Davis, California 95616
²⁰Lawrence Berkeley National Laboratory, Berkeley, California 94720
²¹University of California, Los Angeles, California 90095
²²Indiana University, Bloomington, Indiana 47408
²³Shandong University, Qingdao, Shandong 266237
²⁴Fudan University, Shanghai, 200433
²⁵University of Science and Technology of China, Hefei, Anhui 230026
²⁶Tsinghua University, Beijing 100084
²⁷University of California, Berkeley, California 94720
²⁸University of Heidelberg, Heidelberg 69120, Germany
²⁹NRC "Kurchatov Institute", Institute of High Energy Physics, Protvino 142281
³⁰Wayne State University, Detroit, Michigan 48201
³¹Indian Institute of Science Education and Research (IISER), Berhampur 760010, India
³²Kent State University, Kent, Ohio 44242
³³Rice University, Houston, Texas 77251
³⁴University of Tsukuba, Tsukuba, Ibaraki 305-8571, Japan
³⁵University of Illinois at Chicago, Chicago, Illinois 60607
³⁶Lehigh University, Bethlehem, Pennsylvania 18015
³⁷University of Calabria & INFN-Cosenza, Italy
³⁸National Cheng Kung University, Tainan 70101
³⁹Purdue University, West Lafayette, Indiana 47907
⁴⁰Southern Connecticut State University, New Haven, Connecticut 06515
⁴¹Central China Normal University, Wuhan, Hubei 430079
⁴²Temple University, Philadelphia, Pennsylvania 19122
⁴³Valparaiso University, Valparaiso, Indiana 46383
⁴⁴Indian Institute of Science Education and Research (IISER) Tirupati, Tirupati 517507, India
⁴⁵Institute of Modern Physics, Chinese Academy of Sciences, Lanzhou, Gansu 730000
⁴⁶National Institute of Science Education and Research, HBNI, Jatni 752050, India
⁴⁷Rutgers University, Piscataway, New Jersey 08854
⁴⁸ELTE Eötvös Loránd University, Budapest, Hungary H-1117
⁴⁹University of Texas, Austin, Texas 78712
⁵⁰Warsaw University of Technology, Warsaw 00-661, Poland
⁵¹Max-Planck-Institut für Physik, Munich 80805, Germany
⁵²Creighton University, Omaha, Nebraska 68178
⁵³Indian Institute Technology, Patna, Bihar 801106, India
⁵⁴Ball State University, Muncie, Indiana, 47306
⁵⁵Huzhou University, Huzhou, Zhejiang 313000
⁵⁶Michigan State University, East Lansing, Michigan 48824
⁵⁷Argonne National Laboratory, Argonne, Illinois 60439
⁵⁸South China Normal University, Guangzhou, Guangdong 510631
⁵⁹Frankfurt Institute for Advanced Studies FIAS, Frankfurt 60438, Germany

We present the first measurements of transverse momentum spectra of π^\pm , K^\pm , $p(\bar{p})$ at midrapidity ($|y| < 0.1$) in U+U collisions at $\sqrt{s_{NN}} = 193$ GeV with the STAR detector at the Relativistic Heavy Ion Collider (RHIC). The centrality dependence of particle yields, average transverse momenta, particle ratios and kinetic freeze-out parameters are discussed. The results are compared with the published results from Au+Au collisions at $\sqrt{s_{NN}} = 200$ GeV in STAR. The results are also compared to those from A Multi Phase Transport (AMPT) model.

I. INTRODUCTION

The experimental need to study the medium formed in high-energy heavy-ion collisions is called for by theoretical predictions from Quantum Chromodynamics (QCD) [1–10]. One of the main motivations lies in the exploration of the QCD phase diagram [11–13]. The phase diagram in temperature

(T) and baryon chemical potential (μ_B) can be accessed experimentally by varying the collision energy of heavy-ion collisions [14–17]. Earlier experimental results from the Relativistic Heavy Ion Collider (RHIC) confirmed the presence of a hot and dense deconfined medium dominated by partonic degrees of freedom [2, 6–9]. The fluid of deconfined quarks and gluons, called the Quark Gluon Plasma (QGP), undergoes a transition to a gas of hadrons. The measurements that evidence

the formation of the QGP are: suppression of high transverse momentum (p_T) hadron production in nucleus-nucleus collisions relative to p+p collisions [18–29], the relatively large value of elliptic flow (v_2) associated with light quark as well as strange quark carrying hadrons, and the inequality between baryon and meson v_2 at intermediate p_T [30–40].

A complementary approach to study heavy ion collisions at RHIC relies on varying the colliding system type, shape and geometry. The initial stage dynamics is expected to be reflected in the final state. Hence, the initial geometry and shape of the colliding nuclei could provide insights into the physics of the bulk properties of the system [41–51]. With such a motivation [52–57], the Solenoidal Tracker At RHIC (STAR) took data in the year 2012 from uranium on uranium (U+U) collisions at $\sqrt{s_{NN}} = 193$ GeV because Uranium nucleus (${}_{92}\text{U}^{238}$) is prolate in contrast to the nearly spherical Gold nucleus (${}_{79}\text{Au}^{197}$).

The prolate shape of the uranium nucleus is associated with a long or major axis and a short or minor axis. Due to this, very interesting orientations of the two colliding nuclei are possible even for most central U+U collisions [44]. Among all possible random orientations, three particular orientations are worth noting. Those are termed as : tip-tip, body-body and body-tip. Tip-tip refers to the configuration where the major axes of the two colliding nuclei lie parallel to the beam axis resulting in the tips of both nuclei colliding head-on. Body-body collisions are those with the major axes perpendicular to the beam axis and the minor axes parallel to the beam axis. The body-tip orientation of the two colliding nuclei is straightforward from the above two explanations. Model calculations suggest that a tip-tip collision leads to the production of the highest particle multiplicity [43, 44], and the maximum transverse particle density in U+U collisions could be 6% – 35% higher than in Au+Au collisions [52, 55, 56]. With all possible orientations combined, U+U collisions will still likely give higher energy density and particle multiplicity.

This paper concentrates on the extraction of the bulk properties of the medium in U+U collisions at $\sqrt{s_{NN}} = 193$ GeV, including all orientations of the colliding nuclei. The results presented here include the transverse momentum spectra, the particle yields (dN/dy), the mean transverse momentum ($\langle p_T \rangle$), the particle ratios, and the kinetic freeze-out parameters. These results are obtained for π^\pm , K^\pm and $p(\bar{p})$ as a function of collision centrality. All the results are compared to the corresponding published results of Au+Au collisions at $\sqrt{s_{NN}} = 200$ GeV [58, 59]. A comparative study with AMPT model [60] modified to incorporate the deformation of uranium nucleus [43] is also carried out.

The paper is organized as follows. Section II discusses the STAR detector, event selection, centrality selection, and particle identification (PID) procedures. Section III lists the details of the various correction factors to obtain the transverse momentum spectra of π^\pm , K^\pm and $p(\bar{p})$. The systematic uncertainties associated with the measurements of the yields are discussed in Sec. IV. The results in terms of p_T spectra, dN/dy , $\langle p_T \rangle$, ratios of particle yields, kinetic freeze-out conditions and comparison to model calculations are presented in Sec. V. Finally in Sec. VI a summary is given.

II. EXPERIMENT AND DATA ANALYSIS

A. The STAR Detector

The results presented in this paper are from the analysis of data collected with the STAR detector at RHIC [61] for U+U collisions at $\sqrt{s_{NN}} = 193$ GeV in 2012. The data are obtained with a minimum bias trigger using the Vertex Position Detectors (VPDs) [62, 63] and Zero Degree Calorimeters (ZDCs) [64, 65]. The two VPDs are located at $4.24 < |\eta| < 5.1$. The VPDs also provide the start time of the collision and the position of the collision vertex along the beam direction. The ZDCs are a pair of hadronic calorimeters placed on either side of the beam pipe at a distance of 18 m from the center of the STAR detector that detect hadrons emerging at small angles ($\theta < 2$ mrad) with respect to the beam axis. Particle tracking is done using the cylindrical Time Projection Chamber (TPC) [66] that is filled with P10 gas (90% argon and 10% methane) and maintained at a pressure of 2 mbar above atmospheric pressure. It functions in a uniform 0.5 T magnetic field that is parallel to the beam axis. The TPC has full 2π azimuthal coverage for a pseudorapidity range $|\eta| < 1.0$. Particles are identified in the TPC by their ionization energy loss per unit track length (dE/dx) along the particle trajectory. The identification of particles at higher momentum was achieved by including the Time Of Flight (TOF) detector [62, 67] that surrounds the TPC cylinder. The TOF uses a Multigap Resistive Plate Chamber (MRPC) technology and has pseudorapidity acceptance of $|\eta| < 0.9$ with full azimuthal coverage [62, 67].

B. Event Selection

The criteria for minimum bias triggered event selection begin with the identification of a primary vertex that is the common point of origin of tracks in an event. Conventionally the Z-axis is taken to be along the beam axis with $z = 0$ at the center of the TPC. The Z position of the primary vertex (V_z) was constrained to be within $|V_z| < 30$ cm. In addition, to reduce pile-up events, a cut of $|V_z - V_{z,VPD}| < 3$ cm is applied. Here $V_{z,VPD}$ is the vertex position along the beam axis measured by the VPD. The plane perpendicular to beam direction defines the transverse x-y direction. The radial vertex position $V_r = \sqrt{V_x^2 + V_y^2}$ was required to be less than 2 cm to avoid including interactions between beam and beam pipe. The number of minimum bias U+U collisions at $\sqrt{s_{NN}} = 193$ GeV analysed after these event selections is about 2.70×10^8 .

C. Centrality Selection

The collision centrality is qualitatively a measure of the overlap of the two colliding nuclei or, correspondingly, a measure of the number of colliding nucleons in the U+U collisions. Quantitatively this is determined based on the measured uncorrected charged particle multiplicity in the TPC over the

TABLE I. The average number of participating nucleons $\langle N_{\text{part}} \rangle$, average number of nucleon-nucleon binary collisions $\langle N_{\text{coll}} \rangle$ and number of events in different collision centrality classes in U+U collisions at $\sqrt{s_{\text{NN}}} = 193$ GeV.

Centrality (%)	$\langle N_{\text{part}} \rangle$	$\langle N_{\text{coll}} \rangle$	Events ($\times 10^6$)
0–5	414.9 ± 6.0	1281.3 ± 100	15.7
5–10	355.4 ± 13.9	1011.0 ± 49.7	17.8
10–20	277.5 ± 12.8	714.1 ± 46.6	36.0
20–30	195.4 ± 13.7	435.9 ± 42.5	35.5
30–40	133.1 ± 13.6	253.5 ± 36.3	35.8
40–50	86.2 ± 12.6	137.4 ± 28.6	35.1
50–60	52.6 ± 10.5	69.3 ± 20.3	33.7
60–70	29.4 ± 8.5	31.8 ± 11.8	33.4
70–80	14.7 ± 5.5	13.2 ± 7.0	25.9

full azimuthal coverage and within $|\eta| < 0.5$. This is termed as Reference Multiplicity or simply Refmult in STAR. The centrality selection is done by comparing the Refmult distribution with a Monte Carlo Glauber model simulation [16, 50, 68]. The MC Glauber model is based on charged particle multiplicity calculated using the two-component model with the number of participants (N_{part}) and number of binary nucleon-nucleon collisions (N_{coll}). In this analysis the minimum bias events are divided into nine centrality classes with each class representing the fraction of the total collision cross section parametrized in terms of the number of participating nucleons in the collision, $\langle N_{\text{part}} \rangle$, with the class 0–5% being the class with the highest particle multiplicities and therefore, the most central class. The nine classes are 0–5%, 5–10%, 10–20%, 20–30%, 30–40%, 40–50%, 50–60%, 60–70% and 70–80%. The events for 70–80% centrality class are corrected for the trigger inefficiencies and 80–100% centrality is not used in our analysis because of its significant trigger bias due to vertex inefficiency at low multiplicities and the contamination from electromagnetic interactions. Using the Glauber Model Monte-Carlo simulations, the average number of participant nucleons $\langle N_{\text{part}} \rangle$ and number of nucleon-nucleon binary collisions $\langle N_{\text{coll}} \rangle$ are evaluated. Systematic uncertainties on the average quantities have been estimated by varying input parameters for the Glauber model as well as the total cross section within $\pm 5\%$ and by using different density profiles for the nucleons in the uranium nuclei used in the simulations discussed in Ref. [41]. These values for U+U collisions at $\sqrt{s_{\text{NN}}} = 193$ GeV along with the corresponding number of events analyzed in each centrality bin are listed in Table I. The U+U collisions allow measurements to a higher value of $\langle N_{\text{part}} \rangle > 400$ than in Au+Au collisions at $\sqrt{s_{\text{NN}}} = 200$ GeV ($\langle N_{\text{part}} \rangle \approx 350$).

D. Track Selection

The quality of the tracks used in the analysis was assured by employing the standard track selection process used in STAR [58, 68, 69]. The track selection criteria are tabulated in Table II. To avoid intermixing of tracks from a secondary vertex, the track must have a Distance of Closest Approach (DCA) to the primary vertex of less than 3 cm. The number

TABLE II. Track selection criteria for the tracks used in the analysis.

Criterion type	Value
$ y $	< 0.1
DCA	< 3 cm
nFitPts	> 25
nFitPts/nFitPoss	> 0.52
ndE/dx	> 15

of TPC points used to fit each track (nFitPts) was required to be greater than 25 out of a possible maximum hits (nFitPoss). The ratio of nFitPts to nFitPoss was required to be greater than 0.52 to exclude split tracks [70]. We also required greater than 15 measured points (ndE/dx) along a track to reliably calculate $\langle dE/dx \rangle$. And, finally, the rapidity window chosen for this analysis is $|y| < 0.1$ in order to have the largest p_T coverage for particle identification.

E. Particle Identification

The $\langle dE/dx \rangle$ measured in the TPC is used for particle identification and is a function of rigidity (momentum-to-charge ratio) as shown in Fig. 1 (left panel). The black solid curves represent the theoretical value of $\langle dE/dx \rangle$ predicted by the Bichsel formula [71]. As visible from the figure, the bands of charged pions, kaons, and protons start merging at higher momentum. Hence, the TPC is extensively used to identify pions, kaons, and protons up to p_T of 0.8, 0.8 and 1.0 GeV/c respectively. To extend particle identification to higher p_T we include TOF information for pions and kaons with p_T above 0.4 GeV/c and for protons above 0.5 GeV/c. The right side of Fig. 1 shows the inverse of the particle velocity $1/\beta$ as a function of p/q . The solid black lines are the mean values of $1/\beta$ for a particular particle species. The various bands remain clearly separated up to momentum of 2 GeV/c.

The raw yields of particles identified using $\langle dE/dx \rangle$ from the TPC are obtained using the z -variable [72] as described in Refs. [58, 69]. For a particular particle type it is given by the mathematical expression

$$z_X = \ln \left(\frac{\langle dE/dx \rangle}{\langle dE/dx \rangle_X^B} \right), \quad (1)$$

where X is the particle type chosen [e^\pm , π^\pm , K^\pm , $p(\bar{p})$ in the present analysis] and $\langle dE/dx \rangle_X^B$ is the theoretical value of $\langle dE/dx \rangle$ obtained from Bichsel formula [71] for the corresponding particle. This gives the z -distribution for each particle species within $|y| < 0.1$ for a given p_T and centrality bin.

The z distribution is then fitted by multiple Gaussians, corresponding to each particle species, as discussed in Ref. [69]. The yields are obtained as the area under the Gaussian function for a particular p_T and centrality bin. To extract the raw yields using the TOF, we employ the same technique as described in Refs. [69, 73]. For each detected particle we calcu-




FIG. 1. (Color online) Left panel: The $\langle dE/dx \rangle$ distribution of charged particles as a function of momentum/charge (p/q) obtained from TPC in U+U collisions at $\sqrt{s_{NN}} = 193$ GeV. The curves represent the expected mean value of $\langle dE/dx \rangle$ for the corresponding particles. Right panel: $1/\beta$ as function of p/q . The curves represent the theoretical values of $1/\beta$ for the corresponding particles.




FIG. 2. The difference between p_T^{REC} and p_T^{MC} as function of p_T^{REC} for pions (left), kaons (middle), protons (right) in 0–5% centrality U+U collisions at $\sqrt{s_{NN}} = 193$ GeV. The errors are statistical only and within symbol size.

late a parameter (m^2) defined as

$$m^2 = p^2 \left(\frac{c^2 T^2}{L^2} - 1 \right), \quad (2)$$

where p , T , L , and c respectively represent the particle momentum, time-of-flight, path length of the particle and the velocity of light. The m^2 distribution in a given p_T and centrality bin is fitted to the predicted m^2 distributions as discussed in Refs. [69, 73]. In this way, the raw yields are extracted in different p_T and centrality bins for each particle species studied here.

III. CORRECTION FACTORS

A. Monte-Carlo Embedding Technique

The Monte-Carlo (MC) embedding technique is briefly described below and can be found in detail in Refs [16, 58, 69].

This technique is used to estimate the track reconstruction efficiency and acceptance of the detector. The MC tracks are generated with flat p_T and y distributions, to ensure equal statistics in each p_T bin. These generated tracks have transverse momentum p_T^{MC} . The number of MC tracks is about 5% of the measured multiplicity in each corresponding p_T^{MC} bin and those MC tracks are mixed with the real data tracks at the detector pixel level and allowed to pass through the response of the STAR detector using the GEANT package [74]. These events with embedded tracks are reconstructed like real events taking into consideration all the detector effects.

B. Energy Loss Correction

In the track reconstruction procedure, the Coulomb scattering and energy loss of a charged particle is corrected by assuming the pion mass for each particle track [58, 69]. Thus, a momentum correction is needed for higher mass particles such

as kaons and protons. This energy loss correction mainly affects the particle yields at low momentum. A correction factor is calculated for kaon and proton tracks from the Monte-Carlo embedding data. The difference between the reconstructed transverse momentum p_T^{REC} and initial transverse momentum p_T^{MC} as a function of p_T^{REC} estimates the energy loss correction track by track. The energy loss correction factor is found to be independent of collision energy and centrality [58, 69]. A plot of this energy loss correction as a function of p_T^{REC} is shown in Fig. 2 for this analysis of U+U collisions at $\sqrt{s_{NN}} = 193$ GeV for pions, kaons, and protons.

C. Tracking Efficiency \times Acceptance

A correction factor taking into account the inefficiency of the detector in reconstructing tracks and its acceptance is applied to the p_T spectra. The MC generated embedding sample provides this correction fraction for each particle species in each centrality. It is given by the ratio of the distribution of reconstructed MC tracks to initial generated MC tracks as a function of p_T^{MC} in the selected rapidity window. The reconstruction efficiency shows a centrality dependence as it increases from central to peripheral collisions. Figure 3 represents efficiency \times acceptance as a function of p_T^{MC} for reconstructed pions, kaons, and protons in 0–5% centrality U+U collisions at $\sqrt{s_{NN}} = 193$ GeV. The raw p_T spectra of each particle species and centrality are corrected for the efficiency \times acceptance.

D. TOF Matching efficiency

The TOF detector surrounds the TPC detector within $|\eta| < 0.9$ and full azimuthal angle. Not all TPC tracks are detected by the TOF detector, hence a track matching correction between TPC and TOF is needed. This correction factor is calculated from data for each particle species. This is obtained from the ratio of the number of tracks detected in the TOF to the number of tracks in the TPC within the acceptance under analysis [69]. This track matching efficiency typically varies from about 60% (for central collisions) to 70% (for peripheral collisions) in U+U collisions at $\sqrt{s_{NN}} = 193$ GeV. The p_T spectra in each centrality are divided by this fraction for each p_T bin.

E. Pion Feed-down Correction

The yield of pions contains contributions from weak decays of K_s^0 and Λ . Also, muons can be misidentified as pions because of their small mass difference. Hence, a correction factor estimating these weak decay contributions and muon contamination to the pion yield is needed [58, 69]. This is obtained from MC simulation, where the events generated from Heavy Ion Jet Interaction (HIJING) model [75] pass through the STAR detector response using the GEANT package [74]. The events are reconstructed in the same manner as that of the

real data. In such simulations we have the initial parent particle information. From the final reconstruction, the secondary pions and muons misidentified as pions are calculated. From the final reconstruction we determine the fraction of the pions that are really secondary pions or muons that are misidentified as pions. Therefore, the pion yields in each p_T bin are reduced by the fraction that are secondary pions and muons identified as pions. The total pion feed-down fraction shows only p_T dependence with no centrality dependence. Its value has a maximum of 15% at $p_T = 0.2$ GeV/c and vanishes above $p_T > 1.2$ GeV/c. These values are similar to those reported in Refs.[58, 69].

F. Proton Background Correction

Energetic particles produced in the collision can interact with the beam pipe and detector material producing secondary protons. Because these secondary protons are produced far from the primary vertex, they appear as a long tail in the DCA distribution of protons. However, anti-protons do not have this background and can therefore be used to estimate the proton background fraction by comparing the DCA distributions of proton and anti-proton from the real data [16, 58, 69, 76]. In this method, protons and anti-protons are selected within $|n_{\sigma_p}| < 2$, where n_{σ_p} is z_X of protons divided by the $\langle dE/dx \rangle$ resolution in the TPC. The proton background fraction is calculated for each centrality and each p_T bin. The proton background fraction is found to decrease with increasing p_T value. It shows a slight variation with collision centrality (decreases towards central collisions). The proton background fraction is maximum for low p_T ($\approx 9\%$ for mid-central collisions) and becomes almost negligible for $p_T > 1.2$ GeV/c. It may be noted that the proton and anti-proton yields reported here are inclusive, i.e., feed-down contributions from weak decays are not corrected for, similar to those measured in Au+Au collisions at $\sqrt{s_{NN}} = 200$ GeV [58, 59, 77].

IV. Systematic Uncertainties

The systematic uncertainties associated with the results include contributions from the event and track selection criteria, the PID procedure, the methods of estimation of correction factors, and the errors associated with extrapolations with different functional fits to p_T spectra. The errors associated with all these sources are calculated and added in quadrature.

The systematic errors associated with event and track selection criteria are estimated by varying one selected criterion while the other criteria are kept at their default values. The event and track selection criteria with default and varied values are: 30 cm (default) $< V_z < 40$ cm; 2 cm $< \text{DCA} < 3$ cm (default); 20 $< \text{nFitPts} < 25$ (default); and 10 $< \text{ndE/dx} < 15$ (default). In the identification of particle type, the fit range of the Gaussian function for dE/dx , and the n_σ criteria used in calculation of predicted m^2 distribution are varied (from $|n_\sigma| < 2$ to $|n_\sigma| < 1$). As another way to obtain the yields, bin




FIG. 3. Tracking efficiency \times acceptance as function of p_T^{MC} for reconstructed pions (left), kaons (middle), and protons (right) obtained from embedding in U+U collisions at $\sqrt{s_{NN}} = 193$ GeV. The errors are statistical only and within symbol size.

TABLE III. Systematic uncertainties related to the yields of π , K , and $p(\bar{p})$ in U+U collisions at $\sqrt{s_{NN}} = 193$ GeV.

	π	K	$p(\bar{p})$
Cuts and PID	4%	5%	7%
Extrapolation	8%	9%	10%
Corrections	5%	5%	5%
Total	10%	11%	13%

counting is used and the resulting difference from the fits is taken as part of the systematic uncertainties.

The procedure for calculation of dN/dy and $\langle p_T \rangle$ involves the fitting of the spectra by a suitable function to extrapolate the spectra to the unmeasured p_T region. The main source of uncertainty on dN/dy and $\langle p_T \rangle$ comes from this extrapolation. In order to estimate the uncertainty associated with extrapolation, different fit functions are used. The default fit functions used for pions, kaons, and protons are respectively Bose-Einstein, m_T exponential and double exponential respectively [69]. For the systematic uncertainty estimation the used functions are p_T exponential, Boltzmann, and m_T exponential for pions, kaons, and protons, respectively [69]. The systematic uncertainties on the values of $\langle p_T \rangle$ for pions, kaons, and protons are estimated to be 9%, 10% and 10%, respectively.

In addition, a 5% estimated systematic uncertainty associated with the calculation of the detector efficiency and acceptance was added in quadrature to the statistical errors on the p_T spectra [58]. The systematic uncertainties associated with the pion feed-down and proton background correction were also estimated. While the contribution from the former is negligible, the latter adds a 6–7% systematic uncertainty to the low p_T part of the proton p_T spectra. The different sources of systematic uncertainties on the particle yields for all centralities are tabulated in Table III.

The systematic uncertainty on the particle ratios are obtained using the systematic uncertainty on the particle yields. The correlated uncertainties, i.e., from efficiency, will cancel out completely in the particle ratios. The systematic uncertainties associated with the extrapolated fits also cancels out

for particle and anti-particle ratios.

The uncertainties associated with the kinetic freeze-out parameters have two contributions which are added in quadrature. One of them is the uncertainty that is obtained during the simultaneous fitting of pion, kaon, and proton spectra along with their point-to-point systematic uncertainties. The second is obtained by varying the fit range of the measured p_T spectra.

V. RESULTS AND DISCUSSIONS

A. Transverse Momentum Spectra

The transverse momentum spectra in U+U collisions at $\sqrt{s_{NN}} = 193$ GeV for π^+ , π^- , K^+ , K^- , p and \bar{p} within $|y| < 0.1$ are shown in Fig. 4. The spectra are shown in nine different centrality classes 0–5%, 5–10%, 10–20%, 20–30%, 30–40%, 40–50%, 50–60%, 60–70% and 70–80%. The data points shown for $0.4 < p_T < 2.0$ GeV/c for pions and kaons and for $0.5 < p_T < 2.0$ GeV/c for protons are obtained using both the TPC and the TOF. In addition, the data points measured using only the TPC are shown for $0.2 < p_T < 0.8$ GeV/c, $0.3 < p_T < 0.8$ GeV/c and $0.5 < p_T < 1.0$ GeV/c regions for pions, kaons, and protons respectively. The $0.4 < p_T < 0.8$ GeV/c, $0.4 < p_T < 0.8$ GeV/c and $0.5 < p_T < 1.0$ GeV/c for pions, kaons, and protons respectively are the overlap region containing data points measured using only the TPC and also the data points using the TPC along with the TOF. For each particle in Fig. 4 the spectral shape in this overlapping p_T range is seen to smoothly join the data obtained using only the TPC (low p_T) and the data obtained using the TOF (high p_T). The curves represent the fit to the spectra, shown only for 0–5% central collisions. The respective functions for pion, kaons and protons are Bose-Einstein, m_T exponential and double exponential [69]. The particle yields (dN/dy) and average transverse momenta ($\langle p_T \rangle$) are obtained from the measured points in the p_T spectra. For the unmeasured p_T regions, the contributions to the yield are extracted by extrapolating the fit functions. The behavior of the p_T spectra changes with centrality. As can be observed from the Fig. 4, the slopes of the




FIG. 4. p_T spectra of π^\pm , K^\pm , $p(\bar{p})$ measured at midrapidity ($|y| < 0.1$) in U+U collisions at $\sqrt{s_{NN}} = 193$ GeV in STAR. Spectra are plotted for nine different centrality classes and some of them are scaled for clarity. The p_T spectra are fitted, with Bose-Einstein function for pions, m_T -exponential for kaons, and double exponential for (anti) protons, shown for 0–5% centrality classes in the figure. The uncertainties represent total systematic and statistical uncertainties added in quadrature and are mostly dominated by systematic errors.

spectra show a gradual flattening as one goes from peripheral to central collisions. This is an indication of stronger radial flow effects for particles with increasing centrality. A mass dependence of the inverse slope of the p_T spectra can also be inferred as it increases with increasing mass of the particle.

B. Average Transverse Momenta

The slopes of the spectra can be quantified by $\langle p_T \rangle$. The $\langle p_T \rangle$ for π^+ , K^+ and p is plotted as a function of $\langle N_{part} \rangle$ in Fig. 5 for U+U collisions at $\sqrt{s_{NN}} = 193$ GeV. For comparison, the published results obtained in Au+Au collisions at $\sqrt{s_{NN}} = 200$ GeV are also shown [58]. Not shown explicitly are the $\langle p_T \rangle$ for π^- , K^- and \bar{p} which show identical behavior. The $\langle p_T \rangle$ increases gradually from peripheral to central collisions as well as from pions to kaons and kaons to protons in U+U collisions at $\sqrt{s_{NN}} = 193$ GeV. The hadron mass dependence of $\langle p_T \rangle$ is indicates radial flow effect, which increases from peripheral to central collisions. The $\langle p_T \rangle$ are similar for collisions between deformed uranium nuclei and spherical gold nuclei for comparable values of $\langle N_{part} \rangle$. Moreover, using U+U collisions the $\langle p_T \rangle$ measurements have been made for the highest $\langle N_{part} \rangle$ values observed at RHIC. The $\langle p_T \rangle$ values are listed in Table IV for U+U collisions at $\sqrt{s_{NN}} = 193$ GeV.

C. Particle Yields

The total yield of a particle in a given centrality is obtained by integrating the spectra over the full p_T range. The yields from the unmeasured region of the spectra are extracted using the functional extrapolation [69]. It is represented by the observable dN/dy within rapidity $|y| < 0.1$. The normalised dN/dy with $\langle N_{part} \rangle / 2$ of π^+ , K^+ , p and \bar{p} measured in U+U collisions at $\sqrt{s_{NN}} = 193$ GeV as a function of $\langle N_{part} \rangle$ is shown in Fig. 6. The results are compared with the published results of Au+Au collisions at $\sqrt{s_{NN}} = 200$ GeV [58]. The values of dN/dy are listed in Table V for U+U collisions at $\sqrt{s_{NN}} = 193$ GeV.

The current data extends the measurements to a higher value of N_{part} , with N_{part} extending above 400. Figure 6 shows that the values of dN/dy per participating nucleon pair for π^+ , K^+ and p increase with average number of participant nucleons in U+U collisions at $\sqrt{s_{NN}} = 193$ GeV. Not shown are the plots for π^- and K^- that show identical behavior to π^+ and K^+ . This signifies particle production having contributions from both soft processes and hard processes involving nucleon-nucleon binary collisions. However the N_{part} dependence is smaller for anti-protons, possibly due to annihilation effects. The values of dN/dy are consistent between U+U collisions at $\sqrt{s_{NN}} = 193$ GeV and Au+Au collisions at $\sqrt{s_{NN}} = 200$ GeV within a similar range of $\langle N_{part} \rangle$ and within the quoted systematic uncertainties. The comparison between U+U and Au+Au collision systems was also reported in Ref. [78] on the production of π^0 and η , where it was found




FIG. 5. $\langle p_T \rangle$ of π^+ , K^+ and p as a function of $\langle N_{part} \rangle$ at midrapidity ($|y| < 0.1$) for U+U collisions at $\sqrt{s_{NN}} = 193$ GeV. The results are compared with Au+Au collisions at $\sqrt{s_{NN}} = 200$ GeV [58]. The uncertainties represent the total systematic and statistical uncertainties added in quadrature, and are dominated by systematic uncertainties.

TABLE IV. $\langle p_T \rangle$ values in GeV/c for π^+ , π^- , K^+ , K^- , p , and \bar{p} from U+U collisions at $\sqrt{s_{NN}} = 193$ GeV within rapidity $|y| < 0.1$. The quoted errors are the total systematic and statistical errors added in quadrature, which are dominated by the systematic errors.

Centrality (%)	π^+	π^-	K^+	K^-	p	\bar{p}
0 – 5	0.449 ± 0.039	0.446 ± 0.038	0.686 ± 0.066	0.685 ± 0.065	1.004 ± 0.101	1.003 ± 0.101
5 – 10	0.445 ± 0.038	0.443 ± 0.038	0.684 ± 0.065	0.682 ± 0.065	0.969 ± 0.094	0.961 ± 0.092
10 – 20	0.442 ± 0.038	0.439 ± 0.037	0.682 ± 0.065	0.680 ± 0.065	0.964 ± 0.093	0.963 ± 0.093
20 – 30	0.435 ± 0.037	0.432 ± 0.036	0.676 ± 0.064	0.677 ± 0.064	0.944 ± 0.089	0.943 ± 0.089
30 – 40	0.429 ± 0.036	0.427 ± 0.036	0.667 ± 0.062	0.666 ± 0.061	0.912 ± 0.083	0.911 ± 0.083
40 – 50	0.421 ± 0.035	0.414 ± 0.034	0.654 ± 0.060	0.656 ± 0.060	0.852 ± 0.073	0.852 ± 0.073
50 – 60	0.413 ± 0.034	0.404 ± 0.033	0.640 ± 0.057	0.643 ± 0.058	0.814 ± 0.066	0.813 ± 0.066
60 – 70	0.403 ± 0.032	0.398 ± 0.032	0.622 ± 0.054	0.623 ± 0.063	0.773 ± 0.060	0.760 ± 0.058
70 – 80	0.396 ± 0.031	0.390 ± 0.030	0.619 ± 0.054	0.617 ± 0.053	0.721 ± 0.052	0.702 ± 0.049

that the yields of these particles show similar suppression in U+U and Au+Au collisions relative to the binary scaled $p + p$ collisions.

D. Particle Ratios

Particle yield ratios are measured in U+U collisions at $\sqrt{s_{NN}} = 193$ GeV and compared with the published results from Au+Au collisions at $\sqrt{s_{NN}} = 200$ GeV [58]. Ratios of anti-particle to particle yields π^-/π^+ , K^-/K^+ and \bar{p}/p as a function of $\langle N_{part} \rangle$ in U+U collisions at $\sqrt{s_{NN}} = 193$ GeV are shown in Fig. 7. As a function of $\langle N_{part} \rangle$, the yield ratios shown are all approximately constant and near unity. For the π^-/π^+ yield ratios, this is an indication that the particle production mechanism for π^+ and π^- does not change significantly from central to peripheral collisions. For the K^-/K^+ yield ratios, this suggests that K^+ and K^- are dominantly produced through a pair production process. The \bar{p}/p yield ratios show a slight increase as the collisions change from central to peripheral, which may be an indication that the baryon stopping power is decreasing toward more peripheral collisions. Within systematic uncertainties, these yield ratios agree with the published results in Au+Au collisions at $\sqrt{s_{NN}} = 200$ GeV [58].

Unlike particle yield ratios for K^+/π^+ , K^-/π^- , p/π^+ and \bar{p}/π^- as a function of $\langle N_{part} \rangle$ in U+U collisions at $\sqrt{s_{NN}} = 193$ GeV are presented in Fig. 8. Earlier published results from STAR in Au+Au collisions at $\sqrt{s_{NN}} = 200$ GeV [58] are also shown for comparison. The K^+/π^+ and K^-/π^- yield ratios gradually increase from peripheral to mid-central collisions and saturate from mid-central to central collisions. It could be due to strangeness equilibrium described in various thermodynamical models [79, 80] as well as due to baryon stopping at midrapidity [81–83]. The p/π^+ and \bar{p}/π^- yield ratios are essentially independent of collision centrality in U+U collisions at $\sqrt{s_{NN}} = 193$ GeV. The behavior of all these unlike particle yield ratios in U+U collisions at $\sqrt{s_{NN}} = 193$ GeV is similar, within the quoted uncertainties, to those from the Au+Au collisions at $\sqrt{s_{NN}} = 200$ GeV [58].

E. Kinetic Freeze-out

The evolution of the system with time in high-energy heavy-ion collisions takes the system through different stages. Typically these stages are the QGP phase, the phase transition/crossover, the hadron gas phase, the chemical freeze-out, and the kinetic freeze-out. This last stage fixes the particles' momenta and they stream freely toward the detector. This




FIG. 6. dN/dy of π^+ , K^+ , p and \bar{p} scaled by $\langle N_{\text{part}} \rangle / 2$ as a function of $\langle N_{\text{part}} \rangle$ at midrapidity ($|y| < 0.1$) for U+U collisions at $\sqrt{s_{\text{NN}}} = 193$ GeV. The results are compared with Au+Au collisions at $\sqrt{s_{\text{NN}}} = 200$ GeV [58]. The uncertainties represent total systematic and statistical uncertainties added in quadrature, dominated by systematic uncertainties.

TABLE V. dN/dy values for π^+ , π^- , K^+ , K^- , p , and \bar{p} from U+U collisions at $\sqrt{s_{\text{NN}}} = 193$ GeV within rapidity $|y| < 0.1$. The quoted errors are the total systematic and statistical errors added in quadrature, which are dominated by the systematic errors.

Centrality (%)	π^+	π^-	K^+	K^-	p	\bar{p}
0–5	375.9 ± 37.6	377.9 ± 37.8	58.3 ± 6.5	54.7 ± 6.1	38.5 ± 5.0	29.2 ± 3.8
5–10	315.4 ± 31.6	316.0 ± 31.6	48.0 ± 5.4	46.0 ± 5.2	30.4 ± 4.0	22.3 ± 2.9
10–20	235.1 ± 23.5	236.4 ± 23.7	36.2 ± 4.1	34.4 ± 3.8	21.4 ± 2.8	16.7 ± 2.2
20–30	157.1 ± 15.7	158.9 ± 15.9	24.9 ± 2.8	23.5 ± 2.6	15.0 ± 2.0	11.7 ± 1.5
30–40	102.4 ± 10.2	103.3 ± 10.3	15.8 ± 1.8	15.3 ± 1.7	9.94 ± 1.29	7.92 ± 1.03
40–50	60.9 ± 6.1	61.9 ± 6.2	9.59 ± 1.07	9.13 ± 1.02	6.21 ± 0.81	5.00 ± 0.65
50–60	35.5 ± 3.6	36.6 ± 3.6	5.28 ± 0.59	4.98 ± 0.56	3.65 ± 0.47	2.97 ± 0.39
60–70	18.5 ± 1.9	18.7 ± 1.9	2.59 ± 0.29	2.37 ± 0.27	1.89 ± 0.25	1.54 ± 0.20
70–80	8.35 ± 0.84	8.51 ± 0.86	1.07 ± 0.12	1.01 ± 0.11	0.856 ± 0.111	0.717 ± 0.093

stage is characterized by the kinetic freeze-out temperature T_k and radial flow velocity β .

The kinetic freeze-out parameters are usually calculated from the hydrodynamics based Blast-Wave model [84]. This model assumes at each point in time a locally thermalised system with temperature T_k and moving with a common radial velocity β . The transverse distribution of particles for such a radially boosted system can be written as,

$$\frac{dN}{p_T dp_T} \propto \int_0^R r dr m_T I_0 \left(\frac{p_T \sinh \rho(r)}{T_k} \right) \times K_1 \left(\frac{m_T \cosh \rho(r)}{T_k} \right), \quad (3)$$

where transverse mass $m_T = \sqrt{p_T^2 + m^2}$, m being the mass

of the hadron; $\rho(r) = \tanh^{-1} \beta$; I_0 and K_1 are the modified Bessel functions. We use a radial flow velocity profile of the form

$$\beta = \beta_s (r/R)^n, \quad (4)$$

where β_s is the surface velocity and r/R is the relative radial position in the thermal source with n being the exponent of the flow velocity profile. The average radial flow velocity $\langle \beta \rangle$ can be calculated as $\langle \beta \rangle = \frac{2}{2+n} \beta_s$.

Using the data shown in Fig. 4, the kinetic freeze-out parameters are extracted separately for the data from each centrality class by applying a simultaneous Blast-wave fit [16, 69] to the π^\pm , K^\pm and $p(\bar{p})$ spectra for a given centrality class. The results for one such fit to the 0-5% centrality class are




FIG. 7. π^-/π^+ , K^-/K^+ and \bar{p}/p ratios at midrapidity ($|y| < 0.1$) as a function of $\langle N_{\text{part}} \rangle$ in U+U collisions at $\sqrt{s_{\text{NN}}} = 193$ GeV. The results are compared with Au+Au collisions at $\sqrt{s_{\text{NN}}} = 200$ GeV [58]. The uncertainties represent total systematic and statistical uncertainties added in quadrature, dominated by systematic uncertainties.




FIG. 8. K^+/π^+ , K^-/π^- , p/π^+ and \bar{p}/π^- particle yield ratios at midrapidity ($|y| < 0.1$) as a function of $\langle N_{\text{part}} \rangle$ in U+U collisions at $\sqrt{s_{\text{NN}}} = 193$ GeV. The results are compared with Au+Au collisions at $\sqrt{s_{\text{NN}}} = 200$ GeV [58]. The uncertainties represent the total systematic and statistical uncertainties added in quadrature, dominated by systematic uncertainties.

shown in Fig. 9, illustrated as overlaid on the π^+ , K^+ and p spectra. Several general observations can be made. The Blast-wave model is very sensitive to the p_T range of the spectrum used in the fit [85]. Specifically, the low p_T part of the pion spectra are mostly affected by resonance decays. Thus, the pion spectra are fitted for $p_T > 0.5$ GeV/c. In addition, the Blast-wave model does not fit the high p_T part of the spectra [86].

The kinetic freeze-out parameters obtained from the Blast-wave fits to the U+U collisions at $\sqrt{s_{\text{NN}}} = 193$ GeV are com-

pared with those reported for the Au+Au collisions at $\sqrt{s_{\text{NN}}} = 200$ GeV for a similar p_T range [58, 69] and are shown in Fig. 10. As mentioned earlier, the pion spectra are fitted for p_T greater than 0.5 GeV/c, while kaon and proton spectra are fitted from the lowest available p_T . Typically high p_T values are limited below 1.2 GeV/c. From the left panel showing T_k vs. $\langle N_{\text{part}} \rangle$ it can be seen that T_k decreases with $\langle N_{\text{part}} \rangle$, while the middle panel shows $\langle \beta \rangle$ increasing with $\langle N_{\text{part}} \rangle$. This suggests higher rate of expansion as the collision centrality increases. For peripheral collisions, this suggest a shorter-lived




FIG. 9. Simultaneous Blast-wave fit to the π^\pm , K^\pm , $p(\bar{p})$ p_T spectra in U+U collisions at $\sqrt{s_{NN}} = 193$ GeV in 0-5% centrality class within $|y| < 0.1$. For clarity only distributions for positively charged particles are shown. The uncertainties represent the total systematic and statistical uncertainties added in quadrature.

fireball [87]. The right panel shows the correlation between T_k and $\langle\beta\rangle$ such that an increase in T_k corresponds to a decrease in $\langle\beta\rangle$ and vice-versa. In all three panels it is seen that the values obtained from U+U collisions at $\sqrt{s_{NN}} = 193$ GeV are consistent within errors with those reported for the Au+Au collisions at $\sqrt{s_{NN}} = 200$ GeV [58]. The kinetic freeze-out parameters in U+U collisions at $\sqrt{s_{NN}} = 193$ GeV are listed in Table VI.

TABLE VI. Kinetic freeze-out parameters in U+U collisions at $\sqrt{s_{NN}} = 193$ GeV. The quoted errors are statistical and systematic errors added in quadrature.

Centrality	T_k (MeV)	$\langle\beta\rangle$	n	χ^2/ndf
0–5%	92 ± 12	0.570 ± 0.041	0.970 ± 0.074	0.097
5–10%	94 ± 12	0.567 ± 0.040	0.947 ± 0.152	0.043
10–20%	99 ± 12	0.553 ± 0.037	0.987 ± 0.153	0.042
20–30%	105 ± 11	0.529 ± 0.037	1.035 ± 0.183	0.049
30–40%	110 ± 10	0.493 ± 0.036	1.180 ± 0.210	0.054
40–50%	114 ± 11	0.457 ± 0.031	1.449 ± 0.229	0.052
50–60%	121 ± 11	0.420 ± 0.030	1.564 ± 0.272	0.061
60–70%	128 ± 10	0.362 ± 0.021	2.292 ± 0.269	0.062
70–80%	134 ± 10	0.305 ± 0.015	3.158 ± 0.554	0.102

F. AMPT Model Comparison

In this section, a detailed comparison of the STAR results for U+U collisions at $\sqrt{s_{NN}} = 193$ GeV is done with the AMPT model calculations (version 2.25t7d and 2.26t9) [60]. The comparison of the AMPT model with results from Au+Au collisions at $\sqrt{s_{NN}} = 200$ GeV has been done in Ref. [88]. Such a comparison allows us to constrain the relevant partonic cross sections for the specific reactions.

In the AMPT model, the initial particle distribution is generated by the HIJING model [75]. For the results presented here, the string melting version of the AMPT model is used which incorporated partonic and hadronic interactions. To

TABLE VII. Parameter values in Lund string fragmentation and parton scattering cross section for the three sets of AMPT data used in this paper.

Parameter	Cross section σ		
	1.5 mb	3 mb	10 mb
a	0.5	0.55	2.2
b (GeV^{-2})	0.9	0.15	0.5
α_s	0.33	0.33	0.47
μ (fm^{-1})	3.2	2.265	1.8

incorporate the deformation of uranium nuclei, the AMPT model has been modified, as described in Ref. [43] and briefly discussed below. The nucleon density distribution is parametrized as a deformed Woods-Saxon profile [89]

$$\rho = \frac{\rho_0}{1 + \exp([r - R]/d)} \quad (5)$$

$$R' = R [1 + \beta_2 Y_2^0(\theta) + \beta_4 Y_4^0(\theta)] \quad (6)$$

where ρ_0 is the normal nuclear density, R is the radius of the nucleus, $Y_l^m(\theta)$ denotes the spherical harmonics, θ is the polar angle and d is the surface diffuseness parameter. For the uranium nucleus $R = 6.81$ fm, $d = 0.55$ fm and the surface deformation parameters $\beta_2 = 0.28$ and $\beta_4 = 0.093$ are taken from Ref. [43]. Various other published results having studies performed using different Woods-Saxon profiles can be found in Refs. [50, 90–92]. In Ref. [90] optimization of these parameters was done and the optimized parameters obtained were $R = 6.86$ fm, $d = 0.42$ fm, $\beta_2 = 0.265$, and $\beta_4 = 0$. It was found that for a given R , d and β_2 are anti-correlated. The tuning of parameter β_4 did not affect the results. Using the new parameters, the maximum $dN/d\eta$ value changed only slightly. In Ref. [91], it was found that there was negligible change in the N_{part} values with respect to the old parameters. N_{coll} values changed little but were within the uncertainties. Hence, the different sets of parameters would lead to similar results on the particle production. We have used three different calculations of AMPT calculated data [88], obtained from three different partonic cross sections, for which the initial parameter settings are taken from Refs. [93, 94] and are shown in Table VII. Here, a and b are the parameters in the Lund string fragmentation function [95]: $f(z) \propto z^{-1}(1-z)^1 \exp(-bm_\perp^2/z)$, where z is the light-cone momentum of the produced hadron with transverse mass m_\perp with respect to that of the fragmenting string. Further, based on the Schwinger mechanism for particle production in a strong field, the production probability is proportional to $\exp(-\pi m_\perp^2/\kappa)$, where κ is the string tension, i.e., energy in a unit length of string, and $\kappa \propto [b(2+a)]^{-1}$.

The parton scattering cross section is $\sigma \approx 9\pi\alpha_s^2/(2\mu^2)$, where α_s is the QCD coupling constant and μ is the gluon screening mass in QGP. The three calculated data sets are denoted as AMPTv1 1.5 mb, AMPTv1 10 mb, and AMPTv2 3 mb here and in the following three sections. The statistics analyzed here in the AMPTv1 1.5 mb, AMPTv2 3 mb set is about 6 M and in the AMPTv1 10 mb set is about 0.4 M.




FIG. 10. Left panel: T_k as a function of $\langle N_{\text{part}} \rangle$. Middle panel: $\langle \beta \rangle$ as a function of $\langle N_{\text{part}} \rangle$. Right panel: variation of T_k with $\langle \beta \rangle$. The midrapidity ($|y| < 0.1$) results from U+U collisions at $\sqrt{s_{\text{NN}}} = 193$ GeV are compared with Au+Au collisions at $\sqrt{s_{\text{NN}}} = 200$ GeV [58]. The uncertainties represent the total systematic and statistical uncertainties added in quadrature, dominated by systematic uncertainties.




FIG. 11. $\langle p_T \rangle$ of π^+ , K^+ and p at midrapidity ($|y| < 0.1$) as a function of $\langle N_{\text{part}} \rangle$ obtained for the three sets of AMPT calculated data (AMPTv1 1.5 mb, AMPTv2 3 mb and AMPTv1 10 mb) are shown along with the STAR results obtained in U+U collisions at $\sqrt{s_{\text{NN}}} = 193$ GeV. The width of the bands corresponds to the statistical uncertainty associated with the calculation using the AMPT model.

1. Mean p_T Comparison

The $\langle p_T \rangle$ of π^+ , K^+ and p as a function of $\langle N_{\text{part}} \rangle$ for the three sets of AMPT calculated data are compared with STAR results from U+U collisions at $\sqrt{s_{\text{NN}}} = 193$ GeV in Fig. 11. The $\langle p_T \rangle$ values of all particles are underestimated by AMPTv1 1.5 mb and AMPTv1 10 mb models. The AMPTv2 3 mb model describes the data from mid-central to central collisions for π^+ . For K^+ and p , the AMPTv2 3 mb model describes the data within uncertainties for all centralities.

2. dN/dy Comparison

The yields, dN/dy scaled by $\langle N_{\text{part}} \rangle / 2$ for the three calculations of AMPT data are plotted as a function of $\langle N_{\text{part}} \rangle$ as shown in Fig. 12. For comparison, the measured values obtained from U+U collisions at $\sqrt{s_{\text{NN}}} = 193$ GeV are also shown. For pions and kaons, the AMPTv1 10 mb case shows decidedly better agreement with the measured values. For the protons it is seen that agreement between the AMPT calculated values and the measurements depends on the central-

ity of the collisions with the AMPTv1 10 mb values agreeing better toward peripheral collisions while the AMPTv1 1.5 mb values agree better toward more central collisions. On the other hand, in case of Au+Au collisions at $\sqrt{s_{\text{NN}}} = 200$ GeV, the calculations from the AMPT 1.5 mb case show a better agreement than the AMPT 10 mb case with the charged particle multiplicity density at midrapidity as in Ref. [93].

3. Particle Ratios Comparison

The various anti-particle to particle ratios as a function of $\langle N_{\text{part}} \rangle$ obtained from measured data are compared to these ratios that were calculated in the three AMPT data calculations as shown in Fig. 13. It is observed that these ratios have similar values in all the AMPT cases and agree with STAR data across all centrality classes.

Ratios of unlike particles obtained from the three AMPT data calculations are compared with the STAR results and presented in Fig. 14. The K^+/π^+ and K^-/π^- ratios for the AMPTv1 10 mb and AMPTv2 3mb cases are in good agreement with the STAR data within errors while the AMPT 1.5




FIG. 12. dN/dy for π^+ , K^+ and p scaled by $\langle N_{\text{part}} \rangle / 2$ at midrapidity ($|y| < 0.1$) as a function of $\langle N_{\text{part}} \rangle$ obtained for three calculations of AMPT data (AMPTv1 1.5 mb, AMPTv2 3 mb, and AMPTv1 10 mb) are shown along with the STAR results obtained in U+U collisions at $\sqrt{s_{\text{NN}}} = 193$ GeV. The width of the bands corresponds to the statistical uncertainty associated with the calculation using the AMPT model.




FIG. 13. π^-/π^+ , K^-/K^+ and \bar{p}/p particle yield ratios at midrapidity ($|y| < 0.1$) as a function of $\langle N_{\text{part}} \rangle$ obtained for three calculations of AMPT data (AMPTv1 1.5 mb, AMPTv2 3 mb, and AMPTv1 10 mb) are shown along with the STAR results obtained in U+U collisions at $\sqrt{s_{\text{NN}}} = 193$ GeV. The width of the bands corresponds to the statistical uncertainty associated with the calculation using the AMPT model.

mb case gives values significantly below the STAR data. For the p/π^+ , \bar{p}/π^- cases neither AMPT data set agrees with the data over all centrality classes. The AMPTv1 10 mb data set agrees better for peripheral collisions while the AMPTv1 1.5 mb and AMPTv2 3mb data set agrees better toward central collisions.

VI. SUMMARY

Various basic bulk observables regarding identified particle production in U+U collisions at $\sqrt{s_{\text{NN}}} = 193$ GeV have been presented. The transverse momentum spectra of π^\pm , K^\pm and $p(\bar{p})$ in midrapidity ($|y| < 0.1$) have been measured for nine centrality classes : 0-5%, 5-10%, 10-20%, 20-30%, 30-40%, 40-50%, 50-60%, 60-70% and 70-80%. Other extracted observables from p_T spectra such as average transverse momentum ($\langle p_T \rangle$), particle yields (dN/dy), particle ratios, and kinetic freeze-out properties are also presented as functions of collision centrality or $\langle N_{\text{part}} \rangle$. These observables are compared with the corresponding results from Au+Au collisions

at $\sqrt{s_{\text{NN}}} = 200$ GeV and the AMPT model modified to incorporate the deformation of uranium nucleus.

The mean $\langle p_T \rangle$ values for π , K , and p increase from peripheral to central collisions in U+U collisions at $\sqrt{s_{\text{NN}}} = 193$ GeV. This is an indication of increasing radial flow effects in more central collisions. The increase in $\langle p_T \rangle$ from π to K and to p indicates that the radial flow effect increases with particle mass.

The integrated particle yields dN/dy in midrapidity $|y| < 0.1$ of π^\pm , K^\pm , and p do not scale with $\langle N_{\text{part}} \rangle$, but rather they slowly increase from peripheral to central collisions in U+U collisions at $\sqrt{s_{\text{NN}}} = 193$ GeV. This indicates that, at this energy, the particle production mechanism has contributions from hard processes involving nucleon-nucleon binary collisions. In contrast, \bar{p} shows no such dependence with centrality of the collision.

Anti-particle to particle yield ratios are close to unity in U+U collisions at $\sqrt{s_{\text{NN}}} = 193$ GeV, indicating pair production is the dominant mechanism of particle production at maximum RHIC energy. Their values are also close to those obtained from Au+Au collisions at $\sqrt{s_{\text{NN}}} = 200$ GeV. The




FIG. 14. K^+/π^+ , K^-/π^- , p/π^+ and \bar{p}/π^- particle yield ratios at midrapidity ($|y| < 0.1$) as a function of $\langle N_{\text{part}} \rangle$ obtained for three calculations of AMPT data (AMPTv1 1.5 mb, AMPTv2 3 mb, and AMPTv1 10 mb) are shown along with the STAR results obtained in U+U collisions at $\sqrt{s_{\text{NN}}} = 193$ GeV. The width of the bands corresponds to the statistical uncertainty associated with the calculation using the AMPT model.

K^+/π^+ and K^-/π^- yield ratio increases from peripheral to mid-central collision after which they saturate up to central collisions. The p/π^+ and \bar{p}/π^- ratios do not show any significant centrality dependence in U+U collisions at $\sqrt{s_{\text{NN}}} = 193$ GeV.

The kinetic freeze-out parameters are obtained from the simultaneous Blast-wave fit to π^\pm , K^\pm , and $p(\bar{p})$ p_T spectra in U+U collisions at $\sqrt{s_{\text{NN}}} = 193$ GeV. The kinetic freeze-out temperature T_k decreases from peripheral to central collisions while the average flow velocity $\langle \beta \rangle$ increases from peripheral to central collisions. This indicates a large radial flow effects for central collisions and a short-lived fireball in peripheral collisions. The extracted T_k and $\langle \beta \rangle$ values for similar $\langle N_{\text{part}} \rangle$ values are consistent between Au+Au collisions at $\sqrt{s_{\text{NN}}} = 200$ GeV and U+U collisions at $\sqrt{s_{\text{NN}}} = 193$ GeV.

A detailed comparison of integrated particle yields, mean transverse momentum, and particle yield ratios from STAR results in U+U collisions at $\sqrt{s_{\text{NN}}} = 193$ GeV is carried out with three cases of AMPT string melting corresponding to parton-parton interaction cross sections of 1.5, 3, and 10 mb that was also modified to incorporate the deformation in uranium nucleus. $\langle p_T \rangle$ obtained from the AMPTv1 1.5 mb and AMPTv1 10 mb have lower values than STAR results for all centrality classes while AMPTv2 3 mb could describe pions from mid-central to central collisions and kaons and protons for all centralities. The dN/dy from the AMPTv1 10 mb data are closer to the STAR results for pions and kaons than the AMPTv1 1.5 mb and AMPTv2 3 mb data. However, for protons, dN/dy in central collisions are better explained by the

AMPTv1 1.5 mb data but for peripheral collisions there is better agreement with the AMPTv1 10 mb data. Anti-particle to particle yield ratios are successfully explained by all of the AMPT models implying that they are insensitive to these changes in the parton-parton cross sections. The K/π yield ratios from the AMPTv1 10 mb and AMPTv2 3 mb cases agree with the measurements for all centrality classes, whereas the results from the AMPTv1 1.5 mb case underpredict the STAR results. In case of p/π ratios, AMPTv1 1.5 mb and AMPTv2 3 mb cases more closely explains the STAR results toward central collisions while the AMPTv1 10 mb case seems to be closer to STAR results for peripheral collisions. It can be observed that the three AMPT model cases do not describe all the observables presented here at all $\langle N_{\text{part}} \rangle$ values consistently. The AMPTv2 3 mb model having smaller a and b values describes the $\langle p_T \rangle$ of data better than others for the π^\pm , K^\pm , and $p(\bar{p})$. The AMPTv1 10 mb model having largest cross section, describes the dN/dy of π^\pm , K^\pm , and $p(\bar{p})$ better than the other cases. The antiparticle to particle ratios are described by all the three model cases. These findings call for further investigation into the AMPT model in terms of the particle production mechanism and the collision dynamics.

We have compared the results from U+U collisions to those from Au+Au collisions at similar center-of-mass energies, but the colliding nuclei have different geometrical shapes and hence different initial collision conditions. The presented physical observables: dN/dy , $\langle p_T \rangle$, particle ratios, T_k and $\langle \beta \rangle$ are consistent between U+U and Au+Au for a similar range of $\langle N_{\text{part}} \rangle$ values. This might suggest that, when all the dif-

ferent initial state orientations of the colliding uranium nuclei are combined in the analysis, the resulting final state values approximate what is observed from colliding spherically symmetric nuclei. In other words, these observables are governed by the $\langle N_{\text{part}} \rangle$ for this analysis. In future experiments involving colliding uranium nuclei, it would be interesting to measure the different final state observables for identified orientations of the colliding ellipsoidal uranium nuclei. It has been suggested that this might be done for the most central collisions where the different collision configurations can be selected by a data driven method using selections based on the probability distribution of the charged particle multiplicity and the event anisotropy for U+U collisions [96]. In addition, as discussed in Ref. [44], the spectator neutron measurements can also be used to select certain initial configurations.

VII. ACKNOWLEDGEMENTS

We thank Jun Xu and Zi-Wei Lin for discussions related to the AMPT results in the paper. We thank the RHIC Operations

Group and RCF at BNL, the NERSC Center at LBNL, and the Open Science Grid consortium for providing resources and support. This work was supported in part by the Office of Nuclear Physics within the U.S. DOE Office of Science, the U.S. National Science Foundation, National Natural Science Foundation of China, Chinese Academy of Science, the Ministry of Science and Technology of China and the Chinese Ministry of Education, the Higher Education Sprout Project by Ministry of Education at NCKU, the National Research Foundation of Korea, Czech Science Foundation and Ministry of Education, Youth and Sports of the Czech Republic, Hungarian National Research, Development and Innovation Office, New National Excellency Programme of the Hungarian Ministry of Human Capacities, Department of Atomic Energy and Department of Science and Technology of the Government of India, the National Science Centre and WUT ID-UB of Poland, the Ministry of Science, Education and Sports of the Republic of Croatia, German Bundesministerium für Bildung, Wissenschaft, Forschung und Technologie (BMBF), Helmholtz Association, Ministry of Education, Culture, Sports, Science, and Technology (MEXT) and Japan Society for the Promotion of Science (JSPS).

-
- [1] E. V. Shuryak, Phys. Rept. **61**, 71 (1980).
 - [2] M. Gyulassy and L. McLerran, Nucl. Phys. A **750**, 30 (2005).
 - [3] J. P. Blaizot and E. Iancu, Phys. Rept. **359**, 355 (2002).
 - [4] A. Bazavov *et al.*, Phys. Rev. D **85**, 054503 (2012).
 - [5] Y. Aoki, G. Endrodi, Z. Fodor, S. D. Katz and K. K. Szabo, Nature **443**, 675 (2006).
 - [6] I. Arsene *et al.* [BRAHMS Collaboration], Nucl. Phys. A **757**, 1 (2005).
 - [7] K. Adcox *et al.* [PHENIX Collaboration], Nucl. Phys. A **757**, 184 (2005).
 - [8] B. B. Back *et al.* [PHOBOS Collaboration], Nucl. Phys. A **757**, 28 (2005).
 - [9] J. Adams *et al.* [STAR Collaboration], Nucl. Phys. A **757**, 102 (2005).
 - [10] D. Kharzeev and M. Nardi, Phys. Lett. B **507**, 121 (2001).
 - [11] K. Rajagopal and F. Wilczek, In *At the frontier of particle physics, Handbook of QCD*, edited by M. Shifman (World Scientific, Singapore), Vol. 3. [hep-ph/0011333].
 - [12] E. Laermann and O. Philipsen, Ann. Rev. Nucl. Part. Sci. **53**, 163 (2003).
 - [13] M. A. Stephanov, PoS LAT **2006**, 024 (2006).
 - [14] STAR Note, <https://drupal.star.bnl.gov/STAR/starnotes/public/sn0493>.
 - [15] STAR Note, BES-II white paper, <https://drupal.star.bnl.gov/STAR/starnotes/public/sn0598>.
 - [16] B. I. Abelev *et al.* [STAR Collaboration], Phys. Rev. C **81**, 024911 (2010).
 - [17] P. Braun-Munzinger, J. Stachel, Nature **448**, 302-309, (2007).
 - [18] C. Adler *et al.* [STAR Collaboration], Phys. Rev. Lett. **89**, 202301 (2002).
 - [19] C. Adler *et al.* [STAR Collaboration], Phys. Rev. Lett. **90**, 082302 (2003).
 - [20] J. Adams *et al.* [STAR Collaboration], Phys. Rev. Lett. **91**, 172302 (2003).
 - [21] J. Adams *et al.* [STAR Collaboration], Phys. Rev. Lett. **92**, 052302 (2004).
 - [22] B. I. Abelev *et al.* [STAR Collaboration], Phys. Rev. Lett. **97**, 152301 (2006).
 - [23] B. I. Abelev *et al.* [STAR Collaboration], Phys. Lett. B **655**, 104 (2007).
 - [24] J. Adams *et al.* [STAR Collaboration], Phys. Rev. Lett. **97**, 162301 (2006).
 - [25] S. S. Adler *et al.* [PHENIX Collaboration], Phys. Rev. Lett. **91**, 072301 (2003).
 - [26] I. Arsene *et al.* [BRAHMS Collaboration], Phys. Rev. Lett. **91**, 072305 (2003).
 - [27] B. B. Back *et al.* [PHOBOS Collaboration], Phys. Lett. B **578**, 297 (2004).
 - [28] K. Aamodt *et al.* [ALICE Collaboration], Phys. Lett. B **696**, 30 (2011).
 - [29] S. Chatrchyan *et al.* [CMS Collaboration], Eur. Phys. J. C **72**, 1945 (2012).
 - [30] K. H. Ackermann *et al.* [STAR Collaboration], Phys. Rev. Lett. **86**, 402 (2001).
 - [31] C. Adler *et al.* [STAR Collaboration], Phys. Rev. Lett. **87**, 182301 (2001).
 - [32] C. Adler *et al.* [STAR Collaboration], Phys. Rev. Lett. **89**, 132301 (2002).
 - [33] J. Adams *et al.* [STAR Collaboration], Phys. Rev. Lett. **92**, 062301 (2004).
 - [34] J. Adams *et al.* [STAR Collaboration], Phys. Rev. Lett. **95**, 122301 (2005).
 - [35] B. I. Abelev *et al.* [STAR Collaboration], Phys. Rev. Lett. **99**, 112301 (2007).
 - [36] L. Adamczyk *et al.* [STAR Collaboration], Phys. Rev. Lett. **116**, 062301 (2016).
 - [37] L. Adamczyk *et al.* [STAR Collaboration], Phys. Rev. Lett. **118**, 212301 (2017).
 - [38] B. Alver *et al.* [PHOBOS Collaboration], Phys. Rev. Lett. **98**, 242302 (2007).

- [39] S. S. Adler *et al.* [PHENIX Collaboration], Phys. Rev. Lett. **91**, 182301 (2003).
- [40] B. B. Abelev *et al.* [ALICE Collaboration], J. High Energy Phys. **06** (2015) 190.
- [41] H. Masui, B. Mohanty and N. Xu, Phys. Lett. B **679**, 440 (2009).
- [42] M. R. Haque, M. Nasim and B. Mohanty, Phys. Rev. C **84**, 067901 (2011).
- [43] M. R. Haque, Z. W. Lin and B. Mohanty, Phys. Rev. C **85**, 034905 (2012).
- [44] V. Bairathi, M. R. Haque and B. Mohanty, Phys. Rev. C **91**, no. 5, 054903 (2015).
- [45] L. Adamczyk *et al.* [STAR Collaboration], Phys. Rev. Lett. **118**, 012301 (2017).
- [46] G. Agakishiev *et al.* [STAR Collaboration], Phys. Rev. Lett. **108**, 072301 (2012).
- [47] A. Adare *et al.* [PHENIX Collaboration], Phys. Rev. Lett. **98**, 162301 (2007).
- [48] A. Adare *et al.* [PHENIX Collaboration], Phys. Rev. Lett. **101**, 162301 (2008).
- [49] A. Adare *et al.* [PHENIX Collaboration], Phys. Rev. Lett. **121**, 222301 (2018).
- [50] L. Adamczyk *et al.* [STAR Collaboration], Phys. Rev. Lett. **115**, 222301 (2015).
- [51] J. Adam *et al.* [STAR Collaboration], Phys. Rev. Lett. **121**, 132301 (2018).
- [52] U. W. Heinz and A. Kuhlman, Phys. Rev. Lett. **94**, 132301 (2005).
- [53] A. J. Kuhlman and U. W. Heinz, Phys. Rev. C **72**, 037901 (2005).
- [54] A. Kuhlman, U. W. Heinz and Y. V. Kovchegov, Phys. Lett. B **638**, 171 (2006).
- [55] C. Nepali, G. Fai and D. Keane, Phys. Rev. C **73**, 034911 (2006).
- [56] C. Nepali, G. Fai and D. Keane, Phys. Rev. C **76**, 051902(R) (2007); **76**, 069903(E) (2007).
- [57] T. Hirano, P. Huovinen and Y. Nara, Phys. Rev. C **83**, 021902(R) (2011).
- [58] B. I. Abelev *et al.* [STAR Collaboration], Phys. Rev. C **79**, 034909 (2009).
- [59] J. Adams *et al.* [STAR Collaboration], Phys. Rev. Lett. **92**, 112301 (2004).
- [60] Z. W. Lin, C. M. Ko, B. A. Li, B. Zhang and S. Pal, Phys. Rev. C **72**, 064901 (2005).
- [61] K. H. Ackermann *et al.* [STAR Collaboration], Nucl. Instrum. Meth. A **499**, 624 (2003).
- [62] W. J. Llope *et al.*, Nucl. Instrum. Meth. A **522**, 252 (2004).
- [63] W. J. Llope *et al.*, Nucl. Instrum. Meth. A **759**, 23 (2014).
- [64] C. Adler, A. Denisov, E. Garcia, M. J. Murray, H. Strobele and S. N. White, Nucl. Instrum. Meth. A **470**, 488 (2001).
- [65] C. Adler, A. Denisov, E. Garcia, M. Murray, H. Strobele and S. White, Nucl. Instrum. Meth. A **499**, 433 (2003).
- [66] M. Anderson *et al.*, Nucl. Instrum. Meth. A **499**, 659 (2003).
- [67] J. Wu *et al.*, Nucl. Instrum. Meth. A **538**, 243 (2005).
- [68] L. Adamczyk *et al.* [STAR Collaboration], Phys. Rev. C **94**, no. 6, 064904 (2016).
- [69] L. Adamczyk *et al.* [STAR Collaboration], Phys. Rev. C **96**, no. 4, 044904 (2017).
- [70] M. Calderón de la Barca Sánchez, Ph.D. thesis (Yale University, 2001) (unpublished), arXiv:nucl-ex/**0111004**.
- [71] H. Bichsel, Nucl. Instrum. Meth. A **562**, 154 (2006).
- [72] M. Aguilar-Benitez *et al.*, Z. Phys. C **50**, 405 (1991).
- [73] M. Shao, O. Y. Barannikova, X. Dong, Y. Fisyak, L. Ruan, P. Sorensen and Z. Xu, Nucl. Instrum. Meth. A **558**, 419 (2006).
- [74] V. Fine and P. Nevski, Proc. CHEP **2000**, 143 (2000).
- [75] X. N. Wang and M. Gyulassy, Phys. Rev. D **44**, 3501 (1991).
- [76] C. Adler *et al.* [STAR Collaboration], Phys. Rev. Lett. **86**, 4778 (2001); **90**, 119903(E) (2003).
- [77] C. Adler *et al.* [STAR Collaboration], Phys. Rev. Lett. **87**, 262302 (2001).
- [78] U. Acharya *et al.* [PHENIX Collaboration], Phys. Rev. C **102**, no.6, 064905 (2020).
- [79] M. Kaneta and N. Xu, arXiv:nucl-th/0405068.
- [80] J. Cleymans, B. Kampfer, M. Kaneta, S. Wheaton and N. Xu, Phys. Rev. C **71**, 054901 (2005).
- [81] F. q. Wang, Phys. Lett. B **489**, 273 (2000).
- [82] F. q. Wang and N. Xu, Phys. Rev. C **61**, 021904 (2000).
- [83] P. Braun-Munzinger, J. Cleymans, H. Oeschler and K. Redlich, Nucl. Phys. A **697**, 902 (2002).
- [84] E. Schnedermann, J. Sollfrank and U. W. Heinz, Phys. Rev. C **48**, 2462 (1993).
- [85] B. Abelev *et al.* [ALICE Collaboration], Phys. Rev. C **88**, 044910 (2013).
- [86] G. Wilk and Z. Wlodarczyk, Phys. Rev. Lett. **84**, 2770 (2000).
- [87] U. W. Heinz, arXiv:hep-ph/0407360.
- [88] A. Nandi, L. Kumar and N. Sharma, Phys. Rev. C **102**, no.2, 024902 (2020).
- [89] K. Hagino, N. W. Lwin and M. Yamagami, Phys. Rev. C **74**, 017310 (2006).
- [90] Q. Y. Shou, Y. G. Ma, P. Sorensen, A. H. Tang, F. Videbæk and H. Wang, Phys. Lett. B **749**, 215-220 (2015).
- [91] A. Adare *et al.* [PHENIX Collaboration], Phys. Rev. C **93**, no.3, 034903 (2016).
- [92] A. Adare *et al.* [PHENIX Collaboration], Phys. Rev. C **93**, no.2, 024901 (2016).
- [93] J. Xu and C. M. Ko, Phys. Rev. C **84**, 014903 (2011).
- [94] G. L. Ma and Z. W. Lin, Phys. Rev. C **93**, no.5, 054911 (2016)
- [95] Z. w. Lin, S. Pal, C. M. Ko, B. A. Li and B. Zhang, Phys. Rev. C **64**, 011902 (2001).
- [96] A. Goldschmidt, Z. Qiu, C. Shen and U. Heinz, Phys. Rev. C **92**, no. 4, 044903 (2015).