

# Directed Flow of Identified Particles in Au + Au Collisions at $\sqrt{s_{\text{NN}}} = 200$ GeV at RHIC

Adamczyk,<sup>9</sup> G. Agakishiev,<sup>19</sup> M. M. Aggarwal,<sup>30</sup> Z. Ahammed,<sup>48</sup> A. V. Alakhverdyants,<sup>19</sup> I. Alekseev,<sup>17</sup> J. Alford,<sup>20</sup> B. D. Anderson,<sup>20</sup> C. D. Anson,<sup>28</sup> D. Arkhipkin,<sup>2</sup> G. S. Averichev,<sup>19</sup> J. Balewski,<sup>24</sup> Banerjee,<sup>48</sup> Z. Barnovska,<sup>12</sup> D. R. Beavis,<sup>2</sup> R. Bellwied,<sup>44</sup> M. J. Betancourt,<sup>24</sup> R. R. Betts,<sup>8</sup> A. Bhasin,<sup>18</sup> A. K. Bhati,<sup>30</sup> H. Bichsel,<sup>50</sup> J. Bielcik,<sup>11</sup> J. Bielcikova,<sup>12</sup> L. C. Bland,<sup>2</sup> I. G. Bordyuzhin,<sup>17</sup> W. Borowski,<sup>41</sup> J. Bouchet,<sup>20</sup> A. V. Brandin,<sup>27</sup> S. G. Brovko,<sup>4</sup> E. Bruna,<sup>52</sup> S. Bueltmann,<sup>29</sup> I. Bunzarov,<sup>19</sup> T. P. Burton,<sup>2</sup> J. Butterworth,<sup>36</sup> X. Z. Cai,<sup>40</sup> H. Caines,<sup>52</sup> M. Calderón de la Barca Sánchez,<sup>4</sup> D. Cebra,<sup>4</sup> R. Cendejas,<sup>5</sup> M. C. Cervantes,<sup>42</sup> P. Chaloupka,<sup>12</sup> S. Chattopadhyay,<sup>48</sup> H. F. Chen,<sup>38</sup> J. H. Chen,<sup>40</sup> J. Y. Chen,<sup>7</sup> L. Chen,<sup>7</sup> J. Cheng,<sup>45</sup> M. Cherney,<sup>10</sup> A. Chikanian,<sup>52</sup> W. Christie,<sup>2</sup> P. Chung,<sup>12</sup> J. Chwastowski,<sup>9</sup> M. J. M. Coddington,<sup>42</sup> R. Corliss,<sup>24</sup> J. G. Cramer,<sup>50</sup> H. J. Crawford,<sup>3</sup> X. Cui,<sup>38</sup> A. Davila Leyva,<sup>43</sup> L. C. De Silva,<sup>44</sup> R. R. Debebe,<sup>2</sup> T. G. Dedovich,<sup>19</sup> J. Deng,<sup>39</sup> R. Derradi de Souza,<sup>6</sup> S. Dhamija,<sup>16</sup> L. Didenko,<sup>2</sup> F. Ding,<sup>4</sup> P. Djawotho,<sup>42</sup> X. Dong,<sup>23</sup> J. L. Drachenberg,<sup>42</sup> J. E. Draper,<sup>4</sup> C. M. Du,<sup>22</sup> L. E. Dunkelberger,<sup>5</sup> J. C. Dunlop,<sup>2</sup> L. G. Efimov,<sup>19</sup> M. Elnimr,<sup>51</sup> J. Engelage,<sup>3</sup> G. Eppley,<sup>36</sup> L. Eun,<sup>23</sup> O. Evdokimov,<sup>8</sup> R. Fatemi,<sup>21</sup> J. Fedorisin,<sup>19</sup> R. G. Fersch,<sup>21</sup> P. Filip,<sup>19</sup> E. Finch,<sup>52</sup> Y. Fisyak,<sup>2</sup> C. A. Gagliardi,<sup>42</sup> D. R. Gangadharan,<sup>28</sup> F. Geurts,<sup>36</sup> S. Gliske,<sup>1</sup> Y. N. Gorbunov,<sup>10</sup> O. G. Grebenyuk,<sup>23</sup> D. Grosnick,<sup>47</sup> S. Gupta,<sup>18</sup> W. Guryn,<sup>2</sup> B. Haag,<sup>4</sup> O. Hajkova,<sup>11</sup> A. Hamed,<sup>42</sup> L-X. Han,<sup>40</sup> J. W. Harris,<sup>52</sup> J. P. Hays-Wehle,<sup>24</sup> S. Heppelmann,<sup>31</sup> A. Hirsch,<sup>33</sup> G. W. Hoffmann,<sup>43</sup> D. J. Hofman,<sup>8</sup> S. Horvat,<sup>52</sup> B. Huang,<sup>38</sup> H. Z. Huang,<sup>5</sup> P. Huck,<sup>7</sup> T. J. Humanic,<sup>28</sup> L. Huo,<sup>42</sup> G. Igo,<sup>5</sup> W. W. Jacobs,<sup>16</sup> C. Jena,<sup>14</sup> J. Joseph,<sup>20</sup> E. G. Judd,<sup>3</sup> S. Kabana,<sup>41</sup> K. Kang,<sup>45</sup> J. Kapitan,<sup>12</sup> K. Kauder,<sup>8</sup> H. W. Ke,<sup>7</sup> D. Keane,<sup>20</sup> A. Kechechyan,<sup>19</sup> A. Kesich,<sup>4</sup> D. Kettler,<sup>50</sup> D. P. Kikola,<sup>33</sup> J. Kiryluk,<sup>23</sup> A. Kisiel,<sup>49</sup> V. Kizka,<sup>19</sup> S. R. Klein,<sup>23</sup> D. D. Koetke,<sup>47</sup> T. Kollegger,<sup>13</sup> J. Konzer,<sup>33</sup> I. Koralt,<sup>29</sup> L. Koroleva,<sup>17</sup> W. Korsch,<sup>21</sup> L. Kotchenda,<sup>27</sup> P. Kravtsov,<sup>27</sup> K. Krueger,<sup>1</sup> L. Kumar,<sup>20</sup> M. A. C. Lamont,<sup>2</sup> J. M. Landgraf,<sup>2</sup> S. LaPointe,<sup>51</sup> J. Lauret,<sup>2</sup> A. Lebedev,<sup>2</sup> R. Lednicky,<sup>19</sup> J. H. Lee,<sup>2</sup> W. Leight,<sup>24</sup> M. J. LeVine,<sup>2</sup> C. Li,<sup>38</sup> L. Li,<sup>43</sup> W. Li,<sup>40</sup> X. Li,<sup>33</sup> X. Li,<sup>39</sup> Y. Li,<sup>45</sup> Z. M. Li,<sup>7</sup> L. M. Lima,<sup>37</sup> M. A. Lisa,<sup>28</sup> F. Liu,<sup>7</sup> T. Ljubicic,<sup>2</sup> W. J. Llope,<sup>36</sup> R. S. Longacre,<sup>2</sup> Y. Lu,<sup>38</sup> X. Luo,<sup>7</sup> A. Luszczak,<sup>9</sup> G. L. Ma,<sup>40</sup> Y. G. Ma,<sup>40</sup> D. P. Mahapatra,<sup>14</sup> R. Majka,<sup>52</sup> O. I. Mall,<sup>4</sup> S. Margetis,<sup>20</sup> C. Markert,<sup>43</sup> H. Masui,<sup>23</sup> H. S. Matis,<sup>23</sup> D. McDonald,<sup>36</sup> T. S. McShane,<sup>10</sup> S. Mioduszewski,<sup>42</sup> M. K. Mitrovski,<sup>2</sup> Y. Mohammed,<sup>42</sup> B. Mohanty,<sup>48</sup> B. Morozov,<sup>17</sup> M. G. Munhoz,<sup>37</sup> M. K. Mustafa,<sup>33</sup> M. Naglis,<sup>23</sup> B. K. Nandi,<sup>15</sup> Md. Nasim,<sup>48</sup> T. K. Nayak,<sup>48</sup> L. V. Nogach,<sup>32</sup> G. Odyniec,<sup>23</sup> A. Ogawa,<sup>2</sup> K. Oh,<sup>34</sup> A. Ohlson,<sup>52</sup> V. Okorokov,<sup>27</sup> E. W. Oldag,<sup>43</sup> R. A. N. Oliveira,<sup>37</sup> D. Olson,<sup>23</sup> M. Pachr,<sup>11</sup> B. S. Page,<sup>16</sup> S. K. Pal,<sup>48</sup> Pan,<sup>5</sup> Y. Pandit,<sup>20</sup> Y. Panebratsev,<sup>19</sup> T. Pawlak,<sup>49</sup> B. Pawlik,<sup>9</sup> H. Pei,<sup>8</sup> C. Perkins,<sup>3</sup> W. Peryt,<sup>49</sup> P. Pile,<sup>2</sup> M. Planinic,<sup>53</sup> J. Pluta,<sup>49</sup> D. Plyku,<sup>29</sup> N. Poljak,<sup>53</sup> J. Porter,<sup>23</sup> A. M. Poskanzer,<sup>23</sup> C. B. Powell,<sup>23</sup> D. Prindle,<sup>50</sup> C. Pruneau,<sup>51</sup> N. K. Pruthi,<sup>30</sup> M. Przybycien,<sup>9</sup> P. R. Pujahari,<sup>15</sup> J. Putschke,<sup>51</sup> H. Qiu,<sup>22</sup> R. Raniwala,<sup>35</sup> S. Raniwala,<sup>35</sup> R. L. Ray,<sup>43</sup> R. Redwine,<sup>24</sup> R. Reed,<sup>4</sup> C. K. Riley,<sup>52</sup> H. G. Ritter,<sup>23</sup> J. B. Roberts,<sup>36</sup> O. V. Rogachevskiy,<sup>19</sup> J. L. Romero,<sup>4</sup> L. Ruan,<sup>2</sup> J. Rusnak,<sup>12</sup> N. R. Sahoo,<sup>48</sup> I. Sakrejda,<sup>23</sup> S. Salur,<sup>23</sup> J. Sandweiss,<sup>52</sup> E. Sangaline,<sup>4</sup> A. Sarkar,<sup>15</sup> J. Schambach,<sup>43</sup> R. P. Scharenberg,<sup>33</sup> A. M. Schmah,<sup>23</sup> N. Schmitz,<sup>25</sup> T. R. Schuster,<sup>13</sup> J. Seele,<sup>24</sup> J. Seger,<sup>10</sup> P. Seyboth,<sup>25</sup> N. Shah,<sup>5</sup> E. Shaliev,<sup>19</sup> M. Shao,<sup>38</sup> B. Sharma,<sup>30</sup> M. Sharma,<sup>51</sup> S. S. Shi,<sup>7</sup> Q. Y. Shou,<sup>40</sup> E. P. Sichtermann,<sup>23</sup> R. N. Singaraju,<sup>48</sup> M. J. Skoby,<sup>33</sup> N. Smirnov,<sup>52</sup> D. Solanki,<sup>35</sup> P. Sorensen,<sup>2</sup> U. G. deSouza,<sup>37</sup> H. M. Spinka,<sup>1</sup> B. Srivastava,<sup>33</sup> T. D. S. Stanislaus,<sup>47</sup> S. G. Steadman,<sup>24</sup> J. R. Stevens,<sup>16</sup> R. Stock,<sup>13</sup> M. Strikhanov,<sup>27</sup> B. Stringfellow,<sup>33</sup> A. A. P. Suaide,<sup>37</sup> M. C. Suarez,<sup>8</sup> M. Sumner,<sup>12</sup> X. M. Sun,<sup>23</sup> Y. Sun,<sup>38</sup> Z. Sun,<sup>22</sup> B. Surrow,<sup>24</sup> D. N. Svirida,<sup>17</sup> T. J. M. Symons,<sup>23</sup> A. Szanto de Toledo,<sup>37</sup> J. Takahashi,<sup>6</sup> A. H. Tang,<sup>2</sup> Z. Tang,<sup>38</sup> L. H. Tarini,<sup>51</sup> T. Tarnowsky,<sup>26</sup> D. Thein,<sup>43</sup> J. H. Thomas,<sup>23</sup> J. Tian,<sup>40</sup> A. R. Timmins,<sup>44</sup> D. Tlusty,<sup>12</sup> M. Tokarev,<sup>19</sup> T. A. Trainor,<sup>50</sup> S. Trentalange,<sup>5</sup> R. E. Tribble,<sup>42</sup> P. Tribedy,<sup>48</sup> B. A. Trzeciak,<sup>49</sup> O. D. Tsai,<sup>5</sup> J. Turnau,<sup>9</sup> T. Ullrich,<sup>2</sup> D. G. Underwood,<sup>1</sup> G. Van Buren,<sup>2</sup> G. van Nieuwenhuizen,<sup>24</sup> J. A. Vanfossen, Jr.,<sup>20</sup> R. Varma,<sup>15</sup> G. M. S. Vasconcelos,<sup>6</sup> F. Videbæk,<sup>2</sup> Y. P. Vijoyi,<sup>48</sup> S. Vokal,<sup>19</sup> S. A. Voloshin,<sup>51</sup> A. Vossen,<sup>16</sup> M. Wada,<sup>43</sup> F. Wang,<sup>33</sup> G. Wang,<sup>5</sup> H. Wang,<sup>26</sup> J. S. Wang,<sup>22</sup> Q. Wang,<sup>33</sup> X. L. Wang,<sup>38</sup> Y. Wang,<sup>45</sup> G. Webb,<sup>21</sup> J. C. Webb,<sup>2</sup> G. D. Westfall,<sup>26</sup> C. Whitten Jr.,<sup>5</sup> H. Wieman,<sup>23</sup> S. W. Wissink,<sup>16</sup> R. Witt,<sup>46</sup> W. Witzke,<sup>21</sup> Y. F. Wu,<sup>7</sup> Z. Xiao,<sup>45</sup> W. Xie,<sup>33</sup> K. Xin,<sup>36</sup> H. Xu,<sup>22</sup> N. Xu,<sup>23</sup> Q. H. Xu,<sup>39</sup> W. Xu,<sup>5</sup> Y. Xu,<sup>38</sup> Z. Xu,<sup>2</sup> L. Xue,<sup>40</sup> Y. Yang,<sup>22</sup> Y. Yang,<sup>7</sup> P. Yepes,<sup>36</sup> Y. Yi,<sup>33</sup> K. Yip,<sup>2</sup>

I-K. Yoo,<sup>34</sup> M. Zawisza,<sup>49</sup> H. Zbroszczyk,<sup>49</sup> J. B. Zhang,<sup>7</sup> S. Zhang,<sup>40</sup> W. M. Zhang,<sup>20</sup> X. P. Zhang,<sup>45</sup>  
 Y. Zhang,<sup>38</sup> Z. P. Zhang,<sup>38</sup> F. Zhao,<sup>5</sup> J. Zhao,<sup>40</sup> C. Zhong,<sup>40</sup> X. Zhu,<sup>45</sup> Y. H. Zhu,<sup>40</sup> and Y. Zoukarnieva<sup>19</sup>

(STAR Collaboration)

- <sup>1</sup>Argonne National Laboratory, Argonne, Illinois 60439, USA  
<sup>2</sup>Brookhaven National Laboratory, Upton, New York 11973, USA  
<sup>3</sup>University of California, Berkeley, California 94720, USA  
<sup>4</sup>University of California, Davis, California 95616, USA  
<sup>5</sup>University of California, Los Angeles, California 90095, USA  
<sup>6</sup>Universidade Estadual de Campinas, Sao Paulo, Brazil  
<sup>7</sup>Central China Normal University (HZNU), Wuhan 430079, China  
<sup>8</sup>University of Illinois at Chicago, Chicago, Illinois 60607, USA  
<sup>9</sup>Krakow Universities and Institute  
<sup>10</sup>Creighton University, Omaha, Nebraska 68178, USA  
<sup>11</sup>Czech Technical University in Prague, FNSPE, Prague, 115 19, Czech Republic  
<sup>12</sup>Nuclear Physics Institute AS CR, 250 68 Řež/Prague, Czech Republic  
<sup>13</sup>University of Frankfurt, Frankfurt, Germany  
<sup>14</sup>Institute of Physics, Bhubaneswar 751005, India  
<sup>15</sup>Indian Institute of Technology, Mumbai, India  
<sup>16</sup>Indiana University, Bloomington, Indiana 47408, USA  
<sup>17</sup>Alikhanov Institute for Theoretical and Experimental Physics, Moscow, Russia  
<sup>18</sup>University of Jammu, Jammu 180001, India  
<sup>19</sup>Joint Institute for Nuclear Research, Dubna, 141 980, Russia  
<sup>20</sup>Kent State University, Kent, Ohio 44242, USA  
<sup>21</sup>University of Kentucky, Lexington, Kentucky, 40506-0055, USA  
<sup>22</sup>Institute of Modern Physics, Lanzhou, China  
<sup>23</sup>Lawrence Berkeley National Laboratory, Berkeley, California 94720, USA  
<sup>24</sup>Massachusetts Institute of Technology, Cambridge, MA 02139-4307, USA  
<sup>25</sup>Max-Planck-Institut für Physik, Munich, Germany  
<sup>26</sup>Michigan State University, East Lansing, Michigan 48824, USA  
<sup>27</sup>Moscow Engineering Physics Institute, Moscow Russia  
<sup>28</sup>Ohio State University, Columbus, Ohio 43210, USA  
<sup>29</sup>Old Dominion University, Norfolk, VA, 23529, USA  
<sup>30</sup>Panjab University, Chandigarh 160014, India  
<sup>31</sup>Pennsylvania State University, University Park, Pennsylvania 16802, USA  
<sup>32</sup>Institute of High Energy Physics, Protvino, Russia  
<sup>33</sup>Purdue University, West Lafayette, Indiana 47907, USA  
<sup>34</sup>Pusan National University, Pusan, Republic of Korea  
<sup>35</sup>University of Rajasthan, Jaipur 302004, India  
<sup>36</sup>Rice University, Houston, Texas 77251, USA  
<sup>37</sup>Universidade de Sao Paulo, Sao Paulo, Brazil  
<sup>38</sup>University of Science & Technology of China, Hefei 230026, China  
<sup>39</sup>Shandong University, Jinan, Shandong 250100, China  
<sup>40</sup>Shanghai Institute of Applied Physics, Shanghai 201800, China  
<sup>41</sup>SUBATECH, Nantes, France  
<sup>42</sup>Texas A&M University, College Station, Texas 77843, USA  
<sup>43</sup>University of Texas, Austin, Texas 78712, USA  
<sup>44</sup>University of Houston, Houston, TX, 77204, USA  
<sup>45</sup>Tsinghua University, Beijing 100084, China  
<sup>46</sup>United States Naval Academy, Annapolis, MD 21402, USA  
<sup>47</sup>Valparaiso University, Valparaiso, Indiana 46383, USA  
<sup>48</sup>Variable Energy Cyclotron Centre, Kolkata 700064, India  
<sup>49</sup>Warsaw University of Technology, Warsaw, Poland  
<sup>50</sup>University of Washington, Seattle, Washington 98195, USA  
<sup>51</sup>Wayne State University, Detroit, Michigan 48201, USA  
<sup>52</sup>Yale University, New Haven, Connecticut 06520, USA  
<sup>53</sup>University of Zagreb, Zagreb, HR-10002, Croatia

STAR's measurements of directed flow ( $v_1$ ) around midrapidity for  $\pi^\pm$ ,  $K^\pm$ ,  $K_S^0$ ,  $p$  and  $\bar{p}$  in Au + Au collisions at  $\sqrt{s_{NN}} = 200$  GeV are presented. A negative  $v_1(y)$  slope is observed for most of produced particles ( $\pi^\pm$ ,  $K^\pm$ ,  $K_S^0$  and  $\bar{p}$ ). The proton  $v_1(y)$  slope is found to be much closer to zero compared to antiprotons. A sizable difference is seen between  $v_1$  of protons and antiprotons in 5-30% central collisions. The  $v_1$  excitation function is presented. Comparisons to model calculations

(RQMD, UrQMD, AMPT, QGSM with parton recombination, and a hydrodynamics model with a tilted source) are made. Anti-flow alone cannot explain the centrality dependence of the difference between the  $v_1(y)$  slopes of protons and antiprotons.

PACS numbers: 25.75.Ld

The BNL Relativistic Heavy Ion Collider (RHIC) was built to study a new form of matter known as the Quark Gluon Plasma (QGP) [1], which existed in the universe shortly after the Big-Bang. At RHIC, two nuclei are collided at near light-speed, and the collision produces thousands of particles due to the significant energy deposited. The collective motion of the produced particles can be characterized [2] by Fourier coefficients,

$$v_n = \langle \cos n(\phi - \psi) \rangle \quad (1)$$

where  $n$  denotes the harmonic,  $\phi$  and  $\psi$  denote the azimuthal angle of an outgoing particle and reaction plane, respectively. The reaction plane is defined by the collision axis and the line connecting the centers of two nuclei. Thus far, five of these coefficients have been measured and found to be non-zero at RHIC. They are directed flow  $v_1$ , elliptic flow  $v_2$ , triangular flow  $v_3$ , the 4<sup>th</sup> order harmonic flow  $v_4$  and the 6<sup>th</sup> order harmonic flow  $v_6$ . This paper will focus on the directed flow, the first Fourier coefficient.

Directed flow describes the sideward motion of produced particles in ultra-relativistic nuclear collisions. It is believed to be generated during the nuclear passage time before the thermalization happens, thus it carries early information from the collision [3–6]. The shape of directed flow at midrapidity may be modified by the collective expansion and reveal a signature of a possible phase transition from normal nuclear matter to a QGP [7–9]. It is argued that directed flow, as an odd function of rapidity ( $y$ ), may exhibit a small slope (flatness) at midrapidity due to a strong expansion of the fireball being tilted away from the collision axis. Such tilted expansion gives rise to anti-flow [7] or a 3<sup>rd</sup> flow [8] component (not the third flow harmonic). The anti-flow (or the 3<sup>rd</sup> flow component) is perpendicular to the source surface, and is in the opposite direction to the bouncing-off motion of nucleons. If the tilted expansion is strong enough, it can even overcome the bouncing-off motion and results in a negative  $v_1(y)$  slope at midrapidity, potentially producing a wiggle-like structure in  $v_1(y)$ . Note that although calculations [7, 8] for both anti-flow and 3<sup>rd</sup> flow component are made for collisions at SPS energies where the first order phase transition to a QGP is believed to be the most relevant [9], the direct cause of the negative slope is the strong, tilted expansion, which is also important at RHIC's top energies. Indeed hydrodynamic calculations [10] for Au + Au collisions at  $\sqrt{s_{NN}} = 200$  GeV with a tilted

source as the initial condition can give a similar negative  $v_1(y)$  slope as that found in data. A wiggle structure is also seen in the Relativistic Quantum Molecular Dynamics (RQMD) model [11], and it is attributed to baryon stopping together with a positive space-momentum correlation. In this picture, no phase transition is needed, and pions and nucleons flow in opposite directions. To distinguish between baryon stopping and anti-flow, it is desirable to measure the  $v_1(y)$  for identified particles and compare the sign of their slopes at midrapidity. In particular, the observation of a centrality dependence of proton  $v_1(y)$  may reveal the character of a possible first order phase transition [9]. It is expected that in very peripheral collisions, protons flow in the same direction as spectators. In mid-central collisions, if there is a phase transition, the proton  $v_1(y)$  slope at midrapidity may change sign and become negative. Eventually the slope diminishes in central collisions due to the symmetry of the collisions.

At low energies, the E895 collaboration has shown that  $K_S^0$  has a negative  $v_1(y)$  slope around midrapidity [12], while  $\Lambda$  and protons have positive slopes [13]. This is explained by a repulsive kaon-nucleon potential and an attractive  $\Lambda$ -nucleon potential. The NA49 collaboration [14] has measured  $v_1$  for pions and protons, and a negative  $v_1(y)$  slope is observed by the standard event plane method. The three-particle correlation method  $v_1\{3\}$  [15], which is believed to be less sensitive to non-flow effects, gives a negative slope too, but with a larger statistical error. The non-flow effects are correlations among particles that are not related to the reaction plane, including the quantum Hanbury Brown-Twiss correlation [16], resonance decays [17] and so on. At top RHIC energies,  $v_1$  has been studied mostly for charged particles by both the STAR and the PHOBOS collaborations [18–21]. It is found that  $v_1$  in the forward region follows the limiting fragmentation hypothesis [22], and  $v_1$  as a function of pseudorapidity ( $\eta$ ) depends only on the incident energy, but not on the size of the colliding system at a given centrality. Such system size independence of  $v_1$  can be explained by the hydrodynamic calculation with a tilted initial condition [10]. The systematic study of  $v_1$  for identified particles at RHIC did not begin until recently because it is more challenging for two reasons: 1)  $v_1$  for some identified particles (for example, protons) is much smaller than that of all charged particles, 2) more statistics are needed to determine  $v_1$  for identified particles other than pions.

54 million events from Au + Au collisions at  $\sqrt{s_{NN}} = 200$  GeV have been used in this study, all taken by a minimum-bias trigger with the STAR detector during RHIC's seventh run in year 2007. The main trigger detector used is the Vertex Position Detector (VPD) [23]. The centrality definition of an event was based on the number of charged tracks in the Time Projection Chamber (TPC) [24] with track quality cuts:  $|\eta| < 0.5$ , a Distance of Closest Approach (DCA) to the vertex less than 3 cm, and 10 or more fit points. In the analysis, events are required to have the vertex  $z$  within 30 cm from the center of the TPC, and additional weight is assigned to each event in the analysis accounting for the non-uniform VPD trigger efficiency in the vertex  $z$  direction for different centrality classes. The event plane angle is determined from the sideward deflection of spectator neutrons measured by STAR's Shower Maximum Detector inside the Zero Degree Calorimeters (ZDC-SMDs). Such sideward deflection of spectator neutrons is expected to happen in the reaction plane rather than participant plane, since the ZDC-SMDs are located close to beam rapidity. Being 6 units in  $\eta$  away from midrapidity, ZDC-SMDs also allow a measurement of  $v_1$  with minimal contribution from non-flow correlations. The description of measuring  $v_1$  using the ZDC-SMDs event plane can be found in [20]. Particle Identification (PID) of charged particles is achieved by measuring ionization energy loss ( $dE/dx$ ) inside STAR's TPC, together with the measurement of the momentum ( $p$ ) via TPC tracking. Track quality cuts are the same as used in [25]. In addition, the transverse momentum  $p_T$  for protons is required to be larger than 400 MeV/ $c$ , and DCA is required to be less than 1 cm in order to avoid including background protons which are from knock-out/nuclear interactions of pions with inner detector material. The same cuts are applied to antiprotons as well to ensure a fair comparison with protons. The high-end of the  $p_T$  cut is 1 GeV/ $c$  where protons and pions have the same energy loss in the TPC and thus become indistinguishable. For pions and kaons,  $p_T$  range is 0.15 - 0.75 GeV/ $c$  and 0.2 - 0.6 GeV/ $c$ , respectively.  $K_S^0(\rightarrow \pi^+\pi^-)$  are topologically reconstructed by their charged daughter tracks inside the TPC [26].

Results presented in the following figures contain only statistical errors. Results for pions, protons and antiprotons are not corrected for the feeddown from weak decay particles. The major systematic error in determining the slope of  $v_1(y)$  for identified particles is from the particle misidentification, which was evaluated by varying the  $dE/dx$  cut. Another systematic error comes from the non-uniform  $p_T$  acceptance, as  $v_1(y)$  is obtained by integrating  $v_1$  over the  $p_T$  acceptance which itself depends on the rapidity. This effect is non-negligible for protons and antiprotons




FIG. 1:  $v_1$  for  $\pi^\pm$ ,  $K^\pm$ ,  $K_S^0$  (left panel),  $p$  and  $\bar{p}$  (right panel) as a function of rapidity for 10-70% Au + Au collisions at  $\sqrt{s_{NN}} = 200$  GeV. The lines present the linear fit to the  $\pi^\pm$ ,  $K^\pm$ ,  $K_S^0$ ,  $p$  and  $\bar{p}$ 's  $v_1(y)$  respectively. Data points around  $y = 0.29$  are slightly shifted horizontally to avoid overlapping.

at large rapidity. It is estimated by taking the difference between slopes fitted with points integrated with  $p_T$  acceptance at midrapidity and at large rapidity. In addition, some of the observed protons have originated from interactions between the produced particles and the detector material, and such effect has also been taken into consideration. The total systematic uncertainty is obtained by adding uncertainties mentioned above in quadrature. There are also common systematic errors that should be applied to all particles: the uncertainty due to the first order event plane determination, which was estimated to be  $\sim 10\%$  (relative error) [20], and the uncertainty due to centrality selection, which was estimated to be  $\sim 4\%$  (relative error) by comparing our charged  $v_1(\eta)$  slope to that from the RHIC run in 2004. Other systematic errors have been evaluated to be negligible.

In Fig. 1,  $v_1(y)$  of  $\pi^\pm$ ,  $K^\pm$ ,  $K_S^0$ ,  $p$ , and  $\bar{p}$  are presented for centrality 10-70%. Following convention, the sign of spectator  $v_1$  in the forward region is chosen to be positive, to which the measured sign of  $v_1$  for particles of interest is only relative. Fitting with a linear function, the slopes are  $-0.15 \pm 0.05(\text{stat}) \pm 0.08(\text{sys})(\%)$  for the protons,  $-0.46 \pm 0.06(\text{stat}) \pm 0.04(\text{sys})(\%)$  for the antiprotons,  $-0.27 \pm 0.01(\text{stat}) \pm 0.01(\text{sys})(\%)$  for the pions,  $-0.02 \pm 0.11(\text{stat}) \pm 0.04(\text{sys})(\%)$  for the kaons and  $-0.17 \pm 0.02(\text{stat}) \pm 0.04(\text{sys})(\%)$  for the  $K_S^0$ . The relative 10% common systematic error for all particles is not listed here. The  $v_1(y)$  slope for the produced particle types ( $\pi^\pm$ ,  $K^\pm$ ,  $K_S^0$  and  $\bar{p}$ ) are mostly found to be negative at mid-rapidity, which is consistent with the anti-flow picture. In particu-




FIG. 2: Model calculations of pion (left panel) and proton (right panel)  $v_1(y)$  for 10-70% Au + Au collisions at  $\sqrt{s_{NN}} = 200$  GeV. QGSM\* model presents the basic Quark-Gluon String model with parton recombination [29]. Hydro\* model presents the hydrodynamic expansion from a tilted source [10].

lar, kaons are less sensitive to shadowing effects due to the small kaon-nucleon cross section, yet it shows a negative slope. This is again consistent with the anti-flow picture. Interestingly,  $v_1(y)$  for protons exhibits a clearly flatter shape than that for antiprotons. While mass may contribute to the difference in slope between pions and protons/antiprotons, it cannot explain the difference in slope observed for antiprotons and protons. Indeed, the observed  $v_1$  for protons is a convolution of directed flow of produced protons with that of transported protons (from the original projectile and target nuclei), so the flatness of inclusive proton  $v_1(y)$  around midrapidity could be explained by the negative flow of produced protons being compensated by the positive flow of protons transported from spectator rapidity, as a feature expected in the anti-flow picture.

In Fig. 2, pion and proton  $v_1(y)$  are plotted together with five model calculations, namely, RQMD [11], UrQMD [27], AMPT [28], QGSM with parton recombination [29], and slopes from an ideal hydrodynamic calculation with a tilted source [10]. The model calculations are performed in the same  $p_T$  acceptance and centrality as the data. The RQMD and AMPT model calculations predict the wrong sign and wrong magnitude of pion  $v_1(y)$ , respectively, while the RQMD and the UrQMD predict the wrong magnitude of proton  $v_1(y)$ . None of these models can describe  $v_1(y)$  for pions and protons simultaneously.

In Fig. 3, the slope of  $v_1(y)$  at midrapidity is presented as a function of centrality for protons, antiprotons, and charged pions. Two observations are noteworthy: i) the hydrodynamic model with tilted




FIG. 3: Charged pions (solid stars), protons (solid circles) and antiprotons (solid squares)  $v_1(y)$  slope ( $dv_1/dy$ ) at midrapidity as a function of centrality for Au + Au collisions at  $\sqrt{s_{NN}} = 200$  GeV.




FIG. 4: Proton  $v_1(y')$  slope ( $dv_1/dy'$ ) at midrapidity as a function of center of mass collision energy, where  $y' = y/y_{beam}$ .

source (which is a characteristic of anti-flow) does not predict the difference in  $v_1(y)$  between particle species [30]. ii) If the difference between  $v_1$  of protons and antiprotons is caused by anti-flow alone, then such difference is expected to be accompanied by strongly negative  $v_1$  slopes. In data, the large difference between proton and antiproton  $v_1$  slopes is seen in the 5-30% centrality range, while strongly negative  $v_1$  slopes are found for protons, antiprotons and charged pions in a different centrality range (30-80%). Both observations suggest that additional mechanisms, besides anti-flow, are needed to explain the centrality dependence of the difference between the  $v_1(y)$  slopes of protons and antiprotons.

The excitation function of proton  $v_1(y')$  slope  $F$  ( $= dv_1/dy'$  at midrapidity) is presented in Fig 4. Values for  $F$  are extracted via a polynomial fit of the form  $Fy' + Cy'^3$ , where  $y' = y/y_{beam}$  for which spec-

tators are normalized at  $\pm 1$ . The proton  $v_1(y')$  slope decreases rapidly with increasing energy, reaching zero around  $\sqrt{s_{\text{NN}}} = 9$  GeV. Its sign changes to negative as shown by the data point at  $\sqrt{s_{\text{NN}}} = 17$  GeV, measured by the NA49 experiment [14]. A similar trend has been observed at low energies with a slightly different quantity  $d\langle p_x \rangle / dy'$  [31, 32]. The energy dependence of  $v_1(y')$  slope for protons is driven by two factors, i) the increase in the number of produced protons over transported protons with increasing energy, and ii) the  $v_1$  of both produced and transported protons at different energies. The negative  $v_1(y')$  slope for protons around midrapidity at SPS energies cannot be explained by transport model calculations like UrQMD [33] and AMPT [28], but is predicted by hydro calculations [7, 8]. The present data indicate that the proton  $v_1$  slope remains close to zero at  $\sqrt{s_{\text{NN}}} = 200$  GeV as observed at  $\sqrt{s_{\text{NN}}} = 9$  GeV and  $\sqrt{s_{\text{NN}}} = 17$  GeV heavy ion collisions. Our measurement offers a unique check of the validity of a tilted expansion at RHIC top energy.

In summary, STAR's measurements of directed flow of pions, kaons, protons, and antiprotons for Au + Au collisions at  $\sqrt{s_{\text{NN}}} = 200$  GeV are presented. In the range of 10-70% central collisions,  $v_1(y)$  slopes of pions, kaons ( $K_S^0$ ), and antiprotons are found to be mostly negative at mid-rapidity. However, protons exhibit a clearly flatter shape than antiprotons. A sizable difference is seen between  $v_1$  of protons and antiprotons in 5-30% central collisions. Comparison to models (RQMD, UrQMD, AMPT, QGSM with parton recombination, and hydrodynamics with a

tilted source) is made. None of the models explored can describe  $v_1(y)$  for pions and protons simultaneously. Additional mechanisms besides the anti-flow are needed to explain the centrality dependence of the difference between the  $v_1(y)$  slopes of protons and antiprotons. Our measurement indicates that proton's  $v_1(y)$  slope remains close to zero for Au + Au collisions at  $\sqrt{s_{\text{NN}}} = 200$  GeV. These new measurements on the particle species and centrality dependence of  $v_1(y)$  provides a check for the validity of a tilted expansion at RHIC top energy.

### Acknowledgments

We thank the RHIC Operations Group and RCF at BNL, the NERSC Center at LBNL and the Open Science Grid consortium for providing resources and support. This work was supported in part by the Offices of NP and HEP within the U.S. DOE Office of Science, the U.S. NSF, the Sloan Foundation, the DFG cluster of excellence 'Origin and Structure of the Universe' of Germany, CNRS/IN2P3, FAPESP CNPq of Brazil, Ministry of Ed. and Sci. of the Russian Federation, NNSFC, CAS, MoST, and MoE of China, GA and MSMT of the Czech Republic, FOM and NWO of the Netherlands, DAE, DST, and CSIR of India, Polish Ministry of Sci. and Higher Ed., Korea Research Foundation, Ministry of Sci., Ed. and Sports of the Rep. Of Croatia, and RosAtom of Russia.

- 
- [1] BRAHMS, PHENIX, PHOBOS, and STAR Collaboration, *Nucl. Phys. A* **757** Issues 1-2 (2005).
  - [2] A. M. Poskanzer and S. A. Voloshin, *Phys. Rev. C* **58**, 1671 (1998).
  - [3] E. Schnedermann and U. Heinz, *Phys. Rev. Lett.* **69**, 2908 (1992).
  - [4] D. E. Kahana *et al.*, *Phys. Rev. Lett.* **74**, 4404 (1995).
  - [5] J. Barrette *et al.* (E877 Collaboration), *Phys. Rev. Lett.* **73**, 2532 (1994).
  - [6] I. G. Bearden *et al.* (NA44 Collaboration), *Phys. Rev. Lett.* **78**, 2080 (1997).
  - [7] J. Brachmann *et al.*, *Phys. Rev. C* **61**, 024909 (2000).
  - [8] L. P. Csernai and D. Röhrich, *Phys. Lett. B* **458**, 454 (1999).
  - [9] H. Stöcker, *Nucl. Phys. A* **750**, 121 (2005).
  - [10] P. Bożek and I. Wyskiel, *Phys. Rev. C* **81**, 054902 (2010).
  - [11] R. J. Snellings *et al.*, *Phys. Rev. Lett.* **84**, 2803 (2000).
  - [12] P. Chung *et al.* (E895 Collaboration), *Phys. Rev. Lett.* **85**, 940 (2000).
  - [13] P. Chung *et al.* (E895 Collaboration), *Phys. Rev. Lett.* **86**, 2533 (2001).
  - [14] C. Alt *et al.* (NA49 Collaboration), *Phys. Rev. C* **68**, 034903 (2003).
  - [15] N. Borghini, P. M. Dinh and J.-Y. Ollitrault, *Phys. Rev. C* **66**, 014905 (2002).
  - [16] P. M. Dinh, N. Borghini and J.-Y. Ollitrault, *Phys. Lett. B* **477**, 51 (2000).
  - [17] N. Borghini, P. M. Dinh and J.-Y. Ollitrault, *Phys. Rev. C* **62**, 034902 (2000).
  - [18] J. Adams *et al.* (STAR Collaboration), *Phys. Rev. Lett.* **92**, 062301 (2004).
  - [19] B. B. Back *et al.* (PHOBOS Collaboration), *Phys. Rev. Lett.* **97**, 012301 (2006).
  - [20] J. Adams *et al.* (STAR Collaboration), *Phys. Rev. C* **73**, 034903 (2006).
  - [21] B. I. Abelev *et al.* (STAR Collaboration), *Phys. Rev. Lett.* **101**, 252301 (2008).
  - [22] J. Benecke *et al.*, *Phys. Rev.* **188**, 2159 (1969).
  - [23] W. J. Llope *et al.*, *Nucl. Instrum. Methods A* **522**, 252 (2004).

- [24] M. Anderson *et al.*, *Nucl. Instrum. Method. A* **499**, 659 (2003).
- [25] J. Adams *et al.* (STAR Collaboration), *Phys. Rev. C* **72**, 014904 (2005).
- [26] C. Adler *et al.* (STAR Collaboration), *Phys. Rev. Lett.* **89**, 132301 (2002); J. Adams *et al.* (STAR Collaboration), *Phys. Rev. Lett.* **92**, 052302 (2004).
- [27] M. Bleicher and H. Stöcker, *Phys. Lett. B* **526**, 309 (2002).
- [28] J. Y. Chen *et al.*, *Phys. Rev. C* **81**, 014904 (2010).
- [29] J. Bleibel *et al.*, *Phys. Rev. C* **76**, 024912 (2007).
- [30] P. Božek, private communication, 2010.
- [31] H. Liu *et al.* (E895 Collaboration), *Phys. Rev. Lett.* **84**, 5488 (2000).
- [32] J. Barrette *et al.* (E877 Collaboration), *Phys. Rev. C* **56**, 3254 (1997); *Phys. Rev. C* **55**, 1420 (1997).
- [33] H. Petersen *et al.*, *Phys. Rev. C* **74**, 064908 (2006).