



Letter

Search for the Chiral Magnetic Effect with charge-dependent azimuthal correlations in Xe–Xe collisions at $\sqrt{s_{\text{NN}}} = 5.44$ TeV

ALICE Collaboration ^{*}

ARTICLE INFO

Editor: M. Doser

 Dataset link: <https://www.hepdata.net/record/ins2172062>

ABSTRACT

Charge-dependent two- and three-particle correlations measured in Xe–Xe collisions at $\sqrt{s_{\text{NN}}} = 5.44$ TeV are presented. Results are obtained for charged particles in the pseudorapidity range $|\eta| < 0.8$ and transverse momentum interval $0.2 \leq p_{\text{T}} < 5.0$ GeV/ c for different collision centralities. The three-particle correlator $\gamma_{\alpha\beta} \equiv \langle \cos(\varphi_{\alpha} + \varphi_{\beta} - 2\Psi_2) \rangle$, calculated for different combinations of charge sign α and β , is expected to be sensitive to the presence of the Chiral Magnetic Effect (CME). Its magnitude is similar to the one observed in Pb–Pb collisions in contrast to a smaller CME signal in Xe–Xe collisions than in Pb–Pb collisions predicted by Monte Carlo (MC) calculations including a magnetic field induced by the spectator protons. These observations point to a large non-CME contribution to the correlator. Furthermore, the charge dependence of $\gamma_{\alpha\beta}$ can be described by a blast wave model calculation that incorporates background effects and by the Anomalous Viscous Fluid Dynamics model with values of the CME signal consistent with zero. The Xe–Xe and Pb–Pb results are combined with the expected CME signal dependence on the system size from the MC calculations including a magnetic field to obtain the fraction of CME contribution in $\gamma_{\alpha\beta}$, f_{CME} . The CME fraction is compatible with zero for the 30% most central events in both systems and then becomes positive. This yields an upper limit of 2% (3%) and 25% (32%) at 95% (99.7%) confidence level for the CME signal contribution to $\gamma_{\alpha\beta}$ in the 0–70% Xe–Xe and Pb–Pb collisions, respectively.

1. Introduction

The theory of the strong interaction applied to many-body systems predicts that, at sufficiently high densities and temperatures, the protons and neutrons that compose ordinary matter melt into a plasma where quarks and gluons are no longer confined into hadrons. This hot and dense state of matter is called the quark–gluon plasma (QGP) [1]. The transition from normal hadronic matter to a QGP is supported by Quantum Chromodynamics (QCD) calculations on the lattice [2–5], where it is found to occur at a temperature of about 155 MeV, and at an energy density ϵ of about 0.5 GeV/ fm^3 [6–8]. Collisions between heavy ions accelerated to ultrarelativistic energies can produce the necessary conditions for such a transition to take place [9–11].

Heavy-ion collisions may also allow us to access novel QCD phenomena associated with parity violation in strong interactions [12–20]. Theoretical expectations indicate that the interaction of quarks with gluonic fields describing transitions between topologically different QCD vacuum states changes the quark chirality and leads to a local chiral imbalance. In the presence of the strong magnetic field produced by the colliding ions [21–23], this leads to a charge separation (electric current) relative to the reaction plane, the plane defined by the impact

parameter and the beam axis. This phenomenon is known as the Chiral Magnetic Effect (CME) [20].

The effects from local parity violation are quantified via the coefficient $a_{1,\alpha}$ in a Fourier decomposition of the particle azimuthal distribution [24,25]

$$\frac{dN}{d\Delta\varphi_{\alpha}} \sim 1 + 2v_{1,\alpha} \cos(\Delta\varphi_{\alpha}) + 2a_{1,\alpha} \sin(\Delta\varphi_{\alpha}) + 2v_{2,\alpha} \cos(2\Delta\varphi_{\alpha}) + \dots, \quad (1)$$

where $\Delta\varphi_{\alpha} = \varphi_{\alpha} - \Psi_{\text{RP}}$, φ_{α} is the azimuthal angle of the particle of charge α (+, –), and Ψ_{RP} is the reaction plane angle. The coefficients $v_{n,\alpha}$ characterise the anisotropic flow, i.e., the azimuthal anisotropies in particle production relative to Ψ_{RP} due to initial spatial asymmetries of the collision. The degree of overlap between the two colliding nuclei is estimated by the centrality, with low percentage values corresponding to head-on collisions. The first- and second-order flow coefficients ($v_{1,\alpha}$ and $v_{2,\alpha}$) are called directed and elliptic flow, respectively. Since $a_{1,\alpha}$ changes sign from event to event and the average $\langle a_{1,\alpha} \rangle$ over many events is equal to zero, one can only measure $\langle a_{1,\alpha}^2 \rangle$ or $\langle a_{1,+} a_{1,-} \rangle$ that can be accessed through azimuthal correlation techniques. Thus the CME is expected to have an experimentally accessible signal imprinted in the azimuthal correlations between two particles relative to

^{*} E-mail address: alice-publications@cern.ch.

the reaction plane [25] of the form $\gamma_{\alpha\beta} \equiv \langle \cos(\varphi_\alpha + \varphi_\beta - 2\Psi_{\text{RP}}) \rangle$. The charge-dependent difference of $\gamma_{\alpha\beta}$ is commonly used to search for the CME. In practice, the reaction plane angle is estimated by constructing the second harmonic symmetry plane angle Ψ_2 using azimuthal particle distributions [26], which is why $\gamma_{\alpha\beta}$ is often referred to as a three-particle correlator. The $\gamma_{\alpha\beta}$ correlator measures the difference between the correlations projected onto the reaction plane and perpendicular to it. The contributions from correlations in- and out-of-plane can also be evaluated by measuring the charge-dependent two-particle correlator $\delta_{\alpha\beta} \equiv \langle \cos(\varphi_\alpha - \varphi_\beta) \rangle$.

The first experimental results in Au–Au collisions at a centre-of-mass energy per nucleon–nucleon collision $\sqrt{s_{\text{NN}}} = 200$ GeV at the Relativistic Heavy-Ion Collider (RHIC) [27,28] were compatible with initial expectations for the existence of the CME. The subsequent first measurements at the Large Hadron Collider (LHC) in Pb–Pb collisions at $\sqrt{s_{\text{NN}}} = 2.76$ TeV [29] showed a surprising agreement with the results at lower energies, despite the differences in magnitude of the magnetic field [21–23]. Considering that the charged-particle density, $dN_{\text{ch}}/d\eta$, at the LHC is about three times larger than at RHIC [30,31], any signal due to CME will be considerably diluted since it is expected to follow a $1/(dN_{\text{ch}}/d\eta)$ scaling [19]. This effect will be referred to as dilution in the following. The similarity of the two measurements was indicative of the existence of background effects, coming mostly from “flowing clusters” – charge-dependent correlations modified by elliptic flow [25,32–34]. It was shown in Refs. [35,36] that the local charge conservation coupled to the anisotropic expansion of the medium could explain most if not all the measurements.

To study background effects, the CMS Collaboration performed measurements of charge-dependent correlations in p–Pb collisions at $\sqrt{s_{\text{NN}}} = 5.02$ TeV [37] and the STAR Collaboration in p–Au and d–Au collisions at $\sqrt{s_{\text{NN}}} = 0.2$ TeV [38]. The results suggest that these correlations are similar to those measured in peripheral Pb–Pb and Au–Au collisions. These results might further indicate the dominance of background effects in peripheral collisions where there is no strong correlation between the magnetic field direction and the orientation of the medium via Ψ_{RP} .

These measurements highlighted the need to identify ways of isolating the CME signal from the background. A first attempt was presented by the ALICE Collaboration in Ref. [39] using the Event Shape Engineering method [40]. This method utilises the fluctuations in the shape of the initial state of the system and allows one to select events with the same centrality but different initial geometry, thus varying the background contributions. The study sets an upper limit of 26–33% at 95% confidence level for the CME signal contribution to the charge dependence of $\gamma_{\alpha\beta}$ in the 10–50% centrality interval. A similar study was performed by the CMS Collaboration [41] and the results agree with the measurements in Ref. [39]. A recent study by the ALICE Collaboration [42] found that charge-dependent correlations relative to the higher harmonic symmetry planes can be used as a proxy for the background, assuming that the correlations relative to Ψ_2 and Ψ_3 can be factorised. An upper limit of 15–18% at 95% confidence level for the CME signal has been reported for the 0–40% centrality interval, consistent with previous measurements.

Another approach to address the large backgrounds experimentally is to compare measurements performed in collision systems where the CME contribution is expected to vary significantly, while the background is similar. The STAR Collaboration has recently reported the results of the CME search in an analysis of the three-particle correlator $\gamma_{\alpha\beta}$ measured in collisions of isobar $^{96}_{44}\text{Ru}$ – $^{96}_{44}\text{Ru}$ and $^{96}_{40}\text{Zr}$ – $^{96}_{40}\text{Zr}$ nuclei at $\sqrt{s_{\text{NN}}} = 200$ GeV [43]. No anticipated CME signature (i.e., a larger magnitude of $\gamma_{\alpha\beta}$ in Ru–Ru than in Zr–Zr collisions due to a larger magnetic field in the former) was observed in that analysis. However, a quantitative analysis taking into account the small geometrical differences between the isobar nuclei is needed for the interpretation of the measurements. One can also try to separate the CME signal and background by comparing the results from Pb–Pb and Xe–Xe collisions at the LHC

since the differences in v_2 are typically within 10% in the 5–70% centrality interval [44] but the magnetic field is expected to be significantly larger in Pb–Pb collisions [45], leading to an increase in the CME contribution.

In this article, measurements of charge-dependent azimuthal correlations from Xe–Xe collisions at $\sqrt{s_{\text{NN}}} = 5.44$ TeV are presented. The results are compared with earlier measurements in Pb–Pb collisions at $\sqrt{s_{\text{NN}}} = 5.02$ TeV [42] and calculations from a blast wave parameterisation that incorporate background effects and from the Anomalous Viscous Fluid Dynamics (AVFD) model [46–48]. Furthermore, Monte Carlo (MC) simulations of the magnetic field induced by spectator protons with different initial conditions are used to evaluate the expected change in the CME signal between the Xe–Xe and Pb–Pb collisions. This change is then employed to estimate the fraction of the CME signal in both collision systems.

2. Analysis details

The data set used for these measurements was recorded with the ALICE detector during the 2017 Xe–Xe run at $\sqrt{s_{\text{NN}}} = 5.44$ TeV. A detailed overview of the ALICE detector and its performance are available in Refs. [49,50]. The Inner Tracking System (ITS) [51], the Time Projection Chamber (TPC) [52], the V0 [53], and the Zero Degree Calorimeter (ZDC) [54], the main subsystems used in this analysis, are briefly described in the following. The ITS and TPC cover the full azimuth within the pseudorapidity range $|\eta| < 0.9$. The ITS consists of six layers of silicon detectors and is employed for tracking, vertex reconstruction, and event selection. The TPC is used to reconstruct charged-particle tracks and to identify particles via specific energy loss, dE/dx . The V0 detector, two arrays of 32 scintillator tiles covering $-3.7 < \eta < -1.7$ (V0C) and $2.8 < \eta < 5.1$ (V0A), is used for triggering, event selection, and the determination of centrality [55] and symmetry plane Ψ_2 . Both V0 detectors are segmented in four rings in the radial direction with each ring divided into eight sectors in the azimuthal direction. Two tungsten-quartz neutron ZDCs, installed 112.5 meters from the interaction point on each side, are also used for event selection.

The trigger conditions and the event selection criteria can be found in Ref. [56]. Beam-induced background and pileup events are removed using an offline event selection, employing information from the V0, ZDC, and tracking detectors. The primary vertex position is determined from tracks reconstructed in the ITS and TPC as described in Ref. [50]. Approximately 10^6 Xe–Xe events in the 0–70% centrality interval, with a primary vertex position within ± 10 cm from the nominal interaction point along the beam direction, are used in the analysis. The centrality of the collision is estimated from the energy deposition measured in the V0 detector [55].

The charged-particle tracks reconstructed using the ITS and TPC within $|\eta| < 0.8$ and $0.2 \leq p_{\text{T}} < 5.0$ GeV/c are used to measure the charge-dependent correlations. Each track is required to have a minimum number of 70 space points (out of a maximum of 159) with a χ^2 per TPC space point lower than 4, to cross at least 70 TPC read-out rows, and to have the ratio between the number of crossed rows and the number of findable space points in the TPC larger than 0.8. The selected tracks are also required to have at least 2 ITS hits and a χ^2 per ITS hit smaller than 36. In addition, tracks are selected with a distance of closest approach (DCA) to the reconstructed vertex position smaller than 3.2 cm and 2.4 cm in the longitudinal direction (z) and transverse plane (xy), respectively. These selection criteria reduce the contamination from secondary charged particles (i.e., particles originating from weak decays, conversions, and secondary hadronic interactions in the detector material) and fake tracks (random associations of space points) and ensure a track momentum resolution better than 4% in the considered p_{T} interval [56]. The charged-particle track reconstruction efficiency is estimated from simulations with the HIJING event generator [57,58] combined with the GEANT3 transport model [59]. These simulations include a detailed description of the detector response. The

p_T averaged charge-dependent correlations are corrected for track reconstruction efficiency.

The charge-dependent correlations are measured using two- and three-particle correlators expressed as

$$\delta_{\alpha\beta} \equiv \langle \cos(\varphi_\alpha - \varphi_\beta) \rangle = \langle \cos(\Delta\varphi_\alpha) \cos(\Delta\varphi_\beta) \rangle + \langle \sin(\Delta\varphi_\alpha) \sin(\Delta\varphi_\beta) \rangle \quad (2)$$

$$= \langle v_{1,\alpha} v_{1,\beta} \rangle + B_{\text{in}} + \langle a_{1,\alpha} a_{2,\beta} \rangle + B_{\text{out}},$$

$$\gamma_{\alpha\beta} \equiv \langle \cos(\varphi_\alpha + \varphi_\beta - 2\Psi_2) \rangle \quad (3)$$

$$= \langle \cos(\Delta\varphi_\alpha) \cos(\Delta\varphi_\beta) \rangle - \langle \sin(\Delta\varphi_\alpha) \sin(\Delta\varphi_\beta) \rangle$$

$$= \langle v_{1,\alpha} v_{1,\beta} \rangle + B_{\text{in}} - \langle a_{1,\alpha} a_{2,\beta} \rangle - B_{\text{out}},$$

where $\Delta\varphi_{\alpha(\beta)} = \varphi_{\alpha(\beta)} - \Psi_2$, and B_{in} and B_{out} denote background contributions projected onto Ψ_2 and perpendicular to it, respectively. The term $\langle v_{1,\alpha} v_{1,\beta} \rangle$ is expected to have negligible charge dependence at midrapidity [60]. In addition, $\langle v_1 \rangle$ at midrapidity is zero for a symmetric collision. While $\gamma_{\alpha\beta}$ suppresses background contributions at the level of v_2 (i.e., the relative difference between the particle production in-plane and out-of-plane), $\delta_{\alpha\beta}$ is dominated by short-range correlations unrelated to Ψ_2 (“non-flow”), such as inter-jet correlations and resonance decays.

The orientation of the symmetry plane Ψ_2 is estimated from the azimuthal distribution of the energy deposition measured by the VOA detector, with the x and y components given by

$$Q_{2,x} = \sum_j w_j \cos(2\varphi_j), \quad Q_{2,y} = \sum_j w_j \sin(2\varphi_j), \quad (4)$$

where the index j runs over the 32 sectors of the VOA detector, φ_j is the azimuthal angle of sector j defined by the geometric centre, and w_j is the amplitude of the measured signal in that sector. The symmetry plane resolution is calculated from correlations between the symmetry planes determined with the TPC, the VOA, and the V0C detectors [26]. The effect of the decorrelation of Ψ_2 between mid and forward pseudorapidity has been estimated to be less than 3% for v_2 [61]. Any non-uniform detector response is taken into account by adjusting the components of the \mathbf{Q}_2 vector using a recentering procedure (i.e., subtraction of the \mathbf{Q}_2 vector averaged over many events from the \mathbf{Q}_2 vector of each event) [62]. The non-flow contributions to the charge-dependent azimuthal correlations are greatly suppressed by the large pseudorapidity separation between the TPC and the VOA ($|\Delta\eta| > 2.0$).

The absolute systematic uncertainties were estimated from the variation of the results with different event and track-selection criteria. The event selection contributions were determined by varying the range of the reconstructed collision vertex position from the nominal interaction point along the beam direction, estimating centrality from the number of hits in the first or second layer of the ITS, and imposing stricter pileup rejection criteria than the default selection. Systematic uncertainties related to track selection criteria were evaluated by changing the ITS hit requirements, varying the minimum number of TPC space points, changing the minimum number of crossed TPC readout rows and the ratio between the number of crossed rows and the number of findable space points in the TPC, rejecting tracks close to the TPC sector boundaries to which the sensitive readout rows do not extend, and comparing any differences between results with only positive and only negative charges for pairs of particles with same charge. Finally, changes of the results due to uncertainties in the tracking efficiency arising from an imperfect description in the simulation of the relative abundances of different particle species and their different reconstruction efficiencies [63] were considered as part of the systematic uncertainties. The largest contribution to the systematic uncertainties for $\gamma_{\alpha\beta}$ and $\delta_{\alpha\beta}$ is given by the centrality estimation and track-selection criteria, respectively. The systematic uncertainties are evaluated for each centrality interval. The different sources are assumed uncorrelated and are added in quadrature as an estimate of the total systematic uncertainties if their deviations from the nominal values are significant according to the Bar-

Table 1

Summary of absolute systematic uncertainties on the charge-dependent correlations. The uncertainties depend on centrality, whose minimum and maximum values are listed here.

	Opposite charge	Same charge
$\delta_{\alpha\beta}$	$(6.8 - 33) \times 10^{-5}$	$(3.8 - 13) \times 10^{-5}$
$\gamma_{\alpha\beta}$	$(1.0 - 8.3) \times 10^{-5}$	$(1.4 - 5.9) \times 10^{-5}$

low criterion [64]. The resulting systematic uncertainties increase from central to peripheral collisions and are summarised in Table 1.

3. Results

Fig. 1 compares the $\delta_{\alpha\beta}$ and $\gamma_{\alpha\beta}$ correlators for same- and opposite-charge pairs in Xe–Xe collisions at $\sqrt{s_{\text{NN}}} = 5.44$ TeV to those measured in Pb–Pb collisions at $\sqrt{s_{\text{NN}}} = 5.02$ TeV [42] as a function of centrality and average charged-particle multiplicity density $\langle dN_{\text{ch}}/d\eta \rangle$ at midrapidity [65,66]. The results for same-charge pairs denote the average between pairs of particles with only positive and only negative charges since the two combinations are consistent within statistical uncertainties. Both correlators exhibit strong dependence on the charge-sign combination and qualitatively similar centrality dependence in the two systems. For $\delta_{\alpha\beta}$, the magnitude of the same- and opposite-charge pair correlations is positive and increases from central to peripheral collisions. In contrast to the CME expectation, the correlation for the opposite-charge pairs is stronger than for the same-charge combinations, indicating that background dominates these measurements. For $\gamma_{\alpha\beta}$, the magnitude of opposite-charge pair correlations is close to zero within uncertainties for most of the centrality intervals, while it decreases from central to peripheral collisions becoming more negative for same-charge pairs. Thus, the correlation of opposite-charge pairs is weaker than for same-charge pairs. This ordering is compatible with a charge separation with respect to the reaction plane expected in the presence of the CME.

The $\delta_{\alpha\beta}$ for same-charge pairs shows small (if any) differences between Xe–Xe and Pb–Pb collisions within uncertainties, while the correlations for opposite-charge pairs have larger magnitude in Xe–Xe collisions in the 10–70% centrality interval. The $\gamma_{\alpha\beta}$ for same- and opposite-charge pairs from Xe–Xe and Pb–Pb collisions have similar magnitudes within uncertainties in the 0–10% and 50–70% centrality ranges, while the correlations are stronger in Xe–Xe collisions in the 10–50% centrality interval. These observations can be attributed to the different number of particles produced in the collision within a given centrality interval between the two systems that dilutes the correlations. This is supported by the fact that the $\delta_{\alpha\beta}$ and $\gamma_{\alpha\beta}$ for same- and opposite-charge pairs from Xe–Xe collisions are consistent within uncertainties with the corresponding Pb–Pb results when reported as a function of $\langle dN_{\text{ch}}/d\eta \rangle$.

The $\gamma_{\alpha\beta}$ correlator is also investigated as a function of the pseudorapidity difference $\Delta\eta = |\eta_\alpha - \eta_\beta|$, the transverse momentum difference $\Delta p_T = |p_{T,\alpha} - p_{T,\beta}|$, and the average transverse momentum $\langle p_T \rangle = (p_{T,\alpha} + p_{T,\beta})/2$ of the pair. Fig. 2 presents these results for same- and opposite-charge pairs in the 20–30% centrality interval compared to measurements performed in 30–40% Pb–Pb collisions [42]. Different Xe–Xe and Pb–Pb centrality intervals are selected since they have similar transverse densities ($1/S dN_{\text{ch}}/d\eta \sim 10 \text{ fm}^{-2}$ with S being the transverse area) and transverse sizes ($R = \sqrt{S/\pi} \sim 4 \text{ fm}$) [44], thus the contribution from dilution effects is comparable. In addition, the value of v_2/ε_2 (ε_2 is the second-order eccentricity coefficient and characterises the elliptic shape of the initial geometry) and the influence of radial flow are similar in the two systems for these centrality classes [44,56]. The opposite-charge pair correlations from Xe–Xe collisions show a weak dependence on $\Delta\eta$ and $\langle p_T \rangle$, while they increase with increasing Δp_T of the pair. The correlations for the same-charge pairs do not exhibit any significant dependence on Δp_T and $\langle p_T \rangle$ within uncertainties. These




Fig. 1. The $\delta_{\alpha\beta}$ (top panels) and $\gamma_{\alpha\beta}$ (bottom panels) correlators as a function of centrality (left panels) and charged-particle density [65,66] (right panels) for pairs of particles with same (closed markers) and opposite (open markers) charges from Xe–Xe collisions at $\sqrt{s_{NN}} = 5.44$ TeV (red circles) compared to Pb–Pb collisions at $\sqrt{s_{NN}} = 5.02$ TeV (black squares) [42]. The Pb–Pb points are slightly shifted along the horizontal axis for better visibility in the left panels. Bars (boxes) denote statistical (systematic) uncertainties.




Fig. 2. The dependence of $\gamma_{\alpha\beta}$ on the pseudorapidity difference $|\eta_\alpha - \eta_\beta|$ (left panel), the transverse momentum difference $|\rho_{T_\alpha} - \rho_{T_\beta}|$ (middle panel), and the average transverse momentum $(p_{T_\alpha} + p_{T_\beta})/2$ (right panel) for pairs of particles with same (closed markers) and opposite (open markers) charges from 20–30% Xe–Xe collisions at $\sqrt{s_{NN}} = 5.44$ TeV (red circles) compared to results from 30–40% Pb–Pb collisions at $\sqrt{s_{NN}} = 5.02$ TeV (black squares) [42]. The Pb–Pb and Xe–Xe same-charge points are slightly shifted along the horizontal axis for better visibility in all panels. Bars (boxes) denote statistical (systematic) uncertainties.

correlations show a strong dependence on $\Delta\eta$ with a width of approximately one unit in pseudorapidity difference. The Xe–Xe and Pb–Pb results are compatible within uncertainties demonstrating similar behaviour of this observable despite the differences in the magnetic field.

To get insight into the origin of the charge-dependent effects observed in Xe–Xe collisions, two different approaches were investigated. The first one relies on a blast wave (BW) model based on a parameterisation from Ref. [67]. Input parameters of the model are tuned to describe the p_T spectra [68] and the p_T -differential v_2 values [69] of charged pions and kaons, as well as of protons and antiprotons, measured in the same collision system and centre-of-mass energy. To account for the main source of background in the measurements reported in this article, the model was further extended by including effects from local charge conservation (LCC). This was done by generating sources uniformly at the surface of the ellipse that surrounds the centre of the system and allowing them to decay into particles with opposite charge. In this extension of the BW model, the number of sources that emit oppositely-charged pairs is tuned separately for each centrality interval to reproduce the centrality dependence of $\Delta\delta_{\alpha\beta} \equiv \delta_{\alpha\beta}^{\text{opp.}} - \delta_{\alpha\beta}^{\text{same}}$, the correlator that is mainly sensitive to background effects. This is illustrated in the upper panel of Fig. 3, where the BW curve is represented

by the blue, solid line that goes through the experimental data points. The tuned model is then used to extract the expectation for the centrality dependence of $\Delta\gamma_{\alpha\beta} \equiv \gamma_{\alpha\beta}^{\text{opp.}} - \gamma_{\alpha\beta}^{\text{same}}$, shown in the lower panel of Fig. 3. The width of the band reflects the uncertainty obtained by propagating the corresponding uncertainties of the model parameters using a sub-sampling method. It can be seen that the BW model describes fairly well the measured data points for all centrality intervals. This is in contrast to the picture that emerged in Pb–Pb collisions, where following a similar procedure the same model underestimated the measurements of $\Delta\gamma_{\alpha\beta}$ by as much as $\approx 40\%$ [42].

Additional insight can be obtained by comparing the results with calculations from the Anomalous Viscous Fluid Dynamics (AVFD) model [46–48]. It relies on a Glauber model description of the initial state of the collision and accounts for the development of the early-stage electromagnetic fields as well as for the propagation of anomalous fermion currents. The expanding medium is described using a 2+1 dimensional viscous hydrodynamics (VISH2+1) code [70] which is coupled to a hadron cascade model (UrQMD) [71,72]. Within AVFD, the final-state CME signal induced by the initial chirality imbalance is controlled by the axial current density n_5/s . At the same time, the relevant background in AVFD is governed by the amount of positive and nega-




Fig. 3. Centrality evolution of the difference between opposite- and same-charge pair correlations for $\delta_{\alpha\beta}$ (top panel) and $\gamma_{\alpha\beta}$ (bottom panel) compared to model calculations: blast wave (BW) parameterisation [67] coupled to local charge conservation (LCC) effects (blue curves) and Anomalous Viscous Fluid Dynamics (AVFD) [46,47] (green curves). The BW+LCC model is tuned to reproduce the centrality dependence of $\Delta\delta_{\alpha\beta}$, while AVFD is tuned to describe simultaneously the centrality dependence of $\Delta\delta_{\alpha\beta}$ and $\Delta\gamma_{\alpha\beta}$. Bars (boxes) denote statistical (systematic) uncertainties on the data points, while the thickness of the curves represents the uncertainties on the model calculations.

tive charged partners emitted from the same fluid element relative to the total multiplicity of the event, i.e., the LCC percentage. The model was first calibrated to describe the centrality dependence of both the charged-particle density [66] and the elliptic flow [44]. As a second step, the dependence of $\Delta\delta_{\alpha\beta}$ and $\Delta\gamma_{\alpha\beta}$ on both the CME signal and the background was determined, by analysing samples with either increasing values of n_s/s or LCC percentage, respectively [45]. These results made it possible to extract the combination of n_s/s and LCC percentage that describes the data and led to a quantitative simultaneous description of the centrality dependence of $\Delta\delta_{\alpha\beta}$ and $\Delta\gamma_{\alpha\beta}$ [45]. This is illustrated in the two panels of Fig. 3. Also in this case, the width of the band reflects the uncertainty obtained by propagating the corresponding uncertainties of the model parameters using a sub-sampling




Fig. 4. Difference between opposite- and same-charge pair correlations for $\gamma_{\alpha\beta}$ divided by v_2 [44] as a function of centrality (left) and charged-particle density (right) compared to results from Pb–Pb collisions at $\sqrt{s_{NN}} = 5.02$ TeV [42]. The Pb–Pb points are slightly shifted along the horizontal axis for better visibility in the left panel. Bars (boxes) denote statistical (systematic) uncertainties.

method. The experimental data points can be described by large values of LCC contribution, between 40% and 60% for peripheral and more central Xe–Xe collisions, respectively. In addition, the values of n_s/s extracted from this procedure did not exhibit any significant centrality dependence and were compatible with zero within uncertainties [45]. Similar to the conclusion extracted from the BW model, the study within the AVFD framework indicates that the experimental measurements in Xe–Xe collisions are dominated by background.

The charge separation effect can be further studied using the difference between opposite- and same-charge pair correlations $\Delta\gamma_{\alpha\beta}$ and the CME signal expectations from MC calculations employed as guidance. A comparison between $\Delta\gamma_{\alpha\beta}$ divided by v_2 [44] in Xe–Xe collisions and that measured in Pb–Pb collisions [42] is presented as a function of centrality and charged-particle density [65,66] in Fig. 4. The value of $\Delta\gamma_{\alpha\beta}/v_2$ is positive for all centralities and its magnitude increases from central to peripheral collisions. Furthermore, it is slightly higher in Xe–Xe than Pb–Pb collisions in the 10–60% centrality interval. However, the Xe–Xe and Pb–Pb data points fall approximately onto the same curve when reported as a function of $\langle dN_{ch}/d\eta \rangle$.

The expected centrality dependence of the CME signal in Xe–Xe and Pb–Pb collisions is estimated from MC Glauber [73] simulations including a magnetic field [39]. In these calculations, the ^{208}Pb nucleus is spherical while the ^{129}Xe nucleus is deformed with a deformation parameter $\beta_2 = 0.18 \pm 0.02$ [55]. The centrality classes are determined from the multiplicity of charged particles in the acceptance of the V0 detector, which is generated according to a negative binomial distribution with parameters taken from Ref. [74]. The magnetic field is calculated at the origin from the number of spectator protons using Eq. (A.6) from Ref. [19] with the proper time $\tau = 0.1$ fm/c. Fig. 5 shows the centrality dependence of $\langle (\epsilon B)^2 \cos(2(\Psi_B - \Psi_2)) \rangle$ (Ψ_B is the direction of the magnetic field \vec{B}), which is the expected CME signal contribution in $\gamma_{\alpha\beta}$, for the two collision systems. The expected CME signal is stronger in Pb–Pb than Xe–Xe collisions in a given centrality interval due to the smaller magnetic field strength and a larger decorrelation between Ψ_B and Ψ_2 in Xe–Xe collisions. Similar results are found using T_RENTo [75] initial conditions. This observation coupled to the agreement of $\Delta\gamma_{\alpha\beta}$ between the two collision systems (see Fig. 4) points to a large background contribution to $\gamma_{\alpha\beta}$ in Xe–Xe collisions.

The large differences in the CME signal expectations and the small variations in v_2 ($< 10\%$ for the 5–70% centrality interval) [44] between the two collision systems can be used to disentangle the potential CME signal from the background in Xe–Xe and Pb–Pb collisions. Assuming that both the CME signal and the background scale with $dN_{ch}/d\eta$, the charge dependence of $\gamma_{\alpha\beta}$ for the two collision systems can be ex-




Fig. 5. The expected CME signal as a function of centrality from MC Glauber simulations for Xe-Xe and Pb-Pb collisions [73] (see text for details).

pressed using a two-component approach similar to the one proposed in Ref. [76]

$$\Gamma^{\text{Xe-Xe}} = sB^{\text{Xe-Xe}} + bv_2^{\text{Xe-Xe}}, \quad (5)$$

$$\Gamma^{\text{Pb-Pb}} = sB^{\text{Pb-Pb}} + bv_2^{\text{Pb-Pb}}, \quad (6)$$

where $\Gamma \equiv \gamma_{\alpha\beta} dN_{\text{ch}}/d\eta$, $B \equiv \langle (eB)^2 \cos(2(\Psi_B - \Psi_2)) \rangle$, and v_2 is taken from Ref. [44] and $\gamma_{\alpha\beta}^{\text{Pb-Pb}}$ from Ref. [42]. The s and b parameters quantify the signal and background contributions, respectively, and do not depend on collision system within a given centrality interval as a result of the assumption that both scale with $dN_{\text{ch}}/d\eta$. While the $dN_{\text{ch}}/d\eta$ scaling is expected for b since it is dominated by flowing clusters [25], the domains responsible for s can be considered small and thus they can be regarded as “usual” clusters which scale with $dN_{\text{ch}}/d\eta$. This scaling is further supported by the AVFD calculations performed for Pb-Pb and Xe-Xe collisions [45]. The s and b parameters can be used to calculate the fractions of the CME signal (denoted as f_{CME}) in Xe-Xe and Pb-Pb collisions as

$$f_{\text{CME}} = \frac{sB}{sB + bv_2}. \quad (7)$$

The smaller CME signal in Xe-Xe collisions also results in a tighter limit on f_{CME} in Xe-Xe than in Pb-Pb collisions. It is worth noting that the CME fractions in the two collision systems are correlated because both are calculated with the same s and b parameters, extracted from the data using Eqs. (5) and (6).

Fig. 6 presents the centrality dependence of f_{CME} in Xe-Xe and Pb-Pb collisions for the two models used in this study. The uncertainties in the CME fractions are obtained adding in quadrature the statistical and systematic uncertainties in Eqs. (5) and (6). The f_{CME} does not depend significantly on the proper time used to calculate the magnetic field since varying the value from 0.1 fm/c to 0.01 fm/c, 0.5 fm/c, and 1 fm/c yields similar CME fractions. Furthermore, the dependence on the centrality range used in Pb-Pb collisions has been studied. The analysis was performed using only a single centrality interval of Pb-Pb collisions and all Xe-Xe centrality classes. For all eight Pb-Pb centrality intervals, a good agreement is found with the nominal f_{CME} . The f_{CME} is compatible with zero up to 30% centrality in both systems and then becomes positive for midcentral and peripheral collisions with larger values in Pb-Pb than in Xe-Xe. The CME fraction for the 0–30% centrality interval in Pb-Pb collisions agrees with the one reported in Ref. [42]. Fitting the data points in the centrality range 0–70% with a constant function neglecting any centrality dependence gives $f_{\text{CME}} = -0.003 \pm 0.010$ ($f_{\text{CME}} = -0.001 \pm 0.012$) and $f_{\text{CME}} = 0.147 \pm 0.061$ ($f_{\text{CME}} = 0.150 \pm 0.062$) for MC Glauber ($T_{\text{R}}\text{ENTo}$) initial conditions in Xe-Xe and Pb-Pb collisions, respectively. These results are consistent with zero CME fraction in Xe-Xe collisions and correspond to upper limits on f_{CME} of 2% (3%) and 25% (32%) at 95% (99.7%) confidence level for the 0–70% centrality interval in Xe-Xe and Pb-Pb collisions, respectively. The limits are estimated assuming Gaussian uncertainties.

4. Summary

The charge-dependent two- and three-particle correlators $\delta_{\alpha\beta}$ and $\gamma_{\alpha\beta}$ have been measured in Xe-Xe collisions at $\sqrt{s_{\text{NN}}} = 5.44$ TeV. The charge dependence of these correlators is strongly correlated with centrality, increasing from central to peripheral collisions, and is qualitatively similar to those reported in Pb-Pb collisions. The difference between the Xe-Xe and Pb-Pb results mostly arises from dilution effects since the data points from both collision systems fall approximately onto the same curve when presented as a function of charged-particle density. Monte Carlo simulations with different initial state models predict a significantly larger magnitude of the CME signal in Pb-Pb than Xe-Xe collisions, which implies that the dominant contribution to $\gamma_{\alpha\beta}$ in Xe-Xe collisions is due to background effects. The magnitude of the charge dependence of $\gamma_{\alpha\beta}$ is described over the entire centrality range by a blast wave parameterisation that incorporates local charge conservation tuned to reproduce the components of the background. This magnitude is also reproduced by Anomalous Viscous Fluid Dynamics calculations with large contributions from local charge conservation effects and values of the CME signal close to zero, thus indicating that the background is the dominant contribution to the three-particle correlator.




Fig. 6. Centrality dependence of the CME fraction extracted using Eq. (7) with the expected CME signal from MC Glauber [73] (closed markers) and $T_{\text{R}}\text{ENTo}$ [75] (open markers) models (see text for details). The $T_{\text{R}}\text{ENTo}$ points are slightly shifted along the horizontal axis for better visibility.

tor. In order to get a quantitative estimate of the signal and background contributions, the measured values of $\gamma_{\alpha\beta}$ in Xe–Xe and Pb–Pb collisions are compared using a two-component approach. This procedure allows one to estimate the fraction of the CME signal in both collision systems. Averaging over the 0–70% centrality interval, an upper limit of 2% (3%) and 25% (32%) is estimated at 95% (99.7%) confidence level for the CME contribution to the charge dependence of $\gamma_{\alpha\beta}$ in Xe–Xe and Pb–Pb collisions, respectively.

Declaration of competing interest

The authors declare that they have no known competing financial interests or personal relationships that could have appeared to influence the work reported in this paper.

Data availability

This manuscript has associated data in a HEPData repository at: <https://www.hepdata.net/record/ins2172062>.

Acknowledgements

The ALICE Collaboration would like to thank all its engineers and technicians for their invaluable contributions to the construction of the experiment and the CERN accelerator teams for the outstanding performance of the LHC complex. The ALICE Collaboration gratefully acknowledges the resources and support provided by all Grid centres and the Worldwide LHC Computing Grid (WLCG) collaboration. The ALICE Collaboration acknowledges the following funding agencies for their support in building and running the ALICE detector: A. I. Alikhanyan National Science Laboratory (Yerevan Physics Institute) Foundation (ANSI), State Committee of Science and World Federation of Scientists (WFS), Armenia; Austrian Academy of Sciences, Austrian Science Fund (FWF): [M 2467-N36] and Nationalstiftung für Forschung, Technologie und Entwicklung, Austria; Ministry of Communications and High Technologies, National Nuclear Research Center, Azerbaijan; Conselho Nacional de Desenvolvimento Científico e Tecnológico (CNPq), Financiadora de Estudos e Projetos (Finep), Fundação de Amparo à Pesquisa do Estado de São Paulo (FAPESP) and Universidade Federal do Rio Grande do Sul (UFRGS), Brazil; Bulgarian Ministry of Education and Science, within the National Roadmap for Research Infrastructures 2020–2027 (object CERN), Bulgaria; Ministry of Education of China (MOEC), Ministry of Science & Technology of China (MSTC) and National Natural Science Foundation of China (NSFC), China; Ministry of Science and Education and Croatian Science Foundation, Croatia; Centro de Aplicaciones Tecnológicas y Desarrollo Nuclear (CEADEN), Cubaenergía, Cuba; The Ministry of Education, Youth and Sports of the Czech Republic, Czech Republic; The Danish Council for Independent Research | Natural Sciences, the Villum Fonden and Danish National Research Foundation (DNRF), Denmark; Helsinki Institute of Physics (HIP), Finland; Commissariat à l'Énergie Atomique (CEA) and Institut National de Physique Nucléaire et de Physique des Particules (IN2P3) and Centre National de la Recherche Scientifique (CNRS), France; Bundesministerium für Bildung und Forschung (BMBF) and GSI Helmholtzzentrum für Schwerionenforschung GmbH, Germany; General Secretariat for Research and Technology, Ministry of Education, Research and Religions, Greece; National Research, Development and Innovation Office, Hungary; Department of Atomic Energy, Government of India (DAE), Department of Science and Technology, Government of India (DST), University Grants Commission, Government of India (UGC) and Council of Scientific and Industrial Research (CSIR), India; National Research and Innovation Agency - BRIN, Indonesia; Istituto Nazionale di Fisica Nucleare (INFN), Italy; Japanese Ministry of Education, Culture, Sports, Science and Technology (MEXT) and Japan Society for the Promotion of Science (JSPS) KAKENHI, Japan; Consejo Nacional de Ciencia (CONACYT) y Tecnología, through Fondo de Cooperación Internacional en

Ciencia y Tecnología (FONCICYT) and Dirección General de Asuntos del Personal Académico (DGAPA), Mexico; Nederlandse Organisatie voor Wetenschappelijk Onderzoek (NWO), Netherlands; The Research Council of Norway, Norway; Commission on Science and Technology for Sustainable Development in the South (COMSATS), Pakistan; Pontificia Universidad Católica del Perú, Peru; Ministry of Education and Science, National Science Centre and WUT ID-UB, Poland; Korea Institute of Science and Technology Information and National Research Foundation of Korea (NRF), Republic of Korea; Ministry of Education and Scientific Research, Institute of Atomic Physics, Ministry of Research and Innovation and Institute of Atomic Physics and Universitatea Nationala de Stiinta si Tehnologie Politehnica Bucuresti, Romania; Ministry of Education, Science, Research and Sport of the Slovak Republic, Slovakia; National Research Foundation of South Africa, South Africa; Swedish Research Council (VR) and Knut and Alice Wallenberg Foundation (KAW), Sweden; European Organization for Nuclear Research, Switzerland; Suranaree University of Technology (SUT), National Science and Technology Development Agency (NSTDA) and National Science, Research and Innovation Fund (NSRF via PMU-B B05F650021), Thailand; Turkish Energy, Nuclear and Mineral Research Agency (TENMAK), Turkey; National Academy of Sciences of Ukraine, Ukraine; Science and Technology Facilities Council (STFC), United Kingdom; National Science Foundation of the United States of America (NSF) and United States Department of Energy, Office of Nuclear Physics (DOE NP), United States of America. In addition, individual groups or members have received support from: Marie Skłodowska Curie, European Research Council, Strong 2020 - Horizon 2020 (grant nos. 950692, 824093, 896850), European Union; ICSC - Centro Nazionale di Ricerca in High Performance Computing, Big Data and Quantum Computing, European Union - NextGenerationEU; Academy of Finland (Center of Excellence in Quark Matter) (grant nos. 346327, 346328), Finland; Programa de Apoyos para la Superación del Personal Académico, UNAM, Mexico.

References

- [1] E.V. Shuryak, Quark-gluon plasma and hadronic production of leptons, photons and pions, *Phys. Lett. B* 78 (1978) 150.
- [2] F. Karsch, E. Laermann, A. Peikert, Quark mass and flavor dependence of the QCD phase transition, *Nucl. Phys. B* 605 (2001) 579–599, arXiv:hep-lat/0012023.
- [3] F. Karsch, E. Laermann, A. Peikert, The pressure in two flavor, (2+1)-flavor and three flavor QCD, *Phys. Lett. B* 478 (2000) 447–455, arXiv:hep-lat/0002003.
- [4] F. Karsch, Lattice results on QCD thermodynamics, *Nucl. Phys. A* 698 (2002) 199–208, arXiv:hep-ph/0103314.
- [5] C.R. Allton, et al., The QCD thermal phase transition in the presence of a small chemical potential, *Phys. Rev. D* 66 (2002) 074507, arXiv:hep-lat/0204010.
- [6] A. Bazavov, et al., Equation of state and QCD transition at finite temperature, *Phys. Rev. D* 80 (2009) 014504, arXiv:0903.4379 [hep-lat].
- [7] A. Bazavov, et al., The chiral and deconfinement aspects of the QCD transition, *Phys. Rev. D* 85 (2012) 054503, arXiv:1111.1710 [hep-lat].
- [8] S. Borsanyi, et al., The QCD equation of state with dynamical quarks, *J. High Energy Phys.* 11 (2010) 077, arXiv:1007.2580 [hep-lat].
- [9] CMS Collaboration, S. Chatrchyan, et al., Measurement of the pseudorapidity and centrality dependence of the transverse energy density in PbPb collisions at $\sqrt{s_{NN}} = 2.76$ TeV, *Phys. Rev. Lett.* 109 (2012) 152303, arXiv:1205.2488 [nucl-ex].
- [10] ALICE Collaboration, J. Adam, et al., Direct photon production in Pb–Pb collisions at $\sqrt{s_{NN}} = 2.76$ TeV, *Phys. Lett. B* 754 (2016) 235–248, arXiv:1509.07324 [nucl-ex].
- [11] ALICE Collaboration, J. Adam, et al., Measurement of transverse energy at midrapidity in Pb–Pb collisions at $\sqrt{s_{NN}} = 2.76$ TeV, *Phys. Rev. C* 94 (2016) 034903, arXiv:1603.04775 [nucl-ex].
- [12] T. Lee, A theory of spontaneous T violation, *Phys. Rev. D* 8 (1973) 1226–1239.
- [13] T. Lee, G. Wick, Vacuum stability and vacuum excitation in a spin 0 field theory, *Phys. Rev. D* 9 (1974) 2291–2316.
- [14] P. Morley, I. Schmidt, Strong P, CP, T violations in heavy ion collisions, *Z. Phys. C* 26 (1985) 627.
- [15] D. Kharzeev, R. Pisarski, M.H. Tytgat, Possibility of spontaneous parity violation in hot QCD, *Phys. Rev. Lett.* 81 (1998) 512–515, arXiv:hep-ph/9804221.
- [16] D. Kharzeev, R.D. Pisarski, Pionic measures of parity and CP violation in high-energy nuclear collisions, *Phys. Rev. D* 61 (2000) 111901, arXiv:hep-ph/9906401.
- [17] D.E. Kharzeev, Topology, magnetic field, and strongly interacting matter, *Annu. Rev. Nucl. Part. Sci.* 65 (2015) 193–214, arXiv:1501.01336 [hep-ph].
- [18] D. Kharzeev, A. Zhitnitsky, Charge separation induced by P-odd bubbles in QCD matter, *Nucl. Phys. A* 797 (2007) 67–79, arXiv:0706.1026 [hep-ph].

- [19] D.E. Kharzeev, L.D. McLerran, H.J. Warringa, The effects of topological charge change in heavy ion collisions: 'event by event P and CP violation', Nucl. Phys. A 803 (2008) 227–253, arXiv:0711.0950 [hep-ph].
- [20] K. Fukushima, D.E. Kharzeev, H.J. Warringa, The chiral magnetic effect, Phys. Rev. D 78 (2008) 074033, arXiv:0808.3382 [hep-ph].
- [21] V. Skokov, A. Illarionov, V. Toneev, Estimate of the magnetic field strength in heavy-ion collisions, Int. J. Mod. Phys. A 24 (2009) 5925–5932, arXiv:0907.1396 [nucl-th].
- [22] A. Bzdak, V. Skokov, Event-by-event fluctuations of magnetic and electric fields in heavy ion collisions, Phys. Lett. B 710 (2012) 171–174, arXiv:1111.1949 [hep-ph].
- [23] W.-T. Deng, X.-G. Huang, Event-by-event generation of electromagnetic fields in heavy-ion collisions, Phys. Rev. C 85 (2012) 044907, arXiv:1201.5108 [nucl-th].
- [24] S. Voloshin, Y. Zhang, Flow study in relativistic nuclear collisions by Fourier expansion of azimuthal particle distributions, Z. Phys. C 70 (1996) 665–672, arXiv:hep-ph/9407282.
- [25] S.A. Voloshin, Parity violation in hot QCD: how to detect it, Phys. Rev. C 70 (2004) 057901, arXiv:hep-ph/0406311.
- [26] A.M. Poskanzer, S.A. Voloshin, Methods for analyzing anisotropic flow in relativistic nuclear collisions, Phys. Rev. C 58 (1998) 1671–1678, arXiv:nucl-ex/9805001.
- [27] STAR Collaboration, B.I. Abelev, et al., Azimuthal charged-particle correlations and possible local strong parity violation, Phys. Rev. Lett. 103 (2009) 251601, arXiv:0909.1739 [nucl-ex].
- [28] STAR Collaboration, B.I. Abelev, et al., Observation of charge-dependent azimuthal correlations and possible local strong parity violation in heavy ion collisions, Phys. Rev. C 81 (2010) 054908, arXiv:0909.1717 [nucl-ex].
- [29] ALICE Collaboration, B. Abelev, et al., Charge separation relative to the reaction plane in Pb–Pb collisions at $\sqrt{s_{NN}} = 2.76$ TeV, Phys. Rev. Lett. 110 (2013) 012301, arXiv:1207.0900 [nucl-ex].
- [30] ALICE Collaboration, K. Aamodt, et al., Charged-particle multiplicity density at mid-rapidity in central Pb–Pb collisions at $\sqrt{s_{NN}} = 2.76$ TeV, Phys. Rev. Lett. 105 (2010) 252301, arXiv:1011.3916 [nucl-ex].
- [31] ALICE Collaboration, K. Aamodt, et al., Centrality dependence of the charged-particle multiplicity density at mid-rapidity in Pb–Pb collisions at $\sqrt{s_{NN}} = 2.76$ TeV, Phys. Rev. Lett. 106 (2011) 032301, arXiv:1012.1657 [nucl-ex].
- [32] F. Wang, Effects of cluster particle correlations on local parity violation observables, Phys. Rev. C 81 (2010) 064902, arXiv:0911.1482 [nucl-ex].
- [33] A. Bzdak, V. Koch, J. Liao, Remarks on possible local parity violation in heavy ion collisions, Phys. Rev. C 81 (2010) 031901, arXiv:0912.5050 [nucl-th].
- [34] J. Liao, V. Koch, A. Bzdak, Charge separation effect in relativistic heavy ion collisions, Phys. Rev. C 82 (Nov 2010) 054902, <https://link.aps.org/doi/10.1103/PhysRevC.82.054902>.
- [35] S. Schlichting, S. Pratt, Charge conservation at energies available at the BNL relativistic heavy ion collider and contributions to local parity violation observables, Phys. Rev. C 83 (2011) 014913, arXiv:1009.4283 [nucl-th].
- [36] S. Pratt, S. Schlichting, S. Gavin, Effects of momentum conservation and flow on angular correlations at RHIC, Phys. Rev. C 84 (2011) 024909, arXiv:1011.6053 [nucl-th].
- [37] CMS Collaboration, V. Khachatryan, et al., Observation of charge-dependent azimuthal correlations in p -Pb collisions and its implication for the search for the chiral magnetic effect, Phys. Rev. Lett. 118 (2017) 122301, arXiv:1610.00263 [nucl-ex].
- [38] STAR Collaboration, J. Adam, et al., Charge-dependent pair correlations relative to a third particle in $p + Au$ and $d + Au$ collisions at RHIC, Phys. Lett. B 798 (2019) 134975, arXiv:1906.03373 [nucl-ex].
- [39] ALICE Collaboration, S. Acharya, et al., Constraining the magnitude of the chiral magnetic effect with event shape engineering in Pb–Pb collisions at $\sqrt{s_{NN}} = 2.76$ TeV, Phys. Lett. B 777 (2018) 151–162, arXiv:1709.04723 [nucl-ex].
- [40] J. Schukraft, A. Timmins, S.A. Voloshin, Ultra-relativistic nuclear collisions: event shape engineering, Phys. Lett. B 719 (2013) 394–398, arXiv:1208.4563 [nucl-ex].
- [41] CMS Collaboration, A.M. Sirunyan, et al., Constraints on the chiral magnetic effect using charge-dependent azimuthal correlations in p Pb and PbPb collisions at the CERN large hadron collider, Phys. Rev. C 97 (2018) 044912, arXiv:1708.01602 [nucl-ex].
- [42] ALICE Collaboration, S. Acharya, et al., Constraining the chiral magnetic effect with charge-dependent azimuthal correlations in Pb–Pb collisions at $\sqrt{s_{NN}} = 2.76$ and 5.02 TeV, J. High Energy Phys. 09 (2020) 160, arXiv:2005.14640 [nucl-ex].
- [43] STAR Collaboration, M. Abdallah, et al., Search for the chiral magnetic effect with isobar collisions at $\sqrt{s_{NN}} = 200$ GeV by the STAR collaboration at the BNL relativistic heavy ion collider, Phys. Rev. C 105 (2022) 014901, arXiv:2109.00131 [nucl-ex].
- [44] ALICE Collaboration, S. Acharya, et al., Anisotropic flow in Xe–Xe collisions at $\sqrt{s_{NN}} = 5.44$ TeV, Phys. Lett. B 784 (2018) 82–95, arXiv:1805.01832 [nucl-ex].
- [45] P. Christakoglou, S. Qiu, J. Staa, Systematic study of the chiral magnetic effect with the AVFD model at LHC energies, Eur. Phys. J. C 81 (2021) 717, arXiv:2106.03537 [nucl-th].
- [46] S. Shi, Y. Jiang, E. Lilleskov, J. Liao, Anomalous chiral transport in heavy ion collisions from anomalous-viscous fluid dynamics, Ann. Phys. 394 (2018) 50–72, arXiv:1711.02496 [nucl-th].
- [47] Y. Jiang, S. Shi, Y. Yin, J. Liao, Quantifying the chiral magnetic effect from anomalous-viscous fluid dynamics, Chin. Phys. C 42 (2018) 011001, arXiv:1611.04586 [nucl-th].
- [48] S. Shi, H. Zhang, D. Hou, J. Liao, Signatures of chiral magnetic effect in the collisions of isobars, Phys. Rev. Lett. 125 (2020) 242301, arXiv:1910.14010 [nucl-th].
- [49] ALICE Collaboration, K. Aamodt, et al., The ALICE experiment at the CERN LHC, J. Instrum. 3 (2008) S08002.
- [50] ALICE Collaboration, B.B. Abelev, et al., Performance of the ALICE experiment at the CERN LHC, Int. J. Mod. Phys. A 29 (2014) 1430044, arXiv:1402.4476 [nucl-ex].
- [51] ALICE Collaboration, K. Aamodt, et al., Alignment of the ALICE inner tracking system with cosmic-ray tracks, J. Instrum. 5 (2010) P03003, arXiv:1001.0502 [physics.ins-det].
- [52] J. Alme, et al., The ALICE TPC, a large 3-dimensional tracking device with fast read-out for ultra-high multiplicity events, Nucl. Instrum. Methods A 622 (2010) 316–367, arXiv:1001.1950 [physics.ins-det].
- [53] ALICE Collaboration, E. Abbas, et al., Performance of the ALICE VZERO system, J. Instrum. 8 (2013) P10016, arXiv:1306.3130 [nucl-ex].
- [54] R. Arnaldi, et al., The zero degree calorimeters for the ALICE experiment, Nucl. Instrum. Methods A 581 (2007) 397–401, Nucl. Instrum. Methods A 604 (2009) 765 (Erratum).
- [55] ALICE Collaboration, S. Acharya, et al., Centrality determination using the Glauber model in Xe–Xe collisions at $\sqrt{s_{NN}} = 5.44$ TeV, ALICE-PUBLIC-2018-003, 2018, pp. 1–23, <http://cds.cern.ch/record/2315401>.
- [56] ALICE Collaboration, S. Acharya, et al., Transverse momentum spectra and nuclear modification factors of charged particles in Xe–Xe collisions at $\sqrt{s_{NN}} = 5.44$ TeV, Phys. Lett. B 788 (2019) 166–179, arXiv:1805.04399 [nucl-ex].
- [57] X.-N. Wang, M. Gyulassy, HIJING: a Monte Carlo model for multiple jet production in pp, pA and AA collisions, Phys. Rev. D 44 (1991) 3501–3516.
- [58] M. Gyulassy, X.-N. Wang, HIJING 1.0: a Monte Carlo program for parton and particle production in high-energy hadronic and nuclear collisions, Comput. Phys. Commun. 83 (1994) 307, arXiv:nucl-th/9502021.
- [59] R. Brun, et al., GEANT Detector Description and Simulation Tool, CERN-W5013 1, 1994, p. 1, <https://cds.cern.ch/record/1082634>.
- [60] U. Gürsoy, D. Kharzeev, E. Marcus, K. Rajagopal, C. Shen, Charge-dependent flow induced by magnetic and electric fields in heavy ion collisions, Phys. Rev. C 98 (2018) 055201, arXiv:1806.05288 [hep-ph].
- [61] ATLAS Collaboration, G. Aad, et al., Longitudinal flow decorrelations in Xe+Xe collisions at $\sqrt{s_{NN}} = 5.44$ TeV with the ATLAS detector, Phys. Rev. Lett. 126 (2021) 122301, arXiv:2001.04201 [nucl-ex].
- [62] I. Selyuzhenkov, S. Voloshin, Effects of non-uniform acceptance in anisotropic flow measurement, Phys. Rev. C 77 (2008) 034904, arXiv:0707.4672 [nucl-th].
- [63] ALICE Collaboration, S. Acharya, et al., Transverse momentum spectra and nuclear modification factors of charged particles in pp, p–Pb and Pb–Pb collisions at the LHC, J. High Energy Phys. 11 (2018) 013, arXiv:1802.09145 [nucl-ex].
- [64] R. Barlow, Systematic errors: facts and fictions, in: Conference on Advanced Statistical Techniques in Particle Physics, 7, 2002, pp. 134–144, arXiv:hep-ex/0207026.
- [65] ALICE Collaboration, J. Adam, et al., Centrality dependence of the charged-particle multiplicity density at midrapidity in Pb–Pb collisions at $\sqrt{s_{NN}} = 5.02$ TeV, Phys. Rev. Lett. 116 (2016) 222302, arXiv:1512.06104 [nucl-ex].
- [66] ALICE Collaboration, S. Acharya, et al., Centrality and pseudorapidity dependence of the charged-particle multiplicity density in Xe–Xe collisions at $\sqrt{s_{NN}} = 5.44$ TeV, Phys. Lett. B 790 (2019) 35–48, arXiv:1805.04432 [nucl-ex].
- [67] F. Retiere, M.A. Lisa, Observable implications of geometrical and dynamical aspects of freeze out in heavy ion collisions, Phys. Rev. C 70 (2004) 044907, arXiv:nucl-th/0312024.
- [68] ALICE Collaboration, S. Acharya, et al., Production of pions, kaons, (anti-)protons and ϕ mesons in Xe–Xe collisions at $\sqrt{s_{NN}} = 5.44$ TeV, arXiv:2101.03100 [nucl-ex].
- [69] ALICE Collaboration, S. Acharya, et al., Anisotropic flow of identified hadrons in Xe–Xe collisions at $\sqrt{s_{NN}} = 5.44$ TeV, J. High Energy Phys. 10 (2021) 152, arXiv:2107.10592 [nucl-ex].
- [70] H. Song, U.W. Heinz, Suppression of elliptic flow in a minimally viscous quark-gluon plasma, Phys. Lett. B 658 (2008) 279–283, arXiv:0709.0742 [nucl-th].
- [71] S.A. Bass, et al., Microscopic models for ultrarelativistic heavy ion collisions, Prog. Part. Nucl. Phys. 41 (1998) 255–369, arXiv:nucl-th/9803035.
- [72] M. Bleicher, et al., Relativistic hadron–hadron collisions in the ultrarelativistic quantum molecular dynamics model, J. Phys. G 25 (1999) 1859–1896, arXiv:hep-ph/9909407.
- [73] M.L. Miller, K. Reygers, S.J. Sanders, P. Steinberg, Glauber modeling in high energy nuclear collisions, Annu. Rev. Nucl. Part. Sci. 57 (2007) 205–243, arXiv:nucl-ex/0701025.
- [74] ALICE Collaboration, S. Acharya, et al., Centrality determination in heavy ion collisions, ALICE-PUBLIC-2018-011, 2018, pp. 1–28, <https://cds.cern.ch/record/2636623>.
- [75] J.S. Moreland, J.E. Bernhard, S.A. Bass, Alternative ansatz to wounded nucleon and binary collision scaling in high-energy nuclear collisions, Phys. Rev. C 92 (2015) 011901, arXiv:1412.4708 [nucl-th].
- [76] W.-T. Deng, X.-G. Huang, G.-L. Ma, G. Wang, Test the chiral magnetic effect with isobaric collisions, Phys. Rev. C 94 (2016) 041901, arXiv:1607.04697 [nucl-th].

ALICE Collaboration

S. Acharya ^{126, [id](#)}, D. Adamová ^{86, [id](#)}, A. Adler ⁷⁰, G. Aglieri Rinella ^{32, [id](#)}, M. Agnello ^{29, [id](#)}, N. Agrawal ^{51, [id](#)}, Z. Ahammed ^{134, [id](#)}, S. Ahmad ^{15, [id](#)}, S.U. Ahn ^{71, [id](#)}, I. Ahuja ^{37, [id](#)}, A. Akindinov ^{140, [id](#)}, M. Al-Turany ^{97, [id](#)}, D. Aleksandrov ^{140, [id](#)}, B. Alessandro ^{56, [id](#)}, H.M. Alfanda ^{6, [id](#)}, R. Alfaro Molina ^{67, [id](#)}, B. Ali ^{15, [id](#)}, A. Alici ^{25, [id](#)}, N. Alizadehvandchali ^{115, [id](#)}, A. Alkin ^{32, [id](#)}, J. Alme ^{20, [id](#)}, G. Alocco ^{52, [id](#)}, T. Alt ^{64, [id](#)}, I. Altsybeev ^{140, [id](#)}, J.R. Alvarado ^{44, [id](#)}, M.N. Anaam ^{6, [id](#)}, C. Andrei ^{45, [id](#)}, A. Andronic ^{125, [id](#)}, V. Anguelov ^{94, [id](#)}, F. Antinori ^{54, [id](#)}, P. Antonioli ^{51, [id](#)}, N. Apadula ^{74, [id](#)}, L. Aphecetche ^{103, [id](#)}, H. Appelshäuser ^{64, [id](#)}, C. Arata ^{73, [id](#)}, S. Arcelli ^{25, [id](#)}, M. Aresti ^{52, [id](#)}, R. Arnaldi ^{56, [id](#)}, I.C. Arsene ^{19, [id](#)}, M. Arslanok ^{137, [id](#)}, A. Augustinus ^{32, [id](#)}, R. Averbeck ^{97, [id](#)}, M.D. Azmi ^{15, [id](#)}, A. Badalà ^{53, [id](#)}, J. Bae ^{104, [id](#)}, Y.W. Baek ^{40, [id](#)}, X. Bai ^{119, [id](#)}, R. Bailhache ^{64, [id](#)}, Y. Bailung ^{48, [id](#)}, A. Balbino ^{29, [id](#)}, A. Baldisseri ^{129, [id](#)}, B. Balis ^{2, [id](#)}, D. Banerjee ^{4, [id](#)}, Z. Banoo ^{91, [id](#)}, R. Barbera ^{26, [id](#)}, F. Barile ^{31, [id](#)}, L. Barioglio ^{95, [id](#)}, M. Barlou ⁷⁸, G.G. Barnaföldi ^{46, [id](#)}, L.S. Barnby ^{85, [id](#)}, V. Barret ^{126, [id](#)}, L. Barreto ^{110, [id](#)}, C. Bartels ^{118, [id](#)}, K. Barth ^{32, [id](#)}, E. Bartsch ^{64, [id](#)}, N. Bastid ^{126, [id](#)}, S. Basu ^{75, [id](#)}, G. Batigne ^{103, [id](#)}, D. Battistini ^{95, [id](#)}, B. Batyunya ^{141, [id](#)}, D. Bauri ⁴⁷, J.L. Bazo Alba ^{101, [id](#)}, I.G. Bearden ^{83, [id](#)}, C. Beattie ^{137, [id](#)}, P. Becht ^{97, [id](#)}, D. Behera ^{48, [id](#)}, I. Belikov ^{128, [id](#)}, A.D.C. Bell Hechavarria ^{125, [id](#)}, F. Bellini ^{25, [id](#)}, R. Bellwied ^{115, [id](#)}, S. Belokurova ^{140, [id](#)}, V. Belyaev ^{140, [id](#)}, G. Bencedi ^{46, [id](#)}, S. Beole ^{24, [id](#)}, A. Bercuci ^{45, [id](#)}, Y. Berdnikov ^{140, [id](#)}, A. Berdnikova ^{94, [id](#)}, L. Bergmann ^{94, [id](#)}, M.G. Besoiu ^{63, [id](#)}, L. Betev ^{32, [id](#)}, P.P. Bhaduri ^{134, [id](#)}, A. Bhasin ^{91, [id](#)}, M.A. Bhat ^{4, [id](#)}, B. Bhattacharjee ^{41, [id](#)}, L. Bianchi ^{24, [id](#)}, N. Bianchi ^{49, [id](#)}, J. Bielčík ^{35, [id](#)}, J. Bielčíková ^{86, [id](#)}, J. Biernat ^{107, [id](#)}, A.P. Bigot ^{128, [id](#)}, A. Bilandzic ^{95, [id](#)}, G. Biro ^{46, [id](#)}, S. Biswas ^{4, [id](#)}, N. Bize ^{103, [id](#)}, J.T. Blair ^{108, [id](#)}, D. Blau ^{140, [id](#)}, M.B. Blidaru ^{97, [id](#)}, N. Bluhme ³⁸, C. Blume ^{64, [id](#)}, G. Boca ^{21,55, [id](#)}, F. Bock ^{87, [id](#)}, T. Bodova ^{20, [id](#)}, A. Bogdanov ¹⁴⁰, S. Boi ^{22, [id](#)}, J. Bok ^{58, [id](#)}, L. Boldizsár ^{46, [id](#)}, A. Bolozdynya ^{140, [id](#)}, M. Bombara ^{37, [id](#)}, P.M. Bond ^{32, [id](#)}, G. Bonomi ^{133,55, [id](#)}, H. Borel ^{129, [id](#)}, A. Borissov ^{140, [id](#)}, A.G. Borquez Carcamo ^{94, [id](#)}, H. Bossi ^{137, [id](#)}, E. Botta ^{24, [id](#)}, Y.E.M. Bouziani ^{64, [id](#)}, L. Bratrud ^{64, [id](#)}, P. Braun-Munzinger ^{97, [id](#)}, M. Bregant ^{110, [id](#)}, M. Broz ^{35, [id](#)}, G.E. Bruno ^{96,31, [id](#)}, M.D. Buckland ^{23, [id](#)}, D. Budnikov ^{140, [id](#)}, H. Buesching ^{64, [id](#)}, S. Bufalino ^{29, [id](#)}, O. Bugnon ¹⁰³, P. Buhler ^{102, [id](#)}, Z. Buthelezi ^{68,122, [id](#)}, S.A. Bysiak ¹⁰⁷, M. Cai ^{6, [id](#)}, H. Caines ^{137, [id](#)}, A. Caliva ^{97, [id](#)}, E. Calvo Villar ^{101, [id](#)}, J.M.M. Camacho ^{109, [id](#)}, P. Camerini ^{23, [id](#)}, F.D.M. Canedo ^{110, [id](#)}, S.L. Cantway ^{137, [id](#)}, M. Carabas ^{113, [id](#)}, A.A. Carballo ^{32, [id](#)}, F. Carnesecchi ^{32, [id](#)}, R. Caron ^{127, [id](#)}, L.A.D. Carvalho ^{110, [id](#)}, J. Castillo Castellanos ^{129, [id](#)}, F. Catalano ^{24,29, [id](#)}, C. Ceballos Sanchez ^{141, [id](#)}, I. Chakaberia ^{74, [id](#)}, P. Chakraborty ^{47, [id](#)}, S. Chandra ^{134, [id](#)}, S. Chapeland ^{32, [id](#)}, M. Chartier ^{118, [id](#)}, S. Chattopadhyay ^{134, [id](#)}, S. Chattopadhyay ^{99, [id](#)}, T. Cheng ^{97,6, [id](#)}, C. Cheshkov ^{127, [id](#)}, B. Cheynis ^{127, [id](#)}, V. Chibante Barroso ^{32, [id](#)}, D.D. Chinellato ^{111, [id](#)}, E.S. Chizzali ^{95, [id](#)}, II, J. Cho ^{58, [id](#)}, S. Cho ^{58, [id](#)}, P. Chochula ^{32, [id](#)}, P. Christakoglou ^{84, [id](#)}, C.H. Christensen ^{83, [id](#)}, P. Christiansen ^{75, [id](#)}, T. Chujo ^{124, [id](#)}, M. Ciaccio ^{29, [id](#)}, C. Cicalo ^{52, [id](#)}, F. Cindolo ^{51, [id](#)}, M.R. Ciupek ⁹⁷, G. Clai ^{51,III, [id](#)}, F. Colamaria ^{50, [id](#)}, J.S. Colburn ¹⁰⁰, D. Colella ^{96,31, [id](#)}, M. Colocci ^{32, [id](#)}, M. Concas ^{56, [id](#)}, G. Conesa Balbastre ^{73, [id](#)}, Z. Conesa del Valle ^{130, [id](#)}, G. Contin ^{23, [id](#)}, J.G. Contreras ^{35, [id](#)}, M.L. Coquet ^{129, [id](#)}, T.M. Cormier ^{87,1, [id](#)}, P. Cortese ^{132,56, [id](#)}, M.R. Cosentino ^{112, [id](#)}, F. Costa ^{32, [id](#)}, S. Costanza ^{21,55, [id](#)}, C. Cot ^{130, [id](#)}, J. Crkovská ^{94, [id](#)}, P. Crochet ^{126, [id](#)}, R. Cruz-Torres ^{74, [id](#)}, E. Cuautle ⁶⁵, P. Cui ^{6, [id](#)}, A. Dainese ^{54, [id](#)}, M.C. Danisch ^{94, [id](#)}, A. Danu ^{63, [id](#)}, P. Das ^{80, [id](#)}, P. Das ^{4, [id](#)}, S. Das ^{4, [id](#)}, A.R. Dash ^{125, [id](#)}, S. Dash ^{47, [id](#)}, A. De Caro ^{28, [id](#)}, G. de Cataldo ^{50, [id](#)}, J. de Cuveland ³⁸, A. De Falco ^{22, [id](#)}, D. De Gruttola ^{28, [id](#)}, N. De Marco ^{56, [id](#)}, C. De Martin ^{23, [id](#)}, S. De Pasquale ^{28, [id](#)}, S. Deb ^{48, [id](#)}, R.J. Debski ^{2, [id](#)}, K.R. Deja ¹³⁵, R. Del Grande ^{95, [id](#)}, L. Dello Stritto ^{28, [id](#)}, W. Deng ^{6, [id](#)}, P. Dhankher ^{18, [id](#)}, D. Di Bari ^{31, [id](#)}, A. Di Mauro ^{32, [id](#)}, R.A. Diaz ^{141,7, [id](#)}, T. Dietel ^{114, [id](#)}, Y. Ding ^{127,6, [id](#)}, R. Divià ^{32, [id](#)}, D.U. Dixit ^{18, [id](#)}, Ø. Djuvsland ²⁰, U. Dmitrieva ^{140, [id](#)}, A. Dobrin ^{63, [id](#)}, B. Dönigus ^{64, [id](#)}, J.M. Dubinski ^{135, [id](#)}, A. Dubla ^{97, [id](#)}, S. Dudi ^{90, [id](#)}, P. Dupieux ^{126, [id](#)}, M. Durkac ¹⁰⁶, N. Dzalaiova ¹², T.M. Eder ^{125, [id](#)}, R.J. Ehlers ^{87, [id](#)}, V.N. Eikeland ²⁰, F. Eisenhut ^{64, [id](#)}, D. Elia ^{50, [id](#)}, B. Erasmus ^{103, [id](#)}, F. Ercolessi ^{25, [id](#)}, F. Erhardt ^{89, [id](#)}, M.R. Ersdal ²⁰, B. Espagnon ^{130, [id](#)}, G. Eulisse ^{32, [id](#)}, D. Evans ^{100, [id](#)}, S. Evdokimov ^{140, [id](#)}, L. Fabbietti ^{95, [id](#)}, M. Faggin ^{27, [id](#)},

J. Faivre ^{73, [id](#)}, F. Fan ^{6, [id](#)}, W. Fan ^{74, [id](#)}, A. Fantoni ^{49, [id](#)}, M. Fasel ^{87, [id](#)}, P. Fecchio ²⁹, A. Feliciello ^{56, [id](#)},
 G. Feofilov ^{140, [id](#)}, A. Fernández Téllez ^{44, [id](#)}, L. Ferrandi ^{110, [id](#)}, M.B. Ferrer ^{32, [id](#)}, A. Ferrero ^{129, [id](#)},
 C. Ferrero ^{56, [id](#), [IV](#)}, A. Ferretti ^{24, [id](#)}, V.J.G. Feuillard ^{94, [id](#)}, V. Filova ^{35, [id](#)}, D. Finogeev ^{140, [id](#)}, F.M. Fionda ^{52, [id](#)},
 F. Flor ^{115, [id](#)}, A.N. Flores ^{108, [id](#)}, S. Foertsch ^{68, [id](#)}, I. Fokin ^{94, [id](#)}, S. Fokin ^{140, [id](#)}, E. Fragiaco ^{57, [id](#)}, E. Frajna ^{46, [id](#)},
 U. Fuchs ^{32, [id](#)}, N. Funicello ^{28, [id](#)}, C. Furget ^{73, [id](#)}, A. Furs ^{140, [id](#)}, T. Fusayasu ^{98, [id](#)}, J.J. Gaardhøje ^{83, [id](#)},
 M. Gagliardi ^{24, [id](#)}, A.M. Gago ^{101, [id](#)}, C.D. Galvan ^{109, [id](#)}, D.R. Gangadharan ^{115, [id](#)}, P. Ganoti ^{78, [id](#)},
 C. Garabatos ^{97, [id](#)}, T. García Chávez ^{44, [id](#)}, E. Garcia-Solis ^{9, [id](#)}, K. Garg ^{103, [id](#)}, C. Gargiulo ^{32, [id](#)}, K. Garner ¹²⁵,
 P. Gasik ^{97, [id](#)}, A. Gautam ^{117, [id](#)}, M.B. Gay Ducati ^{66, [id](#)}, M. Germain ^{103, [id](#)}, C. Ghosh ¹³⁴, M. Giacalone ^{25, [id](#)},
 P. Giubellino ^{97, [id](#), [56, \[id\]\(#\)](#)}, P. Giubilato ^{27, [id](#)}, A.M.C. Glaenger ^{129, [id](#)}, P. Glässel ^{94, [id](#)}, E. Glimos ^{121, [id](#)}, D.J.Q. Goh ⁷⁶,
 V. Gonzalez ^{136, [id](#)}, L.H. González-Trueba ^{67, [id](#)}, M. Gorgon ^{2, [id](#)}, S. Gotovac ³³, V. Grabski ^{67, [id](#)},
 L.K. Graczykowski ^{135, [id](#)}, E. Grecka ^{86, [id](#)}, A. Grelli ^{59, [id](#)}, C. Grigoras ^{32, [id](#)}, V. Grigoriev ^{140, [id](#)}, S. Grigoryan ^{141, [id](#), [1](#)},
 F. Grosa ^{32, [id](#)}, J.F. Grosse-Oetringhaus ^{32, [id](#)}, R. Grosso ^{97, [id](#)}, D. Grund ^{35, [id](#)}, G.G. Guardiano ^{111, [id](#)},
 R. Guernane ^{73, [id](#)}, M. Guilbaud ^{103, [id](#)}, K. Gulbrandsen ^{83, [id](#)}, T. Gündem ^{64, [id](#)}, T. Gunji ^{123, [id](#)}, W. Guo ^{6, [id](#)},
 A. Gupta ^{91, [id](#)}, R. Gupta ^{91, [id](#)}, L. Gyulai ^{46, [id](#)}, M.K. Habib ⁹⁷, C. Hadjidakis ^{130, [id](#)}, F.U. Haider ^{91, [id](#)},
 H. Hamagaki ^{76, [id](#)}, A. Hamdi ^{74, [id](#)}, M. Hamid ⁶, Y. Han ^{138, [id](#)}, R. Hannigan ^{108, [id](#)}, M.R. Haque ^{135, [id](#)},
 J.W. Harris ^{137, [id](#)}, A. Harton ^{9, [id](#)}, H. Hassan ^{87, [id](#)}, D. Hatzifotiadou ^{51, [id](#)}, P. Hauer ^{42, [id](#)}, L.B. Havener ^{137, [id](#)},
 S.T. Heckel ^{95, [id](#)}, E. Hellbär ^{97, [id](#)}, H. Helstrup ^{34, [id](#)}, M. Hemmer ^{64, [id](#)}, T. Herman ^{35, [id](#)}, G. Herrera Corral ^{8, [id](#)},
 F. Herrmann ¹²⁵, S. Herrmann ^{127, [id](#)}, K.F. Hetland ^{34, [id](#)}, B. Heybeck ^{64, [id](#)}, H. Hillemanns ^{32, [id](#)}, C. Hills ^{118, [id](#)},
 B. Hippolyte ^{128, [id](#)}, B. Hofman ^{59, [id](#)}, B. Hohlweger ^{84, [id](#)}, G.H. Hong ^{138, [id](#)}, M. Horst ^{95, [id](#)}, A. Horzyk ^{2, [id](#)},
 R. Hosokawa ¹⁴, Y. Hou ^{6, [id](#)}, P. Hristov ^{32, [id](#)}, C. Hughes ^{121, [id](#)}, P. Huhn ⁶⁴, L.M. Huhta ^{116, [id](#)}, T.J. Humanic ^{88, [id](#)},
 A. Hutson ^{115, [id](#)}, D. Hutter ^{38, [id](#)}, J.P. Iddon ^{118, [id](#)}, R. Ilkaev ¹⁴⁰, H. Ilyas ^{13, [id](#)}, M. Inaba ^{124, [id](#)}, G.M. Innocenti ^{32, [id](#)},
 M. Ippolitov ^{140, [id](#)}, A. Isakov ^{86, [id](#)}, T. Isidori ^{117, [id](#)}, M.S. Islam ^{99, [id](#)}, M. Ivanov ^{97, [id](#)}, M. Ivanov ¹², V. Ivanov ^{140, [id](#)},
 M. Jablonski ^{2, [id](#)}, B. Jacak ^{74, [id](#)}, N. Jacazio ^{32, [id](#)}, P.M. Jacobs ^{74, [id](#)}, S. Jadlovska ¹⁰⁶, J. Jadlovsky ¹⁰⁶,
 S. Jaelani ^{82, [id](#)}, L. Jaffe ³⁸, C. Jahnke ^{111, [id](#)}, M.J. Jakubowska ^{135, [id](#)}, M.A. Janik ^{135, [id](#)}, T. Janson ⁷⁰, M. Jercic ⁸⁹,
 S. Jia ^{10, [id](#)}, A.A.P. Jimenez ^{65, [id](#)}, F. Jonas ^{87, [id](#), [125, \[id\]\(#\)](#)}, J.M. Jowett ^{32, [id](#), [97, \[id\]\(#\)](#)}, J. Jung ^{64, [id](#)}, M. Jung ^{64, [id](#)}, A. Junique ^{32, [id](#)},
 A. Jusko ^{100, [id](#)}, J. Kaewjai ¹⁰⁵, P. Kalinak ^{60, [id](#)}, A.S. Kalteyer ^{97, [id](#)}, A. Kalweit ^{32, [id](#)}, V. Kaplin ^{140, [id](#)}, A. Karasu
 Uysal ^{72, [id](#), [V](#)}, D. Karatovic ^{89, [id](#)}, O. Karavichev ^{140, [id](#)}, T. Karavicheva ^{140, [id](#)}, P. Karczmarczyk ^{135, [id](#)},
 E. Karpechev ^{140, [id](#)}, M.J. Karwowska ^{32, [id](#), [135, \[id\]\(#\)](#)}, U. Keschull ^{70, [id](#)}, R. Keidel ^{139, [id](#)}, D.L.D. Keijdener ⁵⁹, M. Keil ^{32, [id](#)},
 B. Ketzer ^{42, [id](#)}, A.M. Khan ^{6, [id](#)}, S. Khan ^{15, [id](#)}, A. Khanzadeev ^{140, [id](#)}, Y. Kharlov ^{140, [id](#)}, A. Khatun ^{117, [id](#), [15, \[id\]\(#\)](#)},
 A. Khuntia ^{107, [id](#)}, M.B. Kidson ¹¹⁴, B. Kileng ^{34, [id](#)}, B. Kim ^{16, [id](#)}, C. Kim ^{16, [id](#)}, D.J. Kim ^{116, [id](#)}, E.J. Kim ^{69, [id](#)},
 J. Kim ^{138, [id](#)}, J.S. Kim ^{40, [id](#)}, J. Kim ^{94, [id](#)}, J. Kim ^{69, [id](#)}, M. Kim ^{18, [id](#), [94, \[id\]\(#\)](#)}, S. Kim ^{17, [id](#)}, T. Kim ^{138, [id](#)}, K. Kimura ^{92, [id](#)},
 S. Kirsch ^{64, [id](#)}, I. Kisel ^{38, [id](#)}, S. Kiselev ^{140, [id](#)}, A. Kisiel ^{135, [id](#)}, J.P. Kitowski ^{2, [id](#)}, J.L. Klay ^{5, [id](#)}, J. Klein ^{32, [id](#)},
 S. Klein ^{74, [id](#)}, C. Klein-Bösing ^{125, [id](#)}, M. Kleiner ^{64, [id](#)}, T. Klemenz ^{95, [id](#)}, A. Kluge ^{32, [id](#)}, A.G. Knospe ^{115, [id](#)},
 C. Kobdaj ^{105, [id](#)}, T. Kollegger ⁹⁷, A. Kondratyev ^{141, [id](#)}, N. Kondratyeva ^{140, [id](#)}, E. Kondratyuk ^{140, [id](#)}, J. König ^{64, [id](#)},
 S.A. Königstorfer ^{95, [id](#)}, P.J. Konopka ^{32, [id](#)}, G. Kornakov ^{135, [id](#)}, M. Korwieser ^{95, [id](#)}, S.D. Koryciak ^{2, [id](#)},
 A. Kotliarov ^{86, [id](#)}, V. Kovalenko ^{140, [id](#)}, M. Kowalski ^{107, [id](#)}, V. Kozuharov ^{36, [id](#)}, I. Králik ^{60, [id](#)}, A. Kravčáková ^{37, [id](#)},
 L. Kreis ⁹⁷, M. Krivda ^{100, [id](#), [60, \[id\]\(#\)](#)}, F. Krizek ^{86, [id](#)}, K. Krizkova Gajdosova ^{35, [id](#)}, M. Kroesen ^{94, [id](#)}, M. Krüger ^{64, [id](#)},
 D.M. Krupova ^{35, [id](#)}, E. Kryshen ^{140, [id](#)}, V. Kučera ^{32, [id](#)}, C. Kuhn ^{128, [id](#)}, P.G. Kuijer ^{84, [id](#)}, T. Kumaoka ¹²⁴,
 D. Kumar ¹³⁴, L. Kumar ^{90, [id](#)}, N. Kumar ⁹⁰, S. Kumar ^{31, [id](#)}, S. Kundu ^{32, [id](#)}, P. Kurashvili ^{79, [id](#)}, A. Kurepin ^{140, [id](#)},
 A.B. Kurepin ^{140, [id](#)}, A. Kuryakin ^{140, [id](#)}, S. Kushpil ^{86, [id](#)}, J. Kvapil ^{100, [id](#)}, M.J. Kweon ^{58, [id](#)}, J.Y. Kwon ^{58, [id](#)},
 Y. Kwon ^{138, [id](#)}, S.L. La Pointe ^{38, [id](#)}, P. La Rocca ^{26, [id](#)}, Y.S. Lai ⁷⁴, A. Lakrathok ¹⁰⁵, M. Lamanna ^{32, [id](#)},
 R. Langoy ^{120, [id](#)}, P. Larionov ^{32, [id](#)}, E. Laudi ^{32, [id](#)}, L. Lautner ^{32, [id](#), [95, \[id\]\(#\)](#)}, R. Lavicka ^{102, [id](#)}, T. Lazareva ^{140, [id](#)},

R. Lea ^{133,55, [id](#)}, H. Lee ^{104, [id](#)}, G. Legras ^{125, [id](#)}, J. Lehrbach ^{38, [id](#)}, R.C. Lemmon ^{85, [id](#)}, I. León Monzón ^{109, [id](#)}, M.M. Lesch ^{95, [id](#)}, E.D. Lesser ^{18, [id](#)}, M. Lettrich ⁹⁵, P. Lévai ^{46, [id](#)}, X. Li ¹⁰, X.L. Li ⁶, J. Lien ^{120, [id](#)}, R. Lietava ^{100, [id](#)}, B. Lim ^{24,16, [id](#)}, S.H. Lim ^{16, [id](#)}, V. Lindenstruth ^{38, [id](#)}, A. Lindner ⁴⁵, C. Lippmann ^{97, [id](#)}, A. Liu ^{18, [id](#)}, D.H. Liu ^{6, [id](#)}, J. Liu ^{118, [id](#)}, I.M. Lofnes ^{20, [id](#)}, C. Loizides ^{87, [id](#)}, S. Lokos ^{107, [id](#)}, J. Lömker ^{59, [id](#)}, P. Loncar ^{33, [id](#)}, J.A. Lopez ^{94, [id](#)}, X. Lopez ^{126, [id](#)}, E. López Torres ^{7, [id](#)}, P. Lu ^{97,119, [id](#)}, J.R. Luhder ^{125, [id](#)}, M. Lunardon ^{27, [id](#)}, G. Luparello ^{57, [id](#)}, Y.G. Ma ^{39, [id](#)}, A. Maevskaya ¹⁴⁰, M. Mager ^{32, [id](#)}, T. Mahmoud ⁴², A. Maire ^{128, [id](#)}, M.V. Makariev ^{36, [id](#)}, M. Malaev ^{140, [id](#)}, G. Malfattore ^{25, [id](#)}, N.M. Malik ^{91, [id](#)}, Q.W. Malik ¹⁹, S.K. Malik ^{91, [id](#)}, L. Malinina ^{141, [id](#), I,VIII}, D. Mal'Kevich ^{140, [id](#)}, D. Mallick ^{80, [id](#)}, N. Mallick ^{48, [id](#)}, G. Mandaglio ^{30,53, [id](#)}, V. Manko ^{140, [id](#)}, F. Manso ^{126, [id](#)}, V. Manzari ^{50, [id](#)}, Y. Mao ^{6, [id](#)}, G.V. Margagliotti ^{23, [id](#)}, A. Margotti ^{51, [id](#)}, A. Marín ^{97, [id](#)}, C. Markert ^{108, [id](#)}, P. Martinengo ^{32, [id](#)}, J.L. Martinez ¹¹⁵, M.I. Martínez ^{44, [id](#)}, G. Martínez García ^{103, [id](#)}, S. Masciocchi ^{97, [id](#)}, M. Maserà ^{24, [id](#)}, A. Masoni ^{52, [id](#)}, L. Massacrier ^{130, [id](#)}, A. Mastroserio ^{131,50, [id](#)}, O. Matonoha ^{75, [id](#)}, P.F.T. Matuoka ¹¹⁰, A. Matyja ^{107, [id](#)}, C. Mayer ^{107, [id](#)}, A.L. Mazuecos ^{32, [id](#)}, F. Mazzaschi ^{24, [id](#)}, M. Mazzilli ^{32, [id](#)}, J.E. Mdhului ^{122, [id](#)}, A.F. Mechler ⁶⁴, Y. Melikyan ^{43,140, [id](#)}, A. Menchaca-Rocha ^{67, [id](#)}, E. Meninno ^{102, [id](#)}, A.S. Menon ^{115, [id](#)}, M. Meres ^{12, [id](#)}, S. Mhlanga ^{114,68}, Y. Miake ¹²⁴, L. Micheletti ^{56, [id](#)}, L.C. Migliorin ¹²⁷, D.L. Mihaylov ^{95, [id](#)}, K. Mikhaylov ^{141,140, [id](#)}, A.N. Mishra ^{46, [id](#)}, D. Miśkowiec ^{97, [id](#)}, A. Modak ^{4, [id](#)}, A.P. Mohanty ^{59, [id](#)}, B. Mohanty ⁸⁰, M. Mohisin Khan ^{15, [id](#), VI}, M.A. Molander ^{43, [id](#)}, Z. Moravcova ^{83, [id](#)}, C. Mordasini ^{95, [id](#)}, D.A. Moreira De Godoy ^{125, [id](#)}, I. Morozov ^{140, [id](#)}, A. Morsch ^{32, [id](#)}, T. Mrnjavac ^{32, [id](#)}, V. Muccifora ^{49, [id](#)}, S. Muhuri ^{134, [id](#)}, J.D. Mulligan ^{74, [id](#)}, A. Mulliri ^{22, [id](#)}, M.G. Munhoz ^{110, [id](#)}, R.H. Munzer ^{64, [id](#)}, H. Murakami ^{123, [id](#)}, S. Murray ^{114, [id](#)}, L. Musa ^{32, [id](#)}, J. Musinsky ^{60, [id](#)}, J.W. Myrcha ^{135, [id](#)}, B. Naik ^{122, [id](#)}, A.I. Nambrath ^{18, [id](#)}, B.K. Nandi ^{47, [id](#)}, R. Nania ^{51, [id](#)}, E. Nappi ^{50, [id](#)}, A.F. Nassirpour ^{75, [id](#)}, A. Nath ^{94, [id](#)}, C. Nattrass ^{121, [id](#)}, M.N. Naydenov ^{36, [id](#)}, A. Neagu ¹⁹, A. Negru ¹¹³, L. Nellen ^{65, [id](#)}, S.V. Nesbo ³⁴, G. Neskovic ^{38, [id](#)}, D. Nesterov ^{140, [id](#)}, B.S. Nielsen ^{83, [id](#)}, E.G. Nielsen ^{83, [id](#)}, S. Nikolaev ^{140, [id](#)}, S. Nikulin ^{140, [id](#)}, V. Nikulin ^{140, [id](#)}, F. Noferini ^{51, [id](#)}, S. Noh ^{11, [id](#)}, P. Nomokonov ^{141, [id](#)}, J. Norman ^{118, [id](#)}, N. Novitzky ^{124, [id](#)}, P. Nowakowski ^{135, [id](#)}, A. Nyanin ^{140, [id](#)}, J. Nystrand ^{20, [id](#)}, M. Ogino ^{76, [id](#)}, A. Ohlson ^{75, [id](#)}, V.A. Okorokov ^{140, [id](#)}, J. Oleniacz ^{135, [id](#)}, A.C. Oliveira Da Silva ^{121, [id](#)}, M.H. Oliver ^{137, [id](#)}, A. Onnerstad ^{116, [id](#)}, C. Oppedisano ^{56, [id](#)}, A. Ortiz Velasquez ^{65, [id](#)}, J. Otwinowski ^{107, [id](#)}, M. Oya ⁹², K. Oyama ^{76, [id](#)}, Y. Pachmayer ^{94, [id](#)}, S. Padhan ^{47, [id](#)}, D. Pagano ^{133,55, [id](#)}, G. Paić ^{65, [id](#)}, S. Paisano-Guzmán ^{44, [id](#)}, A. Palasciano ^{50, [id](#)}, S. Panebianco ^{129, [id](#)}, H. Park ^{124, [id](#)}, H. Park ^{104, [id](#)}, J. Park ^{58, [id](#)}, J.E. Parkkila ^{32, [id](#)}, R.N. Patra ⁹¹, B. Paul ^{22, [id](#)}, H. Pei ^{6, [id](#)}, T. Peitzmann ^{59, [id](#)}, X. Peng ^{6, [id](#)}, M. Pennisi ^{24, [id](#)}, L.G. Pereira ^{66, [id](#)}, D. Peresunko ^{140, [id](#)}, G.M. Perez ^{7, [id](#)}, S. Perrin ^{129, [id](#)}, Y. Pestov ¹⁴⁰, V. Petráček ^{35, [id](#)}, V. Petrov ^{140, [id](#)}, M. Petrovici ^{45, [id](#)}, R.P. Pezzi ^{103,66, [id](#)}, S. Piano ^{57, [id](#)}, M. Pikna ^{12, [id](#)}, P. Pillot ^{103, [id](#)}, O. Pinazza ^{51,32, [id](#)}, L. Pinsky ¹¹⁵, C. Pinto ^{95, [id](#)}, S. Pisano ^{49, [id](#)}, M. Płoskoń ^{74, [id](#)}, M. Planinic ⁸⁹, F. Pliquett ⁶⁴, M.G. Poghosyan ^{87, [id](#)}, B. Polichtchouk ^{140, [id](#)}, S. Politano ^{29, [id](#)}, N. Poljak ^{89, [id](#)}, A. Pop ^{45, [id](#)}, S. Porteboeuf-Houssais ^{126, [id](#)}, V. Pozdniakov ^{141, [id](#)}, K.K. Pradhan ^{48, [id](#)}, S.K. Prasad ^{4, [id](#)}, S. Prasad ^{48, [id](#)}, R. Preghenella ^{51, [id](#)}, F. Prino ^{56, [id](#)}, C.A. Pruneau ^{136, [id](#)}, I. Pshenichnov ^{140, [id](#)}, M. Puccio ^{32, [id](#)}, S. Pucillo ^{24, [id](#)}, Z. Pugelova ¹⁰⁶, S. Qiu ^{84, [id](#)}, L. Quaglia ^{24, [id](#)}, R.E. Quishpe ¹¹⁵, S. Ragoni ^{14,100, [id](#)}, A. Rakotozafindrabe ^{129, [id](#)}, L. Ramello ^{132,56, [id](#)}, F. Rami ^{128, [id](#)}, T.A. Rancien ⁷³, M. Rasa ^{26, [id](#)}, S.S. Räsänen ^{43, [id](#)}, R. Rath ^{51, [id](#)}, M.P. Rauch ^{20, [id](#)}, I. Ravasenga ^{84, [id](#)}, K.F. Read ^{87,121, [id](#)}, C. Reckziegel ^{112, [id](#)}, A.R. Redelbach ^{38, [id](#)}, K. Redlich ^{79, [id](#), VII}, C.A. Reetz ^{97, [id](#)}, H.D. Regules-Medel ⁴⁴, A. Rehman ²⁰, F. Reidt ^{32, [id](#)}, H.A. Reme-Ness ^{34, [id](#)}, Z. Rescakova ³⁷, K. Reygers ^{94, [id](#)}, A. Riabov ^{140, [id](#)}, V. Riabov ^{140, [id](#)}, R. Ricci ^{28, [id](#)}, M. Richter ^{19, [id](#)}, A.A. Riedel ^{95, [id](#)}, W. Riegler ^{32, [id](#)}, C. Ristea ^{63, [id](#)}, M. Rodríguez Cahuantzi ^{44, [id](#)}, S.A. Rodríguez Ramírez ^{44, [id](#)}, K. Røed ^{19, [id](#)}, R. Rogalev ^{140, [id](#)}, E. Rogochaya ^{141, [id](#)}, T.S. Rogoschinski ^{64, [id](#)}, D. Rohr ^{32, [id](#)}, D. Röhrich ^{20, [id](#)}, P.F. Rojas ⁴⁴, S. Rojas Torres ^{35, [id](#)}, P.S. Rokita ^{135, [id](#)}, G. Romanenko ^{141, [id](#)}, F. Ronchetti ^{49, [id](#)}, A. Rosano ^{30,53, [id](#)},

E.D. Rosas⁶⁵, K. Roslon^{135, ^{id}}, A. Rossi^{54, ^{id}}, A. Roy^{48, ^{id}}, S. Roy^{47, ^{id}}, N. Rubini^{25, ^{id}}, D. Ruggiano^{135, ^{id}}, R. Rui^{23, ^{id}}, B. Rumyantsev¹⁴¹, P.G. Russek^{2, ^{id}}, R. Russo^{84, ^{id}}, A. Rustamov^{81, ^{id}}, E. Ryabinkin^{140, ^{id}}, Y. Ryabov^{140, ^{id}}, A. Rybicki^{107, ^{id}}, H. Ryttonen^{116, ^{id}}, W. Rzesza^{135, ^{id}}, O.A.M. Saarimaki^{43, ^{id}}, R. Sadek^{103, ^{id}}, S. Sadhu^{31, ^{id}}, S. Sadovsky^{140, ^{id}}, J. Saetre^{20, ^{id}}, K. Šafařík^{35, ^{id}}, S.K. Saha^{4, ^{id}}, S. Saha^{80, ^{id}}, B. Sahoo^{47, ^{id}}, R. Sahoo^{48, ^{id}}, S. Sahoo⁶¹, D. Sahu^{48, ^{id}}, P.K. Sahu^{61, ^{id}}, J. Saini^{134, ^{id}}, K. Sajdakova³⁷, S. Sakai^{124, ^{id}}, M.P. Salvan^{97, ^{id}}, S. Sambyal^{91, ^{id}}, I. Sanna^{32,95, ^{id}}, T.B. Saramela¹¹⁰, D. Sarkar^{136, ^{id}}, N. Sarkar¹³⁴, P. Sarma^{41, ^{id}}, V. Sarritzu^{22, ^{id}}, V.M. Sarti^{95, ^{id}}, M.H.P. Sas^{137, ^{id}}, J. Schambach^{87, ^{id}}, H.S. Scheid^{64, ^{id}}, C. Schiaua^{45, ^{id}}, R. Schicker^{94, ^{id}}, A. Schmah⁹⁴, C. Schmidt^{97, ^{id}}, H.R. Schmidt⁹³, M.O. Schmidt^{32, ^{id}}, M. Schmidt⁹³, N.V. Schmidt^{87, ^{id}}, A.R. Schmier^{121, ^{id}}, R. Schotter^{128, ^{id}}, A. Schröter^{38, ^{id}}, J. Schukraft^{32, ^{id}}, K. Schwarz⁹⁷, K. Schweda^{97, ^{id}}, G. Scioli^{25, ^{id}}, E. Scomparin^{56, ^{id}}, J.E. Seger^{14, ^{id}}, Y. Sekiguchi¹²³, D. Sekihata^{123, ^{id}}, I. Selyuzhenkov^{97,140, ^{id}}, S. Senyukov^{128, ^{id}}, J.J. Seo^{58, ^{id}}, D. Serebryakov^{140, ^{id}}, L. Šerkšnytė^{95, ^{id}}, A. Sevcenco^{63, ^{id}}, T.J. Shaba^{68, ^{id}}, A. Shabetai^{103, ^{id}}, R. Shahoyan³², A. Shangaraev^{140, ^{id}}, A. Sharma⁹⁰, B. Sharma^{91, ^{id}}, D. Sharma^{47, ^{id}}, H. Sharma^{107, ^{id}}, M. Sharma^{91, ^{id}}, S. Sharma^{76, ^{id}}, S. Sharma^{91, ^{id}}, U. Sharma^{91, ^{id}}, A. Shatat^{130, ^{id}}, O. Sheibani¹¹⁵, K. Shigaki^{92, ^{id}}, M. Shimomura⁷⁷, J. Shin¹¹, S. Shirinkin^{140, ^{id}}, Q. Shou^{39, ^{id}}, Y. Sibiriak^{140, ^{id}}, S. Siddhanta^{52, ^{id}}, T. Siemiarczuk^{79, ^{id}}, T.F. Silva^{110, ^{id}}, D. Silvermyr^{75, ^{id}}, T. Simantathammakul¹⁰⁵, R. Simeonov^{36, ^{id}}, B. Singh⁹¹, B. Singh^{95, ^{id}}, R. Singh^{80, ^{id}}, R. Singh^{91, ^{id}}, R. Singh^{48, ^{id}}, S. Singh^{15, ^{id}}, V.K. Singh^{134, ^{id}}, V. Singhal^{134, ^{id}}, T. Sinha^{99, ^{id}}, B. Sitar^{12, ^{id}}, M. Sitta^{132,56, ^{id}}, T.B. Skaali¹⁹, G. Skorodumovs^{94, ^{id}}, M. Slupecki^{43, ^{id}}, N. Smirnov^{137, ^{id}}, R.J.M. Snellings^{59, ^{id}}, E.H. Solheim^{19, ^{id}}, J. Song^{115, ^{id}}, A. Songmoolnak¹⁰⁵, F. Soramel^{27, ^{id}}, R. Spijkers^{84, ^{id}}, I. Sputowska^{107, ^{id}}, J. Staa^{75, ^{id}}, J. Stachel^{94, ^{id}}, I. Stan^{63, ^{id}}, P.J. Steffanic^{121, ^{id}}, S.F. Stiefelmaier^{94, ^{id}}, D. Stocco^{103, ^{id}}, I. Storehaug^{19, ^{id}}, P. Stratmann^{125, ^{id}}, S. Strazzi^{25, ^{id}}, C.P. Stylianidis⁸⁴, A.A.P. Suaide^{110, ^{id}}, C. Suire^{130, ^{id}}, M. Sukhanov^{140, ^{id}}, M. Suljic^{32, ^{id}}, R. Sultanov^{140, ^{id}}, V. Sumberia^{91, ^{id}}, S. Sumowidagdo^{82, ^{id}}, S. Swain⁶¹, I. Szarka^{12, ^{id}}, S.F. Taghavi^{95, ^{id}}, G. TAILLEPIED^{97, ^{id}}, J. Takahashi^{111, ^{id}}, G.J. Tambave^{20, ^{id}}, S. Tang^{126,6, ^{id}}, Z. Tang^{119, ^{id}}, J.D. Tapia Takaki^{117, ^{id}}, N. Tapus¹¹³, L.A. Tarasovicova^{125, ^{id}}, M.G. Tarzila^{45, ^{id}}, G.F. Tassielli^{31, ^{id}}, A. Tauro^{32, ^{id}}, G. Tejada Muñoz^{44, ^{id}}, A. Telesca^{32, ^{id}}, L. Terlizzi^{24, ^{id}}, C. Terrevoli^{115, ^{id}}, G. Tersimonov³, S. Thakur^{4, ^{id}}, D. Thomas^{108, ^{id}}, A. Tikhonov^{140, ^{id}}, A.R. Timmins^{115, ^{id}}, M. Tkacik¹⁰⁶, T. Tkacik^{106, ^{id}}, A. Toia^{64, ^{id}}, R. Tokumoto⁹², N. Topilskaya^{140, ^{id}}, M. Toppi^{49, ^{id}}, F. Torres-Acosta¹⁸, T. Tork^{130, ^{id}}, A.G. Torres Ramos^{31, ^{id}}, A. Trifiró^{30,53, ^{id}}, A.S. Triolo^{30,53, ^{id}}, S. Tripathy^{51, ^{id}}, T. Tripathy^{47, ^{id}}, S. Trogolo^{32, ^{id}}, V. Trubnikov^{3, ^{id}}, W.H. Trzaska^{116, ^{id}}, T.P. Trzcinski^{135, ^{id}}, A. Tumkin^{140, ^{id}}, R. Turrisi^{54, ^{id}}, T.S. Tveter^{19, ^{id}}, K. Ullaland^{20, ^{id}}, B. Ulukutlu^{95, ^{id}}, A. Uras^{127, ^{id}}, M. Urioni^{55,133, ^{id}}, G.L. Usai^{22, ^{id}}, M. Vala³⁷, N. Valle^{21, ^{id}}, L.V.R. van Doremalen⁵⁹, C. Van Hulse^{130, ^{id}}, M. van Leeuwen^{84, ^{id}}, C.A. van Veen^{94, ^{id}}, R.J.G. van Weelden^{84, ^{id}}, P. Vande Vyvre^{32, ^{id}}, D. Varga^{46, ^{id}}, Z. Varga^{46, ^{id}}, M. Vasileiou^{78, ^{id}}, A. Vasiliev^{140, ^{id}}, O. Vázquez Doce^{49, ^{id}}, O. Vazquez Rueda^{115,75, ^{id}}, V. Vechernin^{140, ^{id}}, E. Vercellin^{24, ^{id}}, S. Vergara Limón⁴⁴, L. Vermunt^{97, ^{id}}, R. Vértesi^{46, ^{id}}, M. Verweij^{59, ^{id}}, L. Vickovic³³, Z. Vilakazi¹²², O. Villalobos Baillie^{100, ^{id}}, G. VINO^{50, ^{id}}, A. Vinogradov^{140, ^{id}}, T. Virgili^{28, ^{id}}, V. Vislavicius⁸³, A. Vodopyanov^{141, ^{id}}, B. Volkel^{32, ^{id}}, M.A. Völkl^{94, ^{id}}, K. Voloshin¹⁴⁰, S.A. Voloshin^{136, ^{id}}, G. Volpe^{31, ^{id}}, B. von Haller^{32, ^{id}}, I. Vorobyev^{95, ^{id}}, N. Vozniuk^{140, ^{id}}, J. Vrláková^{37, ^{id}}, C. Wang^{39, ^{id}}, D. Wang³⁹, Y. Wang^{39, ^{id}}, A. Wegrzynek^{32, ^{id}}, F.T. Weiglhofer³⁸, S.C. Wenzel^{32, ^{id}}, J.P. Wessels^{125, ^{id}}, J. Wiechula^{64, ^{id}}, J. Wikne^{19, ^{id}}, G. Wilk^{79, ^{id}}, J. Wilkinson^{97, ^{id}}, G.A. Willems^{125, ^{id}}, B. Windelband^{94, ^{id}}, M. Winn^{129, ^{id}}, J.R. Wright^{108, ^{id}}, W. Wu³⁹, Y. Wu^{119, ^{id}}, R. Xu^{6, ^{id}}, A. Yadav^{42, ^{id}}, A.K. Yadav^{134, ^{id}}, S. Yalcin^{72, ^{id}}, Y. Yamaguchi^{92, ^{id}}, S. Yang²⁰, S. Yano^{92, ^{id}}, Z. Yin^{6, ^{id}}, I.-K. Yoo^{16, ^{id}}, J.H. Yoon^{58, ^{id}}, S. Yuan²⁰, A. Yuncu^{94, ^{id}}, V. Zaccolo^{23, ^{id}}, C. Zampolli^{32, ^{id}}, F. Zanone^{94, ^{id}}, N. Zardoshti^{32,100, ^{id}}, A. Zarochentsev^{140, ^{id}}, P. Závada^{62, ^{id}}, N. Zaviyalov¹⁴⁰,

M. Zhalov^{140, [id](#)}, B. Zhang^{6, [id](#)}, L. Zhang^{39, [id](#)}, M. Zhang⁶, S. Zhang^{39, [id](#)}, X. Zhang^{6, [id](#)}, Y. Zhang¹¹⁹,
 Z. Zhang^{6, [id](#)}, M. Zhao^{10, [id](#)}, V. Zhrebchevskii^{140, [id](#)}, Y. Zhi¹⁰, D. Zhou^{6, [id](#)}, Y. Zhou^{83, [id](#)}, J. Zhu^{97,6, [id](#)}, Y. Zhu⁶,
 S.C. Zugravel^{56, [id](#)}, N. Zurlo^{133,55, [id](#)}

¹ A.I. Alikhanyan National Science Laboratory (Yerevan Physics Institute) Foundation, Yerevan, Armenia

² AGH University of Krakow, Cracow, Poland

³ Bogolyubov Institute for Theoretical Physics, National Academy of Sciences of Ukraine, Kiev, Ukraine

⁴ Bose Institute, Department of Physics and Centre for Astroparticle Physics and Space Science (CAPSS), Kolkata, India

⁵ California Polytechnic State University, San Luis Obispo, CA, United States

⁶ Central China Normal University, Wuhan, China

⁷ Centro de Aplicaciones Tecnológicas y Desarrollo Nuclear (CEADEN), Havana, Cuba

⁸ Centro de Investigación y de Estudios Avanzados (CINVESTAV), Mexico City and Mérida, Mexico

⁹ Chicago State University, Chicago, IL, United States

¹⁰ China Institute of Atomic Energy, Beijing, China

¹¹ Chungbuk National University, Cheongju, Republic of Korea

¹² Comenius University Bratislava, Faculty of Mathematics, Physics and Informatics, Bratislava, Slovak Republic

¹³ COMSATS University Islamabad, Islamabad, Pakistan

¹⁴ Creighton University, Omaha, NE, United States

¹⁵ Department of Physics, Aligarh Muslim University, Aligarh, India

¹⁶ Department of Physics, Pusan National University, Pusan, Republic of Korea

¹⁷ Department of Physics, Sejong University, Seoul, Republic of Korea

¹⁸ Department of Physics, University of California, Berkeley, CA, United States

¹⁹ Department of Physics, University of Oslo, Oslo, Norway

²⁰ Department of Physics and Technology, University of Bergen, Bergen, Norway

²¹ Dipartimento di Fisica, Università di Pavia, Pavia, Italy

²² Dipartimento di Fisica dell'Università and Sezione INFN, Cagliari, Italy

²³ Dipartimento di Fisica dell'Università and Sezione INFN, Trieste, Italy

²⁴ Dipartimento di Fisica dell'Università and Sezione INFN, Turin, Italy

²⁵ Dipartimento di Fisica e Astronomia dell'Università and Sezione INFN, Bologna, Italy

²⁶ Dipartimento di Fisica e Astronomia dell'Università and Sezione INFN, Catania, Italy

²⁷ Dipartimento di Fisica e Astronomia dell'Università and Sezione INFN, Padova, Italy

²⁸ Dipartimento di Fisica 'E.R. Caianiello' dell'Università and Gruppo Collegato INFN, Salerno, Italy

²⁹ Dipartimento DISAT del Politecnico and Sezione INFN, Turin, Italy

³⁰ Dipartimento di Scienze MIFT, Università di Messina, Messina, Italy

³¹ Dipartimento Interateneo di Fisica 'M. Merlin' and Sezione INFN, Bari, Italy

³² European Organization for Nuclear Research (CERN), Geneva, Switzerland

³³ Faculty of Electrical Engineering, Mechanical Engineering and Naval Architecture, University of Split, Split, Croatia

³⁴ Faculty of Engineering and Science, Western Norway University of Applied Sciences, Bergen, Norway

³⁵ Faculty of Nuclear Sciences and Physical Engineering, Czech Technical University in Prague, Prague, Czech Republic

³⁶ Faculty of Physics, Sofia University, Sofia, Bulgaria

³⁷ Faculty of Science, P.J. Šafárik University, Košice, Slovak Republic

³⁸ Frankfurt Institute for Advanced Studies, Johann Wolfgang Goethe-Universität Frankfurt, Frankfurt, Germany

³⁹ Fudan University, Shanghai, China

⁴⁰ Gangneung-Wonju National University, Gangneung, Republic of Korea

⁴¹ Gauhati University, Department of Physics, Guwahati, India

⁴² Helmholtz-Institut für Strahlen- und Kernphysik, Rheinische Friedrich-Wilhelms-Universität Bonn, Bonn, Germany

⁴³ Helsinki Institute of Physics (HIP), Helsinki, Finland

⁴⁴ High Energy Physics Group, Universidad Autónoma de Puebla, Puebla, Mexico

⁴⁵ Horia Hulubei National Institute of Physics and Nuclear Engineering, Bucharest, Romania

⁴⁶ HUN-REN Wigner Research Centre for Physics, Budapest, Hungary

⁴⁷ Indian Institute of Technology Bombay (IIT), Mumbai, India

⁴⁸ Indian Institute of Technology Indore, Indore, India

⁴⁹ INFN, Laboratori Nazionali di Frascati, Frascati, Italy

⁵⁰ INFN, Sezione di Bari, Bari, Italy

⁵¹ INFN, Sezione di Bologna, Bologna, Italy

⁵² INFN, Sezione di Cagliari, Cagliari, Italy

⁵³ INFN, Sezione di Catania, Catania, Italy

⁵⁴ INFN, Sezione di Padova, Padova, Italy

⁵⁵ INFN, Sezione di Pavia, Pavia, Italy

⁵⁶ INFN, Sezione di Torino, Turin, Italy

⁵⁷ INFN, Sezione di Trieste, Trieste, Italy

⁵⁸ Inha University, Incheon, Republic of Korea

⁵⁹ Institute for Gravitational and Subatomic Physics (GRASP), Utrecht University/Nikhef, Utrecht, Netherlands

⁶⁰ Institute of Experimental Physics, Slovak Academy of Sciences, Košice, Slovak Republic

⁶¹ Institute of Physics, Homi Bhabha National Institute, Bhubaneswar, India

⁶² Institute of Physics of the Czech Academy of Sciences, Prague, Czech Republic

⁶³ Institute of Space Science (ISS), Bucharest, Romania

⁶⁴ Institut für Kernphysik, Johann Wolfgang Goethe-Universität Frankfurt, Frankfurt, Germany

⁶⁵ Instituto de Ciencias Nucleares, Universidad Nacional Autónoma de México, Mexico City, Mexico

⁶⁶ Instituto de Física, Universidade Federal do Rio Grande do Sul (UFRGS), Porto Alegre, Brazil

⁶⁷ Instituto de Física, Universidad Nacional Autónoma de México, Mexico City, Mexico

⁶⁸ iThemba LABS, National Research Foundation, Somerset West, South Africa

⁶⁹ Jeonbuk National University, Jeonju, Republic of Korea

⁷⁰ Johann-Wolfgang-Goethe Universität Frankfurt Institut für Informatik, Fachbereich Informatik und Mathematik, Frankfurt, Germany

⁷¹ Korea Institute of Science and Technology Information, Daejeon, Republic of Korea

⁷² KTO Karatay University, Konya, Turkey

⁷³ Laboratoire de Physique Subatomique et de Cosmologie, Université Grenoble-Alpes, CNRS-IN2P3, Grenoble, France

⁷⁴ Lawrence Berkeley National Laboratory, Berkeley, CA, United States

- ⁷⁵ Lund University Department of Physics, Division of Particle Physics, Lund, Sweden
⁷⁶ Nagasaki Institute of Applied Science, Nagasaki, Japan
⁷⁷ Nara Women's University (NWU), Nara, Japan
⁷⁸ National and Kapodistrian University of Athens, School of Science, Department of Physics, Athens, Greece
⁷⁹ National Centre for Nuclear Research, Warsaw, Poland
⁸⁰ National Institute of Science Education and Research, Homi Bhabha National Institute, Jatni, India
⁸¹ National Nuclear Research Center, Baku, Azerbaijan
⁸² National Research and Innovation Agency – BRIN, Jakarta, Indonesia
⁸³ Niels Bohr Institute, University of Copenhagen, Copenhagen, Denmark
⁸⁴ Nikhef, National institute for subatomic physics, Amsterdam, Netherlands
⁸⁵ Nuclear Physics Group, STFC Daresbury Laboratory, Daresbury, United Kingdom
⁸⁶ Nuclear Physics Institute of the Czech Academy of Sciences, Husinec-Řež, Czech Republic
⁸⁷ Oak Ridge National Laboratory, Oak Ridge, TN, United States
⁸⁸ Ohio State University, Columbus, OH, United States
⁸⁹ Physics department, Faculty of science, University of Zagreb, Zagreb, Croatia
⁹⁰ Physics Department, Panjab University, Chandigarh, India
⁹¹ Physics Department, University of Jammu, Jammu, India
⁹² Physics Program and International Institute for Sustainability with Knotted Chiral Meta Matter (SKCM2), Hiroshima University, Hiroshima, Japan
⁹³ Physikalisches Institut, Eberhard-Karls-Universität Tübingen, Tübingen, Germany
⁹⁴ Physikalisches Institut, Ruprecht-Karls-Universität Heidelberg, Heidelberg, Germany
⁹⁵ Physik Department, Technische Universität München, Munich, Germany
⁹⁶ Politecnico di Bari and Sezione INFN, Bari, Italy
⁹⁷ Research Division and ExtreMe Matter Institute EMMI, GSI Helmholtzzentrum für Schwerionenforschung GmbH, Darmstadt, Germany
⁹⁸ Saga University, Saga, Japan
⁹⁹ Saha Institute of Nuclear Physics, Homi Bhabha National Institute, Kolkata, India
¹⁰⁰ School of Physics and Astronomy, University of Birmingham, Birmingham, United Kingdom
¹⁰¹ Sección Física, Departamento de Ciencias, Pontificia Universidad Católica del Perú, Lima, Peru
¹⁰² Stefan Meyer Institut für Subatomare Physik (SMI), Vienna, Austria
¹⁰³ SUBATECH, IMT Atlantique, Nantes Université, CNRS-IN2P3, Nantes, France
¹⁰⁴ Sungkyunkwan University, Suwon City, Republic of Korea
¹⁰⁵ Suranaree University of Technology, Nakhon Ratchasima, Thailand
¹⁰⁶ Technical University of Košice, Košice, Slovak Republic
¹⁰⁷ The Henryk Niewodniczanski Institute of Nuclear Physics, Polish Academy of Sciences, Cracow, Poland
¹⁰⁸ The University of Texas at Austin, Austin, TX, United States
¹⁰⁹ Universidad Autónoma de Sinaloa, Culiacán, Mexico
¹¹⁰ Universidade de São Paulo (USP), São Paulo, Brazil
¹¹¹ Universidade Estadual de Campinas (UNICAMP), Campinas, Brazil
¹¹² Universidade Federal do ABC, Santo Andre, Brazil
¹¹³ Universitatea Nationala de Stiinta si Tehnologie Politehnica Bucuresti, Bucharest, Romania
¹¹⁴ University of Cape Town, Cape Town, South Africa
¹¹⁵ University of Houston, Houston, TX, United States
¹¹⁶ University of Jyväskylä, Jyväskylä, Finland
¹¹⁷ University of Kansas, Lawrence, KS, United States
¹¹⁸ University of Liverpool, Liverpool, United Kingdom
¹¹⁹ University of Science and Technology of China, Hefei, China
¹²⁰ University of South-Eastern Norway, Kongsberg, Norway
¹²¹ University of Tennessee, Knoxville, TN, United States
¹²² University of the Witwatersrand, Johannesburg, South Africa
¹²³ University of Tokyo, Tokyo, Japan
¹²⁴ University of Tsukuba, Tsukuba, Japan
¹²⁵ Universität Münster, Institut für Kernphysik, Münster, Germany
¹²⁶ Université Clermont Auvergne, CNRS/IN2P3, LPC, Clermont-Ferrand, France
¹²⁷ Université de Lyon, CNRS/IN2P3, Institut de Physique des 2 Infinis de Lyon, Lyon, France
¹²⁸ Université de Strasbourg, CNRS, IPHC UMR 7178, F-67000 Strasbourg, France, Strasbourg, France
¹²⁹ Université Paris-Saclay, Centre d'Etudes de Saclay (CEA), IRFU, Département de Physique Nucléaire (DPhN), Saclay, France
¹³⁰ Université Paris-Saclay, CNRS/IN2P3, IJCLab, Orsay, France
¹³¹ Università degli Studi di Foggia, Foggia, Italy
¹³² Università del Piemonte Orientale, VerCELLI, Italy
¹³³ Università di Brescia, Brescia, Italy
¹³⁴ Variable Energy Cyclotron Centre, Homi Bhabha National Institute, Kolkata, India
¹³⁵ Warsaw University of Technology, Warsaw, Poland
¹³⁶ Wayne State University, Detroit, MI, United States
¹³⁷ Yale University, New Haven, CT, United States
¹³⁸ Yonsei University, Seoul, Republic of Korea
¹³⁹ Zentrum für Technologie und Transfer (ZTT), Worms, Germany
¹⁴⁰ Affiliated with an institute covered by a cooperation agreement with CERN
¹⁴¹ Affiliated with an international laboratory covered by a cooperation agreement with CERN

^I Deceased.

^{II} Also at: Max-Planck-Institut für Physik, Munich, Germany.

^{III} Also at: Italian National Agency for New Technologies, Energy and Sustainable Economic Development (ENEA), Bologna, Italy.

^{IV} Also at: Dipartimento DET del Politecnico di Torino, Turin, Italy.

^V Also at: Yildiz Technical University, Istanbul, Türkiye.

^{VI} Also at: Department of Applied Physics, Aligarh Muslim University, Aligarh, India.

^{VII} Also at: Institute of Theoretical Physics, University of Wrocław, Poland.

^{VIII} Also at: An institution covered by a cooperation agreement with CERN.