

(1975).

<sup>12</sup>M. Lacombe, B. Loiseau, J. M. Richard, R. Vinh Mau, J. Coté, P. Pirès, and R. de Tournreil, Phys. Rev. C 21, 861 (1980).

<sup>13</sup>T. Hamada and I. D. Johnston, Nucl. Phys. 34, 382 (1962).

<sup>14</sup>T. Ueda and A. E. S. Green, Phys. Rev. 174, 1304 (1968).

<sup>15</sup>R. Bryan and B. L. Scott, Phys. Rev. 177, 1435 (1969).

<sup>16</sup>R. de Tournreil and D. W. Sprung, Nucl. Phys. A201, 193 (1973).

<sup>17</sup>I. E. Lagaris and V. R. Pandharipande, Nucl. Phys. A359, 349 (1981).

<sup>18</sup>I. E. Lagaris and V. R. Pandharipande, Nucl. Phys. A359, 331 (1981).

<sup>19</sup>J. L. Friar and J. W. Negele, Adv. Nucl. Phys. 8, 219 (1975).

<sup>20</sup>F. Coester, S. Cohen, B. Day, and C. M. Vincent, Phys. Rev. C 1, 769 (1970).

<sup>21</sup>M. I. Haftel and F. Tabakin, Phys. Rev. C 3, 921 (1971).

<sup>22</sup>I. R. Afnan, D. M. Clement, and F. J. D. Serduke, Nucl. Phys. A170, 625 (1971).

<sup>23</sup>E. L. Lomon, in *Nucleon-Nucleon Interactions—1977*, edited by H. Fearing, D. Measday, and A. Strathlee, AIP Conference Proceedings No. 41 (American Institute of Physics, New York, 1977), p. 554.

<sup>24</sup>J. P. Blaizot, D. Gogny, and B. Grammaticos, Nucl. Phys. A265, 315 (1976).

<sup>25</sup>V. R. Pandharipande, in Proceedings of the International Conference on Recent Progress in Many-Body Theories, Oaxtepec, Mexico, 1981 (to be published).

<sup>26</sup>R. B. Wiringa, to be published.

<sup>27</sup>Y. E. Kim and A. Tubis, Annu. Rev. Nucl. Sci. 24, 69 (1974).

## A Production near Threshold in Central Nucleus-Nucleus Collisions

J. W. Harris, A. Sandoval, R. Stock, and H. Stroebele

*Gesellschaft für Schwerionenforschung, D-6100 Darmstadt, West Germany*

and

R. E. Renfordt

*Philips-Universität Marburg, D-3550 Marburg/Lahn, West Germany*

and

J. V. Geaga,<sup>(a)</sup> H. G. Pugh, and L. S. Schroeder

*Lawrence Berkeley Laboratory, University of California, Berkeley, California 94720*

and

K. L. Wolf

*Argonne National Laboratory, Argonne, Illinois 60439*

and

A. Dacal

*Instituto de Física, University of Mexico, Mexico City 20, Distrito Federal, Mexico*

(Received 23 April 1981)

$\Lambda^1$ s produced in central collisions of  $^{40}\text{Ar} + \text{KCl}$  at 1.8-GeV/u incident energy were detected in a streamer chamber by their charged-particle decay. For central collisions with impact parameters  $b < 2.4$  fm the  $\Lambda$  production cross section is  $7.6 \pm 2.2$  mb. A calculation in which  $\Lambda$  production occurs in the early stage of the collision qualitatively reproduces the results but underestimates the transverse momenta. An average  $\Lambda$  polarization of  $-0.10 \pm 0.05$  is observed.

PACS numbers: 25.70 Bc

In the study of high-energy nucleus-nucleus collisions it is difficult to extract information about the initial stage of the reaction where high baryon densities may occur. Studies of nucleon and cluster emission<sup>1</sup> are consistent with a de-

velopment towards chemical equilibrium in the final stages of the reaction preempting information about the primary stages. In this Letter we report the results of  $\Lambda$  production in central nucleus-nucleus collisions, just above the  $NN$

$\rightarrow \Lambda KN$  threshold.  $\Lambda$  production near threshold is expected to occur either in the first generations of binary nucleon-nucleon collisions or from a localized, collective, high-temperature, high-density system and, in any case, is expected to reflect the first stages of the interaction. Although rescattering of particles in the nuclear medium is expected to be important, the absorption of  $\Lambda$ 's is minimal. Therefore, the study of  $\Lambda$  production would provide a new way to examine the interpenetration phase of nucleus-nucleus collisions.

Strange-baryon production near threshold in nucleus-nucleus collisions is interesting since it provides an opportunity to test the effect of an extended nuclear field on the production mechanism.<sup>2</sup> In the elementary  $NN \rightarrow \Lambda KN$  process, a strange-antistrange quark pair must be produced from the sea.<sup>3</sup> The existence of strange-antistrange quark admixtures<sup>4,5</sup> in the nucleus may enhance the strange-baryon production near threshold.

$\Lambda$  production was studied in central  $^{40}\text{Ar} + \text{KCl}$  collisions at 1.8 GeV/u incident energy using the streamer chamber facility at the Bevalac. The streamer chamber, with its  $4\pi$  solid angle capability, was operated in both inelastic and central trigger modes as previously reported.<sup>6</sup> The central trigger selects events with small total charge in projectile fragments. This trigger, which corresponds to 10% of the reaction cross section ( $\sigma_R = 1.8$  b), is associated with impact




FIG. 1. Results of kinematic fits to the observed neutral strange-particle decays,  $\Lambda \rightarrow p\pi^-$  and  $K^0 \rightarrow \pi^+\pi^-$ , plotted in the invariant mass plane  $m(p\pi^-)$  vs  $m(\pi^+\pi^-)$ .

parameters  $b < 2.4$  fm in a geometric model. The open channels for production of strange particles in this energy regime are  $NN \rightarrow \Lambda KN$  (1.58 GeV),  $NN \rightarrow \Sigma KN$  (1.79 GeV), and  $NN \rightarrow \Lambda KN\pi$  (1.96 GeV). Neutral strange particles were detected by their charged-particle decay in the active volume of the chamber. In order to derive  $\Lambda$  production cross sections and momentum spectra, detection and scanning efficiencies were determined. The detection efficiency is intrinsically a function of the branching ratio for charged-particle decay, the momentum of the decaying neutral particle, its lifetime, and the fiducial volume of the detection apparatus. To eliminate efficiency problems due to the high track density near the primary reaction vertex, only neutrals decaying farther than 10 cm from the target center were accepted. The resultant average efficiency for  $\Lambda$  detection was  $\sim 33\%$ . The neutral strange particles identified by a kinematical fit of the decays  $\Lambda \rightarrow p\pi^-$  and  $K^0 \rightarrow \pi^+\pi^-$  are displayed in the  $m(p\pi^-)$  vs  $m(\pi^+\pi^-)$  invariant mass plane in Fig. 1. There are two distinct bands corresponding to the  $\Lambda$  and  $K^0$  masses. The low number of  $K^0$  decays observed




FIG. 2.  $^{40}\text{Ar} + \text{KCl} \rightarrow \Lambda$  events as a function of  $p_\perp$  and  $p_\parallel$  in the nucleus-nucleus c.m. at 1.8 GeV/u. (a) Scatterplot of the data. The hatched curve shows the region where events were excluded as indicated in the text. The solid curve corresponds to the  $NN \rightarrow \Lambda KN$  kinematic limit. (b) Linear contour plot of the results for the Monte Carlo calculation AA (Fermi motion only). (c) Linear contour plot of the calculation RsII with subsequent  $\Lambda$  rescattering (see text).

is due to the mismatch between the effective fiducial volume of the chamber and the lifetimes of the  $K_S^0$  ( $c\tau=2.68$  cm) and  $K_L^0$  ( $c\tau=1554$  cm). It should be noted that any  $\Lambda$  originating from  $\Sigma^0$  production (100% branching ratio for  $\Sigma^0 \rightarrow \Lambda\gamma$  decay) would be included in these data.<sup>7</sup>

The identified  $\Lambda$ 's are displayed in Fig. 2(a) as a function of  $p_\perp$  and  $p_\parallel$  in the nucleus-nucleus c.m. system. The solid curve in Fig. 2(a) represents the kinematic limit for  $\Lambda$  momenta produced in the elementary  $NN \rightarrow \Lambda KN$  reaction at 1.8 GeV. Most of the  $\Lambda$ 's are observed beyond the kinematic limit indicating the necessity to include Fermi motion and/or collective multiparticle interactions. To investigate further the origin of the observed high  $\Lambda$  momenta, a Monte Carlo phase space calculation incorporating Fermi motion of nucleons in both nuclei was carried out. A Gaussian Fermi momentum density distribution  $f(p) \propto \exp(-p^2/2\sigma^2)$  with  $\sigma=90$  MeV/c was employed<sup>8</sup> and the  $\Lambda$ 's were produced in an elementary  $NN \rightarrow \Lambda KN$  process with a uniform phase space distribution.<sup>9</sup> The results (denoted as AA) after folding in the detection efficiencies are shown in Fig. 2(b). The distribution, which peaks at  $p=0$ , is broadened in phase space but still does not adequately describe the observed spectrum.

Inclusion of  $\Lambda$  rescattering in the nuclear medium was next considered by adopting the simple approach of Randrup.<sup>10</sup> In this model the geometry is assumed to be two partially overlapping spheres of normal nuclear matter density subject to the usual Lorentz contractions. The mean number of  $\Lambda N$  collisions was found to be  $n_0=1.86$  using the  $\Lambda N$  elastic cross section<sup>11</sup>  $\sigma_{\Lambda N}=20$  mb. In each collision the  $\Lambda$  scatters from a baryon whose momentum is obtained from a chosen momentum density distribution. The angular distribution was taken to be isotropic in the  $\Lambda$ -baryon rest frame. The baryon momentum distribution was assumed to be either of two extremes: a thermal "fireball" distribution<sup>12</sup> in the nucleus-nucleus c.m. with  $T_0=120$  MeV (which will be denoted RsI) or a superposition of the initial distributions of baryons in the two incident nuclei represented by two moving (Lorentz contracted) Gaussian Fermi momentum distributions (denoted RsII). Rescattering from the thermal "fireball" distribution (RsI) has little effect on the  $\Lambda$  spectrum. However, rescattering from a superposition of the colliding initial distributions (RsII) shown in Fig. 2(c) depletes events from the central peak and creates peaks near  $p_\parallel = \pm 0.8$

GeV/c. The backward peak at  $p_\parallel = -0.8$  GeV/c is suppressed by the detection efficiency which is included in the calculation. In addition, the  $p_\perp$  is boosted in qualitative agreement with the observations.

A quantitative comparison of the data with results of the Monte Carlo calculations can be found in Table I, where the mean and rms values of  $p_\perp$  and  $|p_\parallel^{c.m.}|$  for the invariant  $(E/2\pi p_\perp)(d^2\sigma/dp_\perp dp_\parallel)$  distributions are presented. The data exhibit a much higher  $\langle p_\perp \rangle$  than predicted by any of the calculations. The RsII calculation which most closely approximates the observed  $\langle p_\perp \rangle$  predicts too large a value for  $\langle |p_\parallel^{c.m.}| \rangle$ . Other possibilities which could influence these results have been explored. One mechanism of importance to  $\Lambda$  production<sup>13</sup> is intermediate  $\Delta$  formation. To test whether the experimental results are sensitive to the process  $NN \rightarrow \Delta N$  followed by  $\Delta N \rightarrow \Lambda KN$ , further calculations including Fermi motion and this two-step process were carried out. The resultant spectrum of  $\Lambda$ 's was found to be similar to that produced in  $NN$  interactions as shown [denoted AA( $\Delta$ )] in row 4 of Table I. A more elaborate cascade calculation is needed; furthermore, additional effects such as high density in the initial stages of the collision either in short-range correlations or hydrodynamic flow may be necessary to describe the observed  $\Lambda$ 's.

The topology for  $\Lambda$ -producing events does not differ from the rest of the central-trigger events. The mean  $\pi^-$  and total charged-particle multiplicities are found to be the same for events with or without  $\Lambda$ 's. In both cases  $\langle n_{\pi^-} \rangle \approx 6$  and  $\langle n_{\text{tot}} \rangle \approx 42$ . The mean  $\Lambda$  multiplicity is found to be  $\langle n_\Lambda \rangle = 0.04$  for the central trigger. Using the measured cross section for central-trigger events,<sup>6</sup>  $\sigma=180$  mb, the  $\Lambda$  cross section in central colli-

TABLE I. Mean and rms values of  $p_\perp$  and  $|p_\parallel^{c.m.}|$  for the invariant  $(E/2\pi p_\perp)(d^2\sigma/dp_\perp dp_\parallel)$  distributions of the data and calculations described in the text.  $\langle p_\perp^2 \rangle$  can be extracted from the table using the relation  $\sigma_{\text{rms}}(p_\perp) = [\langle p_\perp^2 \rangle - \langle p_\perp \rangle^2]^{1/2}$ .

	$\langle p_\perp \rangle$ (GeV/c)	$\sigma_{\text{rms}}(p_\perp)$ (GeV/c)	$\langle  p_\parallel^{c.m.}  \rangle$ (GeV/c)	$\sigma_{\text{rms}}(p_\parallel^{c.m.})$ (GeV/c)
Data	0.493	0.243	0.429	0.264
NN	0.122	0.077	0.117	0.077
AA	0.207	0.150	0.212	0.154
AA( $\Delta$ )	0.227	0.163	0.247	0.179
RsI	0.220	0.180	0.229	0.184
RsII	0.276	0.233	0.641	0.380

sions of  $^{40}\text{Ar} + \text{KCl}$  at 1.8 GeV/u is found to be  $\sigma_{\Lambda} = 7.6 \pm 2.2$  mb. A multiple collision model employing a linear cascade<sup>13,14</sup> with impact parameter selection  $b < 2.4$  fm predicts a  $\Lambda + \Sigma$  cross section of  $\sigma \sim 10$  mb. There is a large uncertainty in the prediction due to the absence of  $pp \rightarrow (\Lambda, \Sigma) + X$  production data at these low energies.

Since  $\Lambda$  decay is self-analyzing for polarization in the decay process  $\Lambda \rightarrow p\pi^-$ , the distribution of decay protons in the  $\Lambda$  decay frame relative to the  $\Lambda$  spin direction<sup>15</sup> is  $dW/d\Omega = (1 + \alpha P \cos\theta)/4\pi$ , where  $P$  is the  $\Lambda$  polarization,  $\alpha = -0.642$ ,<sup>16</sup> and  $\theta$  is the angle between the decay proton and the unit vector normal to the plane. Using the relation<sup>17</sup>  $\alpha P = \langle \cos\theta \rangle / \langle \cos^2\theta \rangle$ , the polarization extracted from the present experiment was found to be  $P = -0.10 \pm 0.05$ . Significant  $\Lambda$  polarization ( $0 > P > -0.3$ ) has been observed in  $p$ -nucleus collisions at incident energies 24–400 GeV.<sup>18</sup> At these higher energies  $|P|$  increases monotonically as a function of transverse momentum and is independent of incident energy. This polarization and its behavior have been described in terms of spin features of the SU(6) quark wave functions<sup>18,19</sup> whereby the spin of the  $\Lambda$  is determined by that of the produced strange quark. Clearly, any interpretation of the  $\Lambda$  polarization observed in the present study is complicated by the rescattering of  $\Lambda$ 's in the surrounding nuclear matter which would be expected to degrade the polarization.

In summary,  $\Lambda$ 's have been studied in collisions of  $^{40}\text{Ar} + \text{KCl}$  at 1.8 GeV/u incident energy just above the  $NN$  threshold. The average  $\Lambda$  polarization was found to be  $P = -0.10 \pm 0.05$ . The  $\Lambda$  production cross section is  $\sigma_{\Lambda} = 7.6 \pm 2.2$  mb for the central-trigger mode. In a Monte Carlo calculation it is necessary to include Fermi motion and rescattering from the initial momentum density distributions to reproduce qualitatively the  $\Lambda$  momentum phase space. This is consistent with the hypothesis that  $\Lambda$ 's are produced in the initial stage of the reaction rather than in the later thermal phase. The high  $\langle p_{\perp} \rangle$  observed in the data cannot be reproduced in the framework of these calculations which are based on a direct  $NN \rightarrow \Lambda KN$  production mechanism in the first binary encounter. Recent Monte Carlo-cascade calculations<sup>20</sup> exhibit a central density pileup of four times normal nuclear matter density in the first few generations of interactions. In such

calculations there is local equilibrium in the high-density region but global nonequilibrium. Such high-density regions may provide a way for producing  $\Lambda$ 's with higher  $p_{\perp}$  than in a direct process.

We wish to thank J. P. Brannigan, the Bevalac operating staff, and the scanning group at Lawrence Berkeley Laboratory for their support. We would also like to thank R. Bock for his continuing interest and support.

This work is supported by the Bundesministerium für Forschung und Technologie, West Germany, and the Director, Office of Energy Research, Division of Nuclear Physics of the Office of High Energy and Nuclear Physics of the U. S. Department of Energy under Contract No. W-7405-ENG-48.

<sup>(a)</sup>Present address: Physics Department, University of California at Los Angeles, Cal. 90024.

<sup>1</sup>A. Sandoval *et al.*, Phys. Rev. C 21, 1321 (1980), and references therein.

<sup>2</sup>J. D. Bjorken and E. A. Paschos, Phys. Rev. 185, 1975 (1969).

<sup>3</sup>M. Holder *et al.*, Phys. Lett. 69B, 377 (1977).

<sup>4</sup>J. G. H. de Groot *et al.*, Z. Phys. C 1, 143 (1979).

<sup>5</sup>Ch. de la Vaissiere *et al.*, Z. Phys. C 1, 3 (1979).

<sup>6</sup>A. Sandoval *et al.*, Phys. Rev. Lett. 45, 874 (1980).

<sup>7</sup> $\Sigma^0$  and  $\Lambda$  production cross sections have been measured in  $pp$  experiments at 2.85 GeV where  $\sigma(\Lambda K^+ p) = 0.051 \pm 0.012$  mb and  $\sigma(\Sigma^0 K^+ p) = 0.013 \pm 0.007$  mb; see R. I. Louttit *et al.*, Phys. Rev. 123, 1465 (1961).

<sup>8</sup>A. S. Goldhaber, Phys. Lett. 53B, 306 (1974).

<sup>9</sup>J. H. Friedman, J. Comput. Phys. 7, 201 (1971).

<sup>10</sup>J. Randrup, Phys. Lett. 99B, 9 (1981).

<sup>11</sup>G. Alexander and U. Karshon, in *Proceedings of the Second International Conference on High Energy Physics and Nuclear Structure, Rehovoth, Israel, 1967* edited by G. Alexander (North-Holland, Amsterdam, 1967), p. 36.

<sup>12</sup>S. Nagamiya, Lawrence Berkeley Laboratory Report No. LBL-12123, unpublished.

<sup>13</sup>J. Randrup and C. M. Ko, Nucl. Phys. A343, 590 (1980).

<sup>14</sup>J. Randrup, private communication.

<sup>15</sup>E. Cummins, Weak Interactions (McGraw-Hill, New York, 1973).

<sup>16</sup>C. Bricman *et al.*, Phys. Lett. 75B, 1 (1978).

<sup>17</sup>M. L. Faccini-Turluer *et al.*, Z. Phys. C 1, 19 (1979).

<sup>18</sup>K. Heller *et al.*, Phys. Rev. Lett. 41, 607 (1978).

<sup>19</sup>B. Andersson *et al.*, Phys. Lett. 85B, 417 (1979).

<sup>20</sup>J. Cugnon *et al.*, Nucl. Phys. A352, 505 (1981).