

INSECTA MUNDI

A Journal of World Insect Systematics

0675

Synonymy and distribution data of *Trypanidius* Blanchard, 1846
(Coleoptera: Cerambycidae: Lamiinae: Acanthocinini) in Cuba

Sergio Devesa

La Iglesia, 4
36988 San Vicente de O Grove
Pontevedra, España

Alejandro Barro

Departamento de Biología Animal y Humana, Facultad de Biología,
Universidad de La Habana, Cuba

Date of issue: December 7, 2018

Sergio Devesa and Alejandro Barro
Synonymy and distribution data of *Trypanidius* Blanchard, 1846
(Coleoptera: Cerambycidae: Lamiinae: Acanthocinini) in Cuba
Insecta Mundi 0675: 1–6

ZooBank Registered: urn:lsid:zoobank.org:pub:E2B9D5DE-7D01-4EBC-961A-ACE7EAB59EA0

Published in 2018 by

Center for Systematic Entomology, Inc.
P.O. Box 141874
Gainesville, FL 32614-1874 USA
<http://centerforsystematicentomology.org/>

Insecta Mundi is a journal primarily devoted to insect systematics, but articles can be published on any non-marine arthropod. Topics considered for publication include systematics, taxonomy, nomenclature, checklists, faunal works, and natural history. *Insecta Mundi* will not consider works in the applied sciences (i.e. medical entomology, pest control research, etc.), and no longer publishes book reviews or editorials. *Insecta Mundi* publishes original research or discoveries in an inexpensive and timely manner, distributing them free via open access on the internet on the date of publication.

Insecta Mundi is referenced or abstracted by several sources, including the Zoological Record and CAB Abstracts. *Insecta Mundi* is published irregularly throughout the year, with completed manuscripts assigned an individual number. Manuscripts must be peer reviewed prior to submission, after which they are reviewed by the editorial board to ensure quality. One author of each submitted manuscript must be a current member of the Center for Systematic Entomology.

Guidelines and requirements for the preparation of manuscripts are available on the *Insecta Mundi* website at <http://centerforsystematicentomology.org/insectamundi/>

Chief Editor: David Plotkin, insectamundi@gmail.com
Assistant Editor: Paul E. Skelley, insectamundi@gmail.com
Head Layout Editor: Robert G. Forsyth
Editorial Board: J. H. Frank, M. J. Paulsen, Michael C. Thomas
Review Editors: Listed on the *Insecta Mundi* webpage

Printed copies (ISSN 0749-6737) annually deposited in libraries

CSIRO, Canberra, ACT, Australia
Museu de Zoologia, São Paulo, Brazil
Agriculture and Agrifood Canada, Ottawa, ON, Canada
The Natural History Museum, London, UK
Muzeum i Instytut Zoologii PAN, Warsaw, Poland
National Taiwan University, Taipei, Taiwan
California Academy of Sciences, San Francisco, CA, USA
Florida Department of Agriculture and Consumer Services, Gainesville, FL, USA
Field Museum of Natural History, Chicago, IL, USA
National Museum of Natural History, Smithsonian Institution, Washington, DC, USA
Zoological Institute of Russian Academy of Sciences, Saint-Petersburg, Russia

Electronic copies (Online ISSN 1942-1354, CDROM ISSN 1942-1362) in PDF format

Printed CD or DVD mailed to all members at end of year. Archived digitally by Portico.
Florida Virtual Campus: <http://purl.fcla.edu/fcla/insectamundi>
University of Nebraska-Lincoln, Digital Commons: <http://digitalcommons.unl.edu/insectamundi/>
Goethe-Universität, Frankfurt am Main: <http://nbn-resolving.de/urn/resolver.pl?urn:nbn:de:hebis:30:3-135240>

Copyright held by the author(s). This is an open access article distributed under the terms of the Creative Commons, Attribution Non-Commercial License, which permits unrestricted non-commercial use, distribution, and reproduction in any medium, provided the original author(s) and source are credited. <http://creativecommons.org/licenses/by-nc/3.0/>

Layout Editor for this article: Robert G. Forsyth

Synonymy and distribution data of *Trypanidius* Blanchard, 1846 (Coleoptera: Cerambycidae: Lamiinae: Acanthocinini) in Cuba

Sergio Devesa

La Iglesia, 4
36988 San Vicente de O Grove
Pontevedra, ESPAÑA
sergio.devesa@gmail.com

Alejandro Barro

Departamento de Biología Animal y Humana, Facultad de Biología,
Universidad de La Habana, CUBA
abarro@fbio.uh.cu

Abstract. *Trypanidius nocturnus* Fisher, 1942 (Coleoptera: Cerambycidae: Lamiinae: Acanthocinini) is synonymized with *Trypanidius insularis* Fisher, 1925. Dorsal habitus photographs of the holotypes and paratypes, as well as male and females of specimens collected in Cuba, Puerto Rico and Hispaniola, are provided. Additionally, collecting data and a distributional map of Cuba are offered.

Key words. West Indies, longhorn woodboring beetles, systematics.

Resumen. *Trypanidius nocturnus* Fisher, 1942 (Coleoptera: Cerambycidae: Lamiinae: Acanthocinini) es sinonimizado con *Trypanidius insularis* Fisher, 1925. Se muestran fotos en vista dorsal de los holotipos y paratipos así como de ejemplares macho y hembra recolectados en Cuba, Puerto Rico y La Española. Adicionalmente, se muestran los datos de colecta y mapa de distribución de los ejemplares colectados en Cuba.

Palabras clave. Antillas, cerambycidos, sistemática.

Introduction

The genus *Trypanidius* Blanchard, 1846 includes 15 species and is widely distributed in the Neotropical region from Central to South America. Three species were described from the Caribbean islands: *T. insularis* Fisher, 1925 from Cuba; *T. nocturnus* Fisher, 1942 from Cuba and Puerto Rico; *T. spilmani spilmani* Villiers, 1980 from Dominica, Martinique and Saint Lucia and *T. spilmani liamaigae* Chalumeau, 1983 from Saint Kitts (Villiers 1980; Chalumeau 1983; Monné 2018).

Fisher (1925: 8) described *T. insularis* based on two specimens: one male (Holotype, collected in Nagua, Oriente, Cuba on July 7, 1922 by S. C. Bruner and C. H. Ballou and deposited in the USNM, No. 28413) (Fig. 1) and one female (Paratype, collected in Guantánamo, Cuba on June 1, 1917 at light by C. T. Ramsden and deposited in the AMNH) (Fig. 2). In his description, Fisher said: “The paratype differs from the type in having the antennae slightly shorter, pubescence above more yellowish brown, black spots on pronotum more distinct, brown pubescence on antennae predominating, black spot on elytra behind middle more sinuate, and the zigzag fascia near apex more confluent and forming a large triangular spot, pubescence beneath sparser and more brownish, with a few white spots toward the sides, and the last abdominal segment longer, narrower at apex, and the dorsal surface only feebly emarginate.” (Fig. 1-2). After its description, no additional information was published for the species.

Fisher (1942: 38) described *T. nocturnus* from a single female specimen, collected in Villalba, Puerto Rico, May 18, 1940 by L. F. Martorell, and deposited in the USNM, No. 55677 (Fig. 3). In his description, Fisher wrote: “This species is allied to *Trypanidius insularis* Fisher described from Cuba, but differs from that species in being more reddish brown and in having a large triangular, yellowish-white pubescent spot covering the apical fourth of each elytron replacing the transverse, white zigzag fascia on the elytron of *insularis*.”

Surprisingly, the same author again described *T. nocturnus* (Fisher 1947: 39) based on two female specimens: the holotype, collected in Villalba, Puerto Rico, May 18, 1940 by L. F. Martorell and deposited in the USNM, No. 55677 and the paratype, collected in Mayaguez, Puerto Rico, September 1938 by J.



1

Nagua, Oriente, Cuba
 Julio 7 de 1922.
 Col. S. C. Bruner y
 C. H. Ballou

Type No.
 28413
 U.S.N.M.

*Trypanidius
 insularis
 Fisher*

Guantanamo, at light
 Cuba 17 June 1917
 Coll. C. T. Ramsden

Paratype

*Trypanidius
 insularis
 Paratype Fisher*



2



3

Villalba, P.R.
 near light
 coll. Mortorell
 (col. light)


*Trypanidius
 nocturnus
 Fisher*

Type No.
 58124
 U.S.N.M.

published number


Mayaguez, P.R.
 Sept. 1938
 Coll. J. Araujo

Paratype
 58124
 U.S.N.M.



4

Figures 1–4. *Trypanidius* spp. 1) *T. insularis* Fisher, 1925, habitus holotype dorsal view (Photo: USNM). 2) *T. insularis*, habitus paratype dorsal view (Photo: AMNH). 3) *T. nocturnus* Fisher, 1942, habitus holotype dorsal view (Photo: USNM). 4) *T. nocturnus* Fisher, 1942, habitus paratype dorsal view (Photo: S. Lingafelter).



Figures 5–10. *Trypanidius insularis* Fisher, 1925: variation in the size and shape of the black and yellowish-white fascia in elytral apical third. 5) Specimen from Cuba, male. 6–7) Specimens from Cuba, females. 8–9) Specimen from the Dominican Republic, female. 8) Dorsal view (Photo: S. Lingafelter). 9) Ventral view (Photo: S. Lingafelter). 10) Specimen from Puerto Rico, female (Photo: S. Lingafelter).

Araujo (Fig. 4). Both specimens are deposited in the USNM, No. 58124; this means that the holotype specimen has two holotype numbers: USNM No. 58124 and USNM No. 55677. Fisher justified the description of the new species as follow: “This species is allied to *Trypanidius insularis* Fisher, but it differs from that species in having the sides of the pronotum more nearly parallel and without a row of coarse punctures along the anterior margin, the antennae and legs densely clothed with short, recumbent, brown pubescence, with a few white hairs intermixed, the underside of the body densely clothed with whitish pubescence and variegated with small, denser pubescent, white spots, and each elytron ornamented with a large triangular yellowish-white pubescent spot covering the apical third and with the tip obtusely produced at the outer margin.”

Fisher used the same character state, presence of a large triangular spot, as a primary diagnostic character in all of these descriptions; this suggests that there is not enough justification to treat *T. insularis* and *T. nocturnus* as distinct species.

Among the other authors who cited these species, Zayas (1975) said that he has only collected *T. nocturnus* in Cuba and suspected that the differences of the elytral fascia between the holotype and the paratype of *T. insularis* represent two different variations of the same species. Carelli et al. (2013), in their review of the genus, did not study any specimen of *T. insularis*; they only examined a photograph of the holotype.

In this paper, we propose the synonymy of *Trypanidius insularis* Fisher, 1925 and *Trypanidius nocturnus* Fisher, 1942.

Materials and Methods

We studied photographs of the types of the species involved in this study (Lingafelter et al. 2014). Color photos of holotypes of both species are also available online on the Cerambycidae of the New World photographic catalog (Bezark 2018) and on the USNM's type database (Lingafelter et al. 2018). Those photos were compared with the specimens studied and with original descriptions. Photographs of additional specimens from Puerto Rico and Hispaniola were studied. Observations were made using an ocular micrometer adapted to an Olympus SZX7 0.8–5.6× stereomicroscope. Photographs were taken with a Canon EOS 5 Mark III camera, equipped with a Canon MP-E 65mm f/2.8 1–5× and EF 100 mm f/2.8 macro IS USM macro lens. Multiple images were stacked with the Zerene Stacker AutoMontage software and processed in Aperture software.

The specimens and images from the following public museums and private collections were examined:

AMNH	American Museum of Natural History, New York, USA
CMNH	Carnegie Museum of Natural History, Pittsburgh, Pennsylvania, USA
CZACC	Colección Zoológica de la Academia de Ciencias de Cuba, I.E.S., Havana, Cuba
CZCTR	Museo de Historia Natural “Charles T. Ramsden”, Santiago de Cuba, Cuba
FZPC	Fernando de Zayas Private Collection, Havana, CUBA
HGPC	Horacio Grillo Private Collection, Universidad Central Las Villas, Cuba
SDPC	Sergio Devesa Private Collection, Pontevedra, Spain
SWLC	Steven W. Lingafelter Collection, Hereford, Arizona, USA
USNM	National Museum of Natural History, Smithsonian Institution, Washington, DC, USA

Results

Trypanidius insularis Fisher 1925

Trypanidius nocturnus Fisher 1942, **new synonym**

Specimens examined. CUBA: (43 specimens: 21 males and 22 females) **CIENFUEGOS Prov.:** Buenos Aires (21°58'00"N/80°9'00"W), may-1952, 1 female (F. de Zayas leg.); jun-1952, 4 females (F. de Zayas leg.); **GRANMA Prov.:** La Platica - P.N. Pico Turquino (20°00'54" N/76°53'28"W), 27-jan-2007,

1 female at light (250W, vapor Hg) (E. Fonseca leg.); Nagua, (20°7'00"N/76°52'00" W), 07-jul-1922, 1 male HOLOTYPE of *T. insularis* (S.C. Bruner y C.H. Ballou leg.); **GUANTÁNAMO Prov.:** Guantánamo (20°8' 40"N/75°12'33"W), 01-jun-1907, 1 female PARATYPE of *T. insularis* (C.T. Ramsden leg.); **ISLA DE LA JUVENTUD Prov.:** Lomas de Sigüanea (21°37'00"N/82°58'00"W), 15-ago-2015, 1 female, pinewood, near Hotel Colony (A. Barro leg.); **SANCTI SPIRITUS Prov.:** Topes de Collantes (21°54'49"N/80°01'19"W), jul-1965, 1 male (F. de Zayas leg.); Topes de Collantes, Estación Biológica (21°56'00"N/79°59'00"W), 25-may-2001, 1 male at light (A. Avila, R. Núñez & A. Lozada leg.); 3-may-2002, 1 male at light (I. Martínez leg.); 4-jun-2002, 1 female (A. Lozada, A. Ávila & R. Núñez leg.); Topes de Collantes, mirador (21°51'50"N/80°01'15"W), jul-2009, 6 males + 2 females at light (S. Devesa leg.); ago-2009, 3 males + 1 female at light (S. Devesa leg.); sep-2009, 2 males + 3 females at light (S. Devesa leg.); oct-2009, 1 female at light (S. Devesa leg.); may-2010, 1 male at light (S. Devesa leg.); jun-2010, 1 female at light (S. Devesa leg.); oct-2010, 1 female at light (S. Devesa leg.); jul-2011, 1 male at light (S. Devesa leg.); **SANTIAGO DE CUBA Prov.:** Loma de Calá a Palma Mocha (20°01'48"N/76°54'36"W), 16-jul-1948, 1 male + 1 female ovipositing on freshly cut tree trunk (F. de Zayas leg.); Loma del Gato (20°00'52"N/76°03'19"W), jun-1956, 1 female (F. de Zayas leg.); Pico Turquino (19°59'00"N/76°50'00"W), jun-1964, 2 males + 3 females at light (F. de Zayas leg.). (Fig. 11). **DOMINICAN REPUBLIC:** Azua, East side of crest, Sierra Martin García, 7 km WNW Barrero, 860 m (18°21'N/70°58'W), 25/26-jul-1992, 1 female in cloud forest adjacent to disturbed forest (C. Young, R. Davidson, S. Thompson and J. Rawlins leg.). **PUERTO RICO:** El Yunque National Park, Mt. Britton (18°17'N/65°47'W), 17-jul-2009, 1 male (J. Anderson & G. Setliff leg.); Villalba, 18-may-1940 (L.F. Martorell leg.); Mayaguez, September-1938 (J. Araujo leg.).

Dimensions in mm (males/females). total length 12.5–14.3/13.0–16.7; pronotum length 2.4–2.7/2.4–3.0; pronotum width 3.7–4.2/3.9–4.8; elytra length 9.2–10.3/9.5–12.6; elytra width at humeri 5.0–5.7/5.2–6.8; total length/pronotal length 5.1–5.6/5.1–5.9; elytral length/elytral width 1.8–1.9/1.7–1.9; pronotal length/pronotal width 0.6–0.7/0.6–0.7.

Geographical distribution. Cuba, Hispaniola, Puerto Rico.



Figure 11. Geographical and spatial distribution of *Trypanidius insularis* Fisher, 1925, in Cuba.

Detailed study of the original descriptions of the two species, examination of the photographs of the holotypes of *T. insularis* (Fig. 1) and *T. nocturnus* (Fig. 3) in the USNM, the paratypes of *T. insularis* (Fig. 2) in the AMNH and *T. nocturnus* (Fig. 4) in the USNM, and 43 specimens collected in Cuba, one specimen from the Dominican Republic (Fig. 8, 9) and one specimen from Puerto Rico (Fig. 10) show that these are the same species. Details of the patterns of pubescence, punctation, color, and other morphology are identical. We conclude that small differences on the pronotum (size of the lateral spine, black spots at apical third, and white pubescence), antennae (size and color of basal annulations on antennomeres) and elytra (shape and dimension of both black spots and yellowish-white fascia at apical

third and sutural area) are intraspecific variation observed in the analyzed specimens. Likewise, the main characteristics previously used to differentiate *T. nocturnus* from *T. insularis* (black spots and a yellowish-white spot in the apical third of the elytra) are present in the description of the paratype of *T. insularis*. The zigzag fascia described in the holotype of *T. insularis* must be considered as a chromatic variation.

Acknowledgments

We thank Dr. Lee Herman for sending the picture of the paratype of *Trypanidius insularis* deposited in the Coleoptera collection of the AMNH. We thank Dr. Steve Lingafelter for sending the picture of the paratype of *Trypanidius nocturnus* deposited in the USNM and the photographs of the specimens from Hispaniola and Puerto Rico. We especially appreciate the critical reading of the manuscript by Allan Carelli Aragao and Steve Lingafelter, as their suggestions contributed greatly to the final product.

Literature Cited

- Bezark, L. G. 2018.** A photographic catalog of the Cerambycidae of the New World. Available at <https://apps2.cdfa.ca.gov/publicApps/plant/bycidDB/wdefault.asp?w=n>. (Last accessed July 12, 2018.)
- Carelli, A., M. L. Monné, and V. S. Machado. 2013.** Taxonomic revision of *Trypanidius* Blanchard, 1846 in South America (Insecta: Coleoptera: Cerambycidae). *Zootaxa* 3691(2): 253–272.
- Chalumeau, F. 1983.** Acanthocinini des Petites Antilles. Nouveaux taxa et observations diverses (Coleoptera: Cerambycidae, Lamiinae). *Nouvelle Revue d'Entomologie*, Paris 13(2): 219–237.
- Fisher, W. S. 1925.** New West Indian Cerambycidae (Coleoptera). Subfamily Lamiinae. *American Museum Novitates* 174: 1–16.
- Fisher, W. S. 1942.** New Coleoptera from Puerto Rico. *Journal of Agriculture of the University of Puerto Rico* 25: 37–39.
- Fisher, W. S. 1947.** New West Indian cerambycid beetles. IV. *Memorias de la Sociedad Cubana de Historia Natural «Felipe Poëy»*, La Habana 19(1): 29–41.
- Lingafelter, S. W., M. A. Monné, C. J. Micheli, and E. H. Nearn. 2018.** Cerambycid primary types of the Smithsonian Institution. Available at <http://smithsoniancerambycidae.com>. (Last accessed July 12, 2018.)
- Lingafelter, S. W., E. H. Nearn, G. L. Tavakilian, M. A. Monné, and M. Biondi. 2014.** Long-horned woodboring beetles (Coleoptera, Cerambycidae and Disteniidae) primary types of the Smithsonian Institution. Smithsonian Institution Scholarly Press; Washington DC. 390 p.
- Monné, M. A. 2018.** Catalogue of the Cerambycidae (Coleoptera) of the Neotropical region. Part II. Subfamily Lamiinae. Available at <http://cerambyxcat.com/>. (Last accessed July 12, 2018.)
- Villiers, A. 1980.** Coléoptères Cerambycidae des Antilles Françaises III. Lamiinae. *Annales de la Société Entomologique de France* (n.s.) 16(4): 541–598.
- Zayas, F. de. 1975.** Revisión de la familia Cerambycidae (Coleoptera, Phytophagoidea). *Academia de Ciencias de Cuba*; Havana, Cuba. 443 p.

Received September 18, 2018; accepted November 2, 2018.
Review editor M. J. Paulsen.